

On the complexity of vertex-coloring edge-weightings

Andrzej Dudek, David Wajc

► **To cite this version:**

Andrzej Dudek, David Wajc. On the complexity of vertex-coloring edge-weightings. *Discrete Mathematics and Theoretical Computer Science, DMTCS*, 2011, Vol. 13 no. 3 (3), pp.45–50. hal-00990502

HAL Id: hal-00990502

<https://hal.inria.fr/hal-00990502>

Submitted on 13 May 2014

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

On the complexity of vertex-coloring edge-weightings

Andrzej Dudek ^{1†} and David Wajc ^{2‡}

¹Department of Mathematics, Western Michigan University, Kalamazoo, MI 49008, USA

²Computer Science Department, Technion Israel Institute of Technology, Haifa 32000, Israel

received 18th March 2010, revised 24th February 2011, accepted 26th October 2011.

Given a graph $G = (V, E)$ and a weight function $w : E \rightarrow \mathbb{R}$, a coloring of vertices of G , induced by w , is defined by $\chi_w(v) = \sum_{e \ni v} w(e)$ for all $v \in V$. In this paper, we show that determining whether a particular graph has a weighting of the edges from $\{1, 2\}$ that induces a proper vertex coloring is NP-complete.

Keywords: vertex-coloring, 1-2-3 conjecture, NP-completeness

1 Introduction

For a given graph $G = (V, E)$, let $w : E \rightarrow \mathbb{R}$ be a weight function. We say that w is *proper* if the coloring of the vertices $\chi_w(v) = \sum_{e \ni v} w(e)$, $v \in V$, is proper. In 2004, Karoński, Łuczak, and Thomason (2004) showed that any graph with no components isomorphic to K_2 has a proper weighting from a finite set of reals. Furthermore, they conjectured that every graph with no components isomorphic to K_2 has a proper weighting from $W = \{1, 2, 3\}$. Addario-Berry, Dalal, McDiarmid, Reed, and Thomason (2007) showed that the above holds if $W = \{1, \dots, 30\}$. This result was improved by Addario-Berry, Dalal, and Reed (2008), who showed that one can take $W = \{1, \dots, 16\}$. Subsequently, Wang and Yu (2008) proved that $W = \{1, \dots, 13\}$ suffices. A recent breakthrough by Kalkowski, Karoński, and Pfender (2010) showed that the set of weights can be as small as $W = \{1, 2, 3, 4, 5\}$.

On the other hand, Addario-Berry, Dalal, and Reed (2008) showed that almost all graphs have a proper weighting from $\{1, 2\}$. In this paper, we show that determining whether a particular graph has a proper weighting of the edges from $\{1, 2\}$ is NP-complete. Consequently, there is no simple characterization of graphs with proper weightings from $\{1, 2\}$, unless $P=NP$. Formally, let

$$1\text{-}2\text{WEIGHT} = \{G : G \text{ is a graph having a proper weighting from } \{1, 2\}\}.$$

[†]Email: andrzej.dudek@wmich.edu.

[‡]Email: sdavidwa@cs.technion.ac.il This work was performed while the author was visiting Carnegie Mellon University.




Fig. 1: A k -disallowing gadget (a) and its symbolic representation (b); a construction of $h(G)$ (c).

Theorem 1.1 *1-2WEIGHT is NP-complete.*

Before we prove this statement, we consider a similar theorem with a somewhat simpler proof, which we use as a template to prove Theorem 1.1. By analogy to 1-2WEIGHT we denote by 0-1WEIGHT the family of graphs with a proper weighting from $\{0, 1\}$ and show the following:

Theorem 1.2 *0-1WEIGHT is NP-complete.*

2 0-1WEIGHT is NP-complete

Here we prove Theorem 1.2.

First note that 0-1WEIGHT is clearly in NP, since one can verify in polynomial time for a given graph whether a weighting of its edges from $\{0, 1\}$ is proper.

Next we consider the well-known NP-hard problem

$$3\text{-COLOR} = \{G : G \text{ is a graph having a proper 3-vertex-coloring}\}.$$

In order to prove that 0-1WEIGHT is NP-hard (and hence NP-complete), we show a reduction from 3-COLOR to 0-1WEIGHT. To this end, we define a polynomial time reduction h , such that $G \in 3\text{-COLOR}$ if and only if $h(G) \in 0\text{-1WEIGHT}$. To achieve this, we need two auxiliary gadgets.

We refer to the first gadget as a **triangle gadget**. This consists of a triangle xab , with x referred to as the *top* and with a and b each having no other coinciding edges. Note that any proper weighting w from $\{0, 1\}$ of a graph with such a triangle must hold $w(xa) \neq w(xb)$; otherwise $\chi_w(a) = w(ab) + w(ax) = w(ba) + w(bx) = \chi_w(b)$. Hence, $\{w(xa), w(xb)\} = \{0, 1\}$ and so every such triangle gadget contributes exactly 1 to $\chi_w(x)$.

The second gadget, called a **k -disallowing gadget**, consists of a *main triangle* vxy with v referred to as the *root* and with x and y each constituting the top of $k - 1$ distinct triangle gadgets (see Figure 1(a)). Note that in any proper weighting w from $\{0, 1\}$, $w(vx) \neq w(vy)$; otherwise, as both $\chi_w(x)$ and $\chi_w(y)$ have $k - 1$ contributed by x and y 's triangles, $\chi_w(x) = w(xv) + w(xy) + k - 1 = w(yv) + w(yx) + k - 1 = \chi_w(y)$. Therefore, if $w(xy) = 0$ then $\{\chi_w(x), \chi_w(y)\} = \{k - 1, k\}$ and, if $w(xy) = 1$ then $\{\chi_w(x), \chi_w(y)\} = \{k, k + 1\}$. In either case, v has one neighbor $z \in \{x, y\}$ with $\chi_w(z) = k$, and consequently, $\chi_w(v) \neq k$ in any proper weighting from $\{0, 1\}$. Also $\{w(vx), w(vy)\} = \{0, 1\}$ and hence this gadget contributes exactly 1 to $\chi_w(v)$.

Now we are ready to show a reduction from 3-COLOR to 0-1WEIGHT, h , such that $G \in 3\text{-COLOR}$ if and only if $h(G) \in 0\text{-1WEIGHT}$. Let $G = (V, E)$ be a graph of order n . We may assume that $n \geq 3$. Otherwise, $n \leq 2$ and G is in 3-COLOR and so it suffices to take as $h(G)$ an empty graph which is trivially in 0-1WEIGHT. For $n \geq 3$ we construct the graph $h(G) = (W, F)$ as follows (see Figure 1(c)). We start with $G = (V, E)$. For each $v \in V$:

- (i) connect v to two new vertices, s and t (distinct for each v);
- (ii) add $n - 1$ new k -disallowing gadgets for all $k \in \{n + 2, n + 3, \dots, 2n\}$ with v as their root.

Clearly, $h(G)$ can be calculated in time polynomial in the size of G .

Fact 2.1 *In $h(G)$ the following holds: any proper weighting w from $\{0, 1\}$ satisfies $\chi_w(v) \in \{n - 1, n, n + 1\}$ for every $v \in V$.*

Proof: Fix $v \in V$. Since $w(vs) + w(vt) \in \{0, 1, 2\}$, v is the endpoint of $\deg(v) \leq n - 1$ edges in V , and v is the root of $(n - 1)$ k -disallowing gadgets (each contributing 1 to $\chi_w(v)$), we have:

$$\chi_w(v) \in \{0, 1, 2\} + \{0, 1, \dots, \deg(v)\} + \{n - 1\} \subseteq \{n - 1, n, \dots, 2n\},$$

where by $A + B$ we mean the set of all sums of an element from A with an element from B . Observing the above and the fact that v is the root of k -disallowing gadgets for all $k \in \{n + 2, \dots, 2n\}$, we find that any proper weighting w from $\{0, 1\}$ satisfies $\chi_w(v) \in \{n - 1, n, n + 1\}$, as claimed. \square

It remains to show that $G \in 3\text{-COLOR}$ if and only if $h(G) \in 0\text{-1WEIGHT}$.

First let us assume that $G \in 3\text{-COLOR}$. That means there exists a proper 3-coloring of G , say $\chi : V \rightarrow \{n - 1, n, n + 1\}$. We define a weighting of the edges of $h(G)$, $w : F \rightarrow \{0, 1\}$ as follows. For all $e \in E$ let $w(e) = 0$. For all $v \in V$, if $\chi(v) = n - 1$ then $w(vs) = w(vt) = 0$; otherwise, if $\chi(v) = n$ then $w(vs) = 1$ and $w(vt) = 0$; and finally, if $\chi(v) = n + 1$ then $w(vs) = w(vt) = 1$. All other edges (parts of gadgets) are weighted as follows: For a triangle gadget xab with root x , $w(xa) = 1, w(xb) = w(ab) = 0$. For a k -disallowing gadget with root v , and main triangle vxy , $w(vx) = w(xy) = 1, w(vy) = 0$, and the weighting of all other triangle gadgets as described above. Note that w is a proper weighting of $h(G)$ (satisfying $\chi_w(v) = \chi(v)$ for all $v \in V$), as required.

Now let us assume that $G \notin 3\text{-COLOR}$. Therefore, for all $\chi : V \rightarrow \{n - 1, n, n + 1\}$, χ is not proper. But, from Fact 2.1, any proper weighting from $\{0, 1\}$ of $h(G)$ satisfies $\chi_w(v) \in \{n - 1, n, n + 1\}$ for all $v \in V$. Thus, there is no such proper weighting and hence $h(G) \notin 0\text{-1WEIGHT}$.

This completes the proof of Theorem 1.2.

3 1-2WEIGHT is NP-complete

The proof of Theorem 1.1 extends the ideas introduced in the proof of Theorem 1.2. Since clearly 1-2WEIGHT is in NP, it remains to show that 1-2WEIGHT is NP-hard. As before, we show a reduction from 3-COLOR to 1-2WEIGHT. To this end, we define a polynomial time reduction f , such that $G \in 3\text{-COLOR}$ if and only if $f(G) \in 1\text{-2WEIGHT}$. Below we define auxiliary gadgets.

As in Section 2, we will use a **triangle gadget**. Now note that every triangle xab , with only x having other adjacent edges (x is referred to as the *top*), contributes exactly 3 to $\chi_w(x)$ in any proper weighting w from $\{1, 2\}$.




Fig. 2: A k -disallowing gadget (a) for even k , (b) for odd k and their symbolic representation (c); a construction of $f(G)$, (d).

Now we define a **2-edge gadget** consisting of the set of vertices $\{x, c, d, e, f\}$ with x and c adjacent and $\{c, d, e, f\}$ spanning a complete graph K_4 . One can check that every proper weighting w from $\{1, 2\}$ of a graph adjacent to such a gadget only at x requires $w(xc) = 2$. We refer to x as the *endpoint*.

We use the above gadgets to construct another gadget, called a **k -disallowing gadget**. As we will see, this gadget has similar properties as its namesake in Section 2. We therefore allow ourselves the re-use of the name for this new, slightly different, gadget. We assume that $k \geq 8$. The k -disallowing gadget contains a *main triangle* vxy with v referred to as the *root*. Moreover, if k is even, x and y each form the endpoint of $(k-6)/2$ edge disjoint 2-edge gadgets and x and y are each tops of distinct triangle gadgets (see Figure 2(a)). If k is odd, x and y each form the endpoint of $(k-3)/2$ edge disjoint 2-edge gadgets (see Figure 2(b)). Note that in any proper weighting w from $\{1, 2\}$, $w(vx) \neq w(vy)$; otherwise, since the weight contributed by gadgets to $\chi_w(x)$ and $\chi_w(y)$ is $k-3$, then $\chi_w(x) = w(xv) + w(xy) + k-3 = w(yv) + w(yx) + k-3 = \chi_w(y)$. Therefore, for any k , if $w(xy) = 1$ then $\{\chi_w(x), \chi_w(y)\} = \{k-1, k\}$ and, if $w(xy) = 2$ then $\{\chi_w(x), \chi_w(y)\} = \{k, k+1\}$. In either case, v has one neighbor $z \in \{x, y\}$ with $\chi_w(z) = k$, and consequently, $\chi_w(v) \neq k$ in any proper weighting from $\{1, 2\}$. Also $\{w(vx), w(vy)\} = \{1, 2\}$, and hence this gadget contributes exactly 3 to $\chi_w(v)$.

Now we are ready to show a polynomial time reduction from 3-COLOR to 1-2WEIGHT, f , such that $G \in 3\text{-COLOR}$ if and only if $f(G) \in 1\text{-2WEIGHT}$. Let $G = (V, E)$ be a graph of order n . As in Section 2, we may assume that $n \geq 3$. We construct the graph $f(G) = (W, F)$ as follows (see Figure 2(d)). We start with $G = (V, E)$. For each $v \in V$:

- (i) connect v to two new vertices s and t (distinct for each v);
- (ii) connect v to all vertices from a new set U_v (distinct for each v) with $|U_v| = n - 1 - \deg(v)$;

(iii) add $n - 1$ new k -disallowing gadgets for all $k \in \{4n + 1, 4n + 2, \dots, 5n - 1\}$ with v as their root.

Clearly, $f(G)$ can be calculated in time polynomial in the size of G .

Fact 3.1 *In $f(G)$ the following holds: any proper weighting w from $\{1, 2\}$ satisfies $\chi_w(v) \in \{4n - 2, 4n - 1, 4n\}$ for every $v \in V$.*

Proof: Fix $v \in V$. Since $w(vs) + w(vt) \in \{2, 3, 4\}$, v is the endpoint of $n - 1$ edges with endpoints in $V \cup U_v$ and v is the root of $(n - 1)$ k -disallowing gadgets (each contributing 3 to $\chi_w(v)$), we have:

$$\chi_w(v) \in \{2, 3, 4\} + \{n - 1, \dots, 2n - 2\} + \{3n - 3\} = \{4n - 2, \dots, 5n - 1\}.$$

Observing the above and the fact that v is the root of k -disallowing gadgets for all $k \in \{4n + 1, 4n + 2, \dots, 5n - 1\}$, we find that any proper weighting w from $\{1, 2\}$ satisfies $\chi_w(v) \in \{4n - 2, 4n - 1, 4n\}$, as claimed. \square

Now we show that $G \in 3\text{-COLOR}$ if and only if $f(G) \in 1\text{-2WEIGHT}$.

First let us assume that $G \in 3\text{-COLOR}$. That means there exists a proper 3-coloring of G , say $\chi : V \rightarrow \{4n - 2, 4n - 1, 4n\}$. We define a weighting of the edges of $f(G)$, $w : F \rightarrow \{1, 2\}$ as follows. For all $e \in E$ let $w(e) = 1$. For all edges $e = vu$ with $v \in V$ and $u \in U_v$ we set $w(e) = 1$. For all $v \in V$, if $\chi(v) = 4n - 2$ then $w(vs) = w(vt) = 1$; otherwise, if $\chi(v) = 4n - 1$ then $w(vs) = 1$ and $w(vt) = 2$; finally, if $\chi(v) = 4n$ then $w(vs) = w(vt) = 2$. All other edges (parts of gadgets) are weighted as follows: For a triangle gadget xab with root x , $w(xa) = 2, w(xb) = w(ab) = 1$. For a 2-gadget defined by $\{x, c, d, e, f\}$ with x adjacent to c , we have $w(xc) = w(cd) = w(ce) = w(de) = w(df) = 2$ and $w(cf) = w(ef) = 1$. For a k -disallowing gadget with root v and main triangle vxy , $w(vx) = w(xy) = 2, w(vy) = 1$, and the weighting of all other gadgets as described above. Note that w is a proper weighting of $f(G)$ (satisfying $\chi_w(v) = \chi(v)$ for all $v \in V$), as required.

Next let us assume that $G \notin 3\text{-COLOR}$. Therefore, for all $\chi : V \rightarrow \{4n - 2, 4n - 1, 4n\}$, χ is not a proper vertex coloring. But, from Fact 3.1, any proper weighting from $\{1, 2\}$ of $f(G)$ satisfies $\chi_w(v) \in \{4n - 2, 4n - 1, 4n\}$ for all $v \in V$. Thus, there is no such proper weighting and hence $f(G) \notin 1\text{-2WEIGHT}$.

This concludes the proof of Theorem 1.1.

4 Concluding remarks

In this paper we showed that determining whether a graph has a proper weighting from either $\{0, 1\}$ or $\{1, 2\}$ is NP-complete. As a matter of fact, these two problems are not the same, in the sense that the corresponding families of graphs 0-1WEIGHT and 1-2WEIGHT are not equal. For example, the graph consisting only of one 2-edge gadget is in 1-2WEIGHT, as seen before, but it is easy to check that it is not in 0-1WEIGHT. Furthermore, we believe that our approach can be generalized to show that determining whether a graph has a proper weighting from $\{a, b\}$ is NP-complete for any different rational numbers a and b . It is not clear if the same would hold for any two distinct irrational numbers.

Acknowledgements

We would like to thank Michał Karoński, who introduced us to the 1-2-3 conjecture. We are also very grateful to the referees for their detailed comments on an earlier version of this paper.

References

- L. Addario-Berry, K. Dalal, C. McDiarmid, B. A. Reed, and A. Thomason. Vertex-colouring edge-weightings. *Combinatorica*, 27(1):1–12, 2007. ISSN 0209-9683. doi: <http://dx.doi.org/10.1007/s00493-007-0041-6>.
- L. Addario-Berry, K. Dalal, and B. A. Reed. Degree constrained subgraphs. *Discrete Appl. Math.*, 156(7):1168–1174, 2008. ISSN 0166-218X. doi: DOI:10.1016/j.dam.2007.05.059. URL <http://www.sciencedirect.com/science/article/B6TYW-4PT2FK1-1/2/f3371b3c9c874a200bf561073f921e12>.
- M. Kalkowski, M. Karoński, and F. Pfender. Vertex-coloring edge-weightings: Towards the 1-2-3-conjecture. *J. Comb. Theory, Ser. B*, 100(3):347–349, 2010.
- M. Karoński, T. Łuczak, and A. Thomason. Edge weights and vertex colours. *J. Combin. Theory Ser. B*, 91:151–157, 2004.
- T. Wang and Q. Yu. On vertex-coloring 13-edge-weighting. *Front. Math. China*, 3(4):581–587, 2008. ISSN 1673-3452 (Print) 1673-3576 (Online). doi: 10.1007/s11464-008-0041-x.