

The b-chromatic number of powers of cycles

Anja Kohl

► **To cite this version:**

Anja Kohl. The b-chromatic number of powers of cycles. *Discrete Mathematics and Theoretical Computer Science, DMTCS*, 2013, Vol. 15 no. 1 (1), pp.147–156. <hal-00990613>

HAL Id: hal-00990613

<https://hal.inria.fr/hal-00990613>

Submitted on 13 May 2014

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The b -chromatic number of powers of cycles

Anja Kohl[†]

Faculty of Information Technology/Mathematics, HTW Dresden, Germany

received 16th May 2011, revised 18th July 2012, accepted 13th March 2013.

A b -coloring of a graph G by k colors is a proper vertex coloring such that each color class contains a color-dominating vertex, that is, a vertex having neighbors in all other $k - 1$ color classes. The b -chromatic number $\chi_b(G)$ is the maximum integer k for which G has a b -coloring by k colors. Let C_n^r be the r th power of a cycle of order n . In 2003, Effantin and Kheddouci established the b -chromatic number $\chi_b(C_n^r)$ for all values of n and r , except for $2r + 3 \leq n \leq 3r$. For the missing cases they presented the lower bound $L := \min\{n - r - 1, r + 1 + \lfloor \frac{n-r-1}{3} \rfloor\}$ and conjectured that $\chi_b(C_n^r) = L$. In this paper, we determine the exact value on $\chi_b(C_n^r)$ for the missing cases. It turns out that $\chi_b(C_n^r) > L$ for $2r + 3 \leq n \leq 2r + 3 + \frac{r-6}{4}$.

Keywords: b -chromatic number, coloring, b -coloring, powers of cycles

1 Introduction

Let $G = (V, E)$ be a simple, undirected graph with vertex set V and edge set E . For $x, y \in V$ we denote by $d(x, y)$ the distance between x and y , which is the number of edges in a shortest (x, y) -path. The r th power of a graph G , written as G^r , is a graph on the same vertex set such that two vertices are joined by an edge if and only if their distance in G is at most r . For $r \geq 1$, let C_n^r and P_n^r denote the r th power of a cycle and a path on n vertices, respectively.

A b -coloring of a graph G by k colors is a proper vertex coloring such that each color class contains a vertex having neighbors in all other $k - 1$ color classes. Such a vertex is called a *color-dominating vertex*. The b -chromatic number $\chi_b(G)$ is the maximum integer k for which G has a b -coloring by k colors. The concept of b -colorings and the b -chromatic number were introduced by Irving and Manlove [5] in 1999 and have already been investigated in more than 50 papers (cf. [1]-[7]).

Since the problem to decide whether $\chi_b(G) \geq K$ for a given graph G and an integer K is \mathcal{NP} -hard in general ([5]), we are interested in exact values on the b -chromatic number for special graphs or graph classes. In 2003, Effantin and Kheddouci [3] determined $\chi_b(P_n^r)$ for all values of n and r . They also investigated powers of cycles C_n^r and obtained the following result:

[†]Email: kohl@informatik.htw-dresden.de.

Theorem 1 (Effantin and Kheddouci, [3]) *The b -chromatic number of C_n^r for $r \geq 1$ is*

$$\chi_b(C_n^r) = \begin{cases} n & \text{if } n \leq 2r + 1, \\ r + 1 & \text{if } n = 2r + 2, \\ r + 1 + \lfloor \frac{n-r-1}{3} \rfloor & \text{if } 3r + 1 \leq n \leq 4r, \\ 2r + 1 & \text{if } n \geq 4r + 1, \end{cases} \quad (1)$$

$$\text{and } \chi_b(C_n^r) \geq \min\{n - r - 1, r + 1 + \lfloor \frac{n-r-1}{3} \rfloor\} \quad \text{if } 2r + 3 \leq n \leq 3r. \quad (2)$$

The determination of $\chi_b(C_n^r)$ for $2r + 3 \leq n \leq 3r$ was posed as an open problem, but the authors conjectured that the lower bound in (2) is the exact value.

In the present paper, we complete Effantin and Kheddouci's work by determining $\chi_b(C_n^r)$ for the missing cases $2r + 3 \leq n \leq 3r$. It turns out that the lower bound in (2) is indeed the exact value on $\chi_b(C_n^r)$ for $2r + 3 + \frac{r-6}{4} < n \leq 3r$, but not for $2r + 3 \leq n \leq 2r + 3 + \frac{r-6}{4}$.

2 Main Result

Theorem 2 *Let C_n^r be the r th power of a cycle of order n with $2r + 3 \leq n \leq 3r$. Moreover, set $l := n - 2(r + 1)$, $R := (r + 1 + 3l) \bmod (5l)$, and $m := \max\{\lfloor \frac{R-2l}{3} \rfloor, 0\}$. Then, $\chi_b(C_n^r) = \frac{3n-R}{5} + m$.*

2.1 Preliminaries

In the following, we only consider powers of cycles C_n^r for which $2r + 3 \leq n \leq 3r$ is satisfied.

Let V be the vertex set of C_n^r and set $l := n - 2(r + 1)$. Then, the complement of C_n^r is an $(l + 1)$ -regular graph and $1 \leq l \leq r - 2$. Moreover, C_n^r has independence number 2.

Let c be a b -coloring of C_n^r by $k := \chi_b(C_n^r)$ colors and let V_1, \dots, V_k be the corresponding color classes such that $1 \leq |V_1| \leq |V_2| \leq \dots \leq |V_k| \leq 2$. By a we denote the number of color classes of cardinality 1. Choose a color-dominating vertex $v_i \in V_i$ for $i = 1, \dots, k$. Let $A := \{v_1, \dots, v_a\}$, $B := \{v_{a+1}, \dots, v_k\}$, and $C := \{w_{a+1}, \dots, w_k\}$ where w_i is the *partner* of v_i , i.e. the vertex in $V_i \setminus \{v_i\}$, for $i \in \{a+1, \dots, k\}$. Vertices from the set A , B , and C are called A -, B -, and C -vertices, respectively.

A *vertex-row* X shall be a set of vertices from V which are consecutive on the underlying cycle C_n . If X only consists of (non-) A -vertices, then we call it a (non-) A -*vertex-row* (analogously for B and C).

The neighborhood $N(x)$ of a vertex $x \in V$ is the set of neighbors of x . Moreover, let $\overline{N}(x) := V \setminus (N(x) \cup \{x\})$ be the *non-neighborhood* of x and for a set $V' \subseteq V$ let $\overline{N}(V') := \bigcup_{x \in V'} \overline{N}(x)$. Note that $|\overline{N}(X)| = l + |X|$ for every vertex-row X .

For a vertex $u \in V$ and a set $V' \subseteq V$ let $d_{C_n}(u, V') := \min\{d_{C_n}(u, v) \mid v \in V'\}$ where $d_{C_n}(u, v)$ denotes the distance of u and v in the underlying C_n .

We refer to Figure 1 for the following considerations.

Let X be an A -vertex-row with $\alpha \geq 1$ vertices. We denote the two subgraphs induced by the vertex-rows between X and $\overline{N}(X)$ in clockwise order and in anti-clockwise order (on the underlying C_n) by $G_r(X)$ and $G_l(X)$, respectively (compare left side of Figure 1). From above we know that $\overline{N}(X)$ contains exactly $l + \alpha$ vertices. Moreover, the set $\overline{N}(\overline{N}(X)) \setminus X$ has then exactly $2l$ vertices, and by symmetry we obtain $|\overline{N}(\overline{N}(X)) \cap V(G_l(X))| = |\overline{N}(\overline{N}(X)) \cap V(G_r(X))| = l$. Since each vertex from $\overline{N}(X)$ is non-adjacent to at least one A -vertex (which corresponds to a color class of cardinality 1), it cannot be color-dominating. So, each vertex from $\overline{N}(X)$ is a C -vertex and therefore has a B -vertex as partner. Let $\overline{N}_p(X)$ be the set of partners of the vertices from $\overline{N}(X)$. Clearly, $\overline{N}_p(X) \subseteq \overline{N}(\overline{N}(X)) \setminus X$.




Fig. 1: Non-neighborhood $\overline{N}(X)$ and extended non-neighborhood $\widehat{N}(X)$ of an A -vertex-row X

Let $v_l(X)$ be the vertex from $\overline{N}_p(X) \cap V(G_l(X))$ that have the largest distance to the set X in C_n , i.e. $d_{C_n}(v_l(X), X) = \max_{v \in \overline{N}_p(X) \cap V(G_l(X))} d_{C_n}(v, X)$. Moreover, set $\beta_l(X) := |\overline{N}_p(X) \cap V(G_l(X))|$.

Analogously, we define $v_r(X)$ and $\beta_r(X)$. From above we know that $\beta_l(X) \leq d_{C_n}(v_l(X), X) \leq |\overline{N}(\overline{N}(X)) \cap V(G_l(X))| = l$ and, analogously, $\beta_r(X) \leq d_{C_n}(v_r(X), X) \leq l$. Along with $\beta_l(X) + \beta_r(X) = |\overline{N}_p(X)| = |\overline{N}(X)| = l + \alpha \geq l + 1$, we further obtain $\beta_l(X), \beta_r(X) \geq 1$. This implies that both vertices $v_l(X)$ and $v_r(X)$ exist.

Since the partners of $v_l(X)$ and $v_r(X)$ are in $\overline{N}(X)$, we know that the sets $\overline{N}(v_l(X))$ and $\overline{N}(X)$ as well as the sets $\overline{N}(v_r(X))$ and $\overline{N}(X)$ intersect, and we can determine that $|\overline{N}(v_l(X)) \setminus \overline{N}(X)| = d_{C_n}(v_l(X), X)$ and $|\overline{N}(v_r(X)) \setminus \overline{N}(X)| = d_{C_n}(v_r(X), X)$. Since each vertex from $\overline{N}(v_l(X))$ is non-adjacent to the color-dominating vertex $v_l(X)$, it cannot be an A -vertex. The same holds for the vertices from $\overline{N}(v_r(X))$. So, altogether, $\overline{N}(v_l(X)) \cup \overline{N}(X) \cup \overline{N}(v_r(X))$ is a non- A -vertex-row with $l + \alpha + d_{C_n}(v_l(X), X) + d_{C_n}(v_r(X), X) \geq l + \alpha + \beta_l(X) + \beta_r(X) = 2(l + \alpha)$ vertices. We call this set the *extended non-neighborhood* of X and denote it by $\widehat{N}(X)$. If X consists of only one vertex x , then we just write $\widehat{N}(x)$ instead of $\widehat{N}(\{x\})$. The same we do with all previously defined notations.

Observation 1 Let X_1, \dots, X_t be a family of $t \geq 1$ disjoint A -vertex-rows (in clockwise order on the C_n) such that $\overline{N}(X_i) \cap \overline{N}(X_{i+1}) \neq \emptyset$ for $i \in \{1, \dots, t-1\}$. Moreover, we assume the given A -vertex-rows to be maximal, i.e. each of it is delimited by non- A -vertices. Then, $\sum_{i=1}^t |X_i| \leq l$.

Proof: Let Y_i be the non- A -vertex-row between X_i and X_{i+1} for $i \in \{1, \dots, t-1\}$ and let Y_0 and Y_t be the vertex-rows with l vertices preceding X_1 and succeeding X_t , respectively. Moreover, set $\mathcal{X} := \bigcup_{i=1}^t X_i$ and $\mathcal{Y} := \bigcup_{i=1}^{t-1} Y_i$. Note that $\mathcal{X} \cup \mathcal{Y}$ is a vertex-row and therefore satisfies $|\overline{N}(\mathcal{X} \cup \mathcal{Y})| = l + |\mathcal{X} \cup \mathcal{Y}|$.

Since $\overline{N}(X_i)$ and $\overline{N}(X_{i+1})$ intersect for each $i \in \{1, \dots, t-1\}$, it follows that $\overline{N}(Y_i) \subset (\overline{N}(X_i) \cup \overline{N}(X_{i+1}))$ and therefore $\overline{N}(\mathcal{Y}) \subset \overline{N}(\mathcal{X})$. Moreover, it implies that $\overline{N}(\mathcal{X})$ is a C -vertex-row. For its cardinality we obtain:

$$\begin{aligned}
|\overline{N}_p(\mathcal{X})| &= |\overline{N}(\mathcal{X})| = |\overline{N}(\mathcal{X}) \cup \overline{N}(\mathcal{Y})| = |\overline{N}(\mathcal{X} \cup \mathcal{Y})| = l + |\mathcal{X} \cup \mathcal{Y}| \\
&= l + \left| \bigcup_{i=1}^t X_i \cup \bigcup_{i=1}^{t-1} Y_i \right| = l + \left| \bigcup_{i=1}^t X_i \right| + \left| \bigcup_{i=1}^{t-1} Y_i \right| = l + \sum_{i=1}^t |X_i| + \sum_{i=1}^{t-1} |Y_i|. \quad (3)
\end{aligned}$$

Moreover, we can easily check that $\overline{N}(\overline{N}(\mathcal{X})) = \bigcup_{i=1}^t X_i \cup \bigcup_{i=0}^t Y_i = \mathcal{X} \cup \bigcup_{i=0}^t Y_i$. Since $\overline{N}_p(\mathcal{X}) \subseteq \overline{N}(\overline{N}(\mathcal{X})) \setminus \mathcal{X}$, we then obtain:

$$|\overline{N}_p(\mathcal{X})| \leq |\overline{N}(\overline{N}(\mathcal{X})) \setminus \mathcal{X}| = \left| \bigcup_{i=0}^t Y_i \right| \leq \sum_{i=0}^t |Y_i| = 2l + \sum_{i=1}^{t-1} |Y_i|. \quad (4)$$

So Inequalities (3) and (4) yield $\sum_{i=1}^t |X_i| \leq l$. \square

Note that for $t = 1$, the previous observation yields that every A -vertex-row contains at most l vertices.

2.2 Proof of Theorem 2

Recall that $k = \chi_b(C_n^r)$, $a = |A|$, and $l = n - 2(r + 1)$. Since the b -coloring by k colors partitions the vertex set into a color classes of cardinality 1 and $k - a$ color classes of cardinality 2, we obtain $n = a + 2(k - a)$ and, thus, $a = 2k - n = 2k - (l + 2r + 2) = 2(k - r - 1) - l$. So, a , l , and n have the same parity. Moreover, according to Inequality (2), $k \geq \min\{n - (r + 1), r + 1 + \lfloor \frac{n - (r + 1)}{3} \rfloor\} = \min\{l + r + 1, r + 1 + \lfloor \frac{l + r + 1}{3} \rfloor\} = r + 1 + \min\{l, \lfloor \frac{l + r + 1}{3} \rfloor\}$ and therefore

$$a + l = 2k - n + l = 2k - 2(r + 1) \geq 2 \min\left\{l, \left\lfloor \frac{l + r + 1}{3} \right\rfloor\right\}. \quad (5)$$

Let Q and R be the quotient and the remainder of the integer division $r + 1 + 3l$ by $5l$, i.e. $r + 1 + 3l = Q \cdot 5l + R$, where $Q \geq 0$ and $0 \leq R < 5l$. Note that then $Q = \lfloor \frac{r + 1 + 3l}{5l} \rfloor$, $R = (r + 1 + 3l) \bmod (5l)$, and $n = l + 2(r + 1) = (2Q - 1)5l + 2R$. Moreover, set $m := \max\{\lfloor \frac{R - 2l}{3} \rfloor, 0\}$.

Case 1. $l + 3 \leq r + 1 < 2l$, i.e. $\frac{5(r + 1)}{2} < n \leq 3r$.

At first we notice that this case is only possible for $l \geq 4$. Moreover, by (5) it follows that

$$a + l \geq 2 \left\lfloor \frac{l + r + 1}{3} \right\rfloor \geq \frac{2(l + r - 1)}{3} = \frac{2r - (l + 2)}{3} + l \geq \frac{l + 2}{3} + l \geq 2 + l. \quad (6)$$

Hence, there exist at least two A -vertices. Choose two vertices $u_1, u_2 \in A$ of maximum distance in C_n . W.l.o.g. u_1 and u_2 are ordered in such a way, that a shortest (u_1, u_2) -path in C_n runs anti-clockwise. Then there are two cases to consider depending on the positions of u_2 and $v_l(u_1)$ (compare Figure 2). From the preliminaries and the premise of the case, we know that the extended non-neighborhood of u_1 has cardinality $|\widehat{N}(u_1)| \geq 2(l + 1) \geq r + 4$. So, since $\widehat{N}(u_1)$ is a vertex-row of at least $r + 4$ vertices, but the maximal vertex-rows induced by $N(u_2)$ have only r vertices, we deduce that u_2 cannot be adjacent to all vertices in $\widehat{N}(u_1)$, and so $\widehat{N}(u_1)$ and $\overline{N}(u_2)$ intersect. Hence, $\widehat{N}(u_1) \cup \overline{N}(u_2)$ is a non- A -vertex-row.




Fig. 2: Extended non-neighborhood $\widehat{N}(u_1)$ and non-neighborhood $\overline{N}(u_2)$ intersect

Subcase 1.1. u_2 and $v_l(u_1)$ are in clockwise order (see left side of Figure 2).

Since $\widehat{N}(u_1) \cup \overline{N}(u_2)$ is a non- A -vertex-row and by the choice of u_1 and u_2 as two A -vertices of maximum distance, there is no A -vertex in $V(G_l(u_2)) \cup \overline{N}(u_2)$ (below the double-line). The remaining vertex set $V(G_r(u_2)) \cup \{u_2\}$ (above the double-line) induces a vertex-row with exactly $r + 1$ vertices. This vertex-row has to contain all a A -vertices and all $l + 1$ B -vertices from $\overline{N}_p(u_1)$. Hence, $r + 1 \geq a + l + 1$. This and the premise of the case yield:

$$a + l \leq r = \frac{(r + 2) + 2(r + 1) - 4}{3} \leq \frac{2l + 2(r + 1) - 4}{3} \leq 2 \left\lfloor \frac{l + r + 1}{3} \right\rfloor. \quad (7)$$

Along with Inequality (6), we obtain $a + l = 2 \lfloor \frac{l+r+1}{3} \rfloor$.

Subcase 1.2. u_2 and $v_l(u_1)$ are in anti-clockwise order (see right side of Figure 2).

By the choice of u_1 and u_2 , there is no A -vertex in $V(G_l(u_2)) \cup \overline{N}(u_2) \cup V(G_r(u_1)) \cup \overline{N}(u_1)$ (below the double-curve). Hence, the remaining vertex set $(V(G_r(u_2)) \cap V(G_l(u_1))) \cup \{u_1, u_2\}$ (above the double-curve) that induces a vertex-row with $1 + d_{C_n}(u_1, u_2)$ vertices has to contain all a A -vertices. Since $d_{C_n}(u_1, u_2) \leq d_{C_n}(u_1, v_l(u_1)) - 1 \leq l - 1$, the non-neighborhoods $\overline{N}(u_1)$ and $\overline{N}(u_2)$ intersect. This implies that $\overline{N}(A) = \overline{N}(u_1) \cup \overline{N}(u_2)$ and therefore $|\overline{N}(A)| = l + 1 + d_{C_n}(u_1, u_2) \geq l + a$. Since $\overline{N}(A) \subseteq C$, $\overline{N}_p(A) \subseteq B$, and $|\overline{N}(A)| = |\overline{N}_p(A)|$, we deduce that $n = |A| + |B| + |C| \geq a + |\overline{N}_p(A)| + |\overline{N}(A)| = a + 2 \cdot |\overline{N}(A)| \geq a + 2(l + a) = 3a + 2l$. Along with $n = l + 2(r + 1)$ we then obtain $a + l \leq \lfloor \frac{2(l+r+1)}{3} \rfloor$. This and Inequality (6) yield

$$2 \left\lfloor \frac{l + r + 1}{3} \right\rfloor \leq a + l \leq \left\lfloor \frac{2(l + r + 1)}{3} \right\rfloor, \quad (8)$$

and since a and l have the same parity implying that $a + l$ is even, we conclude that $a + l = 2 \lfloor \frac{l+(r+1)}{3} \rfloor$.

Hence, in both subcases we obtain $a+l = 2 \lfloor \frac{l+r+1}{3} \rfloor$ which implies that $k = r+1 + \lfloor \frac{l+r+1}{3} \rfloor$. Moreover, since $r + 1 + 3l < 5l$ by the premise of the case, it follows that $R = r + 1 + 3l$ and $m = \lfloor \frac{r+1+l}{3} \rfloor$. Therefore, $k = r + 1 + \lfloor \frac{l+r+1}{3} \rfloor = r + 1 + m = \frac{3(l+2(r+1)) - (r+1+3l)}{5} + m = \frac{3n-R}{5} + m$.

Case 2. $r + 1 \geq 2l \geq 2$, i.e. $2r + 3 \leq n \leq \frac{5(r+1)}{2}$.

The premise yields $r + 1 + 3l = Q \cdot 5l + R \geq 5l$ and therefore $Q \geq 1$.




Fig. 3: Partition of V into $2Q - 1$ bundles of $5l$ vertices each and $2R$ remaining vertices

[“ \geq ”] At first we prove that $k \geq (2Q - 1)3l + R + m = \frac{3n-R}{5} + m$.

In order to do this we construct a b -coloring of C_n^r by $k' := (2Q - 1)3l + R + m$ colors.

Since $n = (2Q - 1)5l + 2R$, we can partition the vertex set V into $2Q - 1$ bundles $\mathcal{B}_1, \dots, \mathcal{B}_{2Q-1}$ of $5l$ vertices each and $2R$ remaining vertices. For $i \in \{1, \dots, 2Q - 1\}$, the bundle \mathcal{B}_i shall consist of a vertex-row X_i with l black vertices, the non-neighborhood $\overline{N}(X_i)$ with $2l$ grey vertices, and the set $\overline{N}_p(X_i) := \overline{N}(\overline{N}(X_i)) \setminus X_i$ with $2l$ black-white vertices, such that the sets X_1, \dots, X_{2Q-1} are positioned on the cycle C_n as described in Figure 3. In order to do this, we start with an arbitrary vertex-row with l vertices as X_1 (on top in the picture). Then the positions of $\overline{N}(X_1)$ and $\overline{N}_p(X_1)$ are already specified. After this, we position X_2 (bottom left) and X_3 (bottom right) leaving out l vertices on each side of $\overline{N}(X_1)$. Then the positions of $\overline{N}(X_2)$ and $\overline{N}_p(X_2)$ as well as $\overline{N}(X_3)$ and $\overline{N}_p(X_3)$ are already specified. If $Q \geq 3$, then we proceed by positioning X_4 (top left) and X_5 (top right) leaving out l vertices on the left side of $\overline{N}(X_3)$ and on the right side of $\overline{N}(X_2)$, respectively. Then also the positions of $\overline{N}(X_4)$ and $\overline{N}_p(X_4)$ as well as $\overline{N}(X_5)$ and $\overline{N}_p(X_5)$ are specified. If $Q \geq 4$, then we position the remaining sets X_6, \dots, X_{2Q-1} in the same manner as we positioned $X_2 - X_5$ using always the order bottom left, bottom right, top left, top right. The $2R$ remaining vertices induce two vertex-rows with R vertices each which we denote by \mathcal{R}_l and \mathcal{R}_r .

Now we color the l vertices from X_1 in clockwise order by colors $1, \dots, l$. Then we color the $2l$ vertices from $\overline{N}(X_1)$ in clockwise order by colors $l + 1, \dots, 3l$, and the same we do for the $2l$ vertices from $\overline{N}(\overline{N}(X_1)) \setminus X_1 = \overline{N}_p(X_1)$. Now all vertices from \mathcal{B}_1 are colored. Next we color the vertices from \mathcal{B}_2 in an analogous way by colors $3l + 1, \dots, 6l$, and we continue until \mathcal{B}_{2Q-1} whose vertices we color by colors $(2Q - 2)3l + 1, \dots, (2Q - 1)3l$. This coloring procedure yields a coloring of $(2Q - 1)5l = n - 2R$ vertices by $(2Q - 1)3l$ colors (compare the left side of Figure 5 for the case $l = Q = 2$). For the coloring of the vertices from $\mathcal{R}_l \cup \mathcal{R}_r$ we have to distinguish the following two cases:




Fig. 4: Partition of the remaining $2R$ vertices for the case $m > 0$

Subcase 2.1. $0 \leq R \leq 2(l + 1)$.

This implies that $m = 0$. We color the vertices from \mathcal{R}_l by colors $(2Q - 1)3l + 1, \dots, (2Q - 1)3l + R$ in clockwise order. The same we do with the vertices from \mathcal{R}_r . Altogether, we obtain a coloring c_b of C_n^r by $(2Q - 1)3l + R = k'$ colors.

Subcase 2.2. $2(l + 1) < R < 5l$.

Here, $m > 0$. We partition $\mathcal{R}_l \cup \mathcal{R}_r$ as depicted in Figure 4 into two bundles $\mathcal{B}_{2Q}, \mathcal{B}_{2Q+1}$ of $2l + 3m$ vertices each and two vertex-rows \mathcal{R}'_l and \mathcal{R}'_r of $R - 2l - 3m$ vertices each. For $i \in \{2Q, 2Q + 1\}$, the bundle \mathcal{B}_i shall consist of a vertex-row X_i with m black vertices, the non-neighborhood $\overline{N}(X_i)$ with $l + m$ grey vertices, and $\overline{N}_p(X_i)$ which is a subset of $l + m$ black-white vertices from the set $\overline{N}(\overline{N}(X_i)) \setminus X_i$.

Now we color the vertices from X_{2Q} in clockwise order by colors $(2Q - 1)3l + 1, \dots, (2Q - 1)3l + m$. Then we color the vertices from $\overline{N}(X_{2Q})$ as well as from $\overline{N}_p(X_{2Q})$ in clockwise order by colors $(2Q - 1)3l + m + 1, \dots, (2Q - 1)3l + 2m + l$. Analogously, we color the vertices from $X_{2Q+1}, \overline{N}(X_{2Q+1})$, and $\overline{N}_p(X_{2Q+1})$ by colors $(2Q - 1)3l + 2m + l + 1, \dots, (2Q - 1)3l + 4m + 2l$. Finally, we color the vertices from \mathcal{R}'_l as well as from \mathcal{R}'_r in clockwise order by colors $(2Q - 1)3l + 4m + 2l + 1, \dots, (2Q - 1)3l + 4m + 2l + (R - 2l - 3m) = (2Q - 1)3l + R + m$ (compare the right side of Figure 5 for the case $l = Q = 2, m = 1$). Altogether, this yields a coloring c_b of C_n^r by $(2Q - 1)3l + R + m = k'$ colors.




Fig. 5: b -coloring by $k' = (2Q - 1)3l + R + m$ colors (here for $l = Q = 2, m = 1$)

It remains to prove that c_b is a b -coloring for both subcases. Consider the Figures 3-5: The black vertices correspond to the color classes of cardinality 1 in the coloring c_b . Moreover, each color class of cardinality 2 consists of a black-white and a grey vertex. Note that in Subcase 2.1 (resp. 2.2) we assume the vertices in R_l (resp. R'_l) to be black-white and the vertices in R_r (resp. R'_r) to be grey.

Because C_n^r has independence number 2, a vertex x from a color class V_j has at least one neighbor in each color class V_g of cardinality 2 for $j \neq g$. Hence, x is color-dominating if and only if x is adjacent to each vertex in a color class V_h of cardinality 1 for $j \neq h$, i.e. if x is adjacent to all (other) black vertices.

Along with the fact that the black vertices are the vertices from $\bigcup_{i=1}^s X_i$ ($s := 2Q - 1$ in Subcase 2.1 and $s := 2Q + 1$ in Subcase 2.2) this implies that x is color-dominating if and only if $x \notin \overline{N}(\bigcup_{i=1}^s X_i) = \bigcup_{i=1}^s \overline{N}(X_i)$. Since all vertices from $\bigcup_{i=1}^s \overline{N}(X_i)$ are grey, we deduce that the black and the black-white vertices are color-dominating and correspond to the A - and the B -vertices, respectively. The grey vertices are then the C -vertices. Moreover, since every color class contains either a black or a black-white vertex it follows that every color class has a color-dominating vertex. Hence, c_b is a proper b -coloring of C_n^r .

For the number k' of colors used in c_b we obtain:

$$\begin{aligned} k' &= (2Q - 1)3l + R + m = r + 1 + Ql + m = \frac{5(r + 1) + (r + 1 + 3l - R)}{5} + m \\ &= \frac{3(l + 2(r + 1)) - R}{5} + m = \frac{3n - R}{5} + m. \end{aligned}$$

This implies that $k \geq k'$ and because of $a = 2k - n$ we further deduce that $a \geq 2k' - n = 2(r + 1 + Ql + m) - (l + 2(r + 1)) = (2Q - 1)l + 2m$.

[“ \leq ”] Now we prove that $k \leq r + 1 + Ql + m = \frac{3n - R}{5} + m$.

Consider the b -coloring c of C_n^r by k colors and suppose that $k > k'$. This is equivalent to $a > (2Q - 1)l + 2m$. Since a and l have the same parity, it suffices to consider $a \geq (2Q - 1)l + 2(m + 1)$. Choose an arbitrary A -vertex-row X with α , $1 \leq \alpha \leq l$, vertices. We refer to Figure 1 for the following considerations.

Let $\beta_l(X)$ and $\beta_r(X)$ be defined as before. In the following, we abbreviate these values by β_l and β_r . Moreover, let $a_l := |A \cap V(G_l(X))|$, $b_l := |(B \setminus \overline{N}_p(X)) \cap V(G_l(X))|$, and $c_l := |C \cap V(G_l(X))|$. Analogously, we define a_r , b_r , and c_r for the subgraph $G_r(X)$. Since $G_l(X)$ and $G_r(X)$ have the same order, we deduce that $|V(G_l(X))| = \beta_l + a_l + b_l + c_l = |V(G_r(X))| = \beta_r + a_r + b_r + c_r$. Every vertex from the set $(B \setminus \overline{N}_p(X)) \cap V(G_l(X))$ has a partner in $C \cap V(G_r(X))$, and vice versa. Hence, $b_l = c_r$. Analogously, we obtain $b_r = c_l$. Moreover, $\beta_l + \beta_r = |\overline{N}_p(X)| = |\overline{N}(X)| = l + \alpha$ and $a = \alpha + a_l + a_r$. Therefore, $\beta_l + a_l = \beta_r + a_r = (l + \alpha - \beta_l) + (a - \alpha - a_l)$ and altogether, $2(\beta_l + a_l) = 2(\beta_r + a_r) = l + a$. By the assumption $a \geq (2Q - 1)l + 2(m + 1)$ we obtain $2(\beta_l + a_l) = 2(\beta_r + a_r) = l + a \geq 2Ql + 2(m + 1)$ and thus, $\beta_l + a_l = \beta_r + a_r \geq Ql + m + 1$. Moreover, since $1 \leq \beta_l, \beta_r \leq l$ (compare the preliminaries), we deduce that $Ql + m + 1 \leq \beta_l + a_l \leq l + a_l$ which yields $a_l \geq (Q - 1)l + m + 1$. Analogously, we obtain $a_r \geq (Q - 1)l + m + 1$.

Let $X_1, \dots, X_{q'}$ be the family of all maximal A -vertex-rows in $V(G_l(X))$ (in clockwise order). For $i \in \{1, \dots, q' - 1\}$ merge the two sets X_i and X_{i+1} if $\overline{N}(X_i) \cap \overline{N}(X_{i+1}) \neq \emptyset$. Continue doing this until the remaining sets, say $\mathcal{X}_1, \dots, \mathcal{X}_q$ (in clockwise order), satisfy $\overline{N}(\mathcal{X}_i) \cap \overline{N}(\mathcal{X}_{i+1}) = \emptyset$ for $i \in \{1, \dots, q - 1\}$. Set $\alpha_i := |\mathcal{X}_i|$ for $i \in \{1, \dots, q\}$. Obviously, $\sum_{i=1}^q \alpha_i = a_l$. Moreover, from Observation 1 we know that $\alpha_i \leq l$ for $i \in \{1, \dots, q\}$. Hence, $ql \geq \sum_{i=1}^q \alpha_i = a_l \geq (Q - 1)l + m + 1 \geq (Q - 1)l + 1$ and therefore $q \geq Q$.

Since every vertex from $\bigcup_{i=1}^q \overline{N}(\mathcal{X}_i)$ is a C -vertex from $V(G_r(X))$ we further deduce that $b_l = c_r \geq |\bigcup_{i=1}^q \overline{N}(\mathcal{X}_i)| = \sum_{i=1}^q |\overline{N}(\mathcal{X}_i)| \geq \sum_{i=1}^q (l + \alpha_i) = a_l + ql \geq a_l + Ql$. Analogously, we can show that $b_r = c_l \geq a_r + Ql$. For the number of vertices n we then obtain:

$$\begin{aligned} n &= |X| + |\overline{N}(X)| + |\overline{N}_p(X)| + a_l + a_r + b_l + b_r + c_l + c_r \\ &= \alpha + 2(l + \alpha) + (a_l + a_r) + 2(b_l + b_r) = a + 2(l + \alpha) + 2(b_l + b_r) \\ &\geq a + 2(l + \alpha) + 2(a_r + a_l + 2Ql) = 3a + 2l + 4Ql \\ &\geq 3(2Q - 1)l + 6(m + 1) + 2l + 4Ql = 10Ql - l + 6(m + 1) \end{aligned} \quad (9)$$

Moreover, since $n = (2Q - 1)5l + 2R$, Inequality (9) yields $R \geq 2l + 3(m + 1)$ and therefore $\lfloor \frac{R-2l}{3} \rfloor \geq m + 1$. This is a contradiction to $m = \max \{ \lfloor \frac{R-2l}{3} \rfloor, 0 \}$.

Therefore, $k \leq k'$, and altogether $k = k' = \frac{3n-R}{5} + m$. \square

3 Conclusion

In [3], Effantin and Kheddouci conjectured that $\chi_b(C_n^r) = \min\{n - (r + 1), r + 1 + \lfloor \frac{n-(r+1)}{3} \rfloor\}$ for $2r + 3 \leq n \leq 3r$. The following observation shows that this conjecture does not hold in general.

Observation 2 Let C_n^r be the r th power of a cycle of order n with $2r + 3 \leq n \leq 3r$, and set $L := \min\{n - (r + 1), r + 1 + \lfloor \frac{n-(r+1)}{3} \rfloor\}$. Then, $\chi_b(C_n^r) = L$ if $n > \frac{9r+6}{4}$ and $\chi_b(C_n^r) > L$ if $n \leq \frac{9r+6}{4}$.

Proof: Recall that $l = n - 2(r + 1)$, $R = (r + 1 + 3l) \bmod (5l)$, $Q = \frac{r+1+3l-R}{5l}$, and $m = \max \{ \lfloor \frac{R-2l}{3} \rfloor, 0 \}$. So we obtain $L = r + 1 + \min\{l, \lfloor \frac{r+1+l}{3} \rfloor\}$.

Case 1. $l + 3 \leq r + 1 < 2l$, i.e. $\frac{5(r+1)}{2} < n \leq 3r$.

Then, $L = r + 1 + \lfloor \frac{r+1+l}{3} \rfloor$. Moreover, we obtain $r + 1 + 3l < 5l$ which implies that $R = r + 1 + 3l$ and $m = \lfloor \frac{R-2l}{3} \rfloor = \lfloor \frac{r+1+l}{3} \rfloor$. Thus, Theorem 2 yields $\chi_b(C_n^r) = \frac{3n-R}{5} + m = \frac{3(l+2(r+1))-(r+1+3l)}{5} + \lfloor \frac{r+1+l}{3} \rfloor = r + 1 + \lfloor \frac{r+1+l}{3} \rfloor = L$.

Case 2. $2l \leq r + 1 < 4l + 3$, i.e. $\frac{9r+6}{4} < n \leq \frac{5(r+1)}{2}$.

Then, $L = r + 1 + l$. Furthermore, it follows that $5l \leq r + 1 + 3l < 7l + 3 = 5l + (2l + 3)$ and therefore $Q = 1$ and $0 \leq R = r + 1 - 2l < 2l + 3$. This yields $m = 0$. Due to Theorem 2, we then obtain $\chi_b(C_n^r) = \frac{3n-R}{5} = \frac{3(l+2(r+1))-(r+1-2l)}{5} = r + 1 + l = L$.

Case 3. $r + 1 \geq 4l + 3 \geq 7$, i.e. $2r + 3 \leq n \leq \frac{9r+6}{4}$.

Then, $L = r + 1 + l$. Moreover, we deduce that $r + 1 + 3l \geq 7l + 3 = 5l + (2l + 3)$ which yields $Q \geq 1$. According to Theorem 2, $\chi_b(C_n^r) = \frac{3n-R}{5} + m = r + 1 + l + \frac{r+1-2l-R}{5} + m = L + \frac{r+1-2l-R}{5} + m$ (note that $\frac{r+1-2l-R}{5}$ is an integer). In order to prove $\chi_b(C_n^r) > L$, we have to show that $\frac{r+1-2l-R}{5} + m \geq 1$.

If $m = 0$, then $R \leq 2l + 2$ and therefore $\frac{r+1-2l-R}{5} = \lceil \frac{r+1-2l-R}{5} \rceil \geq \lceil \frac{r+1-4l-2}{5} \rceil \geq \lceil \frac{1}{5} \rceil = 1$.

If $m > 0$, then $\frac{r+1-2l-R}{5} + m = \frac{r+1-2l-(r+1+3l-5lQ)}{5} + m = (Q - 1)l + m \geq 0 + m \geq 1$. \square

Acknowledgements

The author would like to thank the referees for helpful comments and suggestions.

References

- [1] M. Alkhateeb, A. Kohl, *Upper bounds on the b -chromatic number and results for restricted graph classes*, *Discussiones Mathematicae Graph Theory* 31 (4) (2011), 709-735.
- [2] M. Blidia, F. Maffray, Z. Zemir, *On b -colorings in regular graphs*, *Discrete Appl. Math.* 157 (2009), 1787-1793.
- [3] B. Effantin, H. Kheddouci, *The b -chromatic number of some power graphs*, *DMTCS* 6 (2003), 045-054.
- [4] C.T. Hoang, M. Kouider, *On the b -dominating coloring of graphs*, *Discrete Appl. Math.* 152 (2005), 176-186.
- [5] R.W. Irving, D.F. Manlove, *The b -chromatic number of a graph*, *Discrete Appl. Math.* 91 (1-3) (1999), 127-141.
- [6] M. Kouider, M. Mahéo, *Some bounds for the b -chromatic number of a graph*, *Disc. Math.* 256 (2002), 267-277.
- [7] J. Kratochvíl, Zs. Tuza, M. Voigt, *On the b -chromatic number of graphs*, *WG 2002, LNCS 2573* (2002), 310-320.