

The topology of astrocyte networks controls the propagation of intercellular calcium waves

Jules Lallouette, Maurizio De Pittà, Eshel Ben-Jacob, Hugues Berry

► **To cite this version:**

Jules Lallouette, Maurizio De Pittà, Eshel Ben-Jacob, Hugues Berry. The topology of astrocyte networks controls the propagation of intercellular calcium waves. *BMC Neuroscience*, BioMed Central, 2014, 15 (Suppl 1), P205 (2 p.). <hal-01026523>

HAL Id: hal-01026523

<https://hal.inria.fr/hal-01026523>

Submitted on 21 Jul 2014

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

POSTER PRESENTATION

Open Access


The topology of astrocyte networks controls the propagation of intercellular calcium waves

Jules Lallouette^{1,2*}, Maurizio De Pittà^{1,2,3}, Eshel Ben-Jacob^{3,4}, Hugues Berry^{1,2}

From The Twenty Third Annual Computational Neuroscience Meeting: CNS*2014
Québec City, Canada. 26-31 July 2014

In recent years, astrocytes, one of the main types of glial cells, have been suggested to be active players in neuronal communication. While previously considered to form syn-
cytia with little or no spatial organization, emerging experimental evidence suggests that astrocytes could actually organize into real networks coupled by gap junction

channels, with complex topologies that may depend on the brain region. Intercellular calcium waves (ICW) are considered the main pathway for cell-to-cell signaling in these networks. However, it is still not understood why the extent of these ICW depends on the brain region or the experimental protocol considered. To investigate the



* Correspondence: jules.lallouette@inria.fr

¹EPI Beagle, INRIA Rhône-Alpes, Villeurbanne, France

Full list of author information is available at the end of the article

hypothesis that this variability could actually be linked to the heterogeneous properties of astrocyte networks we studied ICW propagation in biophysically realistic models of three-dimensional astrocyte networks (Figure 1) keeping constant both biophysical properties and spatial distribution of the cells while varying network topology according to different topological schemes (Figure 1D). Numerical simulations revealed that mere changes in network topology could indeed control the extent of ICWs from regenerative ICWs that roughly span the whole network (Figure 1A), to very restricted ones which activate only few tens of astrocytes (Figure 1C). Remarkably, ICW propagation was favored by sparse connectivity (i.e low mean degree) and restriction of cell connections to short distances (i.e large mean-shortest path). Networks with fewer gap junction couplings and stronger distance restrictions on couplings (Figure 1E, *top left quadrant*) supported much larger ICWs than either strongly coupled networks or networks comprising long distance couplings (Figure 1E, *bottom right quadrant*). Our results provide experimentally testable hypotheses to explain several experimental observations and theoretical support to the hypothesis of a functional role for the gap junction couplings in astrocyte networks. In particular, dynamic control of the topology of gap-junction couplings by neuronal activity suggests a novel type of neuron-glia communication.

Authors' details

¹EPI Beagle, INRIA Rhône-Alpes, Villeurbanne, France. ²LIRIS, Université de Lyon, UMR5205 CNRS-INSA, Villeurbanne, France. ³School of Physics and Astronomy, Tel Aviv University, Ramat Aviv, Israel. ⁴Center for Theoretical Biological Physics, Rice University, Houston, TX, USA.

Published: 21 July 2014

doi:10.1186/1471-2202-15-S1-P205

Cite this article as: Lallouette et al.: The topology of astrocyte networks controls the propagation of intercellular calcium waves. *BMC Neuroscience* 2014 15(Suppl 1):P205.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

