

Fourth order energy-preserving locally implicit discretization for linear wave equations

Juliette Chabassier, Sébastien Imperiale

► **To cite this version:**

Juliette Chabassier, Sébastien Imperiale. Fourth order energy-preserving locally implicit discretization for linear wave equations. SECOND FRENCH-RUSSIAN WORKSHOP ON COMPUTATIONAL GEOPHYSICS, Sep 2014, Novosibirsk, Russia. hal-01067115

HAL Id: hal-01067115

<https://hal.inria.fr/hal-01067115>

Submitted on 15 Oct 2014

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Fourth order energy-preserving locally implicit discretization for linear wave equations.

Juliette Chabassier and Sébastien Imperiale

A family of fourth order energy-preserving locally implicit schemes is introduced for linear wave equations. The domain of interest is decomposed into several regions where different fourth order time discretization are used, chosen among a family of implicit or explicit fourth order schemes derived in [1]. The coupling is based on a Lagrangian formulation on the boundaries between the several non conforming meshes of the regions. A global discrete energy is shown to be preserved and leads to global fourth order consistency in time. Numerical results illustrate the good behavior of the schemes and their potential for realistic highly heterogeneous cases or strongly refined geometries (as the Marmousi 2D test case of figure 1), for which using everywhere an explicit scheme can be extremely penalizing because the time step must respect the stability condition adapted to the smallest element or the highest velocities. Accuracy up to fourth order reduces the numerical dispersion inherent to implicit methods used with a large time step, and makes this family of schemes attractive compared to [3] which is only second order accurate in time. The presented technique could be an alternative to local time stepping [2] provided that some limitations are overcome in the future : treatment of dissipative terms, non trivial boundary conditions, coupling with a PML region, fluid structure coupling ...




Figure 1: Marmousi 2D benchmark domain presenting high velocity contrasts and refined geometry.

References

- [1] J. Chabassier, S. Imperiale. Introduction and study of fourth order theta schemes for linear wave equations, *Journal of Computational and Applied Mathematics*, vol. 245, pp 194–212, 2012.
- [2] J. Diaz, M. Grote, Energy conserving explicit local time-stepping for second-order wave equations, *SIAM Journal on Scientific Computing*, vol. 31 (3), pp 1985–2014, 2009.
- [3] S. Descombes, S. Lanteri, L. Moya. Locally implicit time integration strategies in a discontinuous Galerkin method for Maxwell's equations *Inria Research Report*, RR-7983, pp 1–28, 2012.