

Percolation on a non-homogeneous Poisson blob process

Fabio P. Machado

► **To cite this version:**

Fabio P. Machado. Percolation on a non-homogeneous Poisson blob process. Cyril Banderier and Christian Krattenthaler. Discrete Random Walks, DRW'03, 2003, Paris, France. Discrete Mathematics and Theoretical Computer Science, DMTCS Proceedings vol. AC, Discrete Random Walks (DRW'03), pp.171-172, 2003, DMTCS Proceedings. <hal-01183931>

HAL Id: hal-01183931

<https://hal.inria.fr/hal-01183931>

Submitted on 12 Aug 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Percolation on a non-homogeneous Poisson blob process

Fabio P. Machado[†]

Instituto de Matemática e Estatística, Universidade de São Paulo, Rua do Matão 1010, CEP 05508–090, São Paulo SP, Brasil. fmachado@ime.usp.br

We present the main results of a study for the existence of vacant and occupied unbounded connected components in a non-homogeneous Poisson blob process. The method used in the proofs is a multi-scale percolation comparison.

Keywords: Poisson blob model, continuum percolation, phase transition, multi-scale percolation

1 Introduction

One of the most well known examples of phenomena that introduces and motivates the study of continuum percolation is the process of the ground getting wet during a period of rain. At each point hit by a raindrop, one sees a circular wet patch. Right after the rain begins to fall what one sees is a small wet region inside a large dry region. At some instant, so many raindrops have hit the ground that the situation changes from that to a small dry region inside a large wet region. Typically, the parameter in which there is a phase transition behaviour is the density of the raindrops.

Continuum percolation models in which each point of a two-dimensional homogeneous Poisson point process is the centre of a disk of given (or random) radius r , have been extensively studied. In this note we present phase transition results for a sequence of Poisson point process which defines Poisson Boolean models and whose rates depend on the past. In order to prove our results we rely on a multi-scale percolation structure. General reference for percolation and continuum percolation are the books of Grimmett [2] and Meester and Roy [3]. A nice example of the use of multi-scale percolation technique can be found in Fontes *et al* [1].

2 Model and phase transition results

Let $\beta > 0$ be fixed number. Define $A_0 = \emptyset$. Having defined the sets A_0, A_1, \dots, A_n , define the process X_{n+1} as the non-homogeneous Poisson point process with intensity function given by:

$$f_{n+1}(x) = \exp(-\beta|B(x, n+1) \setminus \cup_{k=0}^n A_k|) \quad (1)$$

where $B(a, r)$ is the square of length r having centre at a and $|C|$ is the area (Lebesgue measure) of the set C .

[†]The author is thankful to CNPq (300226/97–7) for financial support.

Let $\{x_i^{(n+1)} : i \geq 1\}$ be the set of points from the process X_{n+1} . Define the set $A_{n+1} = \cup_{i=1}^{\infty} B(x_i^{(n+1)}, n+1)$ as the random set covered by the boxes from the process X_{n+1} . Define the total covered set $A_{\infty} = \cup_{n=0}^{\infty} A_n$.

The fundamental question in continuum percolation theory is about the existence of unbounded connected components. That is why we ask the following questions about the random set A_{∞} and its complement, the set A_{∞}^c . Let A be the component of A_{∞} which contains the origin. If the origin is not contained in A_{∞} , this is the empty set. Define

$$\theta(\beta) = \mathbb{P}_{\beta}(A \text{ is unbounded}). \quad (2)$$

It is clear that $\theta(\beta)$ is a decreasing function of β . Hence define the critical parameter β_c as follows:

$$\beta_c = \sup\{\beta > 0 : \theta(\beta) > 0\}. \quad (3)$$

The following result holds

Theorem 1.

$$0 < \beta_c < \infty.$$

Similar questions can also be asked the complement set of A_{∞} . Define C as the component of $(A_{\infty})^c$ which contain the origin. Define the vacant percolation probability as

$$\theta^*(\beta) = \mathbb{P}_{\beta}(C \text{ is unbounded}). \quad (4)$$

In this case, we have that $\theta^*(\beta)$ is an increasing function of β . Hence define the critical parameter β_c^* as follows:

$$\beta_c^* = \inf\{\beta > 0 : \theta^*(\beta) > 0\}. \quad (5)$$

We also prove the following theorem

Theorem 2.

$$0 < \beta_c^* < \infty.$$

It is clear that X_1 is actually an homogeneous Poisson process with intensity $\exp(-\beta|B(0, 1)|)$. Thus, A_1 will contain the covered set of a Poisson Boolean model with radius random variable being degenerate at $1/2$ and intensity $\exp(-\beta|B(0, 1)|)$. Thus, if $\exp(-\beta|B(0, 1)|) > \lambda_c$, the probability that the origin is contained in an unbounded component of A_1 is positive, where λ_c is the critical intensity of the Poisson Boolean model with radius being degenerate at $1/2$. Therefore, we have $\theta(\beta) > 0$ for this β . Hence, we have that $\beta_c > 0$. A similar argument also holds for β_c^* and we can easily show that, $\beta_c^* > 0$.

This is an announcement of results from a joint work with P. Ferrari, L. Fontes, S. Popov and A. Sarkar. The proofs rely on a multi-scale comparison argument to prove that the probability of certain events related to the existence of an unbounded connected component is exponentially close to 1 for large values of β .

References

- [1] FONTES, L. R.; SCHONMANN, R. H. AND SIDORAVICIUS, V. (2002) Stretched exponential fixation in stochastic Ising models at zero temperature. *Comm. Math. Phys.* 228, no. 3, 495–518.
- [2] GRIMMETT, G. (1999). *Percolation*, Second Edition. Springer, New York.
- [3] MEESTER, R. AND ROY, R. (1996). *Continuum Percolation*. Cambridge University Press, Cambridge.