

Branching rules in the ring of superclass functions of unipotent upper-triangular matrices

Nathaniel Thiem

► **To cite this version:**

Nathaniel Thiem. Branching rules in the ring of superclass functions of unipotent upper-triangular matrices. Krattenthaler, Christian and Strehl, Volker and Kauers, Manuel. 21st International Conference on Formal Power Series and Algebraic Combinatorics (FPSAC 2009), 2009, Hagenberg, Austria. Discrete Mathematics and Theoretical Computer Science, DMTCS Proceedings vol. AK, 21st International Conference on Formal Power Series and Algebraic Combinatorics (FPSAC 2009), pp.859-870, 2009, DMTCS Proceedings. <hal-01185390>

HAL Id: hal-01185390

<https://hal.inria.fr/hal-01185390>

Submitted on 20 Aug 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Branching rules in the ring of superclass functions of unipotent upper-triangular matrices

Nathaniel Thiem

University of Colorado at Boulder

It is becoming increasingly clear that the supercharacter theory of the finite group of unipotent upper-triangular matrices has a rich combinatorial structure built on set-partitions that is analogous to the partition combinatorics of the classical representation theory of the symmetric group. This paper begins by exploring a connection to the ring of symmetric functions in non-commuting variables that mirrors the symmetric group's relationship with the ring of symmetric functions. It then also investigates some of the representation theoretic structure constants arising from the restriction, tensor products and superinduction of supercharacters.

Keywords: set partitions, supercharacters, branching rules, symmetric functions

1 Introduction

The representation theory of the symmetric group S_n —with its connections to partition and tableaux combinatorics—has become a fundamental model in combinatorial representation theory. It has become clear in recent years that the representation theory finite group of unipotent upper-triangular matrix groups $U_n(q)$ can lead to a similarly rich combinatorial theory. While understanding the usual representation theory of $U_n(q)$ is a wild problem, André [1, 2, 3, 4] and Yan [23, 24] constructed a natural approximation to the representation theory that leads to a more computable theory. This approximation (known as a super-representation theory) now relies on set-partition combinatorics in the same way that the representation theory of the symmetric group relies on partition combinatorics.

A fundamental tool in symmetric group combinatorics is the ring of symmetric functions, which encodes the character theory of all symmetric groups simultaneously in a way that polynomial multiplication in the ring of symmetric functions becomes symmetric group induction from Young subgroups. This kind of a relationship has been extended to wreath products and type A finite groups of Lie type (for descriptions see for example [17, 21]). One of the purposes of this paper is to suggest an analogous relationship between the supercharacter theory of $U_n(q)$ and the ring on symmetric functions in non-commuting variables NCSym . In particular, Corollary 3.2 shows that there are a family of algebra isomorphisms from the ring of supercharacters to NCSym , where we replace induction from subgroups with its natural analogue superinduction from subgroups. Unfortunately, there does not yet seem to be a canonical choice (ideally, such a choice would take the Hopf structure of NCSym into account).

The other purpose of this paper is to use the combinatorics of set partitions to supply recursive algorithms for computing restrictions to a family of subgroups called parabolic subgroups. It turns out that if $k \leq n$, then there are many ways in which $U_k(q)$ sits inside $U_n(q)$ as a subgroup. In fact, for every subset $S \subseteq \{1, 2, \dots, n\}$ with k elements, there is a distinct subgroup U_S of $U_n(q)$ isomorphic to $U_k(q)$. The restriction from $U_n(q)$ to U_S depends on S , and Theorem 4.4 sorts out the combinatorial differences for all possible subsets S . This result can then be easily extended to give restriction rules for all parabolic subgroups. These computations require knowledge of tensor product results that were previously done by André [1] for large prime and by Yan [23] for arbitrary primes. For completeness, this paper supplies an alternate proof that relates tensor products to restriction and a generalization of the inflation functor (see Lemma 4.5).

By Frobenius reciprocity we then also obtain the coefficients of superinduction from these subgroups. Corollary 4.10 concludes by stating that superinduced supercharacters from parabolic subgroups are essentially twisted super-permutation characters (again using the generalization of the inflation map). These results give the structure constants for the ring of superclass functions of the finite unipotent upper-triangular groups. However, the underlying coefficient ring is $\mathbb{Z}[q^{-1}]$, unlike in the case of the symmetric group where the ring is \mathbb{Z} .

and

$$S(q) = \bigcup_{n \geq 0} S_n(q), \quad \text{where } S_n(q) = S_{\{1,2,\dots,n\}}(q).$$

Note that if $s_n(q) = |S_n(q)|$, then the generating function

$$\sum_{n \geq 0} s_n(q) \frac{x^n}{n!} = e^{\frac{e^{(q-1)x} - 1}{q-1}}$$

is a q -analogue of the usual exponential generating function of the Bell numbers (where $q = 2$ gives the usual generating function).

Suppose $S \subseteq T \subseteq \{1, 2, \dots, n\}$. Then there is a function

$$\langle \cdot \rangle_T : \left\{ \begin{array}{c} \mathbb{F}_q\text{-labeled} \\ \text{set-partitions of } S \end{array} \right\} \longrightarrow \left\{ \begin{array}{c} \mathbb{F}_q\text{-labeled} \\ \text{set-partitions of } T \end{array} \right\}$$

$$\lambda \qquad \qquad \qquad \mapsto \qquad \qquad \langle \lambda \rangle_T$$

where $\langle \lambda \rangle_T$ is the unique \mathbb{F}_q -labeled set-partition of T with arc set $A(\lambda)$ and labeling function τ_λ . We will use the convention that $\langle \lambda \rangle_n = \langle \lambda \rangle_{\{1,2,\dots,n\}}$.

2.2 Pattern groups

For $n \in \mathbb{Z}_{\geq 1}$, let $U_n(q)$ be the group of $n \times n$ unipotent upper-triangular matrices with entries in \mathbb{F}_q . Given a poset \mathcal{P} of $\{1, 2, \dots, n\}$, the *pattern group* $U_{\mathcal{P}}(q)$ is

$$U_{\mathcal{P}}(q) = \{u \in U_n(q) \mid u_{ij} \neq 0 \text{ implies } i \preceq j \text{ in } \mathcal{P}\}.$$

Remark. If $T_n(q)$ is the group of $n \times n$ diagonal matrices with entries in \mathbb{F}_q^\times , then the set of pattern subgroups of $U_n(q)$ can be characterized as the set of subgroups fixed by the conjugation action of $T_n(q)$ on $U_n(q)$.

Consider the injective map

$$\begin{array}{ccc} S_n & \longrightarrow & \left\{ \begin{array}{c} \text{Posets of} \\ \{1, 2, \dots, n\} \end{array} \right\} \\ K & \mapsto & \mathcal{P}_K \end{array}$$

where $i \prec j$ in \mathcal{P}_K if and only if $i < j$ and both i and j are in the same part of K .

A pattern subgroup $U_{\mathcal{P}}(q)$ is a *parabolic subgroup* of $U_n(q)$ if there exists $K \in S_n$ such that $\mathcal{P} = \mathcal{P}_K$. Note that if $K = K_1 \cup K_2 \cup \dots \cup K_\ell$ is the decomposition of K into parts, then

$$U_{\mathcal{P}_K}(q) \cong U_{|K_1|}(q) \times U_{|K_2|}(q) \times \dots \times U_{|K_\ell|}(q).$$

Thus, the parabolic subgroups of $U_{\mathcal{P}}(q)$ are reminiscent of the Young subgroups of the symmetric groups S_n or parabolic subgroups of a reductive groups of Lie type (such as the general linear group $GL_n(q)$). In fact, we will follow this analogy into the supercharacter theory of $U_n(q)$. To simplify notation, we will typically write

$$U_K(q) = U_{\mathcal{P}_K}(q), \quad \text{for } K \in S_n.$$

Remark. These subgroups are not generally block diagonal. For example,

$$U_{\mathcal{P}_{\{1,3,5\} \cup \{2,4\}}} = \left\{ \left(\begin{array}{ccccc} 1 & 0 & * & 0 & * \\ 0 & 1 & 0 & * & 0 \\ 0 & 0 & 1 & 0 & * \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{array} \right) \mid * \in \mathbb{F}_q \right\} \cong U_3(q) \times U_2(q).$$

However, parabolic subgroups do not include all possible copies of pattern subgroups isomorphic to a direct product of $U_k(q)$'s. For example,

$$U_{\begin{array}{c} 4 \\ 3 \quad 2 \\ 1 \end{array}} = \left\{ \left(\begin{array}{cccc} 1 & 0 & * & * \\ 0 & 1 & 0 & * \\ 0 & 0 & 1 & * \\ 0 & 0 & 0 & 1 \end{array} \right) \mid * \in \mathbb{F}_q \right\} \cong U_3(q) \times U_2(q)$$

is not a parabolic subgroup of $U_4(q)$.

2.3 A supercharacter theory for pattern groups

Given a group G , a supercharacter theory is an approximation to the usual character theory. To be more precise, a supercharacter theory consists of a set of superclasses \mathcal{K} and a set of supercharacters \mathcal{X} , such that

- (a) the set \mathcal{K} is a partition of G such that each part is a union of conjugacy classes,
- (b) the set \mathcal{X} is a set of characters such that each irreducible character appears as the constituent of exactly one supercharacter,
- (c) the supercharacters are constant on superclasses,
- (d) $|\mathcal{K}| = |\mathcal{X}|$,
- (e) the identity element 1 of G is in its own superclass, and the trivial character $\mathbf{1}$ of G is a supercharacter.

This general notion of a supercharacter theory was introduced by Diaconis and Isaacs [12] to generalize work of André and Yan on the character theory of $U_n(q)$.

Remark. The definition includes a reasonable amount of redundancy, as explored in [12, 16].

Diaconis and Isaacs extended the construction of André of a supercharacter theory for $U_n(q)$ to a larger family of groups called algebra groups. We will review the construction for pattern groups (a subset of the set of algebra groups). Let \mathcal{P} be a poset of $\{1, 2, \dots, n\}$ and let

$$\mathfrak{n}_{\mathcal{P}}(q) = U_{\mathcal{P}}(q) - 1,$$

which is an \mathbb{F}_q -algebra.

Fix a nontrivial homomorphism $\vartheta : \mathbb{F}_q^+ \rightarrow \mathbb{C}^\times$. The pattern group $U_{\mathcal{P}}(q)$ acts on the left and right on both $\mathfrak{n}_{\mathcal{P}}(q)$ and the dual space $\mathfrak{n}_{\mathcal{P}}(q)^*$, and the two-sided orbits lead to the sets \mathcal{K} and \mathcal{X} by the following rules. The superclasses are given

$$\begin{array}{ccc} U_{\mathcal{P}}(q) \backslash \mathfrak{n}_{\mathcal{P}}(q) / U_{\mathcal{P}}(q) & \longleftrightarrow & \mathcal{K} \\ U_{\mathcal{P}}(q) X U_{\mathcal{P}}(q) & \mapsto & 1 + U_{\mathcal{P}}(q) X U_{\mathcal{P}}(q), \end{array}$$

and the supercharacters are given by

$$\begin{array}{ccc} U_{\mathcal{P}}(q) \backslash \mathfrak{n}_{\mathcal{P}}(q)^* / U_{\mathcal{P}}(q) & \longleftrightarrow & \mathcal{X} \\ U_{\mathcal{P}}(q) \lambda U_{\mathcal{P}}(q) & \mapsto & \chi^\lambda = \frac{|\lambda U_{\mathcal{P}}(q)|}{|U_{\mathcal{P}}(q) \lambda U_{\mathcal{P}}(q)|} \sum_{\mu \in U_{\mathcal{P}}(q) \lambda U_{\mathcal{P}}(q)} \vartheta \circ \mu. \end{array}$$

The corresponding $U_{\mathcal{P}}$ -modules are given by

$$V^\lambda = \mathbb{C}\text{-span}\{v_\mu \mid \mu \in U_{\mathcal{P}}\lambda\}, \tag{2.1}$$

with action

$$g v_\mu = \vartheta((g\mu)(1-g)) v_{g\mu}, \quad \text{for } g \in U_{\mathcal{P}} \text{ and } \mu \in U_{\mathcal{P}}\lambda.$$

Example. The group $U_n(q)$ was the original motivation for studying supercharacter theories. The following results are due to André, Yan, and Arias-Castro–Diaconis–Stanley. The number of superclasses is

$$|\mathcal{K}| = |\mathcal{X}| = |\mathcal{S}_n(q)|,$$

where, for example,

$$\begin{array}{ccc} \mathcal{S}_n(q) & \longrightarrow & \mathcal{K} \\ \mu & \mapsto & u_\mu, \end{array} \quad \text{and} \quad (u_\mu)_{ij} = \begin{cases} 1, & \text{if } i = j, \\ \tau_\mu(i \frown j), & \text{if } i \frown j \in A(\mu), \\ 0, & \text{otherwise.} \end{cases}$$

The corresponding supercharacter formula for $\lambda, \mu \in \mathcal{S}_n(q)$ is

$$\chi^\lambda(u_\mu) = \begin{cases} \prod_{i \frown l \in A(\lambda)} \frac{q^{l-i-1} \vartheta(\tau_\lambda(i \frown l) \tau_\mu(i \frown l))}{q^{|\{j \frown k \in A(\mu) \mid i < j < k < l\}|}}, & \text{if } i < j < k, i \frown k \in A(\lambda) \\ & \text{implies } i \frown j, j \frown k \notin A(\mu), \\ 0, & \text{otherwise,} \end{cases} \tag{2.2}$$

where $\tau_\mu(i \frown j) = 0$ if $i \frown l \notin A(\mu)$ (see [14] for the corresponding formula for arbitrary pattern groups). Note that the degree of each character is

$$\chi^\lambda(1) = \prod_{i \frown l \in A(\lambda)} q^{l-i-1}. \tag{2.3}$$

It follows directly from the formula that the supercharacters factor nicely

$$\chi^\lambda = \prod_{i \frown l \in A(\lambda)} \chi^{(i \frown l)_n}.$$

It also follows from (2.2) and (2.3) that χ^λ is linear if and only if

$$i \frown k \in A(\lambda) \quad \text{implies} \quad k = i + 1.$$

The set $C(\lambda)$ measures how close the supercharacter χ^λ is to being irreducible. In fact,

$$\langle \chi^\lambda, \chi^\mu \rangle = q^{|C(\lambda)|} \delta_{\lambda\mu}, \tag{2.4}$$

where $\langle \cdot, \cdot \rangle$ is the usual inner product on characters.

Remark. If instead of considering $U_n(q)$ -orbits on $\mathfrak{n}_n(q)$ and $\mathfrak{n}_n(q)^*$, we consider orbits of the full Borel subgroup $B_n(q) = T_n(q)U_n(q)$ on these spaces, then the corresponding supercharacter theory no longer depends on the finite field q . In this case, the combinatorics reduces to considering set-partitions rather than \mathbb{F}_q -labeled set-partitions.

Supercharacters satisfy a variety of nice properties, as described in [12]. The above construction satisfies

- (a) The product of two supercharacters is a $\mathbb{Z}_{\geq 0}$ -linear combination of supercharacters.
- (b) The restriction of a supercharacter from one pattern group to a pattern subgroup is a $\mathbb{Z}_{\geq 0}$ -linear combination of supercharacters.

However, it is not true that the induction functor sends a supercharacter to a $\mathbb{Z}_{\geq 0}$ -linear combination of supercharacters. In fact, an induced supercharacter is generally no longer even a superclass function.

Diaconis and Isaacs therefore define a map superinduction on supercharacters that is adjoint to restriction with respect to the usual inner product on class functions; it turns out that this function averages over superclasses in the same way induction averages over conjugacy classes. In particular, if $H \subseteq G$ are pattern groups (or more generally algebra groups), then *superinduction* is the function

$$\text{SInd} : \left\{ \begin{array}{c} \text{Superclass functions} \\ \text{of } H \\ \chi \end{array} \right\} \longrightarrow \left\{ \begin{array}{c} \text{Superclass functions} \\ \text{of } G \\ \text{SInd}_H^G(\chi), \end{array} \right\}$$

where

$$\text{SInd}_H^G(\chi)(g) = \frac{1}{|G||H|} \sum_{\substack{x, y \in G \\ x(g-1)y+1 \in H}} \chi(x(g-1)y+1), \quad \text{for } g \in G.$$

Unfortunately, while SInd sends superclass functions to superclass functions, it sends supercharacters to $\mathbb{Z}_{\geq 0}[1/q]$ -linear combinations of supercharacters (where q comes from the underlying finite field). In fact, the image is not even generally a character. See also [18] for a further exploration of the relationship between superinduction and induction.

2.4 The ring of symmetric functions in non-commutative variables

Fix a set $X = \{X_1, X_2, \dots\}$ of countably many non-commuting variables. For $K = K_1 \cup K_2 \cup \dots \cup K_\ell \in \mathcal{S}_n$, define the *monomial symmetric function*

$$m_K(X) = \sum_{\substack{k=(k_1, k_2, \dots, k_\ell) \in \mathbb{Z}_{\geq 1}^\ell \\ k_i \neq k_j, 1 \leq i < j \leq \ell}} X_{\pi_1(k)} X_{\pi_2(k)} \cdots X_{\pi_\ell(k)}, \quad \text{where } \pi_j(k) = k_i \text{ if } j \in K_i.$$

The space of symmetric functions in non-commuting variables of homogeneous degree n is

$$\text{NCSym}_n(X) = \mathbb{C}\text{-span}\{m_K(X) \mid K \in \mathcal{S}_n\},$$

and the ring of symmetric functions in non-commuting variables is

$$\text{NCSym} = \bigoplus_{n \geq 0} \text{NCSym}_n(X),$$

where a possible multiplication is given by usual polynomial products. However, note that if $K = \{a_1 < a_2 < \dots < a_m\} \cup \{b_1 < b_2 < \dots < b_n\} \in \mathcal{S}_{m+n}$ with $\sigma = (a_1, a_2, \dots, a_m, b_1, b_2, \dots, b_n)$ the corresponding permutation of $m + 1$ elements, then we could “shuffle” two words according to K ,

$$(X_{i_1} X_{i_2} \dots X_{i_m}) *_K (X_{i_{m+1}} \dots X_{i_{m+n}}) = X_{i_{\sigma^{-1}(1)}} X_{i_{\sigma^{-1}(2)}} \dots X_{i_{\sigma^{-1}(m+n)}}.$$

These operations give a variety of alternate shuffle products for NCSym.

The ring NCSym naturally generalizes the usual ring of symmetric functions [17], but is different from other generalizations such as the ring of noncommutative symmetric functions studied in, for example, [15]. The ring NCSym was introduced by Wolf [22], and further explored by Rosas and Sagan [19]. There has been recent interest in the Hopf structure of NCSym and its Hopf dual – for example, [9, 10]. In particular, [9] show that it has a representation theoretic connection with partition lattice algebras. This paper suggests that the supercharacter theory of $U_n(q)$ also has a representation theoretic connection to NCSym in a way that is more analogous to how the ring of symmetric functions dictates the representation theory of S_n . However, the precise nature of this connection remains open. In particular, it is not clear whether the Hopf structure of NCSym translates naturally into a representation theoretic Hopf structure for the supercharacters of $U_n(q)$.

3 The ring of unipotent superclass functions

This section explores the relationship between NCSym and the space of supercharacters

$$\mathcal{C}(q) = \bigoplus_{n \geq 0} \mathcal{C}_n(q), \quad \text{where } \mathcal{C}_n(q) = \mathbb{C}\text{-span}\{\chi^\lambda \mid \lambda \in \mathcal{S}_n(q)\}.$$

3.1 Parabolic subgroups and set-partition combinatorics

There are different copies of $U_m(q) \times U_n(q)$ as subgroups of $U_{m+n}(q)$ which are not related via an inner automorphism of $U_{m+n}(q)$. In fact, for every $K = K_1 \cup K_2 \in \mathcal{S}_{m+n}$ with $|K_1| = m$ and $|K_2| = n$, $U_{m+n}(q)$ has a parabolic subgroup $U_m(q) \times_K U_n(q) = U_K(q) \cong U_m(q) \times U_n(q)$.


Thus, the space \mathcal{C} has a variety of different products. For $\lambda \in \mathcal{S}_m(q)$, $\mu \in \mathcal{S}_n(q)$, and $K = K_1 \cup K_2 \in \mathcal{S}_{m+n}$ with $|K_1| = m$ and $|K_2| = n$, define

$$\chi^\lambda *_K \chi^\mu = \text{SInd}_{U_m(q) \times_K U_n(q)}^{U_{m+n}(q)} (\chi^\lambda \times \chi^\mu).$$

There is a related map

$$\cup_K : \begin{matrix} \mathcal{S}_m(q) \times \mathcal{S}_n(q) & \longrightarrow & \mathcal{S}_{m+n}(q) \\ (\lambda, \mu) & \longmapsto & \lambda \cup_K \mu, \end{matrix}$$

where $\lambda \cup_K \mu = \lambda' \cup \mu'$ with $\lambda' \in \mathcal{S}_{K_1}(q)$ and $\mu' \in \mathcal{S}_{K_2}(q)$ the same \mathbb{F}_q -labeled set-partitions as λ and μ respectively, but with $\{1, 2, \dots, m\}$ relabeled as K_1 and $\{1, 2, \dots, n\}$ relabeled as K_2 . For example,



It will follow from Corollary 4.10 that $\chi^{\lambda \cup_K \mu}$ is always a nonzero constituent of $\chi^\lambda *_K \chi^\mu$.

3.2 A characteristic map for supercharacters

For $\mu \in \mathcal{S}_n(q)$, let $\kappa_\mu : U_n \rightarrow U_n$ be the superclass characteristic function given by

$$\kappa_\mu(u) = \begin{cases} 1, & \text{if } u \text{ is in the same superclass as } u_\mu, \\ 0, & \text{otherwise,} \end{cases}$$

and

$$z_\mu = \frac{|U_n|}{|U_n(u_\mu - 1)U_n|}$$

Proposition 3.1 For $\mu \in \mathcal{S}_m(q)$ and $\nu \in \mathcal{S}_n(q)$,

$$\text{SInd}_{U_m \times_K U_n}^{U_{m+n}}((z_\mu \kappa_\mu) \otimes (z_\nu \kappa_\nu)) = z_{\mu \cup_K \nu} \kappa_{\mu \cup_K \nu}.$$

Let NCSym be the ring of symmetric functions in non-commuting variables. Let

$$\{p_\lambda \mid \lambda \in \mathcal{S}\}$$

be any basis that satisfies

$$p_\lambda *_K p_\mu = p_{\lambda \cup_K \mu}$$

for all $K = K_1 \cup K_2 \in \mathcal{S}$ with $|K_1| = |\lambda|$ and $|K_2| = |\mu|$. Note that NCSym in fact has several of such bases, such as $\{p_\lambda\}$ in [19] and $\{x_\lambda\}$ in [9].

Corollary 3.2 The function

$$\begin{aligned} \text{ch} : \mathcal{C}(2) &\longrightarrow \text{NCSym} \\ \kappa_\mu &\longmapsto \frac{1}{z_\mu} p_\mu \end{aligned}$$

is an isometric algebra isomorphism.

Questions. This result immediately raises the following questions.

- (1) Does the Hopf algebra structure of NCSym transfer in a representation theoretic way to \mathcal{C} ?
- (2) What is the correct choice of basis p_μ ? In particular, the $\{p_\lambda\}$ of [19] do not seem to give a nice Hopf structure to \mathcal{C} .
- (3) Is there a corresponding NCSym -space for $q > 2$?

Questions (1) and (2) presumably need simultaneous answers, and question (3) suggests there might be an analogue of the ring symmetric functions corresponding to wreath products.

4 Representation theoretic structure constants

This section explores the computation of structure constants in \mathcal{C} . We begin with a family of natural embedding maps of $\mathcal{C}_m(q) \subseteq \mathcal{C}_n(q)$ for $m \leq n$ using a generalization of the inflation functor, and then give algorithms for computing restrictions from $\mathcal{C}_{m+n}(q)$ to $\mathcal{C}_m(q) \otimes \mathcal{C}_n(q)$. To finish the computations we require a method for decomposing tensor products $\mathcal{C}_n(q) \otimes \mathcal{C}_n(q) \rightarrow \mathcal{C}_n(q)$. We conclude with a discussion of the corresponding superinduction coefficients. In this section we will assume a fixed q , and suppress the q from the notation; that is $U_n = U_n(q)$, etc.

4.1 Superinflation of characters

Let $T \subseteq G$ be pattern groups with corresponding algebras \mathfrak{t} and \mathfrak{g} , respectively. There exists a surjective projection

$$\begin{aligned} \pi : \mathfrak{g} = \mathfrak{t} \oplus \mathfrak{t}^\perp &\longrightarrow \mathfrak{t} \\ X + Y &\longmapsto X, \end{aligned}$$

with a corresponding inflation map

$$\begin{aligned} \text{Inf}_\mathfrak{t}^\mathfrak{g} : \mathfrak{t}^* &\longrightarrow \mathfrak{g}^* \\ \mu &\longmapsto \mu \circ \pi. \end{aligned}$$

The *superinflation* map on supermodules is given by

$$\text{Sinf}_T^G : \left\{ \begin{array}{c} \text{Supermodules} \\ \text{of } T \\ V^\mu \end{array} \right\} \begin{array}{l} \longrightarrow \\ \mapsto \end{array} \left\{ \begin{array}{c} \text{Supermodules} \\ \text{of } G \\ V^{\text{Inf}_T^G(\mu)} \end{array} \right\}$$

where supermodules are as in (2.1).

Note that superinflation takes supermodules to supermodules, just as the usual inflation map on characters takes irreducible characters to irreducible characters. Recall, the usual inflation map is constructed from a surjection $\pi : G \rightarrow T$ is given by

$$\text{Inf}_T^G : \left\{ \begin{array}{c} T\text{-modules} \\ V \end{array} \right\} \begin{array}{l} \longrightarrow \\ \mapsto \end{array} \left\{ \begin{array}{c} G\text{-modules} \\ \text{Inf}_T^G(V) \end{array} \right\}$$

where $gv = \pi(g)v$ for $g \in G, v \in \text{Inf}_T^G(V)$. The following proposition says that superinflation is inflation whenever possible.

Proposition 4.1 *Suppose G is a pattern group with pattern subgroups T and H such that $G = T \times H$. Then for any supermodule V^λ of T ,*

$$\text{Sinf}_T^G(V^\lambda) \cong \text{Inf}_T^G(V^\lambda).$$

We will be primarily be interested in the superinflation function between parabolic subgroups of $U_n(q)$. In this case, if $U_K(q) \subseteq U_L(q)$, then

$$\text{Sinf}_{U_K(q)}^{U_L(q)}(\chi^\lambda) = \chi^{(\lambda)_L}.$$

For example,

$$\begin{array}{ccc} U_{\{2,3,5,7\}} & \xrightarrow{\text{Sinf}} & U_{\{1\} \cup \{2,3,5,7\}} & \xrightarrow{\text{Sinf}} & U_7 \\ \chi^\circ \begin{array}{c} \overset{a}{\curvearrowright} \quad \overset{b}{\curvearrowright} \\ \bullet \quad \circ \quad \bullet \quad \circ \end{array} & \mapsto & \chi \begin{array}{c} \overset{a}{\curvearrowright} \quad \overset{b}{\curvearrowright} \\ \bullet \quad \bullet \quad \bullet \quad \circ \end{array} & \mapsto & \chi \begin{array}{c} \overset{a}{\curvearrowright} \quad \overset{b}{\curvearrowright} \\ \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \end{array}$$

Thus, superinflation allows us to embed $C_m(q) \subseteq C_n(q)$ for all $m < n$, although this embedding still depends on the embedding of $U_m(q)$ inside $U_n(q)$.

Remark. While the superinflation function does match up with the usual inflation when possible, it does not generally behave as nicely as the usual inflation function. In particular, it is no longer generally true that $\text{Res}_T^G \circ \text{Sinf}_T^G(\chi) = \chi$ for χ a class function of T . For example,

$$\chi^\circ \begin{array}{c} \overset{a}{\curvearrowright} \quad \overset{b}{\curvearrowright} \\ \bullet \quad \circ \quad \bullet \quad \circ \end{array} (1) = q^1 \neq q^3 = \chi \begin{array}{c} \overset{a}{\curvearrowright} \quad \overset{b}{\curvearrowright} \\ \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} (1).$$

4.2 Restrictions

In this section we give algorithms for computing restrictions between parabolic subgroups of $U_n(q)$. Since supercharacters decompose into tensor products of arcs, for $\lambda \in \mathcal{S}_n(q)$,

$$\chi^\lambda = \prod_{i \stackrel{a}{\sim} l \in A(\lambda)} \chi^{(i \stackrel{a}{\sim} l)_n},$$

our strategy is to compute restrictions to for each $\chi^{(i \stackrel{a}{\sim} l)_n}$. We then use a tensor product result in Section 4.3 to glue back together the resulting restrictions.

We begin with two observations, and then Theorem 4.4 gives a general algorithm. Recall that for $K = K_1 \cup K_2 \cup \dots \cup K_\ell \in \mathcal{S}_n, U_K$ is a subgroup of $U_n(q)$ isomorphic to

$$U_{|K_1|} \times U_{|K_2|} \times \dots \times U_{|K_\ell|}.$$

Proposition 4.2 *Let $U_K \subseteq U_L$ be parabolic subgroups of U_n with $L = L_1 \cup L_2 \cup \dots \cup L_\ell \in \mathcal{S}_n$. Then*

$$\text{Res}_{U_K}^{U_L}(\chi^{\lambda_1} \times \dots \times \chi^{\lambda_\ell}) = \text{Res}_{U_{K_1}}^{U_{L_1}}(\chi^{\lambda_1}) \times \text{Res}_{U_{K_2}}^{U_{L_2}}(\chi^{\lambda_2}) \times \dots \times \text{Res}_{U_{K_\ell}}^{U_{L_\ell}}(\chi^{\lambda_\ell})$$

where U_{K_j} is the parabolic subgroup of U_{L_j} corresponding to the vertices L_j .

The next proposition gives information about each factor in Proposition 4.2.

Proposition 4.3 For $i < l$, $a \in \mathbb{F}_q^\times$ and $K = K_1 \cup K_2 \cup \dots \cup K_\ell \in \mathcal{S}_n$,

$$\text{Res}_{U_K}^{U_n}(\chi^{\langle i \overset{a}{\sim} l \rangle_n}) = \frac{\text{Res}_{U_{K_1}}^{U_n}(\chi^{\langle i \overset{a}{\sim} l \rangle_n})}{q^{|\{i < k < l | k \notin K_1\}|}} \times \frac{\text{Res}_{U_{K_2}}^{U_n}(\chi^{\langle i \overset{a}{\sim} l \rangle_n})}{q^{|\{i < k < l | k \notin K_2\}|}} \times \dots \times \frac{\text{Res}_{U_{K_\ell}}^{U_n}(\chi^{\langle i \overset{a}{\sim} l \rangle_n})}{q^{|\{i < k < l | k \notin K_\ell\}|}}$$

For $S \subseteq \{1, 2, \dots, n\}$, let

$$U_S = \{u \in U_n \mid u_{ij} \neq 0 \text{ implies } i, j \in S\}.$$

Note that while U_S is not itself a parabolic subgroup of U_n , it is isomorphic to the parabolic subgroup $U_{\langle S \rangle_n}$.

Theorem 4.4 Let $S \subseteq \{1, 2, \dots, n\}$. Then for $1 \leq i < l \leq n$ and $a \in \mathbb{F}_q^\times$,

$$\text{Res}_{U_S}^{U_n}(\chi^{\langle i \overset{a}{\sim} l \rangle_n}) = \begin{cases} q^{|\{i < k < l | k \notin S\}|} \chi^{\langle i \overset{a}{\sim} l \rangle_S}, & \text{if } i, l \in S, \\ q^{|\{i < k < l | k \notin S\}|} \left(\mathbf{1} + \sum_{\substack{i < j < l, j \in S \\ b \in \mathbb{F}_q^\times}} \chi^{\langle j \overset{b}{\sim} l \rangle_S} \right), & \text{if } i \notin S, l \in S, \\ q^{|\{i < k < l | k \notin S\}|} \left(\mathbf{1} + \sum_{\substack{i < k < l, k \in S \\ b \in \mathbb{F}_q^\times}} \chi^{\langle i \overset{b}{\sim} k \rangle_S} \right), & \text{if } i \in S, l \notin S, \\ q^{|\{i < k' < l | k' \notin S\}|} \left((|S \cap [i, l]|(q-1) + 1)\mathbf{1} + (q-1) \sum_{\substack{i < j' < k' < l \\ j', k' \in S, c \in \mathbb{F}_q^\times}} \chi^{\langle j' \overset{c}{\sim} k' \rangle_S} \right), & \text{if } i, l \notin S. \end{cases}$$

Example. Let $n = 7$, $j = 2$, $k = 5$, so that

$$U_{[2,5]} = U_{\{2,3,4,5\}} = \left\{ \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & * & * & 0 & 0 & 0 \\ 0 & 0 & 1 & * & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & * & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix} \right\} \subseteq \left\{ \begin{pmatrix} 1 & * & * & * & * & * & * \\ 0 & 1 & * & * & * & * & * \\ 0 & 0 & 1 & * & * & * & * \\ 0 & 0 & 0 & 1 & * & * & * \\ 0 & 0 & 0 & 0 & 1 & * & * \\ 0 & 0 & 0 & 0 & 0 & 1 & * \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix} \right\} = U_7$$

Then

$$\begin{aligned} \text{Res}_{U_{[2,5]}}^{U_7}(\chi^{\langle i \overset{a}{\sim} l \rangle_n}) &= \chi^{\circ \dots \circ \circ \circ \circ \circ \circ} + \sum_{b \in \mathbb{F}_q^\times} \chi^{\langle i \overset{b}{\sim} l \rangle_n} + \sum_{b \in \mathbb{F}_q^\times} \chi^{\langle i \overset{b}{\sim} l \rangle_n} + \sum_{b \in \mathbb{F}_q^\times} \chi^{\langle i \overset{b}{\sim} l \rangle_n} \\ \text{Res}_{U_{[2,5]}}^{U_7}(\chi^{\langle i \overset{a}{\sim} l \rangle_n}) &= q \left((4q-3)\chi^{\circ \dots \circ \circ \circ \circ \circ \circ} + (q-1) \sum_{b \in \mathbb{F}_q^\times} \chi^{\langle i \overset{b}{\sim} l \rangle_n} + (q-1) \sum_{b \in \mathbb{F}_q^\times} \chi^{\langle i \overset{b}{\sim} l \rangle_n} \right. \\ &\quad + (q-1) \sum_{b \in \mathbb{F}_q^\times} \chi^{\langle i \overset{b}{\sim} l \rangle_n} + (q-1) \sum_{b \in \mathbb{F}_q^\times} \chi^{\langle i \overset{b}{\sim} l \rangle_n} + (q-1) \sum_{b \in \mathbb{F}_q^\times} \chi^{\langle i \overset{b}{\sim} l \rangle_n} \\ &\quad \left. + (q-1) \sum_{b \in \mathbb{F}_q^\times} \chi^{\langle i \overset{b}{\sim} l \rangle_n} \right). \end{aligned}$$

4.3 Tensor products

We have seen in the previous section that when we decompose supercharacters into tensor products of irreducible characters, the restriction rules are manageable to compute. This section explains how to glue back together the resulting products of characters. The main result – Corollary 4.6 – has been computed by André for large primes in [1, Lemmas 6–8] and for arbitrary primes by Yan in [23, Propositions 7.2-7.5], but we reprove it here quickly using the machinery developed in this paper.

We begin with a lemma that further establishes the relationship between tensor products and restrictions. For $H \subseteq G$ and χ a superclass function of G , let

$$\text{SinfRes}_H^G(\chi) = \text{Sinf}_H^G \text{Res}_H^G(\chi).$$

Lemma 4.5 For $i < j < k < l$,

$$\begin{aligned} \chi^{\langle i \overset{a}{\sim} k \rangle_n} \otimes \chi^{\langle i \overset{b}{\sim} l \rangle_n} &= \text{SinfRes}_{U_{[i+1,l]}}^{U_n} (\chi^{\langle i \overset{a}{\sim} k \rangle_n}) \otimes \chi^{\langle i \overset{b}{\sim} l \rangle_n}, & a, b \in \mathbb{F}_q^\times, \\ \chi^{\langle i \overset{a}{\sim} l \rangle_n} \otimes \chi^{\langle j \overset{b}{\sim} l \rangle_n} &= \chi^{\langle i \overset{a}{\sim} l \rangle_n} \otimes \text{SinfRes}_{U_{[i,l-1]}}^{U_n} (\chi^{\langle j \overset{b}{\sim} l \rangle_n}), & a, b \in \mathbb{F}_q^\times, \\ \chi^{\langle i \overset{a}{\sim} l \rangle_n} \otimes \chi^{\langle i \overset{-a}{\sim} l \rangle_n} &= \text{SinfRes}_{U_{[i+1,l]}}^{U_n} (\chi^{\langle i \overset{a}{\sim} l \rangle_n}) \otimes \text{SinfRes}_{U_{[i,l-1]}}^{U_n} (\chi^{\langle i \overset{-a}{\sim} l \rangle_n}), & a \in \mathbb{F}_q^\times, \\ \chi^{\langle i \overset{a}{\sim} l \rangle_n} \otimes \chi^{\langle i \overset{b}{\sim} l \rangle_n} &= \chi^{\langle i \overset{a+b}{\sim} l \rangle_n} \otimes \text{SinfRes}_{U_{[i+1,l-1]}}^{U_n} (\chi^{\langle i \overset{a+b}{\sim} l \rangle_n}), & a, b \in \mathbb{F}_q^\times, b \neq -a. \end{aligned}$$

Combining Lemma 4.6 with Theorem 4.4 we obtain the following corollary.

Corollary 4.6 For $i < k, j < l, a, b \in \mathbb{F}_q^\times$, and $\{i, k\} \neq \{j, l\}$,

$$\chi^{\langle i \overset{a}{\sim} k \rangle_n} \otimes \chi^{\langle j \overset{b}{\sim} l \rangle_n} = \begin{cases} \chi^{\langle \{i \overset{a}{\sim} k\} \cup \{j \overset{b}{\sim} l\} \rangle_n}, & \text{if } \{i, k\} \cap \{j, l\} = \emptyset, \\ \chi^{\langle i \overset{a}{\sim} j \overset{b}{\sim} l \rangle_n}, & \text{if } i < j = k < l, \\ \chi^{\langle i \overset{b}{\sim} l \rangle_n} + \sum_{\substack{i < j' < k \\ c \in \mathbb{F}_q^\times}} \chi^{\langle \{j' \overset{c}{\sim} k\} \cup \{i \overset{b}{\sim} l\} \rangle_n}, & \text{if } i = j < k < l, \\ \chi^{\langle i \overset{a}{\sim} l \rangle_n} + \sum_{\substack{j < k' < l \\ c \in \mathbb{F}_q^\times}} \chi^{\langle \{i \overset{a}{\sim} l\} \cup \{j \overset{c}{\sim} k'\} \rangle_n}, & \text{if } i < j < k = l, \end{cases}$$

For $i < l, a, b \in \mathbb{F}_q^\times$,

$$\chi^{\langle i \overset{a}{\sim} l \rangle_n} \otimes \chi^{\langle i \overset{b}{\sim} l \rangle_n} = \begin{cases} \mathbf{1} + \sum_{\substack{i < j' < l \\ c \in \mathbb{F}_q^\times}} \chi^{\langle i \overset{c}{\sim} j' \rangle_n} + \sum_{\substack{i < k' < l \\ c \in \mathbb{F}_q^\times}} \chi^{\langle k' \overset{c}{\sim} l \rangle_n} + \sum_{\substack{i < j', k' < l \\ c, d \in \mathbb{F}_q^\times}} \chi^{\langle \{i \overset{c}{\sim} j'\} \cup \{k' \overset{d}{\sim} l\} \rangle_n}, & \text{if } b = -a, \\ ((l - i - 1)(q - 1) + 1) \chi^{\langle i \overset{a+b}{\sim} l \rangle_n} + (q - 1) \sum_{\substack{i < j' < k' < l \\ c \in \mathbb{F}_q^\times}} \chi^{\langle \{j' \overset{c}{\sim} k'\} \cup \{i \overset{a+b}{\sim} l\} \rangle_n}, & \text{otherwise.} \end{cases}$$

Remark. The coefficients of the tensor products are not understood in general, although it is clear from Corollary 4.6 that they are polynomial in q .

4.4 Superinduction

Let $S \subseteq \{1, 2, \dots, n\}$. If $\mu \in \mathcal{S}_S(q)$ and $\lambda \in \mathcal{S}_n(q)$, then by Frobenius reciprocity,

$$\langle \chi^\lambda, \text{SInd}_{U_S}^{U_n}(\chi^\mu) \rangle_{U_n} = \langle \text{Res}_{U_S}^{U_n}(\chi^\lambda), \chi^\mu \rangle_{U_S}.$$

Thus, if

$$\text{SInd}_{U_S}^{U_n}(\chi^\mu) = \sum_{\nu} a_{\mu}^{\nu} \chi^{\nu} \quad \text{and} \quad \text{Res}_{U_S}^{U_n}(\chi^\lambda) = \sum_{\gamma} b_{\gamma}^{\lambda} \chi^{\gamma},$$

then by (2.4)

$$q^{|C(\lambda)|} a_{\mu}^{\lambda} = q^{|C(\mu)|} b_{\mu}^{\lambda},$$

where $C(\nu)$ is the set of crossings of ν . Therefore,

$$\text{SInd}_{U_S}^{U_n}(\chi^\mu) = \sum_{\nu} a_{\mu}^{\nu} \chi^{\nu} = \sum_{\nu} q^{|C(\mu)| - |C(\nu)|} b_{\mu}^{\nu} \chi^{\nu}.$$

In general, if $U_K \subseteq U_n$ with $K \in \mathcal{S}_n$, then

$$\text{SInd}_{U_K}^{U_n}(\chi^\mu) = \sum_{\nu} q^{|C_K(\mu)| - |C(\nu)|} b_{\mu}^{\nu} \chi^{\nu}, \tag{4.1}$$

where $C_K(\nu)$ is the set of crossings that occur within the same parts of K .

With this discussion, we obtain the following corollary of Sections 4.2 and 4.3. When combined with Corollary 4.10, below, these results give a reasonably direct way to compute superinduction for some cases.

Corollary 4.7 Let $K = \{1, 2, \dots, k\} \cup \{k + 1, k + 2, \dots, n\} \in \mathcal{S}_n$ be a set-partition with two parts. Then

$$\text{SInd}_{U_K}^{U_n}(\mathbf{1}) = \sum_{\substack{\lambda \in \mathcal{S}_n(q) \\ \text{if } i \sim j \in \lambda, \\ \text{then } i \approx j \in K}} q^{-|C(\lambda)|} \chi^\lambda,$$

where $i \sim j$ if and only if i and j are in the same part in K .

Corollary 4.7 has some immediate combinatorial consequences. Let

$$SG_{n \times m} = \{a \in M_{n \times n}(\{0, 1\}) \mid a \text{ has at most one } 1 \text{ and every row and column}\}$$

be the set of $m \times n$ 0-1 matrices with at most one 1 in every row and column. Define statistics for $w \in SG_{m \times n}$

$$\text{ones}(w) = |\{(i, j) \in [1, n] \times [1, m] \mid w_{ij} = 1\}|$$

$$\text{sow}(w) = |\{(j, k) \in [1, n] \times [1, m] \mid w_{jk} = 0, w_{ik} = 1 \text{ for some } i < j \text{ or } w_{jl} = 1 \text{ for some } k < l\}|.$$

For example, if

$$w = \begin{pmatrix} \underline{0} & \underline{1} & \underline{0} & \underline{0} \\ \underline{0} & \underline{0} & \underline{0} & \underline{1} \\ \underline{0} & \underline{0} & \underline{0} & \underline{0} \end{pmatrix}, \quad \text{then} \quad \begin{aligned} \text{ones}(w) &= 2 \\ \text{sow}(w) &= 6. \end{aligned}$$

Corollary 4.8 Let m and n be positive integers. Then

$$(a) \quad q^{mn} = \sum_{w \in SG_{m \times n}} (q - 1)^{\text{ones}(w)} q^{\text{sow}(w)}$$

$$(b) \quad 0 = \sum_{w \in SG_{m \times n}} (-1)^{w_{1n}} (q - 1)^{\text{ones}(w)} q^{\text{sow}(w)}.$$

We conclude with some observations relating superinduction to these superpermutation ‘‘characters.’’ The first corollary follows from Frobenius reciprocity.

Corollary 4.9 Let $H \subseteq G$ be pattern groups, and let $\mu \in (H - 1)$. If $\chi^\mu(1) \text{Sinf}_H^G(\chi^\mu)(h) = \text{Sinf}_H^G(\chi^\mu)(1) \chi^\mu(h)$, for all $h \in H$, then

$$\text{SInd}_H^G(\chi^\mu) = \frac{\chi^\mu(1)}{\text{Sinf}_H^G(\chi^\mu)(1)} \text{Sinf}_H^G(\chi^\mu) \otimes \text{SInd}_H^G(\mathbf{1}).$$

The assumption in Corollary 4.9 is not so unusual. In fact,

Corollary 4.10 Let $U_K \subseteq U_L$ be parabolic subgroups of U_n , where $K = K_1 \cup K_2 \cup \dots \cup K_\ell, L \in \mathcal{S}_n$. Then for $\mu \in \mathcal{S}_{K_1}(q) \times \mathcal{S}_{K_2}(q) \times \dots \times \mathcal{S}_{K_\ell}(q)$,

$$\text{SInd}_{U_K}^{U_L}(\chi^\mu) = \frac{\chi^\mu(1)}{\text{Sinf}_{U_K}^{U_L}(\chi^\mu)(1)} \text{Sinf}_{U_K}^{U_L}(\chi^\mu) \otimes \text{SInd}_{U_K}^{U_L}(\mathbf{1}).$$

Remark. While the assumption in Corollary 4.9 is sufficient, it is not necessary. For example, if

$$H = \left\{ \begin{pmatrix} 1 & * & * \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \right\} \subseteq U_3 = \left\{ \begin{pmatrix} 1 & * & * \\ 0 & 1 & * \\ 0 & 0 & 1 \end{pmatrix} \right\}.$$

then for these groups,

$$\chi^{\begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}} \begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} = 1 \quad \text{and} \quad \text{Sinf}_H^{U_3} \left(\chi^{\begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}} \right) \begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} = 0.$$

However, it remains true that

$$\text{SInd}_H^G \left(\chi^{\begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}} \right) = q^{-1} \text{Sinf}_H^G \left(\chi^{\begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}} \right) \otimes \text{SInd}_H^G(\mathbf{1}).$$

In fact, the conclusion of Corollary 4.9 may be true for all pattern groups; I know of no counter-example.

References

- [1] André, C. “Basic characters of the unitriangular group,” *J. algebra* **175** (1995), 287–319.
- [2] André, C. “Irreducible characters of finite algebra groups,” *Matrices and group representations Coimbra, 1998* Textos Mat. Sér B **19** (1999), 65–80.
- [3] André, C. “The basic character table of the unitriangular group,” *J. algebra* **241** (2001), 437–471.
- [4] André, C. “Basic characters of the unitriangular group (for arbitrary primes),” *Proc. Amer. Math. Soc.* **130** (2002), 1934–1954.
- [5] André, C; Neto, A. “Super-characters of finite unipotent groups of types B_n , C_n and D_n ,” *J. Algebra* **305** (2006), 394–429.
- [6] André, C; Nicolás, A. “Supercharacters of the adjoint group of a finite radical ring,” August 2006 preprint.
- [7] Arregi, J; Vera-Lopez, A. “Computing in unitriangular matrices over finite fields.” *Linear algebra and Appl.* **387** (2004), 193–219.
- [8] Arias-Castro E; Diaconis, P; Stanley, R. “A super-class walk on upper-triangular matrices,” *J. algebra* **278** (2004), 739–765.
- [9] Bergeron, N; Hohlweg, C; Rosas, M; Zabrocki, M. “Grothendieck bialgebras, partition lattices, and symmetric functions in noncommutative variables,” *Electron. J. Combin.* **13** (2006).
- [10] Bergeron, N; Reutenauer, C; Rosas, M; Zabrocki, M. “Invariants and coinvariants of the symmetric groups in noncommuting variables,” *Canad. J. Math.* **60** (2008), 266–296.
- [11] Boyarchenko, M; Drinfeld, V. “A motivated introduction to character sheaves and the orbit method for unipotent groups in positive characteristic.” Preprint (2006), arXiv.0609769.
- [12] Diaconis, P; Isaacs, M. “Supercharacters and superclasses for algebra groups,” *Trans. Amer. Math. Soc.* **360** (2008), 2359–2392.
- [13] Diaconis, P; Saloff-Coste, L. “Comparison techniques for random walk on finite groups,” *Ann. Probab.* **21** (1993), 2131–2156.
- [14] Diaconis, P; Thiem, N. “Supercharacter formulas for pattern groups.” To appear in *Trans. Amer. Math. Soc.*
- [15] Gelfand, I.M; Krob, D; Lascoux, A; Leclerc, B; Retakh, V; Thibon, J-Y. “Noncommutative symmetric functions,” *Adv. Math.* **112** (1995), 218–348.
- [16] Hendrickson, A. *Supercharacter theories of finite cyclic groups*. Unpublished Ph.D. Thesis, Department of Mathematics, University of Wisconsin, 2008.
- [17] Macdonald, I.G. *Symmetric functions and Hall polynomials. Second edition. With Contributions by A. Zelevinsky*. Oxford Mathematical Monographs. Oxford Science Publications. The Clarendon Press, Oxford University Press, New York, 1995.
- [18] Marberg, E; Thiem, N. “Superinduction for pattern groups.” A 2007 preprint, , arXiv:0712.1228.
- [19] Rosas, M; Sagan, B. “Symmetric functions in noncommuting variables,” *Trans. Amer. Math. Soc.* **358** (2004), 215–232.
- [20] Thiem, N; Venkateswaran, V. “Restricting supercharacters of the finite group of unipotent uppertriangular matrices,” A 2007 preprint, arXiv:0712.1237.
- [21] Thiem, N; Vinroot, C.R. “On the characteristic map of finite unitary groups,” *Adv. Math.* **210** (2007), 707–732.
- [22] Wolf, M.C. “Symmetric functions of noncommuting elements,” *Duke Math. J.* **2** (1936), 626–637.
- [23] Yan, N. *Representation theory of the finite unipotent linear groups*, Unpublished Ph.D. Thesis, Department of Mathematics, University of Pennsylvania, 2001.
- [24] Yan, N. “Representations of finite unipotent linear groups by the method of clusters,” 2006 preprint.