

Matroid Polytopes and Their Volumes

Federico Ardila, Carolina Benedetti, Jeffrey Doker

► **To cite this version:**

Federico Ardila, Carolina Benedetti, Jeffrey Doker. Matroid Polytopes and Their Volumes. Krattenthaler, Christian and Strehl, Volker and Kauers, Manuel. 21st International Conference on Formal Power Series and Algebraic Combinatorics (FPSAC 2009), 2009, Hagenberg, Austria. Discrete Mathematics and Theoretical Computer Science, DMTCS Proceedings vol. AK, 21st International Conference on Formal Power Series and Algebraic Combinatorics (FPSAC 2009), pp.77-88, 2009, DMTCS Proceedings. <hal-01185426>

HAL Id: hal-01185426

<https://hal.inria.fr/hal-01185426>

Submitted on 20 Aug 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Matroid Polytopes and Their Volumes

Federico Ardila^{1†}, Carolina Benedetti^{2‡} and Jeffrey Doker³

¹San Francisco, CA, USA, federico@math.sfsu.edu.

²Universidad de Los Andes, Bogotá, Colombia, c.benedetti92@uniandes.edu.co.

³University of California, Berkeley, Berkeley, CA, USA, doker@math.berkeley.edu.

Abstract. We express the matroid polytope P_M of a matroid M as a signed Minkowski sum of simplices, and obtain a formula for the volume of P_M . This gives a combinatorial expression for the degree of an arbitrary torus orbit closure in the Grassmannian $Gr_{k,n}$. We then derive analogous results for the independent set polytope and the associated flag matroid polytope of M . Our proofs are based on a natural extension of Postnikov's theory of generalized permutohedra.

Résumé. On exprime le polytope matroïde P_M d'un matroïde M comme somme signée de Minkowski de simplices, et on obtient une formule pour le volume de P_M . Ceci donne une expression combinatoire pour le degré d'une clôture d'orbite de tore dans la Grassmannienne $Gr_{k,n}$. Ensuite, on déduit des résultats analogues pour le polytope ensemble indépendant et pour le polytope matroïde drapeau associé à M . Nos preuves sont fondées sur une extension naturelle de la théorie de Postnikov de permutoèdres généralisés.

Keywords: Matroid; generalized permutohedron; matroid polytope; Minkowski sum; mixed volume; flag matroid.

[†]NSF grant DMS-0801075

[‡]Proyecto Semilla of the Universidad de Los Andes

1 Introduction

The theory of matroids can be approached from many different points of view; a matroid can be defined as a simplicial complex of independent sets, a lattice of flats, a closure relation, etc. A relatively new point of view is the study of matroid polytopes, which in some sense are the natural combinatorial incarnations of matroids in algebraic geometry and optimization. Our paper is a contribution in this direction.

We begin with the observation that matroid polytopes are members of the family of *generalized permutohedra* (14). With some modifications of Postnikov's beautiful theory, we express the matroid polytope P_M as a signed Minkowski sum of simplices, and use that to give a formula for its volume $\text{Vol}(P_M)$. This is done in Theorems 2.5 and 3.3. Our answers are expressed in terms of the beta invariants of the contractions of M .

Formulas for $\text{Vol}(P_M)$ were given in very special cases by Stanley (17) and Lam and Postnikov (11), and a polynomial time algorithm for finding $\text{Vol}(P_M)$ was constructed by de Loera et. al. (6). One motivation for this computation is the following. The closure of the torus orbit of a point p in the Grassmannian $Gr_{k,n}$ is a toric variety X_p , whose degree is the volume of the matroid polytope P_{M_p} associated to p . Our formula allows us to compute the degree of X_p combinatorially.

One can naturally associate two other polytopes to a matroid M : its independent set polytope and its associated flag matroid polytope. By a further extension of Postnikov's theory, we also write these polytopes as signed Minkowski sums of simplices and give formulas for their volumes. This is the content of Sections 4 and 5.

Throughout the paper we assume familiarity with the basic concepts of matroid theory; for further information we refer the reader to (13).

2 Matroid Polytopes are Generalized Permutohedra

A *generalized permutohedron* is a polytope whose inequality description is of the following form:

$$P_n(\{z_I\}) = \{(t_1, \dots, t_n) \in \mathbb{R}^n : \sum_{i=1}^n t_i = z_{[n]}, \sum_{i \in I} t_i \geq z_I \text{ for all } I \subseteq [n]\}$$

where z_I is a real number for each $I \subseteq [n] := \{1, \dots, n\}$, and $z_\emptyset = 0$. Different choices of z_I can give the same generalized permutohedron: if one of the inequalities does not define a face of $P_n(\{z_I\})$, then we can increase the value of the corresponding z_I without altering the polytope. When we write $P_n(\{z_I\})$, we will always assume that the z_I s are all chosen minimally; *i.e.*, that all the defining inequalities are tight.

The *Minkowski sum* of two polytopes P and Q in \mathbb{R}^n is defined to be $P + Q = \{p + q : p \in P, q \in Q\}$. We say that the *Minkowski difference* of P and Q is $P - Q = R$ if $P = Q + R$.⁽ⁱ⁾ The following lemma shows that generalized permutohedra behave nicely with respect to Minkowski sums.

Lemma 2.1 $P_n(\{z_I\}) + P_n(\{z'_I\}) = P_n(\{z_I + z'_I\})$.

⁽ⁱ⁾ We will only consider Minkowski differences $P - Q$ such that Q is a Minkowski summand of P . More generally, the Minkowski difference of two arbitrary polytopes P and Q in \mathbb{R}^n is defined to be $P - Q = \{r \in \mathbb{R}^n \mid r + Q \subseteq P\}$ (14). It is easy to check that $(Q + R) - Q = R$, so the two definitions agree in the cases that interest us. In this paper, a signed Minkowski sum equality such as $P - Q + R - S = T$ should be interpreted as $P + R = Q + S + T$.

Let Δ be the standard unit $(n - 1)$ -simplex

$$\begin{aligned}\Delta &= \{(t_1, \dots, t_n) \in \mathbb{R}^n : \sum_{i=1}^n t_i = 1, t_i \geq 0 \text{ for all } 1 \leq i \leq n\} \\ &= \text{conv}\{e_1, \dots, e_n\},\end{aligned}$$

where $e_i = (0, \dots, 0, 1, 0, \dots, 0)$ with a 1 in its i th coordinate. As J ranges over the subsets of $[n]$, let Δ_J be the face of the simplex Δ defined by

$$\Delta_J = \text{conv}\{e_i : i \in J\} = P_n(\{z(J)_I\})$$

where $z(J)_I = 1$ if $I \supseteq J$ and $z(J)_I = 0$ otherwise. Lemma 2.1 gives the following proposition.

The next two propositions are due to Postnikov in the case $y_I \geq 0$.

Proposition 2.2 (14, Proposition 6.3) *For any $y_I \geq 0$, the Minkowski sum $\sum y_I \Delta_I$ of dilations of faces of the standard $(n - 1)$ -simplex is a generalized permutohedron. We can write*

$$\sum_{A \subseteq E} y_I \Delta_I = P_n(\{z_I\}),$$

where $z_I = \sum_{J \subseteq I} y_J$ for each $I \subseteq [n]$.

Proposition 2.3 *Every generalized permutohedron $P_n(\{z_I\})$ can be written uniquely as a signed Minkowski sum of simplices, as*

$$P_n(\{z_I\}) = \sum_{I \subseteq [n]} y_I \Delta_I$$

where $y_I = \sum_{J \subseteq I} (-1)^{|I|-|J|} z_J$ for each $I \subseteq [n]$.

Proof: First we need to separate the right hand side into its positive and negative parts. By Proposition 2.2,

$$\sum_{I \subseteq [n]: y_I < 0} (-y_I) \Delta_I = P_n(\{z_I^-\}) \text{ and } \sum_{I \subseteq [n]: y_I \geq 0} y_I \Delta_I = P_n(\{z_I^+\})$$

where $z_I^- = \sum_{J \subseteq I: y_J < 0} (-y_J)$ and $z_I^+ = \sum_{J \subseteq I: y_J \geq 0} y_J$. Now $z_I + z_I^- = z_I^+$ gives

$$P_n(\{z_I\}) + \sum_{I \subseteq [n]: y_I < 0} (-y_I) \Delta_I = \sum_{I \subseteq [n]: y_I \geq 0} y_I \Delta_I,$$

as desired. Uniqueness is clear. \square

Let M be a matroid of rank r on the set E . The *matroid polytope* of M is the polytope P_M in \mathbb{R}^E whose vertices are the indicator vectors of the bases of M . The known description of the polytope P_M by inequalities makes it apparent that it is a generalized permutohedron:

Proposition 2.4 (19) *The matroid polytope of a matroid M on E with rank function r is $P_M = P_E(\{r - r(E - I)\}_{I \subseteq E})$.*

The *beta invariant* (5) of M is a non-negative integer given by

$$\beta(M) = (-1)^{r(M)} \sum_{X \subseteq E} (-1)^{|X|} r(X)$$

which stores significant information about M ; for example, $\beta(M) = 0$ if and only if M is disconnected and $\beta(M) = 1$ if and only if M is series-parallel. If

$$T_M(x, y) = \sum_{A \subseteq E} (x-1)^{r(E)-r(A)} (y-1)^{|A|-r(A)} = \sum_{i,j} b_{ij} x^i y^j$$

is the *Tutte polynomial* (20) of M , then $\beta(M) = b_{10} = b_{01}$ for $|E| \geq 2$.

Our next results are more elegantly stated in terms of the *signed beta invariant* of M , which we define to be

$$\tilde{\beta}(M) = (-1)^{r(M)+1} \beta(M).$$

Theorem 2.5 *Let M be a matroid of rank r on E and let P_M be its matroid polytope. Then*

$$P_M = \sum_{A \subseteq E} \tilde{\beta}(M/A) \Delta_{E-A}. \quad (1)$$

Proof: By Propositions 2.3 and 2.4, $P_M = \sum_{I \subseteq E} y_I \Delta_I$ where

$$\begin{aligned} y_I &= \sum_{J \subseteq I} (-1)^{|I|-|J|} (r - r(E-J)) = - \sum_{J \subseteq I} (-1)^{|I|-|J|} r(E-J) \\ &= - \sum_{E-J \supseteq E-I} (-1)^{|E-J|-|E-I|} (r(E-J) - r(E-I)) \\ &= - \sum_{X \subseteq I} (-1)^{|X|} (r(E-I \cup X) - r(E-I)) \\ &= - \sum_{X \subseteq I} (-1)^{|X|} r_{M/(E-I)}(X) = \tilde{\beta}(M/(E-I)) \end{aligned}$$

as desired. □

Example 2.6 *Let M be the matroid on $E = [4]$ with bases $\{12, 13, 14, 23, 24\}$; its matroid polytope is a square pyramid. Theorem 2.5 gives $P_M = \Delta_{234} + \Delta_{134} + \Delta_{12} - \Delta_{1234}$, as illustrated in Figure 1. The dotted lines in the polytope $\Delta_{234} + \Delta_{134} + \Delta_{12}$ are an aid to visualize the Minkowski difference.*

One way of visualizing the Minkowski sum of two polytopes P and Q is by grabbing a vertex v of Q and then using it to “slide” Q around in space, making sure that v never leaves P . The region that Q sweeps along the way is $P + Q$. Similarly, the Minkowski difference $P - R$ can be visualized by picking a vertex v of R and then “sliding” R around in space, this time making sure that no point in R ever leaves P . The region that v sweeps along the way is $P - R$. This may be helpful in understanding Figure 1.

Some remarks about Theorem 2.5 are in order.




Fig. 1: A matroid polytope as a signed Minkowski sum of simplices.

- Generally most terms in the sum of Theorem 2.5 are zero. The nonzero terms correspond to the *coconnected flats* A , which we define to be the sets A such that M/A is connected. These are indeed flats, since contracting by them must produce a loopless matroid.
- A matroid and its dual have congruent matroid polytopes, and Theorem 2.5 gives different formulas for them. For example $P_{U_{1,3}} = \Delta_{123}$ while $P_{U_{2,3}} = \Delta_{12} + \Delta_{23} + \Delta_{13} - \Delta_{123}$.
- The study of the subdivisions of a matroid polytope into smaller matroid polytopes, originally considered by Lafforgue (10), has recently received significant attention (1; 2; 7; 15). Speyer conjectured (15) that the subdivisions consisting of series-parallel matroids have the largest number of faces in each dimension and proved this (16) for a large and important family of subdivisions: those which arise from a tropical linear space. The important role played by series-parallel matroids is still somewhat mysterious. Theorem 2.5 characterizes series-parallel matroids as those whose matroid polytope has no repeated Minkowski summands. It would be interesting to connect this characterization to matroid subdivisions; this may require extending the theory of mixed subdivisions to signed Minkowski sums.
- Theorem 2.5 provides a geometric interpretation for the beta invariant of a matroid M in terms of the matroid polytope P_M . In Section 5 we see how to extend this to certain families of Coxeter matroids. This is a promising point of view towards the notable open problem (4, Problem 6.16.6) of defining useful enumerative invariants of a Coxeter matroid.

3 The Volume of a Matroid Polytope

Our next goal is to present an explicit combinatorial formula for the volume of an arbitrary matroid polytope. Formulas have been given for very special families of matroids by Stanley (17) and Lam and Postnikov (11). Additionally, a polynomial time algorithm for computing the volume of an arbitrary matroid polytope was recently discovered by de Loera et. al. (6). Let us say some words about the motivation for this question.

Consider the Grassmannian manifold $Gr_{k,n}$ of k -dimensional subspaces in \mathbb{C}^n . Such a subspace can be represented as the row space of a $k \times n$ matrix A of rank k , modulo the left action of GL_k which does not change the row space. The $\binom{n}{k}$ maximal minors of this matrix are the *Plücker coordinates* of the subspace, and they give an embedding of $Gr_{k,n}$ as a projective algebraic variety in $\mathbb{CP}^{\binom{n}{k}-1}$.

Each point p in $Gr_{k,n}$ gives rise to a matroid M_p whose bases are the k -subsets of n where the Plücker coordinate of p is not zero. Gelfand, Goresky, MacPherson, and Serganova (9) first considered the stratification of $Gr_{k,n}$ into *matroid strata*, which consist of the points corresponding to a fixed matroid.

The torus $\mathbb{T} = (\mathbb{C}^*)^n$ acts on \mathbb{C}^n by $(t_1, \dots, t_n) \cdot (x_1, \dots, x_n) = (t_1 x_1, \dots, t_n x_n)$ for $t_i \neq 0$; this action extends to an action of \mathbb{T} on $Gr_{k,n}$. For a point $p \in Gr_{k,n}$, the closure of the torus orbit $X_p = \overline{\mathbb{T} \cdot p}$ is a toric variety which only depends on the matroid M_p of p , and the polytope corresponding to X_p under the moment map is the matroid polytope of M_p (9). Under these circumstances it is known (8) that the volume of the matroid polytope M_p equals the degree of the toric variety X_p as a projective subvariety of $\mathbb{C}\mathbb{P}^{\binom{n}{k}-1}$:

$$\text{Vol } P_{M_p} = \deg X_p.$$

Therefore, by finding the volume of an arbitrary matroid polytope, one obtains a formula for the degree of the toric varieties arising from arbitrary torus orbits in the Grassmannian.

To prove our formula for the volume of a matroid polytope, we first recall the notion of the *mixed volume* $\text{Vol}(P_1, \dots, P_n)$ of n (possibly repeated) polytopes P_1, \dots, P_n in \mathbb{R}^n . All volumes in this section are normalized with respect to the lattice generated by $e_1 - e_2, \dots, e_{n-1} - e_n$ where our polytopes live; so the standard simplex Δ has volume $1/(n-1)!$.

Proposition 3.1 (12) *Let n be a fixed positive integer. There exists a unique function $\text{Vol}(P_1, \dots, P_n)$ defined on n -tuples of polytopes in \mathbb{R}^n , called the mixed volume of P_1, \dots, P_n , such that, for any collection of polytopes Q_1, \dots, Q_m in \mathbb{R}^n and any nonnegative real numbers y_1, \dots, y_m , the volume of the Minkowski sum $y_1 Q_1 + \dots + y_m Q_m$ is the polynomial in y_1, \dots, y_m given by*

$$\text{Vol}(y_1 Q_1 + \dots + y_m Q_m) = \sum_{i_1, \dots, i_n} \text{Vol}(Q_{i_1}, \dots, Q_{i_n}) y_{i_1} \cdots y_{i_n}$$

where the sum is over all ordered n -tuples (i_1, \dots, i_n) with $1 \leq i_r \leq m$.

Proposition 3.1 still holds if some of the y_i s are negative as long as the expression $y_1 Q_1 + \dots + y_m Q_m$ still makes sense, as stated in the following Proposition.

Proposition 3.2 *If $P = y_1 Q_1 + \dots + y_m Q_m$ is a signed Minkowski sum of polytopes in \mathbb{R}^n , then*

$$\text{Vol}(y_1 Q_1 + \dots + y_m Q_m) = \sum_{i_1, \dots, i_n} \text{Vol}(Q_{i_1}, \dots, Q_{i_n}) y_{i_1} \cdots y_{i_n}$$

where the sum is over all ordered n -tuples (i_1, \dots, i_n) with $1 \leq i_r \leq m$.

Proof: We first show that

$$\text{Vol}(A - B) = \sum_{k=0}^n (-1)^k \binom{n}{k} \text{Vol}(A, \dots, A, B, \dots, B) \quad (2)$$

when B is a Minkowski summand of A in \mathbb{R}^n . Let $A - B = C$. By Proposition 3.1, for $t \geq 0$ we have that

$$\text{Vol}(C + tB) = \sum_{k=0}^n \binom{n}{k} \text{Vol}(C, \dots, C, B, \dots, B) t^k =: f(t)$$

and we are interested in computing $\text{Vol}(C) = f(0)$. Invoking Proposition 3.1 again, for $t \geq 0$ we have that

$$\text{Vol}(A + tB) = \sum_{k=0}^n \binom{n}{k} \text{Vol}(A, \dots, A, B, \dots, B) t^k =: g(t). \quad (3)$$

But $A + tB = C + (t+1)B$ and therefore $g(t) = f(t+1)$ for all $t \geq 0$. Therefore $g(t) = f(t+1)$ as polynomials, and $\text{Vol} C = f(0) = g(-1)$. Plugging into (3) gives the desired result.

Having established (2), separate the given Minkowski sum for P into its positive and negative parts as $P = Q - R$, where $Q = x_1 Q_1 + \dots + x_r Q_r$ and $R = y_1 R_1 + \dots + y_s R_s$ with $x_i, y_i \geq 0$. For positive t we can write $Q + tR = \sum x_i Q_i + \sum t y_j R_j$, which gives two formulas for $\text{Vol}(Q + tR)$.

$$\begin{aligned} \text{Vol}(Q + tR) &= \sum_{k=0}^n \binom{n}{k} \text{Vol}(Q, \dots, Q, R, \dots, R) t^k \\ &= \sum_{\substack{1 \leq i_a \leq r \\ 1 \leq j_b \leq s}} \text{Vol}(Q_{i_1}, \dots, Q_{i_{n-k}}, R_{j_1}, \dots, R_{j_k}) x_{i_1} \cdots x_{i_{n-k}} y_{j_1} \cdots y_{j_k} t^k \end{aligned}$$

The last two expressions must be equal as polynomials. A priori, we cannot plug $t = -1$ into this equation; but instead, we can use the formula for $\text{Vol}(Q - R)$ from (2), and then plug in coefficient by coefficient. That gives the desired result. \square

Theorem 3.3 *If a connected matroid M has n elements, then the volume of the matroid polytope P_M is*

$$\text{Vol} P_M = \frac{1}{(n-1)!} \sum_{(J_1, \dots, J_{n-1})} \tilde{\beta}(M/J_1) \tilde{\beta}(M/J_2) \cdots \tilde{\beta}(M/J_{n-1}),$$

summing over the ordered collections of sets $J_1, \dots, J_{n-1} \subseteq [n]$ such that, for any distinct i_1, \dots, i_k , $|J_{i_1} \cap \dots \cap J_{i_k}| < n - k$.

Proof: Postnikov (14, Corollary 9.4) gave a formula for the volume of a (positive) Minkowski sum of simplices. We would like to apply his formula to the signed Minkowski sum in Theorem 2.5, and Proposition 3.2 makes this possible. \square

In Theorem 3.3, the hypothesis that M is connected is needed to guarantee that the matroid polytope P_M has dimension $n - 1$. More generally, if we have $M = M_1 \oplus \dots \oplus M_k$ then $P_M = P_{M_1} \times \dots \times P_{M_k}$ so the $((n - k)$ -dimensional) volume of P_M is $\text{Vol} P_M = \text{Vol} P_{M_1} \cdots \text{Vol} P_{M_k}$.

4 Independent Set Polytopes

In this section we show that our analysis of matroid polytopes can be carried out similarly for the *independent set polytope* I_M of a matroid M , which is the convex hull of the indicator vectors of the independent sets of M . The inequality description of I_M is known to be:

$$I_M = \{(x_1, \dots, x_n) \in \mathbb{R}^n : x_i \geq 0 \text{ for } i \in [n], \sum_{i \in A} x_i \leq r(A) \text{ for all } A \subseteq E\}. \quad (4)$$

The independent set polytope of a matroid is not a generalized permutahedron. Instead, it is a Q -polytope; i.e., a polytope of the form

$$Q_n(\{z_J\}) = \{(t_1, \dots, t_n) \in \mathbb{R}^n : t_i \geq 0 \text{ for all } i \in [n], \sum_{i \in J} t_i \leq z_J \text{ for all } J \subseteq [n]\}$$

where z_J is a non-negative real number for each $J \subseteq [n]$. We can also express these polytopes as signed Minkowski sums of simplices, though the simplices we use are not the Δ_J s, but those of the form

$$\begin{aligned} D_J &= \text{conv}\{0, e_i : i \in J\} \\ &= Q_n(\{d(J)_I\}) \end{aligned}$$

where $d(J)_I = 0$ if $I \cap J = \emptyset$ and $d(J)_I = 1$ otherwise.

The following lemmas on Q -polytopes are proved in a way analogous to the corresponding lemmas for generalized permutahedra as was done in Section 2, and we leave them to the reader.

Lemma 4.1 $Q_n(\{z_J\}) + Q_n(\{z'_J\}) = Q_n(\{z_J + z'_J\})$

Proposition 4.2 For any $y_I \geq 0$ we have

$$\sum_{I \subseteq [n]} y_I D_I = Q_n(\{z_J\})$$

where $z_J = \sum_{I: I \cap J \neq \emptyset} y_I$.

Proposition 4.3 Every Q -polytope $Q_n(\{z_J\})$ can be written uniquely⁽ⁱⁱ⁾ as a signed Minkowski sum of D_I s as

$$Q_n(\{z_J\}) = \sum_{I \subseteq [n]} y_I D_I,$$

where

$$y_J = - \sum_{I \subseteq J} (-1)^{|J|-|I|} z_{[n]-I}.$$

Proof: We need to invert the relation between the y_I s and the z_J s given by $z_J = \sum_{I: I \cap J \neq \emptyset} y_I$. We rewrite this relation as

$$z_{[n]} - z_J = \sum_{I \subseteq [n]-J} y_I$$

and apply inclusion-exclusion. As in Section 2, we first do this in the case $y_I \geq 0$ and then extend it to arbitrary Q -polytopes. \square

Theorem 4.4 Let M be a matroid of rank r on E and let I_M be its independent set polytope. Then

$$I_M = \sum_{A \subseteq E} \tilde{\beta}(M/A) D_{E-A}$$

where $\tilde{\beta}$ denotes the signed beta invariant.

⁽ⁱⁱ⁾ assuming $y_\emptyset = 0$

The great similarity between Theorems 2.5 and 4.4 is not surprising, since P_M is the facet of I_M which maximizes the linear function $\sum_{i \in E} x_i$, and Δ_I is the facet of D_I in that direction as well. In fact we could have first proved Theorem 4.4 and then obtained Theorem 2.5 as a corollary.

Theorem 4.5 *If a connected matroid M has n elements, then the volume of the independent set polytope I_M is*

$$\text{Vol } I_M = \frac{1}{n!} \sum_{(J_1, \dots, J_n)} \tilde{\beta}(M/J_1) \tilde{\beta}(M/J_2) \cdots \tilde{\beta}(M/J_n)$$

where the sum is over all n -tuples (J_1, \dots, J_n) of subsets of $[n]$ such that, for any distinct i_1, \dots, i_k , we have $|J_{i_1} \cap \cdots \cap J_{i_k}| \leq n - k$.

Notice that by Hall's marriage theorem, the condition on the J_i s is equivalent to requiring that $(E - J_1, \dots, E - J_n)$ has a system of distinct representatives (SDR); that is, there are $a_1 \in E - J_1, \dots, a_n \in E - J_n$ with $a_i \neq a_j$ for $i \neq j$.

Proof: By Theorem 4.4 and Proposition 3.1 it suffices to compute the mixed volume $\text{Vol}(D_{A_1}, \dots, D_{A_n})$ for each n -tuple (A_1, \dots, A_n) of subsets of $[n]$. Bernstein's theorem (18) tells us that $\text{Vol}(D_{A_1}, \dots, D_{A_n})$ is the number of isolated solutions in $(\mathbb{C} - \{0\})^n$ of the system of equations:

$$\begin{cases} \beta_{1,0} + \beta_{1,1}t_1 + \beta_{1,2}t_2 + \cdots + \beta_{1,n}t_n & = 0 \\ \beta_{2,0} + \beta_{2,1}t_1 + \beta_{2,2}t_2 + \cdots + \beta_{2,n}t_n & = 0 \\ & \vdots \\ \beta_{n,0} + \beta_{n,1}t_1 + \beta_{n,2}t_2 + \cdots + \beta_{n,n}t_n & = 0 \end{cases}$$

where $\beta_{i,0}$ and $\beta_{i,j}$ are generic complex numbers when $j \in A_i$, and $\beta_{i,j} = 0$ if $j \notin A_i$.

This system of linear equations will have one solution if it is non-singular and no solutions otherwise. Because the $\beta_{i,0}$ are generic, such a solution will be non-zero if it exists. The system is non-singular when the determinant is non-zero, and by genericity that happens when (A_1, \dots, A_n) has an SDR.

We conclude that $\text{Vol}(D_{E-J_1}, \dots, D_{E-J_n})$ is 1 if $(E - J_1, \dots, E - J_n)$ has an SDR and 0 otherwise, and the result follows. \square

Example 4.6 *Let I_M be the independent set polytope of the uniform matroid $U_{2,3}$. We have $I_M = D_{12} + D_{23} + D_{13} - D_{123}$. Theorem 4.5 should confirm that its volume is $\frac{5}{6}$; let us carry out that computation.*

The coconnected flats of M are $1, 2, 3$ and \emptyset and their complements are $\{23, 13, 12, 123\}$. We need to consider the triples of coconnected flats whose complements contain an SDR. Each one of the 24 triples of the form (a, b, c) , where $a, b, c \in [3]$ are not all equal, contributes a summand equal to 1. The 27 permutations of triples of the form (a, b, \emptyset) , contribute a -1 each. The 9 permutations of triples of the form $(a, \emptyset, \emptyset)$ contribute a 1 each. The triple $(\emptyset, \emptyset, \emptyset)$ contributes a -1 . The volume of I_M is then $\frac{1}{6}(24 - 27 + 9 - 1) = \frac{5}{6}$.

5 Truncation Flag Matroids

We will soon see that any flag matroid polytope can also be written as a signed Minkowski sum of simplices Δ_I . We now focus on the particularly nice family of *truncation flag matroids*, introduced by Borovik, Gelfand, Vince, and White, where we obtain an explicit formula for this sum.

The *strong order* on matroids is defined by saying that two matroids M and N on the same ground set E , having respective ranks $r_M < r_N$, are *concordant* if their rank functions satisfy that $r_M(Y) - r_M(X) \leq r_N(Y) - r_N(X)$ for all $X \subset Y \subseteq E$. (4).

Flag matroids are an important family of *Coxeter matroids* (4). There are several equivalent ways to define them; in particular they also have an algebro-geometric interpretation. We proceed constructively. Given pairwise concordant matroids M_1, \dots, M_m on E of ranks $k_1 < \dots < k_m$, consider the collection of flags (B_1, \dots, B_m) , where B_i is a basis of M_i and $B_1 \subset \dots \subset B_m$. Such a collection of flags is called a *flag matroid*, and M_1, \dots, M_m are called the *constituents* of \mathcal{F} .

For each flag $B = (B_1, \dots, B_m)$ in \mathcal{F} let $v_B = v_{B_1} + \dots + v_{B_m}$, where $v_{\{a_1, \dots, a_i\}} = e_{a_1} + \dots + e_{a_i}$. The *flag matroid polytope* is $P_{\mathcal{F}} = \text{conv}\{v_B : B \in \mathcal{F}\}$.

Theorem 5.1 (4, Cor 1.13.5) *If \mathcal{F} is a flag matroid with constituents M_1, \dots, M_k , then $P_{\mathcal{F}} = P_{M_1} + \dots + P_{M_k}$.*

As mentioned above, this implies that every flag matroid polytope is a signed Minkowski sum of simplices Δ_I ; the situation is particularly nice for truncation flag matroids, which we now define.

Let M be a matroid over the ground set E with rank r . Define M_i to be the rank i *truncation* of M , whose bases are the independent sets of M of rank i . One easily checks that the truncations of a matroid are concordant, and this motivates the following definition of Borovik, Gelfand, Vince, and White.

Definition 5.2 (3) *The flag $\mathcal{F}(M)$ with constituents M_1, \dots, M_r is a flag matroid, called the truncation flag matroid or underlying flag matroid of M .*

Our next goal is to present the decomposition of a truncation flag matroid polytope as a signed Minkowski sum of simplices. For that purpose, we define the *gamma invariant* of M to be $\gamma(M) = b_{20} - b_{10}$, where $T_M(x, y) = \sum_{i,j} b_{ij} x^i y^j$ is the Tutte polynomial of M .

Proposition 5.3 *The gamma invariant of a matroid is given by*

$$\gamma(M) = \sum_{I \subseteq E} (-1)^{r-|I|} \binom{r - r(I) + 1}{2}.$$

Unlike the beta invariant, the gamma invariant is not necessarily nonnegative. In fact its sign is not simply a function of $|E|$ and r . For example, $\gamma(U_{k,n}) = -\binom{n-3}{k-1}$, and $\gamma(U_{k,n} \oplus C) = \binom{n-2}{k-1}$ where C denotes a coloop.

As we did with the beta invariant, define the *signed gamma invariant* of M to be $\tilde{\gamma}(M) = (-1)^{r(M)} \gamma(M)$.

Theorem 5.4 *The truncation flag matroid polytope of M can be expressed as:*

$$P_{\mathcal{F}(M)} = \sum_{I \subseteq E} \tilde{\gamma}(M/I) \Delta_{E-I}.$$

Proof: By Theorems 2.5 and 5.1, $P_{\mathcal{F}(M)}$ is

$$\sum_{i=1}^r P_{M_i} = \sum_{i=1}^r \sum_{I \subseteq E} \sum_{J \subseteq I} (-1)^{|I|-|J|} (i - r_i(E - J)) \Delta_I,$$

where $r_i(A) = \min\{i, r(A)\}$ is the rank function of M_i . Then

$$\begin{aligned} P_{\mathcal{F}(M)} &= \sum_{I \subseteq E} \left[\sum_{J \subseteq I} (-1)^{|I|-|J|} \sum_{i=r(E-J)+1}^r (i - r(E - J)) \right] \Delta_I \\ &= \sum_{I \subseteq E} \left[\sum_{J \subseteq I} (-1)^{|I|-|J|} \binom{r - r(E - J) + 1}{2} \right] \Delta_I \\ &= \sum_{I \subseteq E} \left[\sum_{X \subseteq I} (-1)^{|X|} \binom{r_{M/(E-I)} - r_{M/(E-I)}(X) + 1}{2} \right] \Delta_I \\ &= \sum_{I \subseteq E} \tilde{\gamma}(M/(E - I)) \Delta_I \end{aligned}$$

as desired. □

Corollary 5.5 *If a connected matroid M has n elements, then*

$$\text{Vol } P_{\mathcal{F}(M)} = \frac{1}{(n-1)!} \sum_{(J_1, \dots, J_{n-1})} \tilde{\gamma}(M/J_1) \tilde{\gamma}(M/J_2) \cdots \tilde{\gamma}(M/J_{n-1}),$$

summing over the ordered collections of sets $J_1, \dots, J_{n-1} \subseteq [n]$ such that, for any distinct i_1, \dots, i_k , $|J_{i_1} \cap \dots \cap J_{i_k}| < n - k$.

Proof: This follows from Proposition 3.2 and Theorem 5.4. □

Acknowledgements

This work is part of the SFSU-Colombia Combinatorics Initiative. We are very grateful to SFSU and the Universidad de Los Andes for supporting this initiative.

References

- [1] F. Ardila, A. Fink, and F. Rincon. *Valuations for matroid polytope subdivisions*. Canadian Mathematical Bulletin, to appear.
- [2] L. J. Billera, N. Jia, and V. Reiner. *A quasisymmetric function for matroids*. arXiv:math.CO/0606646. Preprint, 2007.

- [3] A. Borovik, I.M. Gelfand, A. Vince, and N. White. *The lattice of flats and its underlying flag matroid polytope*. Ann. Combin. **1** (1997) 17-26.
- [4] A. Borovik, I.M. Gelfand, and N. White. *Coxeter matroids*. Birkhäuser, Boston (2003).
- [5] H. Crapo. *A higher invariant for matroids*. J. Combin. Theory **2** (1967), 406-417.
- [6] J. A. de Loera, D. Haws, and M. Köppe. *Erhart polynomials of matroid polytopes and polymatroids*. arXiv:math.CO/0710.4346v1. Preprint, 2007.
- [7] H. Derksen. *Symmetric and quasisymmetric functions associated to polymatroids*. arXiv:math.CO/0801.4393v1 Preprint, 2008.
- [8] W. Fulton. *Introduction to toric varieties*. Ann. of Math. Stud. 131, Princeton Univ. (1993).
- [9] I.M. Gelfand, M. Goresky, R.D. MacPherson, and V.V. Serganova. *Combinatorial geometries, convex polyhedra, and Schubert cells*. Advances Math. **63** (1987), 301-316.
- [10] L. Lafforgue. *Chirurgie des grassmanniennes*. CRM Monograph Series, **19**. American Mathematical Society, Providence, RI (2003).
- [11] T. Lam and A. Postnikov. *Alcoved polytopes*. Discrete and Computational Geometry **38** (2007), 453-478.
- [12] P. McMullen. *Valuations and Euler-type relations on certain classes of convex polytopes*. Proc. London Math. Soc. **3-35** (1977), 113-135.
- [13] J. G. Oxley. *Matroid theory*. Oxford University Press. New York, 1992.
- [14] A. Postnikov. *Permutohedra, associahedra and beyond*. arXiv:math.CO/0507163 v1. Preprint, 2005.
- [15] D. Speyer. *Tropical linear spaces*. arXiv:math.CO/0410455. Preprint, 2004.
- [16] D. Speyer. *A matroid invariant via the K-theory of the Grassmannian*. arXiv:math.CO/0603551. Preprint, 2006.
- [17] R. Stanley. *Eulerian partition of a unit hypercube*. Higher combinatorics, M. Aigner ed. Reidel Dordrecht, Boston (1977).
- [18] B. Sturmfels. *Solving systems of polynomial equations*. CBMS Regional Conference in Mathematics, **97**. AMS (2002).
- [19] D.J.A. Welsh. *Matroid theory*. Academic Press Inc. London (1976).
- [20] N. White. *Theory of matroids*. Encyclopedia of Mathematics and its Applications, **26**. Cambridge University Press (1986).