

Cover time of a random graph with given degree sequence

Mohammed Abdullah, Colin Cooper, Alan Frieze

► **To cite this version:**

Mohammed Abdullah, Colin Cooper, Alan Frieze. Cover time of a random graph with given degree sequence. 21st International Meeting on Probabilistic, Combinatorial, and Asymptotic Methods in the Analysis of Algorithms (AofA'10), 2010, Vienna, Austria. pp.1-20. hal-01185603

HAL Id: hal-01185603

<https://hal.inria.fr/hal-01185603>

Submitted on 20 Aug 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Cover time of a random graph with given degree sequence

Mohammed Abdullah¹ and Colin Cooper¹ and Alan Frieze^{2†}

¹Department of Computer Science, King's College, University of London, London WC2R 2LS, UK

²Department of Mathematical Sciences, Carnegie Mellon University, Pittsburgh PA15213, USA

In this paper we establish the cover time of a random graph $G(\mathbf{d})$ chosen uniformly at random from the set of graphs with vertex set $[n]$ and degree sequence \mathbf{d} . We show that under certain restrictions on \mathbf{d} , the cover time of $G(\mathbf{d})$ is with high probability asymptotic to $\frac{d-1}{d-2} \frac{\theta}{d} n \log n$. Here θ is the average degree and d is the *effective minimum degree*. The effective minimum degree is the first entry in the sorted degree sequence which occurs order n times.

Keywords: Random walks, random graphs, cover time

1 Introduction

Let $G = (V, E)$ be a connected graph with $|V| = n$ vertices and $|E| = m$ edges. For $v \in V$, let C_v be the expected time taken for a simple random walk \mathcal{W}_v on G starting at v , to visit every vertex of G . The *vertex cover time* C_G of G is defined as $C_G = \max_{v \in V} C_v$. The vertex cover time of connected graphs has been extensively studied. It is a classic result of [1] that $C_G \leq 2m(n-1)$. It was shown by [6], [7], that for any connected graph G , the cover time satisfies $(1 - o(1))n \log n \leq C_G \leq (1 + o(1))\frac{4}{27}n^3$. Between these two extremal examples, the cover time, both exact and asymptotic, has been determined for a number of different classes of random and non-random graphs. The above bounds being general, we can often gain insight into the behavior of random walks by taking into consideration the structural properties of the underlying graphs, and studying how estimates of the the main parameters - including cover time - are affected in light of this knowledge.

One way is to specify the degrees of the vertices and investigate how far this determines the properties of the walk. For any connected graph, [10] gave a cover time upper bound of $4n^2 d_{ave}/d_{min}$, where d_{ave} is the average degree and d_{min} is the minimum degree. For d -regular graphs, [8] improved this to $2n^2$. It is interesting that the vertex degree d does not itself feature in this bound. Although the bound of [10] is clearly $O(n^2)$ when d_{ave}/d_{min} is a constant, the factor of $1/d_{min}$ means that the result is not robust to small numbers of small degree vertices that should have inconsequential effects on the cover time. The paper [5] addresses this to some extent with a result on cyclic cover time. For an optimal cyclic list of all the vertices of a graph, the *cyclic cover time* is the expected time it takes a random walk to travel

[†]Research supported in part by NSF Grant ccf0502793

from vertex to vertex along the list, until it completes a full cycle. This is obviously an upper bound on the cover time. The paper [5] show that the cyclic cover time for a graph is $\Theta(n^2 d_{ave} (d^{-1})_{ave})$, where $(d^{-1})_{ave}$ is the average of the inverse of the degrees. Note, in the case of a graph where the sets of vertices of degree d and $\Delta \gg d$ each make up a constant proportion of the total number of vertices, we get $n^2 d_{ave} (d^{-1})_{ave} = \Omega(n^2 \Delta / d)$

In this paper we extend the above results by studying the cover time of random graphs $\mathcal{G}(\mathbf{d})$ picked uniformly at random (uar) from the set $\mathcal{G}(\mathbf{d})$ of simple graphs with vertex set $V = [n]$ and degree sequence $\mathbf{d} = (d_1, d_2, \dots, d_n)$. We make the following definitions: Let $V_j = \{i \in V : d_i = j\}$ and let $n_j = |V_j|$. Let $\sum_{i=1}^n d_i = 2m$ and let $\theta = 2m/n$ be the average degree.

It seems reasonable to ask how the various entries in the degree sequence affect the cover time. In particular, how much does the cover time depend on the vertices of low degree, and how much on the average degree of the graph? In fact, as in [5], both parameters play a part, as is shown in Theorem 1.

Immediately on fixing the degree sequence, \mathbf{d} , some definitional problems arise, as e.g. there may be just a few low degree vertices spread over a wide range. To get round this, we define an *absolute* minimum degree δ , and an *effective* minimum degree d . The effective minimum degree is the first entry in the sorted degree sequence which occurs order n times. We fix the minimum degree at 3 to ensure the graph is connected (**whp**). Between the minimum degree, and the effective minimum degree, we place an upper bound on the number n_i of vertices of degree i . The bound we choose of $n_i = O(n^{ci/d})$, is not as arbitrary as it looks. Certainly when $c = d/(d-1)$, the effective minimum degree drops below d , so clearly there is some $c < d/(d-1)$ which is critical. Finally, we make some constraints on the average degree and upper tail of the degree sequence, to ensure simple graphs occur with high enough probability in the configuration model. Thus we are left with the following list of conditions.

Let $0 < \alpha < 1$ be constant, $0 < c < 1/8$ be constant and let d be a positive integer. Let $\gamma \rightarrow \infty$ with n . We suppose the degree sequence \mathbf{d} satisfies the following conditions:

- (i) Average degree $\theta = o(\sqrt{\log n})$.
- (ii) Minimum degree $\delta \geq 3$.
- (iii) For $\delta \leq i < d$, $n_i = O(n^{ci/d})$.
- (iv) $n_d = \alpha n + o(n)$. We call d the *effective minimum degree*.
- (v) Maximum degree $\Delta = O(n^{c(d-1)/d})$.
- (vi) Upper tail size $\sum_{j=\gamma\theta}^{\Delta} n_j = O(\Delta)$.

We call a degree sequence \mathbf{d} which satisfies conditions (i)–(vi) *nice*, and apply the same adjective to $\mathcal{G}(\mathbf{d})$.

Theorem 1 *Let $G(\mathbf{d})$ be chosen uar from $\mathcal{G}(\mathbf{d})$, where \mathbf{d} is nice. Then whp*

$$C(G(\mathbf{d})) \sim \frac{d-1}{d-2} \frac{\theta}{d} n \log n.$$

We note that if $d \sim \theta$, i.e. the graph is pseudo-regular, then

$$C(G) \sim \frac{d-1}{d-2} n \log n,$$

which extends the result of [3] for random d -regular graphs.

Structure of the paper

The proof of Theorem 1 is based on an application of (5) below. Put simply, (5) says that, if we ignore which vertices the random walk visits during the mixing time, the probability a vertex v is not visited by step t is asymptotic to $\exp(-\pi_v t/R_v)$. Here $\pi_v = d(v)/2m$ and R_v is the expected number of returns to v during the mixing time, for a walk starting at v . We estimate R_v in Section 4, and describe and prove the required **whp** graph properties in Section 3. The proof that (5) is valid **whp** for $\mathcal{G}(\mathbf{d})$ is similar to proofs in earlier papers and is given in the Appendix. The cover time $C(G)$ is established as follows in Section 5. Firstly a general upper bound is proved in Section 5.1. In Section 5.2 a lower bound is determined by the set of vertices S which maximize $\sum_{v \in S} \exp(-\pi_v t/R_v)$.

2 Estimating first visit probabilities

Convergence of the random walk

In this section G denotes a fixed connected graph with n vertices. A random walk \mathcal{W}_u is started from a vertex u . Let $\mathcal{W}_u(t)$ be the vertex reached at step t , let P be the matrix of transition probabilities of the walk and let $P_u^{(t)}(v) = \mathbf{Pr}(\mathcal{W}_u(t) = v)$. We assume that the random walk \mathcal{W}_u on D is ergodic with stationary distribution π , where $\pi_v = d(v)/(2m)$, and $d(v)$ is the degree of vertex v .

Let

$$d(t) = \max_{u,x \in V} |P_u^{(t)}(x) - \pi_x|,$$

and let T be a positive integer such that for $t \geq T$

$$\max_{u,x \in V} |P_u^{(t)}(x) - \pi_x| \leq n^{-3}. \quad (1)$$

Fix two vertices u, v . Considering the walk \mathcal{W}_v , starting at v , let $r_t = \mathbf{Pr}(\mathcal{W}_v(t) = v)$ be the probability that this walk returns to v at step $t = 0, 1, \dots$. Let

$$R_T(z) = \sum_{j=0}^{T-1} r_j z^j \quad (2)$$

and

$$\lambda = \frac{1}{KT} \quad (3)$$

for a sufficiently large constant K .

For $t \geq T$ let $\mathbf{A}_v(t)$ be the event that \mathcal{W}_u does not visit v in steps $T, T+1, \dots, t$.

Lemma 2 *Suppose that*

(a) *For some constant $\psi > 0$, we have*

$$\min_{|z| \leq 1+\lambda} |R_T(z)| \geq \psi.$$

(b) *$T^2\pi_v = o(1)$ and $T\pi_v = \Omega(n^{-2})$ for all $v \in V$.*

There exists

$$p_v = \frac{\pi_v}{R_v(1 + O(T\pi_v))}, \quad (4)$$

where

$$R_v = R_T(1)$$

is from (2), such that for all $v \in V$ and $t \geq T$,

$$\Pr(\mathbf{A}_v(t)) = \frac{(1 + O(T\pi_v))}{(1 + p_v)^t} + O(T^2\pi_v e^{-\lambda t/2}). \quad (5)$$

3 Required graph properties

3.1 Mixing time

Given a graph G , the *conductance* $\Phi(G)$ of a random walk \mathcal{W}_u on G is defined by

$$\Phi(G) = \min_{\pi(S) \leq 1/2} \frac{e(S : \bar{S})}{d(S)}$$

where $d(S) = \sum_{i \in S} d_i$, and $e(A : B)$ denotes the number of edges with one endpoint in A and the other in B . The lemma below follows from Lemma 10 of the Appendix by applying (8).

Lemma 3 *Let \mathbf{d} be a nice degree sequence and let $G(\mathbf{d})$ be chosen uniformly at random from the $\mathcal{G}(\mathbf{d})$, then whp*

$$\Phi(G) \geq 0.01.$$

Note that $\Phi(G) \geq 0.01$ in Lemma 3 implies $G(\mathbf{d})$ is connected.

We note a result from Sinclair [11], that

$$|P_u^{(t)}(x) - \pi_x| \leq (\pi_x/\pi_u)^{1/2} (1 - \Phi^2/2)^t. \quad (6)$$

Referring to Lemma 3 and (6), if we choose A sufficiently large and

$$T = A \log n \quad (7)$$

then (1) holds.

There is a technical point here. The result (6) assumes that the walk is lazy. A lazy walk moves to a neighbour with probability 1/2 at any step. This assumption halves the conductance. Asymptotically, the cover time, and the value of $R_T(1)$ are also doubled. Otherwise, the lazy assumption has a negligible effect on the analysis. We will ignore this assumption for the rest of the paper, and continue as though there are no lazy steps.

3.2 Structural properties of $G(\mathbf{d})$

We make our calculations in the configuration model, see Bollobás [2]. Let $W = [2m]$ be our set of *configuration points* and let $W_i = [d_1 + \dots + d_{i-1} + 1, d_1 + \dots + d_i]$, $i \in [n]$, partition W . The function $\phi : W \rightarrow [n]$ is defined by $w \in W_{\phi(w)}$. Given a pairing F (i.e. a partition of W into m pairs) we obtain a (multi-)graph G_F with vertex set $[n]$ and an edge $(\phi(u), \phi(v))$ for each $\{u, v\} \in F$. Choosing a pairing F uniformly at random from among all possible pairings of the points of W produces a random (multi-)graph G_F . Let $\nu = \sum_i d_i(d_i - 1)/(2m)$. Assuming that $\Delta = o(m^{1/3})$ (as it will be for nice sequences), the probability that G_F is simple is given by

$$P_S = \Pr(G_F \text{ is simple}) \sim e^{-\frac{\nu}{2} - \frac{\nu^2}{4}}, \quad (8)$$

(see e.g. [9]), and each simple graph $G \in \mathcal{G}(\mathbf{d})$ is equiprobable.

Observe that our assumptions (i)–(vi) that \mathbf{d} is nice imply that $\nu = o(\sqrt{\log n})$. Indeed if $\theta = \sqrt{\log n}/\gamma^3$ where $\gamma \rightarrow \infty$ then

$$\nu \leq \frac{1}{\theta n} \left(\sum_{j=3}^{\gamma\theta} n_j j^2 + \sum_{j=\gamma\theta}^{\Delta} n_j j^2 \right) \leq \frac{1}{\theta n} (n\gamma^2\theta^2 + O(\Delta^3)) = o(\sqrt{\log n}).$$

All the **whp** statements in this paper fail with probability at most $n^{-\Omega(1)}$, whereas P_S in (8) is at least $e^{-o(\log n)}$. This justifies our use of the configuration model.

Let C be a large constant and let

$$\omega = \log \log \log n, \quad \omega' = C \log \log n. \quad (9)$$

We use these values for ω, ω' throughout the paper. A cycle C or path P is *small*, if it has at most $2\omega' + 1$ vertices, otherwise it is *large*.

Let

$$\ell = B \log^2 n \quad (10)$$

for some large constant B . A vertex v is *light* if it has degree at most ℓ , otherwise it is *heavy*. A small path is *light* if all vertices are light. A small cycle is *light* if it has at most one heavy vertex.

For a vertex v , let G_v be the subgraph induced by the set of vertices within a distance ω of v . A vertex v is *locally tree-like* if G_v is a tree. A vertex v is *r-regular*, if it is locally tree-like and each vertex of G_v , (with the possible exception of v), has degree r . A vertex v is *r-compliant*, if there exists a tree subgraph \mathcal{T}_v of G_v rooted at v , in which each vertex of \mathcal{T}_v (with the possible exception of v) has degree r .

Lemma 4 Whp:

- (a) No pair of small light cycles are connected by a small light path.
- (b) No pair of vertices on a small light cycle are joined by a small light path.

Proof We first note a useful inequality. For integer $x > 0$, let $F(2x) = \frac{(2x)!}{2^x x!}$, then

$$\frac{F(\theta n - 2x)}{F(\theta n)} = \frac{(\theta n - 2x)!}{\left(\frac{\theta n}{2} - x\right)! 2^{\frac{\theta n}{2} - x}} \frac{\left(\frac{\theta n}{2}\right)! 2^{\frac{\theta n}{2}}}{(\theta n)!} = \left(\prod_{i=1}^x \theta n - 2i + 1 \right)^{-1} \leq \left(\frac{1}{\theta n - 2x + 1} \right)^x.$$

We prove part **(a)** in detail; the calculations for part **(b)** are similar.

(a) Let μ denote the expected number of light cycle-path-cycle subgraphs consisting of cycles of length a, b joined by a path length c . Then

$$\mu \leq \sum_{a=3}^{2\omega'+1} \sum_{b=3}^{2\omega'+1} \sum_{c=1}^{2\omega'+1} \binom{n}{a} \binom{n}{b} \binom{n}{c} \frac{(a-1)!}{2} \frac{(b-1)!}{2} c! ab \ell^{2(a+b+c-2)} \Delta^6 \frac{\mathcal{F}(\theta n - 2(a+b+c+1))}{\mathcal{F}(\theta n)} \quad (11)$$

Explanation. Choose a vertices for one cycle, b vertices for the other and c vertices for the path. At most one vertex in a cycle is not light, and has degree more than ℓ (and at most Δ). Each light vertex has up to $\ell(\ell-1)$ ways to connect to a neighbour, for a total of (at least) $((a-1) + (b-1) + c)$ light vertices, explaining the exponent of ℓ . The remaining, possibly heavy vertex in each cycle can connect in up to $\Delta(\Delta-1)$ ways to neighbours in the cycle and $\Delta-2$ ways from a cycle to a path. Thus μ is bounded by

$$\begin{aligned} \mu &= \sum_{a=3}^{2\omega'+1} \sum_{b=3}^{2\omega'+1} \sum_{c=1}^{2\omega'+1} n^a n^b n^c \ell^{2(a+b+c-2)} \Delta^6 \left(\frac{1}{\theta n - 12\omega' + 6} \right)^{a+b+c} \\ &\leq \frac{\Delta^6}{\theta n - 12\omega' + 6} \sum_a \sum_b \sum_c \left(\frac{n \ell^2}{\theta n - 12\omega' + 6} \right)^{a+b+c} \\ &= O\left(\frac{\Delta^6 \ell^{12\omega'+6} \omega'^3}{\theta n} \right). \end{aligned}$$

Thus $\Pr(\mu > 0) = o(n^{-\epsilon})$, for some constant $\epsilon > 0$, since $\Delta = O(n^{c(d-1)/d})$ where $c < 1/6$. \square

Lemma 5 Whp:

(a) The number of vertices $v \in V$ that are not d -compliant is at most $n^{4c(d-1)/d}$.

(b) There is no small vertex v , $\delta \leq d(v) < d$ which is not d -compliant.

Proof (a) We lower bound the probability P that v is d -compliant by the success, in the configuration model, of the following process.

Process \mathcal{P} : For $0 \leq i \leq \omega - 1$, and for each vertex w at level i , the first $d-1$ unpaired points of w pair with points of distinct unused vertices u of degree $d(u) \geq d$.

The tree created by process \mathcal{P} involves $N_1 - 1 = d_v \sum_{i=0}^{i=\omega-1} (d-1)^i \leq \Delta(d-1)^\omega$ pairings. Let σ represent the sum of degrees of vertices of degree less than d . Thus

$$P \geq \prod_{i=1}^{N_1} \frac{\theta n - i\Delta - \sigma}{\theta n - 2i + 1} \geq \left(1 - \frac{N_1\Delta + \sigma}{\theta n} \right)^{N_1}.$$

Let X count the number of vertices v that are not d -compliant. Using the inequality $1 - (1-x)^y \leq xy$ for real x, y , $0 \leq x \leq 1, y \geq 1$, we have

$$\mathbf{E}[X] \leq n(1-P) = n \left(1 - \left(1 - \frac{N_1\Delta + \sigma}{\theta n} \right)^{N_1} \right) \leq \frac{N_1(N_1\Delta + \sigma)}{\theta}. \quad (12)$$

We have that $\Delta = O(n^{c(d-1)/d})$, and from $n_i = O(n^{ci/d})$ we find that $\sigma = O(n^{c(d-1)/d})$. Thus

$$E[X] = \tilde{O}(\Delta^3 + \Delta\sigma) = \tilde{O}(n^{3c(d-1)/d}) \leq K \log^L n n^{3c(d-1)/d}$$

for some $K, L > 0$. Then,

$$\Pr(X \geq n^{4c(d-1)/d}) = \tilde{O}(n^{-c(d-1)/d}). \quad (13)$$

(b) In this case we have that the number of small vertices is $O(n^{\frac{c(d-1)}{d}})$ and so $\mathbf{E}[X] = \tilde{O}(n^{\frac{2c(d-1)}{d}-1})$. \square

Lemma 6 Whp: *There are $n^{1-o(1)}$ d -regular vertices $v \in V$ with $d(v) = d$.*

Proof We consider d -regular vertices that have a root vertex v of degree d . Recall that $n_d = |V_d| = \alpha n + o(n)$ for some constant $\alpha > 0$. Let $N_2 = 1 + d(d-1)^\omega$. A d -regular tree of depth ω contains N_2 vertices. We proceed in a similar manner to Lemma 5, and bound the probability P that a vertex v is d -regular by bounding the probability of success of the construction of a d -regular tree in the configuration model.

$$P = \Pr(\text{a vertex } v \text{ is } d\text{-regular}) = \prod_{i=1}^{N_2-1} \frac{d(n_d - i)}{\theta n - 2i + 1} \geq \left(\frac{d n_d - N_2}{\theta n} \right)^{N_2}. \quad (14)$$

Let M count the number of d -regular vertices, then $\mathbf{E}[M] = \mu = n_d P$, and

$$\mu = \mathbf{E}[M] \geq n^{1-o(1)}. \quad (15)$$

To estimate $\mathbf{Var}[M]$, let I_v be the indicator that vertex v is d -regular. We have

$$\mathbf{E}[M^2] = \mu + \sum_{v \in V_d} \sum_{w \in V_d, w \neq v} \mathbf{E}[I_v I_w], \quad (16)$$

and

$$\mathbf{E}[I_v I_w] = \Pr(v, w \text{ are } d\text{-regular}, G_v \cap G_w = \emptyset) + \Pr(v, w \text{ are } d\text{-regular}, G_v \cap G_w \neq \emptyset).$$

Now

$$\Pr(v, w \text{ are } d\text{-regular}, G_v \cap G_w = \emptyset) = \prod_{i=1}^{2N_2-2} \frac{d(n_d - i - 1)}{\theta n - 2i + 1} \leq P^2. \quad (17)$$

For any vertex v , the number of vertices u such that $G_v \cap G_w \neq \emptyset$ is bounded from above by $N_2 + dN_2^2$. Using this and (17), we can bound (16) from above by $\mu + \mu^2 + \mu(N_2 + dN_2^2)$.

By the Chebychev Inequality, for some constant $0 < \tilde{\epsilon} < 1$,

$$\Pr(|M - \mu| > \mu^{\frac{1}{2} + \tilde{\epsilon}}) \leq \frac{\mathbf{Var}[M]}{\mu^{1+2\tilde{\epsilon}}} = \frac{\mathbf{E}[M^2] - \mathbf{E}[M]^2}{\mu^{1+2\tilde{\epsilon}}} \leq \frac{\mu + \mu N_2 + \mu d N_2^2}{\mu^{1+2\tilde{\epsilon}}} = O(n^{-\epsilon}).$$

The lemma now follows from (15). \square

4 Expected number of returns in the mixing time

The local graph Γ_v . For vertex v , inductively define a sub-graph Γ_v of G_v as follows: If $u \in G_v$ is heavy, delete an edge $(u, w) \in G_v$ iff there is no (w, v) -path that is light. After this process is completed, let Γ_v be the connected component of G_v rooted at v . The following lemma is a consequence of this construction.

Lemma 7 *Either Γ_v is a tree, or Γ_v contains a unique light cycle C .*

Denote by Γ_v° the subset of the vertices of Γ_v consisting of pruned heavy vertices, and vertices at distance ω from the root v .

Lemma 8 *Let \mathcal{W}_v^* denote the walk on Γ_v starting at v with Γ_v° made into absorbing states. Let $R_v^* = \sum_{t=0}^{\infty} r_t^*$ where r_t^* is the probability that \mathcal{W}_v^* is at vertex v at time t . There exists a constant $\zeta \in (0, 1)$ such that*

$$R_v = R_v^* + O(\zeta^\omega).$$

Proofs of a lemma similar to Lemma 8 are given in e.g. [3]. For completeness the proof of Lemma 8 is given in the Appendix.

Lemma 9 *Let $G(d)$ be good. For a vertex $v \in V$,*

(a) *If v is d -regular, then $R_v = \frac{d-1}{d-2} + O(\zeta^\omega)$.*

(b) *If v is d -compliant then $R_v \leq \frac{d-1}{d-2}(1+o(1))$.*

(c) *For any v , $R_v \leq \frac{\delta-1}{\delta-2}(1+o(1))$.*

(a) We calculate R_v^* for a walk \mathcal{W}_v^* on an d -regular tree Γ_v with Γ_v° made into absorbing states. For a biased random walk on $(0, 1, \dots, k)$, starting at vertex 1, with absorbing states 0, k , and with transition probabilities at vertices $(1, \dots, k-1)$ of $q = \mathbf{Pr}(\text{move left})$, $p = \mathbf{Pr}(\text{move right})$; then

$$\mathbf{Pr}(\text{absorption at } k) = \frac{(q/p) - 1}{(q/p)^k - 1}. \quad (18)$$

We project \mathcal{W}_v^* onto $(0, 1, \dots, \omega)$ with $p = \frac{d-1}{d}$ and $q = \frac{1}{d}$ giving

$$\mathbf{Pr}(\text{absorption at } \Gamma_v^\circ) = \left(1 - \frac{1}{d-1}\right) \left(1 + O\left(\frac{1}{(d-1)^\omega}\right)\right).$$

Let f_v be the probability of a return to v . Then

$$R_v^* = \frac{1}{1 - f_v} = \frac{d-1}{d-2} + O\left(\frac{1}{(d-1)^\omega}\right) \quad (19)$$

and part (a) of the lemma follows.

(b) If v is d -compliant, we can prune G_v removing edges from each vertex (other than v) until v is d -regular. Treating the edges as having unit resistance, this pruning process cannot decrease the effective

resistance between v and a hypothetical vertex ζ that is connected by a zero-resistance edge to each of the vertices in Γ_v° (and no others). Then by part **(a)** and Rayleigh's monotonicity law part **(b)** of the lemma follows. (Here we are using the fact that the probability of reaching ζ before returning to v is equal to $\frac{1}{d(v)R}$ where R is the effective resistance between v and ζ . Rayleigh's Law states that deleting edges increases R).

(c) All vertices on a path from v to Γ_v° have degree at least δ . Thus in expectation there are at most $(\delta - 1)/(\delta - 2) + o(1)$ returns to v before absorption. If absorption is at distance ω , the arguments in (a), (b) above apply. If not, absorption is at a heavy vertex $u \in \Gamma_v^\circ$, that is at distance less than ω from v . Such a vertex will have at most two paths back (on the unique light cycle). All other paths to v in $G(\mathbf{d})$ go via other heavy vertices. Hence if a particle is at u , with probability at most $2/\ell$ it will enter a path to v in Γ_v and probability at least $1 - 2/\ell$ enter a path in which it will only reach v by going through another vertex in Γ_v° first. Thus the probability of reaching v in time T after having visited a heavy vertex in Γ_v° is at most $O(T/\ell)$. So $\sum_{t=0}^{\omega} r_t - r_t^* = O(\omega T/\ell) = o(1)$. \square

5 Cover time of $G(\mathbf{d})$

5.1 Upper bound on cover time

Let $T_G(u)$ be the time taken by the random walk \mathcal{W}_u to visit every vertex of a connected graph G . Let U_t be the number of vertices of G which have not been visited by \mathcal{W}_u at step t . We note the following:

$$C_u = \mathbf{E}[T_G(u)] = \sum_{t>0} \Pr(T_G(u) \geq t), \quad (20)$$

$$\Pr(T_G(u) \geq t) = \Pr(T_G(u) > t - 1) = \Pr(U_{t-1} > 0) \leq \min\{1, \mathbf{E}[U_{t-1}]\}. \quad (21)$$

Recall from (5) that $\mathbf{A}_s(v)$ is the event that vertex v has not been visited by time s . It follows from (20), (21) that

$$C_u \leq t + 1 + \sum_{s \geq t} \mathbf{E}[U_s] = t + 1 + \sum_v \sum_{s \geq t} \Pr(\mathbf{A}_s(v)). \quad (22)$$

Let $t_0 = \left(\frac{d-1}{d-2}\frac{\theta}{d}\right) n \log n$ and $t_1 = (1 + \epsilon)t_0$, where $\epsilon = o(1)$ is sufficiently large that all inequalities claimed below hold. We will use the notation d_v for $d(v)$. We assume that the high probability claims of Sections 3, 4 hold. In the Appendix, we establish that condition (a) of Lemma 2 holds. Condition (b) of Lemma 2, that $T\pi_v = o(1)$, holds trivially as the maximum degree is n^a , $a < 1$.

Recall from (4) that $p_v = (1 + (T\pi_v))d_v/(\theta n R_v)$. Thus by (5), the probability that \mathcal{W}_u has not visited v during $[T, t]$ is given by

$$\Pr(\mathbf{A}_t(v)) = (1 + o(1))e^{-tp_v} + O(T^2\pi_v e^{-\lambda t/2}) \quad (23)$$

$$= (1 + o(1))e^{-tp_v}. \quad (24)$$

Thus

$$\begin{aligned} \sum_{t \geq t_1} (1 + o(1))e^{-tp_v} &= (1 + o(1))e^{-t_1 p_v} \sum_{t \geq t_1} e^{-(t-t_1)p_v} \\ &\leq 2p_v^{-1} e^{-t_1 p_v} \\ &= O(1) \frac{\theta n R_v}{d_v} \exp \left\{ -(1 + \Theta(\epsilon)) \frac{d_v}{d} \frac{d-1}{d-2} \frac{\log n}{R_v} \right\}. \end{aligned} \quad (25)$$

We consider the following partition of V :

- (i) $V_A = \bigcup_{\delta \leq i < d} V_i$.
- (ii) $V_B = \bigcup_{i \geq \delta} \{v \in V_i : v \text{ is } d\text{-compliant}\}$.
- (iii) $V_C = \bigcup_{i \geq \delta} \{v \in V_i : v \text{ is not } d\text{-compliant}\}$.

Case (i): $\delta \leq d_v < d$.

For these vertices, Γ_v is d -compliant by Lemma 5. Consider vertices in V_i , $i < d$. By Lemma 9 (b), $R_v \leq (1+o(1))\frac{d-1}{d-2}$ so for $v \in V_i$ (25) is bounded by $O(\theta n)n^{-(1+o(1))\frac{i}{d}}$. Recall that $|V_i| = O(n^{ci/d})$ where $c < 1$. Thus

$$\sum_{v \in V_i} \sum_{t \geq t_1} (1+o(1))e^{-tp_v} \leq O(\theta n)n^{ci/d}n^{-(1+o(1))i/d} = o(t_1).$$

Case (ii): $d \leq d_v$, v is d -compliant.

For $v \in V_B$ (25) is bounded by $O(\theta)n^{-\Theta(\epsilon)}$. Therefore

$$\sum_{v \in V_B} \sum_{t \geq t_1} (1+o(1))e^{-tp_v} \leq \sum_{v \in V_B} O(\theta)n^{-\Theta(\epsilon)} = O(\theta n)n^{-\Theta(\epsilon)} = o(t_1).$$

Case (iii): $d \leq d_v$, v is not d -compliant.

For vertices $v \in V_C$ (25) is bounded by $O(\theta n)n^{-(1+\Theta(\epsilon))\frac{\delta-2}{\delta-1}\frac{d-1}{d-2}}$. By Lemma 5, $|V_C| \leq n^{4c(d-1)/d}$ where $4c < 1/2 \leq \frac{d}{d-2}\frac{\delta-2}{\delta-1}$. Hence

$$\begin{aligned} \sum_{v \in V_C} \sum_{t \geq t_1} (1+o(1))e^{-tp_v} &= \sum_{v \in V_C} O(\theta n)n^{-(1+\Theta(\epsilon))\frac{\delta-2}{\delta-1}\frac{d-1}{d-2}} \\ &= O(n^{4c(d-1)/d}\theta n)n^{-(1+\Theta(\epsilon))\frac{\delta-2}{\delta-1}\frac{d-1}{d-2}} \\ &= o(t_1). \end{aligned}$$

In each of the cases above, the term $\sum_v \sum_{s \geq t} \Pr(\mathbf{A}_s(v)) = o(t_1)$ and thus, from (22), $C_u \leq (1+o(1))t_1$ as required. This completes the proof of the upper bound on cover time of $G(\mathbf{d})$. \square

5.2 Lower bound on cover time

Let $t_2 = (1-\epsilon)t_0$, where $\epsilon = o(1)$ is sufficiently large that all inequalities claimed below hold. For vertex u of degree d , we exhibit a set of vertices S such that at time t_2 the probability the set S is covered by the walk \mathcal{W}_u tends to zero. Hence $T_G(u) > t_2$, **whp** which implies that $C_G \geq t_0 - o(t_0)$.

We construct S as follows. Let S_d be the set of d -regular vertices of degree d . Lemma 6 tells us that $|S_d| = n^{1-o(1)}$. Let $\omega' = C \log \log n$ for some large C . Let S be a maximal subset of S_d such that the distance between any two elements of S is least ω' . Thus $|S| = \Omega(n^{1-o(1)}/d^{\omega'})$.

Let $S(t)$ denote the subset of S which has not been visited by \mathcal{W}_u after step t . Let $v \in S$, then

$$\Pr(\mathbf{A}_v(t_2)) = (1+o(1))e^{-t_2 p_v(1-O(p_v))} + o(n^{-2}).$$

Hence

$$\mathbf{E}(|S(t_2)|) \geq (1+o(1))|S|e^{-(1-\epsilon)t_0 p_v} \tag{26}$$

$$= \Omega\left(\frac{n^{\epsilon/2-o(1)}}{d^{\omega'}}\right) \rightarrow \infty. \tag{27}$$

Let $Y_{v,t}$ be the indicator for the event $A_t(v)$. Let $Z = \{v, w\} \subset S$. We will show (below) that that for $v, w \in S$

$$\mathbf{E}(Y_{v,t_2} Y_{w,t_2}) = \frac{1 + O(T\pi_v)}{(1 + p_Z)^{t_2}} + o(n^{-2}), \quad (28)$$

where $p_Z \sim p_v + p_w + o(1/\log n)$. Thus

$$\mathbf{E}(Y_{v,t_2} Y_{w,t_2}) = (1 + o(1))\mathbf{E}(Y_{v,t_2})\mathbf{E}(Y_{w,t_2})$$

which implies

$$\mathbf{E}(|S(t_2)|(|S(t_2)| - 1)) \sim \mathbf{E}(|S(t_2)|)(\mathbf{E}(|S(t_2)|) - 1). \quad (29)$$

It follows from (27) and (29), that

$$\Pr(S(t_2) \neq \emptyset) \geq \frac{\mathbf{E}(|S(t_2)|)^2}{\mathbf{E}(|S(t_2)|^2)} = \frac{1}{\frac{\mathbf{E}(|S(t_2)|(|S(t_2)|-1))}{\mathbf{E}(|S(t_2)|)^2} + \mathbf{E}(|S(t_2)|)^{-1}} = 1 - o(1).$$

Proof of (28). Let \widehat{G} be obtained from G by merging v, w into a single node Z . This node has degree $2d$ and is d -regular. $R_Z = (R_v + R_w)/2 + \rho$ where ρ is the expected number of passages between v, w in T steps. By Lemma 4 the number of light paths between v, w is at most 2. Using arguments similar to Lemma 9, we find $\rho = O(T/(\delta - 1)^{\omega'}) = o(1/\log n)$.

There is a natural measure-preserving mapping from the set of walks in G which start at u and do not visit v or w , to the corresponding set of walks in \widehat{G} which do not visit Z . Thus the probability that \mathcal{W}_u does not visit v or w in steps $T \dots t$ is asymptotically equal to the probability that a random walk $\widehat{\mathcal{W}}_u$ in \widehat{G} which also starts at u does not visit Z in steps $T \dots t$. The detailed argument is given in [4].

We apply Lemma 2 to \widehat{G} . That $\pi_Z = \frac{2d}{\theta_n}$ is clear. Furthermore, the vertex Z is tree-like up to distance ω in \widehat{G} . The derivation of R_Z as in Lemma 9(a) is valid. The fact that the root vertex of the corresponding infinite tree has degree $2d$ does not affect the calculation of R_Z^* . \square

References

- [1] R. Aleliunas, R.M. Karp, R.J. Lipton, L. Lovász and C. Rackoff, Random Walks, Universal Traversal Sequences, and the Complexity of Maze Problems. *Proceedings of the 20th Annual IEEE Symposium on Foundations of Computer Science* (1979) 218-223.
- [2] B. Bollobás, *A probabilistic proof of an asymptotic formula for the number of labelled regular graphs*, *European Journal on Combinatorics* 1 (1980) 311-316.
- [3] C. Cooper and A. M. Frieze. The cover time of random regular graphs, *SIAM Journal on Discrete Mathematics*, 18 (2005) 728-740.
- [4] C. Cooper and A. M. Frieze. The cover time of the giant component of a random graph, *Random Structures and Algorithms*, 32 (2008) 401-439.
- [5] D. Coppersmith, U. Feige, J. Shearer, Random walks on regular and irregular graphs, *SIAM Journal on Discrete Mathematics*, 9(2), 301-308, 1996.

- [6] U. Feige, A tight upper bound for the cover time of random walks on graphs, *Random Structures and Algorithms*, 6 (1995) 51-54.
- [7] U. Feige, A tight lower bound for the cover time of random walks on graphs, *Random Structures and Algorithms*, 6 (1995) 433-438.
- [8] U. Feige, Collecting coupons on trees, and the cover time of random walks, *Computational Complexity*, 6 (1996/1997), 341–356.
- [9] B. McKay and N. Wormald, Asymptotic enumeration by degree sequence of graphs with degrees $o(n^{1/2})$. *Combinatorica*, 11 (1991), 369-382.
- [10] J.D. Kahn, N. Linial, N.Nisan and M.E. Saks, On the cover time of random walks on graphs. *Journal of Theoretical Probability*, 2(1):121-128 January 1989.
- [11] A. Sinclair, Improved bounds for mixing rates of Markov chains and multicommodity flow. *Combinatorics, Probability and Computing*, 1 (1992), 351-370.

Appendix

Proof of conductance bound in Lemma 3

By the conductance of a configuration C , we mean the conductance of a random walk on the underlying multi-graph $M(C)$. It is however, the configurations we sample uar in the proof of Lemma 10.

Lemma 10 *Let $\mathbf{d} = (d_1, d_2, \dots, d_n)$ be a sequence of natural numbers, satisfying $\min d_i \geq 3$ and $\theta \leq n^{1/4}$. With probability $1 - o(n^{-1/9})$ the conductance Φ of a uar sampled configuration $C(\mathbf{d})$ satisfies $\Phi \geq 0.01$.*

Proof Let $F(a) = a!/((a/2)!2^{(a/2)})$. With this notation,

$$\frac{F(b)F(a-b)}{F(a)} = \frac{\binom{a/2}{b/2}}{\binom{a}{b}} = O(1) \left(\frac{b}{a}\right)^{b/2} \left(1 - \frac{b}{a}\right)^{(a-b)/2}. \quad (30)$$

For any $S \subseteq V$ let $d(S)$ denote the sum of the degrees of the vertices of S . A set S is *small* if $d(S) \leq (\theta n)^{1/4}$. A set is *large* if $(\theta n)^{1/4} \leq d(S) \leq \theta n/2$. Let $\beta < 1$ be a positive constant. We choose $\beta = 0.99$.

SMALL SETS ($\delta|S| \leq d(S) \leq (\theta n)^{1/4}$).

Let $N(s, \beta)$ be the expected number of small sets S of size s with at least $\beta d(S)$ induced edges.

$$N(s, \beta) = \sum_S \binom{d(S)}{\beta d(S)} \frac{F(\beta d(S))F(\theta n - \beta d(S))}{F(\theta n)}. \quad (31)$$

Thus using (30), $\delta s \leq d(s) \leq (\theta n)^{1/4}$ and $\delta \geq 3$ we find

$$\begin{aligned} N(s, \beta) &\leq O(1) \sum_S \left(\frac{e}{\beta}\right)^{\beta d(S)} \left(\frac{\beta d(S)}{\theta n}\right)^{\beta d(S)/2} \\ &\leq O(1) \left(\frac{ne}{s} \left(\frac{e^2}{\beta(\theta n)^{3/4}}\right)^{3\beta/2}\right)^s \\ &= O(n^{-(9\beta/8-1)s}). \end{aligned}$$

Thus

$$\sum_{\substack{|S|=s \\ S \text{ Small}}} N(s, \beta) = O(n^{-(9\beta/8-1)s}).$$

LARGE SETS ($(\theta n)^{1/4} \leq d(S) \leq \theta n/2$).

Let $N(s, \beta)$ be the expected number of large sets S of size s inducing at least $\beta d(S)$ edges. As before, $N(s, \beta)$ is given by (31). Let $d(S) = \alpha \theta n$ where $0 < \alpha \leq 1/2$. Let $\varepsilon = 1 - \beta$. We note the following approximation:

$$\binom{d(S)}{\beta d(S)} = \binom{\alpha \theta n}{\beta \alpha \theta n} = \frac{O(1)}{\sqrt{\varepsilon \beta \alpha \theta n}} \frac{1}{\beta^\beta \alpha^\alpha \varepsilon^\varepsilon \theta^{\varepsilon \alpha \theta n}}.$$

Thus

$$N(s, \beta) \leq \sum_S \frac{O(1)}{\sqrt{\varepsilon \beta \alpha \theta n}} \left(\frac{(\alpha \beta)^{\alpha \beta} (1 - \alpha \beta)^{1 - \alpha \beta}}{(\varepsilon^\varepsilon \beta^\beta)^{2\alpha}}\right)^{\frac{\theta n}{2}} = \sum_S f(S). \quad (32)$$

Let $s = cn$. We henceforth assume that we choose the value $\alpha = \alpha^*$ which maximizes $f(S)$ for $|S| = cn$. With this convention we can write

$$N(cn, \beta) \leq \frac{O(1)}{\sqrt{\varepsilon\beta c(1-c)\alpha\theta n^2}} \left(\left(\frac{(\alpha\beta)^{\alpha\beta}(1-\alpha\beta)^{1-\alpha\beta}}{(\varepsilon^\varepsilon\beta\beta)^{2\alpha}} \right)^{\frac{\theta}{2}} \frac{1}{c^c(1-c)^{1-c}} \right)^n. \quad (33)$$

We split the proof for large sets into two parts: Those sets for which $\alpha \leq 1/\theta$ and those for which $1/\theta \leq \alpha \leq 1/2$.

Case of $\alpha \leq 1/\theta$.

We need to remove the dependence on c in the right hand side of the expression (33) for $N(cn, \beta)$. We first deal with the square root term. Since $\frac{1}{n} \leq c \leq \frac{n-1}{n}$, we have that $c(1-c) \geq \frac{n-1}{n^2}$ and so

$$c(1-c)\alpha\theta n^2 \geq \frac{n-1}{n^2}(\theta n)^{1/4}n \geq (\theta n)^{1/4}/2.$$

Therefore, as β, ε are positive constants,

$$\frac{1}{\sqrt{\varepsilon\beta c(1-c)\alpha\theta n^2}} = \frac{O(1)}{(\theta n)^{1/8}}.$$

We next consider the main term of (33). For $0 \leq x \leq 1/2$, the function

$$g(x) = x^x(1-x)^{1-x}$$

satisfies, $g(0) = 1$ and is monotonically decreasing with minimum $g(1/2) = 1/2$.

Since $d(S) \geq 3s$, and $s = cn$, from $d(S) = \alpha\theta n$ we deduce that $c \leq \alpha\theta/3$. As $\alpha \leq 1/\theta$ then $c \leq \alpha\theta/3 \leq 1/3$. Therefore $g(c) \geq g(\alpha\theta/3)$, and we can replace c by $\alpha\theta/3$ in (33). Hence

$$\begin{aligned} N(cn, \beta) &= \frac{O(1)}{(\theta n)^{1/8}} \left(\frac{(\alpha\beta)^{\alpha\beta\theta/2}(1-\alpha\beta)^{1-\alpha\beta\theta/2}}{(\alpha\theta/3)^{\alpha\theta/3}(1-\alpha\theta/3)^{1-\alpha\theta/3}} \frac{(1-\alpha\beta)^{\theta/2-1}}{(\varepsilon^\varepsilon\beta\beta)^{\alpha\theta}} \right)^n \\ &= \frac{O(1)}{(\theta n)^{1/8}} (\phi(\alpha, \beta, \theta))^n. \end{aligned}$$

We next maximize $\phi(\alpha, \beta, \theta)$. Let $h(x, y) = (yx)^x(1-yx)^{1-x}$ for $0 < x, y \leq 1$. Considering $h(x, y)$ as a function of y , there is a unique maximum at $y = 1$, given by

$$\begin{aligned} \frac{\partial}{\partial y} \log(h(x, y)) &= x \left(\frac{1}{y} - \frac{1-x}{1-yx} \right) = 0, \\ \frac{\partial^2}{\partial y^2} \log(h(x, y)) &= -x \left(\frac{1}{y^2} + \frac{x(1-x)}{(1-yx)^2} \right) < 0. \end{aligned}$$

Therefore $h(x, y) < h(x, 1) = g(x)$. So $h(\alpha\beta\theta/2, 2/\theta) < g(\alpha\beta\theta/2) < g(\alpha\theta/3)$. Hence

$$\phi(\alpha, \beta, \theta) \leq \frac{(1-\alpha\beta)^{\theta/2-1}}{(\varepsilon^\varepsilon\beta\beta)^{\alpha\theta}}.$$

We prove below, that

$$\frac{\partial}{\partial \theta} \left\{ \frac{(1 - \alpha\beta)^{\theta/2-1}}{(\varepsilon^\varepsilon \beta^\beta)^{\alpha\theta}} \right\} < 0. \quad (34)$$

Since $\theta \geq \delta \geq 3$, we have that

$$\frac{(1 - \alpha\beta)^{\theta/2-1}}{(\varepsilon^\varepsilon \beta^\beta)^{\alpha\theta}} \leq \frac{e^{-\alpha\beta/2}}{(\varepsilon^\varepsilon \beta^\beta)^{3\alpha}} \leq \lambda^\alpha,$$

where $\lambda < 0.7$, provided $\beta \geq 0.99$.

Now since $\alpha\theta n \geq (\theta n)^{1/4}$ for large sets, and $\theta \leq n^{1/4}$ by conditions of the lemma, we have that $\alpha n \geq n^{1/16}$. Thus

$$\begin{aligned} N(cn, \beta) &= \frac{O(1)}{(\theta n)^{1/8}} (\phi(\alpha, \beta, \theta))^n \\ &= O(\lambda^{n^{1/16}}). \end{aligned}$$

As $s = cn$ can take at most n values we have that $\sum N(cn, \beta) = O(n\lambda^{n^{1/16}})$.

Proof of (34).

$$\frac{\partial}{\partial \theta} \left\{ \frac{(1 - \alpha\beta)^{\theta/2-1}}{(\varepsilon^\varepsilon \beta^\beta)^{\alpha\theta}} \right\} = \frac{1}{1 - \alpha\beta} \left(\frac{(1 - \alpha\beta)^{\frac{1}{2}}}{(\varepsilon^\varepsilon \beta^\beta)^\alpha} \right)^\theta \log \left(\frac{(1 - \alpha\beta)^{\frac{1}{2}}}{(\varepsilon^\varepsilon \beta^\beta)^\alpha} \right).$$

Let

$$f(\alpha, \beta) = \frac{(1 - \alpha\beta)}{(\varepsilon^\varepsilon \beta^\beta)^{2\alpha}}.$$

When $\alpha = 0$, $f(\alpha, \beta) = 1$. We prove that, for $\beta \geq 0.99$, $f(\alpha, \beta) < 1$ for $\alpha > 0$, which will establish the result. Note that

$$\frac{\partial}{\partial \alpha} f(\alpha, \beta) = \frac{-1}{(\varepsilon^\varepsilon \beta^\beta)^{2\alpha}} (\beta + (1 - \alpha\beta) \log(\varepsilon^\varepsilon \beta^\beta)^2). \quad (35)$$

Consider

$$\begin{aligned} \frac{d}{d\beta} \{ \log(\varepsilon^\varepsilon \beta^\beta)^2 + \beta \} &\equiv \frac{d}{d\beta} \{ \log((1 - \beta)^{1-\beta} \beta^\beta)^2 + \beta \} \\ &= 2 \log \left(\frac{\beta}{1 - \beta} \right) + 1. \end{aligned}$$

For $\beta > \frac{1}{2}$, the last line above is positive, and thus $\log(\varepsilon^\varepsilon \beta^\beta)^2 > -\beta$. It follows that (35) is negative, as required.

Case of $1/\theta \leq \alpha \leq 1/2$.

Continuing to evaluate $N(s, \beta)$ as before, and referring to $f(S)$ as given by the right hand side term of (32), let

$$A(\alpha) = \frac{(\alpha\beta)^{\alpha\beta} (1 - \alpha\beta)^{1-\alpha\beta}}{(\varepsilon^\varepsilon \beta^\beta)^{2\alpha}}.$$

Thus

$$\begin{aligned}\log(A(\alpha)) &= (\alpha\beta) \log(\alpha\beta) + (1 - \alpha\beta) \log(1 - \alpha\beta) - 2\alpha \log(\varepsilon^\varepsilon \beta^\beta), \\ \frac{\partial}{\partial \alpha} \log(A(\alpha)) &= \beta \log(\alpha\beta) - \beta \log(1 - \alpha\beta) - 2 \log(\varepsilon^\varepsilon \beta^\beta).\end{aligned}$$

Setting $\frac{\partial}{\partial \alpha} \log(A(\alpha)) = 0$ gives

$$\alpha = \frac{\varepsilon^{2\varepsilon/\beta} \beta}{1 + \varepsilon^{2\varepsilon/\beta} \beta^2}.$$

Let α_0 be the solution to this when $\beta = 0.99$. Thus $\alpha_0 \approx 0.477$. Also,

$$\frac{\partial^2}{\partial \alpha^2} \log(A(\alpha)) = \beta \left(\frac{1}{\alpha} + \frac{\beta}{1 - \alpha\beta} \right) > 0$$

hence the stationary point α_0 is a minima. As $\theta \geq 3$ and by inspection, $A(0.5) < A(1/3)$ then $A(\alpha_0) \leq A(1/\theta)$. We can use $\alpha^* = 1/\theta$ as the value of α maximizing $A(\alpha)$ in the range $1/\theta \leq \alpha \leq 1/2$. It follows that

$$\begin{aligned}\sum_{\substack{S \text{ Large} \\ \alpha \geq 1/\theta}} f(S) &= \left(\frac{1}{\sqrt{\theta n}} \right) 2^n (A(1/\theta))^{\frac{\theta n}{2}} \\ &= O(1) 2^n \left(\frac{(\beta/\theta)^{\frac{\beta}{2}} (1 - \beta/\theta)^{\frac{1}{2}(\theta - \beta)}}{\varepsilon^\varepsilon \beta^\beta} \right)^n.\end{aligned}$$

Let

$$T(\theta) = \left(\frac{\beta}{\theta} \right)^\beta \left(1 - \frac{\beta}{\theta} \right)^{\theta - \beta},$$

then

$$\frac{\partial}{\partial \theta} \log(T(\theta)) = \log \left(\frac{\theta - \beta}{\theta} \right).$$

Thus $T(\theta)$ is monotone decreasing in θ , and so $T(\theta) \leq T(3)$. Finally

$$\begin{aligned}\sum N(s, \beta) &\leq O(n) 2^n \left(\frac{(\beta/3)^{\frac{\beta}{2}} (1 - \beta/3)^{\frac{1}{2}(3 - \beta)}}{\varepsilon^\varepsilon \beta^\beta} \right)^n \\ &= O(n (0.8)^n).\end{aligned}$$

This completes the proof of the lemma. □

Proof of Lemma 8

For convenience, we restate the lemma.

Lemma 11 *Let \mathcal{W}_v^* denote the walk on Γ_v starting at v with Γ_v° made into absorbing states. Let $R_v^* = \sum_{t=0}^{\infty} r_t^*$ where r_t^* is the probability that \mathcal{W}_v^* is at vertex v at time t . There exists a constant $\zeta \in (0, 1)$ such that*

$$R_v = R_v^* + O(\zeta^\omega).$$

Proof We bound $|R_v - R_v^*|$ by using

$$R_v - R_v^* = \left(\sum_{t=0}^{\omega} r_t - r_t^* \right) + \left(\sum_{t=\omega+1}^T r_t - r_t^* \right) - \sum_{t=T+1}^{\infty} r_t^*. \quad (36)$$

Case $t \leq \omega$. When a particle starting from v is absorbed at Γ_v° , this is either at distance ω , or by a heavy vertex u at distance less than ω from v . In the case of a heavy vertex u , by the light cycle condition, there are at most two light paths back to v from u of length at most ω . All other paths of length at most ω go via other heavy vertices. Hence if a particle is at u , with probability at most $2/\ell$ it will enter a light path to v . Thus the probability of reaching v in time ω after having landed on a heavy vertex of Γ_v° is at most $O(\omega/\ell) = o(\zeta^\omega)$. In the alternative case that absorption is at distance ω from v , then for $t < \omega$, $r_t^* = r_t$. Thus we can write

$$\left(\sum_{t=0}^{\omega} r_t - r_t^* \right) = o(\zeta^\omega). \quad (37)$$

Case $\omega + 1 \leq t \leq T$. Using (6) with $x = u = v$ and $\zeta = (1 - \Phi^2/2) < 1$, we have for $t \geq \omega$, that $r_t = \pi_v + O(\zeta^t)$. Since $\Delta = O(n^a)$, $a < 1$, we have $T\pi_v = o(\zeta^\omega)$ and so

$$\sum_{t=\omega+1}^T |r_t - r_t^*| = \sum_{t=\omega+1}^T r_t \leq \sum_{t=\omega+1}^T (\pi_v + \zeta^t) = O(\zeta^\omega). \quad (38)$$

Case $t \geq T + 1$. It remains to estimate $\sum_{t=T+1}^{\infty} r_t^*$. We upper bound r_t^* by a probability σ_t as follows. Assume first that Γ_v is a tree. Consider an unbiased random walk $X_0^{(b)}, X_1^{(b)}, \dots$ starting at $|b| < a \leq \omega$ on the infinite line $(\dots, -a, \dots, -1, 0, 1, \dots, a, \dots)$. $X_m^{(b)}$ is the sum of m independent ± 1 random variables. The central limit theorem implies that there exists a constant $c > 0$ such that

$$\Pr(|X_{ca^2}^{(0)}| < a) \leq e^{-1/2}. \quad (39)$$

Now for any t and b with $|b| < a$, we have

$$\Pr(|X_\tau^{(b)}| < a, \tau = 0, \dots, t) \leq \Pr(|X_\tau^{(0)}| < a, \tau = 0, \dots, t) \quad (40)$$

which is justified with the following game: We have two walks, A and B coupled to each other, with A starting at position 0 and B at position b , which, w.l.o.g, we shall assume is positive. The walk is a simple random walk which comes to a halt when either of the walks hits an absorbing state (that being, $-a$ or a). Since they are coupled, B will win iff they drift $(a - b)$ to the right from 0 and A will win iff they drift $-a$ to the left from 0. Given the symmetry of the walk, B has a higher chance of winning.

For $t > T$, we define σ_t by

$$\sigma_t = \Pr(|X_\tau^{(0)}| < a, \tau = 0, 1, \dots, t) \leq \left(e^{-1/2} \right)^{\lfloor t/(ca^2) \rfloor}. \quad (41)$$

The paths from v to Γ_v° in the tree satisfy $a \leq \omega$, and so

$$\sum_{t=T+1}^{\infty} \sigma_t \leq \sum_{t=T+1}^{\infty} e^{-t/(3c\omega^2)} \leq \frac{e^{-T/(3c\omega^2)}}{1 - e^{-1/(3c\omega^2)}} = O(\omega^2 e^{-\Theta(\frac{\log n}{\omega^2})}) = O(\zeta^\omega)$$

We now turn to the case where Γ_v contains a unique light cycle C . Let x be the furthest vertex of C from v in Γ_v . This is the only possible place where the random walk is more likely to get closer to v at the next step. We can see this by considering the breadth first construction of Γ_v . Thus we can compare our walk with random walk on $[-a, a]$ where there is a unique value $x < a$ such that only at $\pm x$ is the walk more likely to move towards the origin and even then this probability is at most $2/3$. Using results (39), (40) for the unbiased walk on the line, we have

$$\Pr(\exists \tau \leq ca^2 : |X_\tau^{(b)}| \geq x) \geq 1 - e^{-1/2}.$$

The probability the particle walks from x to a without returning to the cycle is at least $1/3(a-x)$. Thus

$$\Pr(\exists \tau \leq ca^2 : |X_{\tau+a-x}^{(b)}| \geq a) \geq (1 - e^{-1/2})/3a \geq \frac{13}{100a},$$

and so

$$\sigma_t = \Pr(|X_\tau^{(0)}| < a, \tau = 0, 1, \dots, t) \leq (1 - 13/(100a))^{\lfloor t/(2ca^2) \rfloor} \leq e^{-t/(20ca^3)}. \quad (42)$$

As $a \leq \omega$,

$$\sum_{t=T+1}^{\infty} \sigma_t \leq \sum_{t=T+1}^{\infty} e^{-t/(20c\omega^3)} \leq \frac{e^{-T/(20c\omega^3)}}{1 - e^{-1/(20c\omega^3)}} = O\left(\omega^3 e^{-O\left(\frac{\log n}{\omega^3}\right)}\right) = O(\zeta^\omega)$$

□

Condition (a) of Lemma 2

Lemma 12 *There exists a constant $\psi > 0$ such that for $|z| \leq 1 + \lambda$, $|R_T(z)| \geq \psi$.*

Proof As in Lemma 8, we consider the walk \mathcal{W}_v^* on Γ_v , starting from v , and with absorption at Γ_v° . For this walk, let β_t be the probability of a first return to v at step t , and let r_t^* be the probability of a return to v at step t .

Let $\beta(z) = \sum_{t=1}^T \beta_t z^t$, let $\alpha(z) = 1/(1 - \beta(z))$, and write $\alpha(z) = \sum_{t=0}^{\infty} \alpha_t z^t$. Thus α_t is the probability of a return to v at time t for a walk \mathcal{W}_v^\dagger , all of whose excursions from v are length at most T . Observe that $\alpha_t \leq r_t^* \leq r_t$. We shall prove below that the radius of convergence of $\alpha(z)$ is at least $1 + \Omega(1/\omega^3)$.

We can write

$$\begin{aligned} R_T(z) &= \alpha(z) + Q(z) \\ &= \frac{1}{1 - \beta(z)} + Q(z), \end{aligned} \quad (43)$$

where $Q(z) = Q_1(z) + Q_2(z)$, and

$$\begin{aligned} Q_1(z) &= \sum_{t=0}^T (r_t - \alpha_t) z^t \\ Q_2(z) &= - \sum_{t=T+1}^{\infty} \alpha_t z^t. \end{aligned}$$

We note that $Q(0) = 0$, $\alpha(0) = 1$ and $\beta(0) = 0$.

We claim that the expression (43) is well defined for $|z| \leq 1 + \lambda$. We will show below that

$$|Q_2(z)| = o(1) \quad (44)$$

for $|z| \leq 1 + 2\lambda$ and thus the radius of convergence of $Q_2(z)$ (and hence $\alpha(z)$) is greater than $1 + \lambda$. This will imply that $|\beta(z)| < 1$ for $|z| \leq 1 + \lambda$. For suppose there exists z_0 such that $|\beta(z_0)| \geq 1$. Then $\beta(|z_0|) \geq |\beta(z_0)| \geq 1$ and we can assume (by scaling) that $\beta(|z_0|) = 1$. We have $\beta(0) < 1$ and so we can assume that $\beta(|z|) < 1$ for $0 \leq |z| < |z_0|$. But as ρ approaches 1 from below, (43) is valid for $z = \rho|z_0|$ and then $|R_T(\rho|z_0|)| \rightarrow \infty$, contradiction.

Recall that $\lambda = 1/KT$. Clearly $\beta(1) \leq 1$ and so for $|z| \leq 1 + \lambda$

$$\beta(|z|) \leq \beta(1 + \lambda) \leq \beta(1)(1 + \lambda)^T \leq e^{1/K}.$$

Using $|1/(1 - \beta(z))| \geq 1/(1 + \beta(|z|))$ we obtain

$$|R_T(z)| \geq \frac{1}{1 + \beta(|z|)} - |Q(z)| \geq \frac{1}{1 + e^{1/K}} - |Q(z)|. \quad (45)$$

We now prove that $|Q(z)| = o(1)$ for $|z| \leq 1 + \lambda$ and the lemma will follow.

Turning our attention first to $Q_1(z)$, we have

$$|Q_1(z)| \leq (1 + \lambda)^T |Q_1(1)| \leq e^{2/K} \sum_{t=0}^T |r_t - \alpha_t|. \quad (46)$$

From (37), (38) of the proof of Lemma 8, we see that $\sum_{t=0}^T |r_t - \alpha_t| = o(1)$, hence $|Q_1(z)| = o(1)$.

We now consider $Q_2(z)$. As in Lemma 8, let r_t^* be the probability that a walk \mathcal{W}_v^* on Γ_v starting at v has not been absorbed at Γ_v° by step t . Then $\alpha_t \leq r_t^* \leq \sigma_t$, so

$$|Q_2(z)| \leq \sum_{t=T+1}^{\infty} \sigma_t |z|^t,$$

In the case where G_v is a tree we can use (41) to prove that the radius of convergence of $Q_2(z)$ is at least $e^{1/(3c\omega^2)} \gg 1 + 2\lambda$. So for $|z| \leq 1 + \lambda$,

$$|Q_2(z)| \leq \sum_{t=T+1}^{\infty} e^{\lambda t - t/(3c\omega^2)} = o(1).$$

In the case that G_v contains a unique cycle, we can use (42) to see that the radius of convergence of $Q_2(z)$ is at least $e^{\frac{1}{20c\omega^3}} \gg 1 + 2\lambda$. So for $|z| \leq 1 + \lambda$,

$$|Q_2(z)| \leq \sum_{t=T+1}^{\infty} e^{\lambda t - t/(20c\omega^3)} = o(1).$$

□

