



Minkowski decompositions of associahedra

Carsten Lange

► **To cite this version:**

Carsten Lange. Minkowski decompositions of associahedra. Bousquet-Mélou, Mireille and Wachs, Michelle and Hultman, Axel. 23rd International Conference on Formal Power Series and Algebraic Combinatorics (FPSAC 2011), 2011, Reykjavik, Iceland. Discrete Mathematics and Theoretical Computer Science, DMTCS Proceedings vol. AO, 23rd International Conference on Formal Power Series and Algebraic Combinatorics (FPSAC 2011), pp.611-622, 2011, DMTCS Proceedings. <hal-01215039>

HAL Id: hal-01215039

<https://hal.inria.fr/hal-01215039>

Submitted on 13 Oct 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Minkowski decompositions of associahedra

Carsten Lange[†]

Freie Universität Berlin, FB Mathematik und Informatik, Berlin, Germany

Abstract. Realisations of associahedra can be obtained from the classical permutahedron by removing some of its facets and the set of facets is determined by the diagonals of certain labeled convex planar n -gons as shown by Hohlweg and Lange (2007). Ardila, Benedetti, and Doker (2010) expressed polytopes of this type as Minkowski sums and differences of scaled faces of a standard simplex and computed the corresponding coefficients y_I by Möbius inversion from the z_I if tight right-hand sides z_I for all inequalities of the permutahedron are assumed. Given an associahedron of Hohlweg and Lange, we first characterise all tight values z_I in terms of non-crossing diagonals of the associated labeled n -gon, simplify the formula of Ardila et al., and characterise the remaining terms combinatorially.

Résumé. Dans un article paru en 2007, Hohlweg et Lange décrivent des associaèdres réalisés à partir du permutoèdre en enlevant certaines de ses facettes. Ces facettes sont déterminées par les diagonales d’une famille de n -gones étiquetés. En 2010, Ardila, Benedetti et Doker ont montré que ces polytopes s’expriment par des sommes et différences de Minkowski de faces pondérées d’un simplexe. De plus, si les coefficients z_I des inégalités décrivant l’associaèdre à partir du permutoèdre sont optimaux, alors les coefficients y_I correspondants sont calculés par une inversion de Möbius. Étant donné un tel associaèdre, nous décrivons d’abord les valeurs optimales z_I en termes de diagonales non croisées d’un certain n -gone étiqueté, ensuite nous simplifions la formule de Ardila et al. pour finalement décrire combinatoirement les termes restants.

Keywords: reflection and Coxeter groups, lattice polytopes, associahedra, Minkowski sums

1 Generalised permutahedra and associahedra

Generalised permutahedra and Minkowski decompositions. A generalised permutahedron according to A. Postnikov is a convex $(n - 1)$ -polytope that has the following description by inequalities

$$P_n(\{z_I\}) := \{\mathbf{x} \in \mathbb{R}^n \mid \sum_{i \in [n]} x_i = z_{[n]} \text{ and } \sum_{i \in I} x_i \geq z_I \text{ for } \emptyset \subset I \subset [n]\},$$

where $[n]$ denotes the set $\{1, 2, \dots, n\}$, [Pos09]. The classical permutahedron, as described for example by G. M. Ziegler, [Zie98], corresponds to the choice $z_I = \frac{|I|(|I|+1)}{2}$ for $\emptyset \subset I \subseteq [n]$. Obviously, some of the above inequalities may be redundant for $P_n(\{z_I\})$ and sufficiently small increases and decreases of z_I of redundant inequalities do not change the set $P_n(\{z_I\})$ unless the inequality $\sum_{i \in I} x_i \geq z_I$ is tight. As described next, all generalised permutahedra have a Minkowski decomposition into faces of a simplex and the coefficients y_I of this decomposition can be computed if *all* (tight) values z_I are known.

[†]Partially supported by the DFG grant (Forschergruppe 565 *Polyhedral Surfaces*).

For any two polytopes P and Q , the Minkowski sum $P + Q$ is defined as $\{p + q \mid p \in P, q \in Q\}$. In contrast, we define the Minkowski difference $P - Q$ of P and Q only if there is a polytope R such that $P = Q + R$. We are interested in decompositions of generalised permutahedra into Minkowski sums and differences of faces of the $(n - 1)$ -dimensional standard simplex

$$\Delta_n = \text{conv}\{e_1, e_2, \dots, e_n\},$$

where e_i is a standard basis vector of \mathbb{R}^n . The faces Δ_I of Δ_n are given by $\text{conv}\{e_i\}_{i \in I}$ for $I \subseteq [n]$. If a polytope P is the Minkowski sum and difference of scaled faces of Δ_n , we say that P has a Minkowski decomposition into faces of the standard simplex. The following two results are key observations.

Lemma 1.1 ([ABD10, Lemma 2.1]) $P_n(\{z_I\}) + P_n(\{z'_I\}) = P_n(\{z_I + z'_I\})$.

If we consider the function $I \mapsto z_I$ that assigns every subset of $[n]$ the corresponding tight value z_I of $P_n(\{z_I\})$, then the Möbius inverse of this function assigns to I the coefficient y_I of a Minkowski decomposition of $P_n(\{z_I\})$ into faces of the standard simplex:

Proposition 1.2 ([ABD10, Proposition 2.3])

Every generalised permutahedron $P_n(\{z_I\})$ can be written uniquely as a Minkowski sum and difference of faces of Δ_n :

$$P_n(\{z_I\}) = \sum_{I \subseteq [n]} y_I \Delta_I$$

where $y_I = \sum_{J \subseteq I} (-1)^{|I \setminus J|} z_J$ for each $I \subseteq [n]$.

Studying properties of generalised permutahedra, Postnikov proved a weaker version of Proposition 1.2 that requires $y_I \geq 0$ for all $I \subseteq [n]$, [Pos09]. Although we can compute the values y_I theoretically if all tight values z_I are known, the formula of Proposition 1.2 is computationally expensive.

We could stop here and be fascinated how the Möbius inversion relates the description by half spaces and Minkowski decompositions. Nevertheless, we go further and study this relationship for certain realisations of associahedra, a certain subclass of generalised permutahedra described in the following subsection. It turns out that the formula of Ardila, Benedetti & Doker can be simplified significantly and that there is a combinatorial interpretation of the terms that remain after this simplification. The simplified formula extracts the combinatorial core data for the Möbius inversion: which subsets J of I are essential to compute y_I . In general, the sets J obtained for the same I but different realisations of an n -dimensional associahedron are different. Nevertheless, the sets J have a simple combinatorial characterisation as shown in Section 3.

Associahedra as generalised permutahedra. Associahedra form a class of combinatorially equivalent simple polytopes and can be realised as generalised permutahedra. They are often defined by specifying their 1-skeleton or graph. A theorem of G. Kalai, [Kal88], implies that the face lattice of an $(n - 1)$ -dimensional associahedron As_{n-1} is in fact completely determined by this graph. Now, the graph of an associahedron is isomorphic to a graph with all triangulations (without new vertices) of a convex plane $(n + 2)$ -gon Q as vertex set and all pairs of distinct triangulations that differ in precisely one proper diagonal⁽ⁱ⁾ as edge set. Alternatively, the edges of As_{n-1} are in bijection with the set of triangulations with

⁽ⁱ⁾ A proper diagonal is a line segment connecting a pair of vertices of Q whose relative interior is contained in the interior of Q . A non-proper diagonal is a diagonal that connects vertices adjacent in ∂Q and a degenerate diagonal is a diagonal where the end-points are equal.

one proper diagonal removed. Similarly, k -faces of As_{n-1} are in bijection to triangulations of Q with k proper diagonals deleted. In particular, the facets of As_{n-1} are in bijection with the proper diagonals of Q . J.-L. Loday published a beautiful construction of associahedra in 2004, [Lod04]. This construction was generalised by C. Hohlweg and C. Lange, [HL07], and explicitly describes realisations of As_{n-1} as generalised permutahedra that depend on combinatorics induced by the choice of a Coxeter element c of the symmetric group Σ_n on n elements. Figure 1 shows two realisations of As_3 for different choices of c . Before explaining how to obtain these realisations, we stop for some general remarks.

S. Fomin and A. Zelevinsky introduced generalised associahedra in the context of cluster algebras of finite type, [FZ03], and it is well-known that associahedra and generalised associahedra associated to cluster algebras of type A are combinatorially equivalent. The construction of [HL07] was generalised by C. Hohlweg, C. Lange, and H. Thomas, [HLT11] to generalised associahedra. Their construction depends also on choosing a Coxeter element c and the normal vectors of the facets are determined by combinatorial properties of c . Since the normal fans of these realisations turn out to be Cambrian fans as described by N. Reading and D. Speyer, [RS09], the obtained realisations are generalised associahedra associated to any given cluster algebra of finite type. N. Reading and D. Speyer conjectured a linear isomorphism between Cambrian fans and g -vector fans associated to cluster algebras of finite type with acyclic initial seed introduced by S. Fomin and A. Zelevinsky, [FZ07]. They proved their conjecture up to an assumption of another conjecture of [FZ07]. S.-W. Yang and A. Zelevinsky gave an alternative proof of the conjecture of Reading and Speyer in [YZ08]. We emphasize in this context that the results of Section 2 and 3 can be read along these lines: the computations of z_I and y_I for fixed I and varying c involve sums over different choices of z_{R_δ} where the choice of δ depends on c . Moreover, the values z_{R_δ} that occur in these sums can be chosen within a large class as described for example in [HLT11]. From this point of view, we suggest that combinatorial properties of the g -vector fan for cluster algebras of finite type A with respect to an acyclic initial seed are reflected by the Minkowski decompositions studied in this manuscript.

We now return to the construction of [HL07] and give a brief outline. The choice of a Coxeter element c corresponds to a partition of $[n]$ into a *down set* D_c and an *up set* U_c :

$$D_c = \{d_1 = 1 < d_2 < \dots < d_\ell = n\} \quad \text{and} \quad U_c = \{u_1 < u_2 < \dots < u_m\}.$$

This partition induces a labeling of Q with label set $[n+1]_0 := [n+1] \cup \{0\}$ as follows. Pick two vertices of Q which are the end-points of a path with $\ell + 2$ vertices on the boundary of Q , label the vertices of this path counter-clockwise increasing using the label set $\overline{D}_c := D_c \cup \{0, n+1\}$ and label the remaining path clockwise increasing using the label set U_c . Without loss of generality, we shall always assume that the label set D_c is on the right-hand side of the diagonal $\{0, n+1\}$ oriented from 0 to $n+1$, see Figure 1 for two examples. We derive the values z_I for some subsets $I \subset [n]$ obtained from this labeled $(n+2)$ -gon Q using proper diagonals of Q as follows. Orient each proper diagonal δ from the smaller to the larger labeled end-point of δ , associate to δ the set R_δ that consists of all labels on the strict right-hand side of δ , and replace the elements 0 and $n+1$ by the smaller respectively larger label of the end-points contained in U_c if possible. For each proper diagonal δ we have $R_\delta \subseteq [n]$ but for $n > 2$, obviously not every subset of $[n]$ is of this type. We set

$$\tilde{z}_I^c := \begin{cases} \frac{|I|(|I|+1)}{2} & \text{if } I = R_\delta \text{ for some proper diagonal } \delta, \\ -\infty & \text{else,} \end{cases}$$

compare Table 1 for the examples of Figure 1. In [HL07] it is shown that $P_n(\{\tilde{z}_I^c\})$ is in fact an associa-




Fig. 1: Two 3-dimensional associahedra $As_3^\xi = P_4(\{\tilde{z}_f^\xi\})$ with vertex coordinates computed for differently chosen Coxeter elements according to [HL07] after application of an orthogonal transformation. The different Coxeter elements are encoded by different labelings of hexagons as indicated.

$D_c = \{1, 3, 4\}$ and $U_c = \{2\}$:

δ	$\{0, 3\}$	$\{0, 4\}$	$\{0, 5\}$	$\{1, 2\}$	$\{1, 4\}$	$\{1, 5\}$	$\{2, 3\}$	$\{2, 4\}$	$\{3, 5\}$
R_δ	$\{1\}$	$\{1, 3\}$	$\{1, 3, 4\}$	$\{2, 3, 4\}$	$\{3\}$	$\{3, 4\}$	$\{1, 2\}$	$\{1, 2, 3\}$	$\{4\}$
\tilde{z}_{R_δ}	1	3	6	6	1	3	3	6	1

$D_c = \{1, 4\}$ and $U_c = \{2, 3\}$:

δ	$\{0, 4\}$	$\{2, 4\}$	$\{3, 4\}$	$\{0, 5\}$	$\{0, 3\}$	$\{1, 2\}$	$\{2, 5\}$	$\{1, 3\}$	$\{1, 5\}$
R_δ	$\{1\}$	$\{1, 2\}$	$\{1, 2, 3\}$	$\{1, 4\}$	$\{1, 3, 4\}$	$\{2, 3, 4\}$	$\{1, 2, 4\}$	$\{3, 4\}$	$\{4\}$
\tilde{z}_{R_δ}	1	3	6	3	6	6	6	3	1

Tab. 1: The tables list R_δ and z_I associated to the proper diagonal δ of a labeled hexagon. The upper table corresponds to the associahedron shown on the left of Figure 1, the bottom one to the one on the right.

hedron of dimension $n - 1$ realised in \mathbb{R}^n for every choice of c . We refer to these realisations as As_{n-1}^c . As shown by C. Hohlweg, C. Lange, and H. Thomas, one also obtains associahedra if the finite values of \tilde{z}_I^c are replaced by the corresponding right-hand sides of a permutahedron obtained as convex hull of a (non-degenerate) Σ_n -orbit of a point of \mathbb{R}^n , where Σ_n acts by permutation of coordinates, [HLT11]. The results of Sections 2 and 3 remain true in this situation.

Some instances of As_{n-1}^c have been studied earlier. For example, the realisations of J.-L. Loday, [Lod04], and of G. Rote, F. Santos, and I. Streinu, [RSS03], related to one-dimensional point configurations, are affinely equivalent to As_{n-1}^c if $U_c = \emptyset$ or $U_c = \{2, 3\}$. Moreover, G. Rote, F. Santos, and I. Streinu point out in Section 5.3 that their realisation is not affinely equivalent to the realisation of F. Chapoton, S. Fomin, and A. Zelevinsky, [CFZ02], which in turn is affinely equivalent to As_{n-1}^c if $U_c = \{2\}$ or $U_c = \{3\}$.

Outline. In this article, we study Minkowski decompositions of a family of realisations of associahedra. In the next section, we explain a combinatorial procedure to compute tight values z_I for the redundant inequalities from the irredundant ones for the realisations of [HL07], compare Theorem 2.8. A key step for the computation is the up and down interval decomposition of I defined in Definition 2.2. In Section 3, we simplify the formula for y_I of Proposition 1.2 and show that most terms in that alternating sum cancel. It turns out that at most four summands remain and they are combinatorially characterised by properties of the up and down interval decomposition of I , see Theorem 3.1 for the precise statement. A full version of this extended abstract that includes all proofs will be published later.

2 Tight values for all z_I^c for As_{n-1}^c

As already mentioned, the facet-defining inequalities for As_{n-1}^c correspond to proper diagonals of Q and these are precisely the irredundant inequalities for the generalised permutahedron $P_n(\{\tilde{z}_I^c\})$. As motivated in the previous section, we want to determine tight values for all z_I , $I \subseteq [n]$, in order to compute the coefficients y_I of the Minkowski decomposition of As_{n-1}^c described by F. Ardila, C. Benedetti, and J. Doker. Since we already know that \tilde{z}_I^c is a tight value if and only if $I = R_\delta$ for some proper diagonal of Q , we now aim for a description of the missing \tilde{z}_I^c . The concept of an up and down interval decomposition induced by the partitioning $D_c \cup U_c$ (or equivalently induced by c) of a given interval $I \subset [n]$ is a key concept that we introduce first.

Definition 2.1 (up and down intervals)

Let $D_c = \{d_1 = 1 < d_2 < \dots < d_\ell = n\}$ and $U_c = \{u_1 < u_2 < \dots < u_m\}$ be the partition of $[n]$ induced by a Coxeter element c .

- (a) A set $S \subseteq [n]$ is a non-empty interval of $[n]$ if $S = \{r, r + 1, \dots, s\}$ for some $0 < r \leq s < n$. We write S as closed interval $[r, s]$ (end-points included) or as open interval $(r - 1, s + 1)$ (end-points not included). An empty interval is an open interval $(k, k + 1)$ for some $1 \leq k < n$.
- (b) A non-empty open down interval is a set $S \subseteq D_c$ such that $S = \{d_r < d_{r+1} < \dots < d_s\}$ for some $1 \leq r \leq s \leq \ell$. We write S as open down interval $(d_{r-1}, d_{s+1})_{D_c}$ where we allow $d_{r-1} = 0$ and $d_{s+1} = n + 1$, i.e. $d_{r-1}, d_{s+1} \in \overline{D}_c$. For $1 \leq r \leq \ell - 1$, we have the empty down interval $(d_r, d_{r+1})_{D_c}$.
- (c) A closed up interval is a non-empty set $S \subseteq U_c$ such that $S = \{u_r < u_{r+1} < \dots < u_s\}$ for some $1 \leq r \leq s \leq m$. We write $[u_r, u_s]_{U_c}$.

We often omit the words *open* and *closed* when we consider down and up intervals. There will not be any ambiguity, because we are not going to deal with closed down intervals and open up intervals. Up intervals are always non-empty, while down intervals may be empty. It turns out to be convenient to distinguish the empty down intervals $(d_r, d_{r+1})_{D_c}$ and $(d_s, d_{s+1})_{D_c}$ if $r \neq s$ although they are equal as sets.

Definition 2.2 (up and down interval decomposition)

Let $D_c = \{d_1 = 1 < d_2 < \dots < d_\ell = n\}$ and $U_c = \{u_1 < u_2 < \dots < u_m\}$ be the partition of $[n]$ induced by a Coxeter element c and $I \subseteq [n]$ be non-empty.

(a) An up and down interval decomposition of type (v, w) of I is a partition of I into disjoint up and down intervals I_1^U, \dots, I_w^U and I_1^D, \dots, I_v^D obtained by the following procedure.

1. Suppose there are \tilde{v} non-empty inclusion maximal down intervals of I denoted by $\tilde{I}_k^D = (\tilde{a}_k, \tilde{b}_k)_{D_c}$, $1 \leq k \leq \tilde{v}$, with $\tilde{b}_k \leq \tilde{a}_{k+1}$ for $1 \leq k < \tilde{v}$. Let $E_i^D = (d_{r_i}, d_{r_i+1})_{D_c}$ denote all empty down intervals with $\tilde{b}_k \leq d_{r_i} < d_{r_i+1} \leq \tilde{a}_{k+1}$ for $0 \leq k \leq \tilde{v}$, $\tilde{b}_0 = 0$, and $\tilde{a}_{\tilde{v}+1} = n + 1$. Denote the open intervals $(\tilde{a}_i, \tilde{b}_i)$ and (d_{r_i}, d_{r_i+1}) of $[n]$ by \tilde{I}_i and E_i respectively.
2. Consider all up intervals of I which are contained in (and inclusion maximal within) some interval \tilde{I}_i or E_i obtained in Step 1 and denote these up intervals by

$$I_1^U = [\alpha_1, \beta_1]_{U_c}, \dots, I_w^U = [\alpha_w, \beta_w]_{U_c}.$$

Without loss of generality, we assume $\alpha_i \leq \beta_i < \alpha_{i+1}$.

3. A down interval $I_i^D = (a_i, b_i)_{D_c}$, $1 \leq i \leq w$, is a down interval obtained in Step 1 that is either a non-empty down interval \tilde{I}_k^D or an empty down interval E_k^D with the additional property that there is some up interval I_j^U obtained in Step 2 such that $I_j^U \subseteq E_k$. Without loss of generality, we assume $b_i \leq a_{i+1}$ for $1 \leq i < w$.

(b) An up and down interval decomposition of type $(1, w)$ is called *nested*. A nested component of I is an inclusion-maximal subset J of I such that the up and down decomposition of J is nested.

The following example illustrates Lemma 2.4 and Lemma 2.5.

Example 2.3

Let $D_c = \{d_1 = 1 < d_2 < \dots < d_\ell = n\}$ and $U_c = \{u_1 < u_2 < \dots < u_m\}$ be the partition of $[n]$ induced by a Coxeter element c . The proper diagonals $\delta = \{a, b\}$, $a < b$, of the labeled polygon Q are in bijection to certain non-empty proper subsets $I \subset [n]$ that have an up and down interval decomposition of type $(1, 0)$, $(1, 1)$, or $(1, 2)$. More precisely, we have

- (a) $R_\delta = (a, b)_{D_c}$ if and only if R_δ has an up and down decomposition of type $(1, 0)$.
- (b) $R_\delta = (0, b)_{D_c} \cup [u_1, a]_{U_c}$ or $R_\delta = (a, n + 1)_{D_c} \cup [b, u_m]_{U_c}$ if and only if R_δ has a decomposition of type $(1, 1)$.
- (c) $R_\delta = (0, n + 1)_{D_c} \cup [u_1, a]_{U_c} \cup [b, u_m]_{U_{p_c}}$ if and only if R_δ has an up and down decomposition of type $(1, 2)$.

Lemma 2.4

Given a partition $[n] = D_c \sqcup U_c$ induced by a Coxeter element c and a non-empty subset $I \subseteq [n]$. Let I have a nested up and down interval decomposition of type $(1, 0)$ with down interval $(a, b)_{D_c}$ and no up interval $[\alpha_i, \beta_i]_{U_c}$. Associate the diagonal $\delta_1 := \{a, b\}$ to I .

- (a) The diagonal δ_1 is a proper diagonal if and only if $I \subset [n]$.
- (b) If the diagonal δ_1 is not proper then $\delta_1 = \{0, n\}$, in particular $U_c = \emptyset$.

(c) $I = R_{\delta_1}$.

Lemma 2.5

Given a partition $[n] = D_c \sqcup U_c$ induced by the Coxeter element c and a non-empty subset $I \subseteq [n]$. Let I have a nested up and down interval decomposition of type $(1, w)$ with $w \geq 1$, down interval $(a, b)_{D_c}$, and up intervals $[\alpha_i, \beta_i]_{U_c}$ for $1 \leq i \leq w$. Associate $w + 1$ diagonals δ_i to I :

$$\delta_1 := \{a, \alpha_1\}, \quad \delta_i := \{\beta_{i-1}, \alpha_i\} \text{ for } 1 < i < w + 1, \text{ and } \quad \delta_{w+1} := \{\beta_w, b\}.$$

Then

- (a) The diagonals δ_i are proper diagonals of Q for $1 < i < w + 1$.
- (b) The diagonal δ_1 is either a proper diagonal or the edge $\{0, u_1\}$ of Q .
- (c) The diagonal δ_{w+1} is either a proper diagonal or the edge $\{u_m, n + 1\}$ of Q .
- (d) The diagonals δ_i are non-crossing.
- (e) Let $W \subseteq [w + 1]$ be the index set of the proper diagonals among $\delta_1, \dots, \delta_{w+1}$ and m be the maximal element of W . Then

$$I = R_{\delta_m} \setminus \left(\bigcup_{i \in W \setminus \{m\}} [n] \setminus R_{\delta_i} \right).$$

Proof: We only prove statement (e). The up and down interval decomposition of I guarantees $I \subseteq R_{\delta_i}$ for $i \in W$. Thus

$$R_{\delta_m} \cap R_{\delta_i} = R_{\delta_m} \setminus ([n] \setminus R_{\delta_i}), \quad \text{for } i \in W \setminus \{m\}$$

and

$$I = \bigcap_{i \in W} R_{\delta_i} = \bigcap_{i \in W \setminus \{m\}} (R_{\delta_m} \cap R_{\delta_i}) = R_{\delta_m} \setminus \left(\bigcup_{i \in W \setminus \{m\}} [n] \setminus R_{\delta_i} \right).$$

□

Lemma 2.6

Given a partition $[n] = D_c \sqcup U_c$ induced by a Coxeter element c . Let I be a non-empty proper subset of $[n]$ with up and down interval decomposition of type (v, w) . Then there are v nested components of type $(1, w_1), \dots, (1, w_v)$ with $w = \sum_{j=1}^v w_j$. For $1 \leq i \leq v$ and $1 \leq j \leq w_i$ let $[\alpha_{i,j}, \beta_{i,j}]_{U_c}$ denote the up intervals nested in the down interval $(a_i, b_i)_{D_c}$ and associate $w_i + 1$ diagonals $\delta_{i,j}$, $1 \leq j \leq w_i + 1$, to each nested component in the same way as in Lemma 2.4 and Proposition 2.5. Then

- (a) The diagonals $\delta_{i,j}$ are non-crossing.
- (b) As in Proposition 2.5 it can happen that we have to adjust to index sets W_1 and W_v to avoid non-proper diagonals.

$$I = \bigcup_{i=1}^v \left(R_{\delta_{i,m_i}} \setminus \left(\bigcup_{j \in W_i \setminus \{m_i\}} [n] \setminus R_{\delta_{i,j}} \right) \right).$$

Proof: There are precisely v nested components, one for every down interval. Moreover, for each up interval $[\alpha, \beta]_{U_c}$ exists a unique down interval $(a, b)_{D_c}$ with $[\alpha, \beta]_{U_c} \subseteq (a, b)$. Depending on the type, apply Lemma 2.4 or Lemma 2.5 to each nested component. Combining the statements gives the claim. □

Definition 2.7 Given a partition $[n] = D_c \sqcup U_c$ induced by a Coxeter element c . Let I be a non-empty proper subset of $[n]$ with up and down interval decomposition of type (v, w) , nested components of type $(1, w_1), \dots, (1, w_v)$ and associated diagonals $\delta_{i,j}$. The subset \mathcal{D}_I of proper diagonals of $\{\delta_{i,j} \mid 1 \leq i \leq v \text{ and } 1 \leq j\}$ is called set of proper diagonals associated to I . Similarly, we say that $\delta \in \mathcal{D}_I$ is a proper diagonal associated to I .

Theorem 2.8

Given a partition $[n] = D_c \sqcup U_c$ induced by a Coxeter element c . Let I be a non-empty proper subset of $[n]$ with up and down interval decomposition of type (v, w) . We use the notation of Lemma 2.6 for its up and down interval decomposition. For non-empty $I \subseteq [n]$ we set

$$z_I^c := \sum_{i=1}^v \left(\sum_{j \in W_i} \tilde{z}_{R_{\delta_{i,j}}}^c - (|W_i| - 1)z_{[n]} \right).$$

Then $P(\{z_I^c\}) = P(\{\tilde{z}_I^c\}) = \text{As}_{n-1}^c$ and all z_I^c are tight.

Proof: The verification of the inequality is a straightforward calculation:

$$\begin{aligned} \sum_{i \in I} x_i &= \sum_{k=1}^v \left(\sum_{i \in R_{\delta_{k,m_k}} \setminus (\cup_{\ell \in W_k \setminus \{m_k\}} [n] \setminus R_{\delta_{k,\ell}})} x_i \right) \\ &= \sum_{k=1}^v \left(\sum_{i \in R_{\delta_{k,m_k}}} x_i - \sum_{\ell \in W_k \setminus \{m_k\}} \sum_{i \in [n] \setminus R_{\delta_{k,\ell}}} x_i \right) \\ &= \sum_{k=1}^v \left(\sum_{i \in R_{\delta_{k,m_k}}} x_i + \sum_{\ell \in W_k \setminus \{m_k\}} \left(- \sum_{i \in [n] \setminus R_{\delta_{k,\ell}}} x_i \right) \right) \\ &\geq \sum_{k=1}^v \left(\tilde{z}_{R_{\delta_{k,w_k}}}^c + \sum_{\ell \in W_k \setminus \{m_k\}} \left(\tilde{z}_{R_{\delta_{k,\ell}}}^c - z_{[n]} \right) \right). \end{aligned}$$

The second equality holds, since the sets $[n] \setminus R_{\delta_{k,\ell}}$ are for fixed k and varying ℓ pairwise disjoint. The last inequality holds, since $\sum_{i \in R_{\delta}} x_i \geq \tilde{z}_{R_{\delta}}^c$ is equivalent to $-\sum_{i \in [n] \setminus R_{\delta}} x_i \geq \tilde{z}_{R_{\delta}}^c - z_{[n]}$ for every proper diagonal δ . Since the irredundant facets of the associahedron As_{n-1}^c correspond to inequalities for proper diagonals δ and these inequalities are tight for As_{n-1}^c , the claim follows. \square

As expected, we have $z_{R_{\delta}}^c = \tilde{z}_{R_{\delta}}^c$ for all proper diagonals δ , since $v = |W_1| = 1$.

3 Combinatorial description and efficient computation of y_I

Proposition 1.2 together with Theorem 2.8 provides a way to compute all Minkowski coefficients y_I since all tight values z_I^c for As_{n-1}^c can be computed. The main goal of this section is to provide the simplified

formula for y_I of Theorem 3.1 which consists of at most four non-zero summand for each $I \subseteq [n]$ and has a combinatorial interpretation.

Throughout this section, we use the following notation and make some general assumptions unless explicitly mentioned otherwise. Let $[n] = D_c \sqcup U_c$ be a partition of $[n]$ induced by a Coxeter element c with $D_c = \{d_1 = 1 < d_2 < \dots < d_\ell = n\}$ and $U_c = \{u_1 < \dots < u_m\}$. Consider a non-empty subset $I \subseteq [n]$ with up and down interval decomposition of type (v, k) . If I has a nested up and down interval decomposition, then in particular $v = 1$ and $I = (a, b)_{D_c} \cup \bigcup_{i=1}^k [\alpha_i, \beta_i]_{U_c}$ with $\alpha_k < \beta_k \leq \alpha_{k+1}$ as before. In this situation, we denote the smallest (respectively largest) element of I by γ (respectively Γ) and consider the diagonals

$$\delta_1 = \{a, b\}, \quad \delta_2 = \{a, \Gamma\}, \quad \delta_3 = \{\gamma, b\}, \quad \text{and} \quad \delta_4 = \{\gamma, \Gamma\}.$$

The subset of proper diagonals of $\{\delta_1, \delta_2, \delta_3, \delta_4\}$ is denoted by \mathcal{D}_I . It is convenient to use the mnemonic δ for diagonals, but unfortunately the meaning of δ_i in this and the preceding section (starting with Lemma 2.4) is not consistent. We now extend our definition of R_δ and z_{R_δ} to non-proper and degenerate diagonals δ . Is $\delta = \{x, y\}$ not a proper diagonal, we set

$$R_\delta := \begin{cases} \emptyset & \text{if } x, y \in \overline{D}_c \\ [n] & \text{otherwise,} \end{cases} \quad \text{and} \quad z_{R_\delta}^c := \begin{cases} 0 & \text{if } R_\delta = \emptyset \\ \frac{n(n+1)}{2} & \text{if } R_\delta = [n]. \end{cases}$$

The main result of this section is

Theorem 3.1

Let I be a non-empty subset of $[n]$ with a nested up and down interval decomposition of type $(1, k)$. Then

$$y_I = \sum_{\delta \in \mathcal{D}_I} (-1)^{|I \setminus R_\delta|} z_{R_\delta}^c.$$

The ideas used to prove Theorem 3.1 also prove the following corollary.

Corollary 3.2

Let I be a non-empty subset of $[n]$ with a nested up and down interval decomposition of type (v, k) and $v > 1$. Then $y_I = 0$.

The rest of this section sketches the proof of Theorem 3.1. We start with some general remarks and then sketch the proof of Theorem 3.1 if $\mathcal{D}_I = \{\delta_1, \delta_2, \delta_3, \delta_4\}$. The same techniques combined with a more detailed statement of Lemma 3.3 settle the remaining cases of Theorem 3.1, see Corollary 3.4.

Combining Proposition 1.2 and Theorem 2.8, we obtain

$$y_I = \sum_{J \subseteq I} (-1)^{|I \setminus J|} z_J^c = \sum_{J \subseteq I} (-1)^{|I \setminus J|} \sum_{i=1}^{v_J} \left(\tilde{z}_{R_{\delta_{i, m_i^J}}}^c + \sum_{j \in W_i^J \setminus \{m_i^J\}} (\tilde{z}_{R_{\delta_{i, j}}}^c - z_{[n]}) \right).$$

This formula is rather awkward at first sight but it turns out that the right-hand side simplifies significantly for our instances As_{n-1}^c . We say that a proper diagonal δ (associated to J) is of type $\tilde{z}_{R_\delta}^c$ (in the expression for y_I), if there exists a set $J \subseteq I$ and an index $i \in [v_J]$ such that $\delta = \delta_{i, m_i^J}^J$. Similarly, we say that a proper diagonal δ (associated to J) is of type $(\tilde{z}_{R_\delta}^c - z_{[n]})$ (in the expression for y_I), if there exists a set J and indices $i \in [v_J]$ and $j \in W_i^J \setminus \{m_i^J\}$ such that $\delta = \delta_{i, j}^J$.




Fig. 2: Here $I = (2, 14)_{D_c} \cup [3, 3]_{U_c} \cup [6, 11]_{U_c}$ with $\gamma = 3$ and $\Gamma = 13$. For $\delta = \{5, 12\}$, the up and down interval decomposition of R_δ is of type $(1, 0)$. Moreover, δ is associated to $\{8, 10\}$, $\{3, 8, 10\}$, $\{3, 8, 10, 13\}$, and $\{8, 10, 13\}$ since $(2, 5) \cap I = \{3\}$ and $(12, 14) \cap I = \{13\}$. Thus the contribution of δ to y_I vanishes. The only diagonals associated to $J \subseteq I$ with up and down interval decomposition of type $(1, 0)$ and non-vanishing contribution to y_I are diagonals associated to only one subset $J \subseteq I$, i.e. $\delta_1 = \{2, 14\}$ and $\delta_2 = \{2, 13\}$ in this example.

Lemma 3.3

Let I be a non-empty proper subset of $[n]$ with up and down interval decomposition of type $(1, k)$.

(a) There is no partition $[n] = D_c \sqcup U_c$ induced by a Coxeter element c and no non-empty $I \subset [n]$ such that \mathcal{D}_I is one of the following sets:

$$\emptyset, \{\delta_2\}, \{\delta_3\}, \{\delta_4\}, \{\delta_1, \delta_2\}, \{\delta_1, \delta_3\}, \{\delta_2, \delta_4\}, \text{ or } \{\delta_3, \delta_4\}.$$

(b) There is a partition $[n] = D_c \sqcup U_c$ induced by a Coxeter element c and a non-empty $I \subset [n]$ such that \mathcal{D}_I is one of the following sets:

$$\{\delta_1\}, \{\delta_1, \delta_4\}, \{\delta_2, \delta_3\}, \{\delta_1, \delta_2, \delta_3\}, \{\delta_1, \delta_2, \delta_4\}, \{\delta_1, \delta_3, \delta_4\}, \{\delta_2, \delta_3, \delta_4\}, \text{ or } \{\delta_1, \delta_2, \delta_3, \delta_4\}.$$

The proof of Part (a) is left to the reader, while a more detailed analysis for Part (b) is implicitly given (and needed) for Corollary 3.4. We now sketch the proof of Theorem 3.1 if $\mathcal{D}_I = \{\delta_1, \delta_2, \delta_3, \delta_4\}$.

Proof: We have to analyse the equation for y_I given above. Let δ be some diagonal $\delta_{i,j}^J$ that occurs on the equation’s right-hand side. In particular, δ is a proper and non-degenerate diagonal, since δ is a diagonal of an up and down interval decomposition. By Example 2.3, the up and down interval decomposition of R_δ is either of type $(1, 0)$, $(1, 1)$ or $(1, 2)$. A good understanding which sets $S \subseteq I$ satisfy $\delta \in \mathcal{D}_S$ and in which cases δ associated to S is of type $\tilde{z}_{R_\delta}^c$ or of type $(\tilde{z}_{R_\delta}^c - z_{[n]})$ in the expression for y_I is essential for the simplification. The complete picture is a case study of the mentioned three cases for R_δ , two of them split further into subcases. In order to illustrate some of the arguments used, we show the simplest case where R_δ has an up and down interval decomposition of type $(1, 0)$.

1. R_δ has up and down decomposition of type $(1, 0)$, see Figure 2.

Then $R_\delta = (\tilde{a}, \tilde{b})_{D_c} \subseteq (a, b)_{D_c}$ and $R_\delta \subseteq I$. Let $S \subseteq I$ be a set with $\delta \in \mathcal{D}_S$. Then $(\tilde{a}, \tilde{b})_{D_c}$ is a nested component of type $(1, 0)$ of S and all other nested components are subsets of $(a, \tilde{a}) \cap I$ and $(\tilde{b}, b) \cap I$. It follows that a set S satisfies $\delta \in \mathcal{D}_S$ if and only if

$$R_\delta \subseteq S \subseteq R_\delta \cup ((a, \tilde{a}) \cap I) \cup ((\tilde{b}, b) \cap I).$$

We now collect all terms for $\tilde{z}_{R_\delta}^c$ in the expression for y_I . Since δ is a proper diagonal, we have $\tilde{z}_{R_\delta}^c \neq 0$ and the resulting alternating sum vanishes if and only if there is more than one term of this

type, that is, if and only if $((a, \tilde{a}) \cap I) \cup ((\tilde{b}, b) \cap I) \neq \emptyset$. If $((a, \tilde{a}) \cap I) \cup ((\tilde{b}, b) \cap I) = \emptyset$, we obtain $(-1)^{|I \setminus R_\delta|} z_{R_\delta}^c$ as contribution for y_I . Moreover, the condition $((a, \tilde{a}) \cap I) \cup ((\tilde{b}, b) \cap I) = \emptyset$ guarantees $\delta \in \mathcal{D}_I$. The diagonal δ_1 is always of this type. Similarly, we have $\delta_2 \in \mathcal{D}_I$ if $\Gamma \in D_c$, $\delta_3 \in \mathcal{D}_I$ if $\gamma \in D_c$, and $\delta_4 \in \mathcal{D}_I$ if $\gamma, \Gamma \in D_c$.

Once the three cases are settled, we have a smaller number of diagonals that have a potential contribution to y_I . We now proceed by another case study and distinguish the four cases

1. $\gamma, \Gamma \in D_c$,
2. $\gamma \in U_c$ and $\Gamma \in D_c$,
3. $\gamma, \Gamma \in U_c$, and
4. $\gamma \in D_c, \Gamma \in U_c$.

Collecting the diagonals in each of these four cases from the previous analysis, additional terms cancel and the claim follows. \square

We end this section listing the precise statements needed to settle the degenerate cases $\mathcal{D}_I \neq \{\delta_1, \delta_2, \delta_3, \delta_4\}$ of Theorem 3.1 in the following corollary.

Corollary 3.4

Let I be non-empty proper subset of $[n]$ with up and down interval decomposition of type $(1, k)$.

(a) Suppose that I satisfies one of the following conditions

- (i) $\mathcal{D}_I = \{\delta_1\}$,
- (ii) $\mathcal{D}_I = \{\delta_1, \delta_3, \delta_4\}$, $(a, b)_D = \{\Gamma\}$, and $\gamma \in U_c$
- (iii) $\mathcal{D}_I = \{\delta_1, \delta_2, \delta_4\}$, $(a, b)_D = \{\gamma\}$, and $\Gamma \in U_c$
- (iv) $\mathcal{D}_I = \{\delta_1, \delta_2, \delta_3\}$ and $(a, b)_D = \{\gamma, \Gamma\}$, or
- (v) $\mathcal{D}_I = \{\delta_2, \delta_3, \delta_4\}$ and $(a, b)_D = \emptyset$.

Then the Minkowski coefficient y_I of As_{n-1}^c is

$$y_I = \sum_{\delta \in \mathcal{D}_I} (-1)^{|I \setminus R_\delta|} z_{R_{\delta_1}}.$$

(b) Suppose that I satisfies one of the following conditions

- (i) $\mathcal{D}_I = \{\delta_1, \delta_4\}$,
- (ii) $\mathcal{D}_I = \{\delta_2, \delta_3\}$,
- (iii) $\mathcal{D}_I = \{\delta_1, \delta_3, \delta_4\}$ and $\bigcup_{i=1}^k [\alpha_i, \beta_i]_{U_c} = \{\Gamma\}$,
- (iv) $\mathcal{D}_I = \{\delta_1, \delta_2, \delta_4\}$ and $\bigcup_{i=1}^k [\alpha_i, \beta_i]_{U_c} = \{\gamma\}$,
- (v) $\mathcal{D}_I = \{\delta_1, \delta_2, \delta_3\}$ and $\bigcup_{i=1}^k [\alpha_i, \beta_i]_{U_c} = \{\gamma, \Gamma\}$.

Then the Minkowski coefficient y_I of As_{n-1}^c is

$$y_I = (-1)^{|\{\gamma, \Gamma\}|} z_{[n]} + \sum_{\delta \in \mathcal{D}_I} (-1)^{|I \setminus R_\delta|} z_{R_{\delta_1}}.$$

References

[ABD10] Federico Ardila, Carolina Benedetti, and Jeffrey Doker. Matroid polytopes and their volumes. *Discrete Comput Geom*, 43:841–854, 2010.

- [CFZ02] Frédéric Chapoton, Sergey Fomin, and Andrei Zelevinsky. Polytopal realizations of generalized associahedra. *Canad Math Bull*, 45:537–566, 2002.
- [FZ03] Sergey Fomin and Andrei Zelevinsky. Y-systems and generalized associahedra. *Ann. of Math.*, 158:977–1018, 2003.
- [FZ07] Sergey Fomin and Andrei Zelevinsky. Cluster algebras iv. coefficients. *Compos Math*, 143:112–164, 2007.
- [HL07] Christophe Hohlweg and Carsten Lange. Realizations of the associahedron and cyclohedron. *Discrete Comput Geom*, 37:517–543, 2007.
- [HLT11] Christophe Hohlweg, Carsten Lange, and Hugh Thomas. Permutahedra and generalized associahedra. *Adv Math*, 226:608–640, 2011.
- [Kal88] Gil Kalai. A simple way to tell a simple polytope from its graph. *J. Combin Theory Ser A*, 49:381–383, 1988.
- [Lod04] Jean-Louis Loday. Realizations of the Stasheff polytope. *Arch Math*, 83:267–278, 2004.
- [Pos09] Alexander Postnikov. Permutahedra, associahedra, and beyond. *Int Math Res Not*, pages 1026–1106, 2009.
- [RS09] Nathan Reading and David Speyer. Cambrian fans. *J Europ Math Soc*, 11:411–447, 2009.
- [RSS03] Günter Rote, Francisco Santos, and Ileana Streinu. *Discrete and Computational Geometry – The Goodman-Pollack Festschrift*, chapter Expansive Motions and the Polytope of Pointed Pseudo-Triangulations, pages 699–736. Number 25 in Algorithms and Combinatorics. Springer-Verlag, Berlin, 2003.
- [YZ08] Shih-Wei Yang and Andrei Zelevinsky. Cluster algebras of finite type via coxeter elements and principal minors. *Transform Groups*, 13:855–895, 2008.
- [Zie98] Günter M. Ziegler. *Lectures on Polytopes*, volume 152 of *Graduate Texts in Mathematics*. Springer-Verlag, Heidelberg, 1998.