

## How often do we reject a superior value? (Extended abstract)

Kamilla Oliver, Helmut Prodinger

► **To cite this version:**

Kamilla Oliver, Helmut Prodinger. How often do we reject a superior value? (Extended abstract). Bousquet-Mélou, Mireille and Wachs, Michelle and Hultman, Axel. 23rd International Conference on Formal Power Series and Algebraic Combinatorics (FPSAC 2011), 2011, Reykjavik, Iceland. Discrete Mathematics and Theoretical Computer Science, DMTCS Proceedings vol. AO, 23rd International Conference on Formal Power Series and Algebraic Combinatorics (FPSAC 2011), pp.741-752, 2011, DMTCS Proceedings. <hal-01215069>

**HAL Id: hal-01215069**

**<https://hal.inria.fr/hal-01215069>**

Submitted on 13 Oct 2015

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# How often do we reject a superior value?

(Extended abstract)

Kamilla Oliver<sup>1†</sup> and Helmut Prodinger<sup>2‡</sup>

<sup>1</sup>Erlangen, Germany

<sup>2</sup>Department of Mathematics, University of Stellenbosch, Stellenbosch, South Africa

**Abstract.** Words  $a_1a_2 \dots a_n$  with independent letters  $a_k$  taken from the set of natural numbers, and a weight (probability) attached via the geometric distribution  $pq^{i-1}$  ( $p + q = 1$ ) are considered. A consecutive record (motivated by the analysis of a skip list structure) can only advance from  $k$  to  $k + 1$ , thus ignoring perhaps some larger (=superior) values. We investigate the number of these rejected superior values. Further, we study the probability that there is a single consecutive maximum and show that (apart from fluctuations) it tends to a constant.

**Résumé.** On considère des mots  $a_1a_2 \dots a_n$  formés de lettres à valeurs entières, tirées de façon indépendante avec une distribution géométrique  $pq^{i-1}$  ( $p + q = 1$ ). Un record  $k + 1$  est dit consécutif si la lettre précédente est  $k$ . La notion est motivée par des considérations algorithmiques. Les autres records sont rejetés. Nous étudions le nombre de records rejetés. Nous étudions aussi la probabilité qu'il y ait un seul maximum consécutif, et montrons qu'elle converge vers une constante, à certaines fluctuations près.

**Keywords:** combinatorics on words, records, generating functions, Rice's method,  $q$ -series

## 1 Introduction

We consider words  $a_1a_2 \dots a_n$  with letters  $a_k$  taken from the set of natural numbers, and a weight (probability) attached to it by saying that the letter  $i \in \mathbb{N}$  occurs with probability  $pq^{i-1}$  ( $p + q = 1$ ) and that the letters are independent. The parameter  $\mathcal{K}(a_1a_2 \dots a_n)$ , which we call the number of *weak consecutive records*, has proved to be essential in the analysis of a skip list structure Louchard and Prodinger (2006).<sup>(i)</sup> The word is scanned from left to right, and assuming that the current record (maximum) is value  $k$ , any letter different from  $k$ ,  $k + 1$  is ignored. If, however, the symbol scanned is one of these, we call it a weak consecutive record, and set the value of the current maximum to it. So, the current record either stays at  $k$  or advances to the next value  $k + 1$ . The skip list version assumes that the first letter of the word defines the first record.

<sup>†</sup>K.O.'s work was partially done when she visited the Department of Mathematics of the University of Stellenbosch. She is thankful for the hospitality encountered there. olikamilla@gmail.com

<sup>‡</sup>hprodinger@sun.ac.za

<sup>(i)</sup> In order to learn more about this structure, the interested reader is invited to consult the earlier paper Louchard and Prodinger (2006), which also appeared in this journal; it also contains many pointers to some interesting earlier papers.

For the sake of clarity, we consider the word 1 3 1 1 2 4 3 5 3 5 1 4 1 2 3 4 6 5 1 and underline each consecutive weak maximum:  $\underline{1} \ 3 \ \underline{1} \ \underline{2} \ 4 \ \underline{3} \ 5 \ \underline{3} \ 5 \ 1 \ \underline{4} \ 2 \ 1 \ 3 \ \underline{4} \ 6 \ \underline{5} \ 1$ . The number of underlined symbols (9 in this case) is the parameter  $\mathcal{K}$  of interest.

In Louchard and Prodinger (2006), the average of the parameter  $\mathcal{K}(n)$  was shown to be (with  $Q = 1/q$ ,  $L = \log Q$ )

$$\mathbb{E}\mathcal{K}(n) = 1 + (Q + 1) \sum_{j=1}^n \binom{n}{j} \frac{(-1)^{j-1} (q; q)_{j-1} p^{j+1} q^j}{1 - q^{j+1}},$$

which was also evaluated asymptotically.

**Theorem 1 [Old theorem]** *The expectation of the  $\mathcal{K}(n)$ -parameter is asymptotic to*

$$\mathbb{E}\mathcal{K}(n) \sim (Q + 1) \log_Q n + \frac{(Q + 1)\gamma}{L} + \frac{Q + 1}{L} \log(p) - (Q + 1)\alpha - \frac{(1 + q)^2}{2pq} + 1 + \delta(\log_Q n).$$

The constant  $\alpha$  is given by

$$\alpha = \sum_{i \geq 1} \frac{q^i}{1 - q^i};$$

$\delta(x)$  is a small periodic function. Its Fourier coefficients could be given in principle.

In Oliver and Prodinger (2009), we investigated the parameter  $\mathcal{M}$ , which is the *maximum of the underlined values*. Now, clearly, for that, we do not need to underline repetitions of the current maximum, as in the instance of the  $\mathcal{K}$ -parameter. So, when our current maximum is  $k - 1$ , we ignore all letters different from  $k$ , and when it occurs (with probability  $pq^{k-1}$ ) we set the current maximum to  $k$ . We obtained explicit and asymptotic enumerations:

**Theorem 2 [Old theorem]** *The average and variance of the  $\mathcal{M}(n)$ -parameter are asymptotically given by*

$$\mathbb{E}\mathcal{M}(n) \sim \log_Q n - \alpha + \frac{\log(p)}{L} + \frac{\gamma}{L} + \frac{1}{2} + \delta_E(\log_Q n),$$

$$\mathbb{V}\mathcal{M}(n) \sim \frac{\pi^2}{6L^2} + \frac{1}{12} - \beta + \delta_V(\log_Q n).$$

Here,  $\delta(x)$  is an unspecified periodic function of period 1 and small amplitude. Its Fourier coefficients could be computed in principle. The poles come from the residues at  $z = \chi_k = \frac{2\pi ik}{L}$ . These expansions follow from explicit expressions for the first and second (factorial) moment:

$$\frac{1}{p} + \sum_{j=1}^{n-1} \binom{n-1}{j} (-1)^{j+1} \frac{(q)_{j-1} q^j p^{j+1}}{1 - q^{j+1}}$$

and

$$\frac{2q}{p^2} + 2 \sum_{j=1}^n \binom{n}{j} (-1)^{j+1} (q)_{j-1} p^{j+1} q^j \left[ -\frac{1}{1 - q^{j+1}} \sum_{m=1}^{j-1} \frac{q^m}{1 - q^m} + \frac{1}{(1 - q^{j+1})^2} \right].$$

□



## 2 The difference between ordinary maxima and consecutive maxima

Since in the present setting the consecutive maximum can only advance by +1, we will miss (reject) some values that are larger than that. In this section, we count the number of rejected values.

Let  $c_k(z, u)$  be the generating function where  $[z^n u^j]c_k(z, u)$  is the probability that a random word of length  $n$  has consecutive maximum equal to  $k$ , and  $j$  better values have been rejected.

### A recursion

$$c_k(z, u) = c_{k-1}(z, u) \frac{zpq^{k-1}}{1 - z[1 + (u-1)q^{k+1} - pq^k]} + \frac{zpq^{k-1}}{1 - z[1 + (u-1)q^{k+1} - pq^k]}.$$

It holds for  $k \geq 1$ , and we assume that  $c_0 = 0$ . It is good to use an abbreviation:

$$\lambda_k := 1 - z[1 + (u-1)q^{k+1} - pq^k].$$

This follows from taking the instance with consecutive maximum  $k-1$ , and attaching the letter  $k$ , followed by an arbitrary sequence of letters different from  $k+1$ ; the large ones are marked by  $u$ . The last terms reflects the situation that  $k$  is the first letter of the word.

Then

$$\frac{c_k(z, u)\lambda_1 \dots \lambda_k}{z^k p^k q^{\binom{k}{2}}} = \frac{c_{k-1}(z, u)\lambda_1 \dots \lambda_{k-1}}{z^{k-1} p^{k-1} q^{\binom{k-1}{2}}} + \frac{\lambda_1 \dots \lambda_{k-1}}{z^{k-1} p^{k-1} q^{\binom{k-1}{2}}}.$$

This first order recursion can now be solved by summation:

$$c_k(z, u) = \frac{z^k p^k q^{\binom{k}{2}}}{\lambda_1 \dots \lambda_k} \sum_{j=0}^{k-1} \frac{\lambda_1 \dots \lambda_j}{z^j p^j q^{\binom{j}{2}}} = \sum_{j=0}^{k-1} \frac{z^{k-j} p^{k-j} q^{\binom{k}{2} - \binom{j}{2}}}{\lambda_{j+1} \dots \lambda_k}.$$

### Expectations

The generating function of the expectations is obtained by differentiation:

$$d_k(z) := \left. \frac{\partial}{\partial u} c_k(z, u) \right|_{u=1},$$

and is finally given by

$$\sum_{k \geq 1} d_k(z),$$

since we must take any final consecutive maximum into account. Now

$$d_k(z) = \sum_{j=0}^{k-1} \frac{z^{k-j} p^{k-j} q^{\binom{k}{2} - \binom{j}{2}}}{\lambda_{j+1} \dots \lambda_k} \Big|_{u=1} \sum_{i=j+1}^k \frac{zq^{i+1}}{1 - z(1 - pq^i)}$$

and

$$\lambda_{j+1} \dots \lambda_k \Big|_{u=1} = (1-w)^{j-k} \prod_{i=j+1}^k (1-wpq^i) = (1-w)^{j-k} (wpq^{j+1}; q)_{k-j}.$$

Rewriting in  $w$ -notation:

$$d_k(z) = \sum_{0 \leq j < i \leq k} \frac{w^{k-j} (-1)^{k-j-1} p^{k-j} q^{\binom{k}{2} - \binom{j}{2}}}{(wpq^{j+1}; q)_{k-j}} \frac{wq^{i+1}}{1-wpq^i}.$$

Summing up leads after a lengthy computation to:

$$\sum_{k \geq 1} d_k(z) = \frac{1}{p} \sum_{j \geq 0} \sum_{i \geq 1} q^{\binom{i}{2}} \frac{(-pq^{j+1}w)^{i+1}}{(wpq^{j+1}; q)_i}.$$

Note (this will be discussed later):

$$\sum_{h \geq 1} \frac{(-q^d w)^h q^{\binom{h}{2}}}{(w; q)_h} = - \sum_{n \geq 1} q^d \frac{(q; q)_{n+d-1}}{(q; q)_d} w^n. \tag{1}$$

Therefore

$$\sum_{k \geq 1} d_k(z) = -\frac{1}{p} \sum_{n \geq 1} (q; q)_{n-1} (wpq)^{n+1} \frac{1}{1-q^{n+1}}.$$

Hence the coefficient of  $w^n$  in this expression is

$$\frac{p^{n-1} q^n}{1-q^n} (q; q)_{n-2}$$

for  $n \geq 2$  and 0 otherwise. And now we rewrite this in terms of coefficients  $z^n$ :

$$[z^n] \sum_{j \geq 1} d_j = (-1)^n [w^n] (1-w)^{n-1} \sum_{j \geq 1} d_j = \sum_{k=1}^{n-1} \binom{n-1}{k} (-1)^{k+1} \frac{p^k q^{k+1}}{1-q^{k+1}} (q; q)_{k-1}.$$

### Second factorial moments

Let

$$f_k(z) := \frac{\partial^2}{\partial u^2} c_k(z, u) \Big|_{u=1}.$$

We have

$$\frac{\partial^2}{\partial u^2} \lambda_{j+1} \dots \lambda_k \Big|_{u=1} = \frac{2}{(1-w)^{j-k} (wpq^{j+1}; q)_{k-j}} \sum_{j+1 \leq h \leq i \leq k} \frac{w^2 q^{h+i+2}}{(1-wpq^h)(1-wpq^i)}.$$

Combined with the rest,

$$f_k(z) = \sum_{0 \leq j < h \leq i \leq k} (-pw)^{k-j} q^{\binom{k}{2} - \binom{j}{2}} \frac{2}{(wpq^{j+1}; q)_{k-j}} \frac{w^2 q^{h+i+2}}{(1-wpq^h)(1-wpq^i)}.$$

And this must be summed (the long computation is not shown here):

$$\sum_{k \geq 1} f_k(z) = -\frac{2q}{p^2} \sum_{N \geq 3} \frac{(q; q)_{N-2} (wpq)^N}{1 - q^N} \sum_{n=1}^{N-2} \frac{1}{1 - q^n}.$$

The coefficient of  $w^N$  is

$$-\frac{2(q; q)_{N-2} p^{N-2} q^{N+1}}{1 - q^N} \sum_{n=1}^{N-2} \frac{1}{1 - q^n}$$

for  $N \geq 1$ , otherwise 0. Now we rewrite this:

$$[z^n] \sum_{k \geq 1} f_k(z) = \sum_{j=1}^{n-1} \binom{n-1}{j} (-1)^j \frac{2(q; q)_{j-1} p^{j-1} q^{j+2}}{1 - q^{j+1}} \sum_{m=1}^{j-1} \frac{1}{1 - q^m}.$$

Let us summarize what we found so far:

**Theorem 3** *The expectation and the second factorial moment of the number of superior elements that we reject when we consider the consecutive maxima instead of the true maxima, assuming random words of length  $n$ , are given by*

$$\sum_{k=1}^{n-1} \binom{n-1}{k} (-1)^{k+1} \frac{p^k q^{k+1}}{1 - q^{k+1}} (q; q)_{k-1}$$

and

$$\sum_{j=1}^{n-1} \binom{n-1}{j} (-1)^j \frac{2(q; q)_{j-1} p^{j-1} q^{j+2}}{1 - q^{j+1}} \sum_{m=1}^{j-1} \frac{1}{1 - q^m}.$$

□

In various places of these computations, the formula (1) was used. It can be shown using Heine's transform Andrews (1976); a proof is included in the full paper.

### Asymptotics

The main ingredient is *Rice's formula* Flajolet and Sedgewick (1995) which allows to write an alternating sum as a contour integral:

$$\sum_{k=1}^n \binom{n}{k} (-1)^k f(k) = -\frac{1}{2\pi i} \int_{\mathcal{C}} \frac{n! \Gamma(-z)}{\Gamma(n+1-z)} f(z) dz.$$

The positively oriented curve  $\mathcal{C}$  encircles the poles  $1, 2, \dots, n$  and no others. (The lower summation index 1 could be replaced by another constant, say  $a$ .) Changing the contour, there are extra residues, which have to be taken into account with the opposite sign, and they constitute the asymptotic expansion of the sum. In our instance, this residue is at 0. There are also contributions from poles at  $\frac{2\pi ik}{L}$ , with  $L = \log Q$ ,  $Q = \frac{1}{q}$ , which we do not compute explicitly. When one collects them, they constitute a (tiny) periodic oscillation.

The function  $f(z)$  extrapolates the sequence  $f(k)$ . This is often obvious, but sometimes requires some work to provide such a function. For our expected value, this discussion leads to

$$\sum_{k=1}^{n-1} \binom{n-1}{k} (-1)^{k+1} \frac{p^k q^{k+1}}{1-q^{k+1}} (q; q)_{k-1} = \frac{1}{2\pi i} \int_{\mathcal{C}} \frac{(n-1)! \Gamma(-z)}{\Gamma(n-z)} \frac{p^z q^{z+1}}{1-q^{z+1}} (q; q)_{z-1} dz.$$

It is not *a priori* clear what  $(q; q)_{z-1} = \frac{(q; q)_z}{1-q^z}$  should be. However, note that

$$(q; q)_z = \frac{(q; q)_{\infty}}{(q; q^{z+1})_{\infty}}.$$

The expansion

$$(q; q)_z \sim 1 - \alpha Lz + \frac{\alpha^2 + \beta}{2} L^2 z^2$$

with

$$\alpha = \sum_{j \geq 1} \frac{q^j}{1-q^j} \quad \text{and} \quad \beta = \sum_{j \geq 1} \frac{q^j}{(1-q^j)^2}$$

is not hard to obtain. Providing this expansion, the rest can be done by a computer, and the expectation comes out as

$$\frac{q}{p} \log_Q n + \frac{q}{p} \log(p) - \frac{q}{p} \alpha + \frac{q\gamma}{pL} - \frac{q(1+q)}{2p^2}.$$

For the second factorial moment, things are more involved, and we must find the continuation of

$$\sum_{m=1}^{j-1} \frac{1}{1-q^m} = j-1 + \sum_{m=1}^{j-1} \frac{q^m}{1-q^m}.$$

But

$$\sum_{m=1}^{j-1} \frac{q^m}{1-q^m} = \sum_{m=1}^j \frac{q^m}{1-q^m} - \frac{q^j}{1-q^j} = \alpha - \sum_{m \geq 1} \frac{q^{m+j}}{1-q^{m+j}} - \frac{q^j}{1-q^j},$$

and thus we may take

$$z-1 + \alpha - \sum_{m \geq 1} \frac{q^{m+z}}{1-q^{m+z}} - \frac{q^z}{1-q^z}.$$

Near  $z = 0$ , this function can be expanded as

$$z-1 + \beta Lz - \frac{1}{Lz} + \frac{1}{2} - \frac{Lz}{12}.$$



Again, the rest can be done by a computer, and the second factorial moment can be obtained. It is not displayed here. From this, the variance is computed by adding the expectation and subtracting the square of the expectation. We summarize the results.

**Theorem 4** *The expectation and the variance of the number of superior elements that we reject when we consider the consecutive maxima instead of the true maxima, assuming random words of length  $n$ , are given by*

$$\text{expectation} \sim \frac{q}{p} \log_Q n + \frac{q}{p} \log(p) - \frac{q}{p} \alpha + \frac{q\gamma}{pL} - \frac{q(1+q)}{2p^2} + \delta.(\log_Q n).$$

and

$$\begin{aligned} \text{variance} \sim & \frac{q \log_Q n}{p^2} \\ & + \frac{q \log(p)}{p^2 L} - \frac{q\alpha}{p^2} + \frac{q\gamma}{p^2 L} - \frac{q^2 \beta}{p^2} + \frac{q^2 \pi^2}{6p^2 L^2} - \frac{2q^2}{p^2 L} \\ & + \frac{q(q^3 - 6 + 16q^2 + q)}{12p^4} + \delta.(\log_Q n). \end{aligned}$$

Here,  $\delta.(x)$  denotes an unspecified tiny periodic function of period 1. It might differ in different places. The leading term in the variance of order  $\log n$  has no fluctuating component, since they cancel out.  $\square$

### 3 Probability of a single (consecutive) winner

#### Exact enumeration

Recall that the generating function of words with consecutive maximum =  $k$  is given by

$$b_k(z) = c_k(z, 1) = \sum_{j=0}^{k-1} \frac{q^{\binom{k}{2} - \binom{j}{2}} (-pw)^{k-j}}{(wpq^{j+1}; q)_{k-j}}.$$

Then

$$e_k(z) = (1 + b_k(z)) \frac{zp^k}{1 - z(1 - pq^k - pq^{k+1})} = - \left( 1 + \sum_{j=0}^{k-1} \frac{q^{\binom{k}{2} - \binom{j}{2}} (-pw)^{k-j}}{(wpq^{j+1}; q)_{k-j}} \right) \frac{wp^k}{1 - wp(1 + q)q^k}$$

is the generating function where the maximum is =  $k + 1$ , and it is a single consecutive maximum. This must be summed over  $k + 1 \geq 1$ , to get the generating function of words with a single consecutive maximum (details are in the full paper):

$$\begin{aligned} \sum_{k \geq 0} e_k(z) &= - \sum_{m \geq 0} (1 + q)^m (wp)^{m+1} \sum_{j \geq 0} q^{j(m+1)} \left( 1 - \sum_{n \geq 1} q^m \frac{(q; q)_{n+m-1}}{(q; q)_m} (wpq^{j+1})^n \right) \\ &=: -A + B. \end{aligned}$$

Then

$$A = \sum_{m \geq 0} (1 + q)^m (wp)^{m+1} \frac{1}{1 - q^{m+1}}$$

and

$$B = \frac{1}{q} \sum_{m \geq 0} (1+q)^m \sum_{n \geq 1} \frac{(q; q)_{n+m-1}}{(q; q)_m} (wpq)^{m+1+n} \frac{1}{1 - q^{m+1+n}}.$$

The coefficient of  $w^N$  in  $A$  is  $\frac{p^N(1+q)^{N-1}}{1-q^N}$  for  $N \geq 1$  and 0 otherwise.

The coefficient of  $w^N$  in  $B$  is

$$p^N q^{N-1} \frac{(q; q)_{N-2}}{1 - q^N} \sum_{m=0}^{N-2} \frac{(1+q)^m}{(q; q)_m}$$

for  $N \geq 2$  and 0 otherwise.

Therefore we get an explicit expression.

**Theorem 5** *The probability that a random word of length  $n$  has a single consecutive maximum is given (exactly) by*

$$\sum_{k=0}^n \binom{n}{k} (-1)^k \frac{p^{k+1}(1+q)^k}{1 - q^{k+1}} + \sum_{k=1}^n \binom{n}{k} (-1)^{k+1} p^{k+1} q^k \frac{(q; q)_{k-1}}{1 - q^{k+1}} \sum_{m=0}^{k-1} \frac{(1+q)^m}{(q; q)_m}.$$

□

### Asymptotics

The first sum is  $O(1/n)$ , so we concentrate on the second one. The continuation of

$$\sum_{m=0}^{k-1} \frac{(1+q)^m}{(q; q)_m}$$

isn't as easy as before, since we cannot push it to infinity, and pulling out dominant terms isn't too obvious either. So we proceed in a different way (computation not shown in this abstract):

$$\sum_{m=0}^{n-1} \frac{(1+q)^m}{(q; q)_m} = \frac{1}{(q; q)_\infty} \sum_{j \geq 0} \frac{(-1)^j q^{\binom{j+1}{2}}}{(q; q)_j} \frac{1 - (1+q)^n q^{jn}}{1 - (1+q)q^j}.$$

Therefore we continue

$$p^{k+1} q^k \frac{(q; q)_{k-1}}{1 - q^{k+1}} \sum_{m=0}^{k-1} \frac{(1+q)^m}{(q; q)_m}$$

via

$$p^{z+1} q^z \frac{(q; q)_{z-1}}{1 - q^{z+1}} \frac{1}{(q; q)_\infty} \sum_{j \geq 0} \frac{(-1)^j q^{\binom{j+1}{2}}}{(q; q)_j} \frac{1 - (1+q)^z q^{jz}}{1 - (1+q)q^j}.$$

At  $z = 0$ , this function is regular, with the value

$$\frac{1}{(q; q)_\infty} \sum_{j \geq 0} \frac{(-1)^j q^{\binom{j+1}{2}}}{(q; q)_j} \frac{j - \log_Q(1+q)}{1 - (1+q)q^j}.$$

This follows from

$$\frac{1 - (1+q)^z q^{jz}}{1 - q^z} \sim \frac{1 - (1+z \log(1+q))(1 - Ljz)}{Lz} \sim j - \log_Q(1+q), \quad (z \rightarrow 0).$$

So, at  $z = 0$ , there is ultimately a simple pole, and, apart from the complicated value, the negative residue is one.

**Theorem 6** *The probability that the consecutive record is single, is asymptotically, as  $n \rightarrow \infty$ , given by*

$$\frac{1}{(q; q)_\infty} \sum_{j \geq 0} \frac{(-1)^j q^{\binom{j+1}{2}}}{(q; q)_j} \frac{j - \log_Q(1+q)}{1 - (1+q)q^j} + \delta(\log_Q n),$$

with a periodic function of period 1 and tiny oscillation  $\delta(x)$ . □

One could also consider a random variable, namely the number of times the consecutive maximum occurs, and compute moments. However, we decided not to go that route here.

## Acknowledgements

We thank Guy Louchard for valuable remarks.

## References

- G. Andrews. *The Theory of Partitions*, volume 2 of *Encyclopedia of Mathematics and its Applications*. Addison–Wesley, 1976.
- G. Andrews, R. Askey, and R. Roy. *Special Functions*, volume 71 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, 1999.
- Y. Baryshnikov, B. Eisenberg, and G. Stengle. A necessary and sufficient condition for the existence of the limiting probability of a tie for first place. *Statist. Probab. Lett.*, 23:203–209, 1995.
- J. J. A. M. Brands, F. W. Steutel, and R. J. G. Wilms. On the number of maxima in a discrete sample. *Statist. Probab. Lett.*, 20:209–217, 1994.
- T. Bruss and C. A. O’Cinneide. On the maximum and its uniqueness for geometric random samples. *J. Appl. Probab.*, 27:598–610, 1990.
- B. Eisenberg, G. Stengle, and G. Strang. The asymptotic probability of a tie for first place. *Ann. Appl. Probab.*, 3:731–745, 1993.

- P. Flajolet and R. Sedgewick. Mellin transforms and asymptotics: Finite differences and Rice's integrals. *Theoretical Computer Science*, 144:101–124, 1995.
- P. Kirschenhofer and H. Prodinger. The number of winners a in discrete geometrically distributed sample. *Annals in Applied Probability*, 6:687–694, 1996.
- G. Louchard and H. Prodinger. Analysis of a new skip list variant. *Discrete Mathematics and Theoretical Computer Science*, proc. AG:365–374, 2006.
- G. Louchard and H. Prodinger. The number of elements close to near-records in geometric samples. *Quaestiones Mathematicae*, 29:447–470, 2006.
- K. Oliver and H. Prodinger. Consecutive records in geometrically distributed words. *submitted*, 2009.
- A. Pakes and F. W. Steutel. On the number of records near the maximum. *Austral. J. Statist.*, 39:179–192, 1997.
- Y. Qi and R. J. G. Wilms. The limit behavior of maxima modulo one and the number of maxima. *Statist. Probab. Lett.*, 34:75–84, 1997.

