

On r -stacked triangulated manifolds

Satoshi Murai, Eran Nevo

► **To cite this version:**

Satoshi Murai, Eran Nevo. On r -stacked triangulated manifolds. 25th International Conference on Formal Power Series and Algebraic Combinatorics (FPSAC 2013), 2013, Paris, France. pp.181-192. hal-01229706

HAL Id: hal-01229706

<https://hal.inria.fr/hal-01229706>

Submitted on 17 Nov 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

On r -stacked triangulated manifolds

Satoshi Murai^{1†} and Eran Nevo^{2‡}

¹ *Department of Mathematical Science, Faculty of Science, Yamaguchi University, 1677-1 Yoshida, Yamaguchi 753-8512, Japan*

² *Department of Mathematics, Ben Gurion University of the Negev, Be'er Sheva 84105, Israel*

Abstract. The notion of r -stackedness for simplicial polytopes was introduced by McMullen and Walkup in 1971 as a generalization of stacked polytopes. In this paper, we define the r -stackedness for triangulated homology manifolds and study their basic properties. In addition, we find a new necessary condition for face vectors of triangulated manifolds when all the vertex links are polytopal.

Résumé. Généralisant les polytopes simpliciaux empilés, McMullen et Walkup ont introduit en 1971 la notion de r -empilement pour les polytopes simpliciaux. Dans cet article, nous définissons la notion de r -empilement pour les variétés homologiques simpliciales et étudions ses propriétés élémentaires. En outre, nous donnons une nouvelle condition pour les f -vecteurs des variétés simpliciales lorsque tous les sommets ont un lien polytopal.

Keywords: stackedness, triangulation, manifold, f -vector, face ring

1 Introduction

A triangulated d -ball is said to be r -stacked if it has no interior faces of dimension $\leq d - r - 1$, and the boundary of an r -stacked d -ball is called an r -stacked $(d-1)$ -sphere. It is known that r -stacked d -balls and $(d-1)$ -spheres with $r < \frac{d}{2}$ have many nice combinatorial properties, and they have been used to obtain several important results on polytopes and triangulated spheres. For example, they appeared in Barnette's lower bound theorem [Ba1, Ba2] and in the generalized lower bound conjecture given by McMullen and Walkup [MW]. They also appeared in the proof of the sufficiency of the famous g -theorem by Billera and Lee [BL] (see [KIL]) as well as in the construction of many non-polytopal triangulated spheres given by Kalai [Ka1]. The purpose of this paper is to extend this notion to triangulated manifolds, and establish their fundamental properties.

Throughout the paper, we fix a field \mathbf{k} . For a simplicial complex Δ and its face $F \in \Delta$, the *link of F in Δ* is the simplicial complex

$$\mathrm{lk}_{\Delta}(F) = \{G \in \Delta : F \cup G \in \Delta \text{ and } F \cap G = \emptyset\}.$$

A simplicial complex Δ of dimension d is said to be a (\mathbf{k}) -homology d -sphere if, for all faces $F \in \Delta$ (including the empty face \emptyset), one has $\beta_i(\mathrm{lk}_{\Delta}(F)) = 0$ for $i \neq d - \#F$ and $\beta_{d-\#F}(\mathrm{lk}_{\Delta}(F)) = 1$, where

[†]Partially supported by KAKENHI #22740018.

[‡]partially supported by Marie Curie grant IRG-#270923 and by ISF grant.

$\beta_i(\Delta) = \dim_{\mathbf{k}} \tilde{H}_i(\Delta; \mathbf{k})$ is the i th Betti number of Δ over \mathbf{k} . A simplicial complex is said to be *pure* if all its facets have the same dimension. A (\mathbf{k} -)homology d -manifold without boundary is a d -dimensional pure simplicial complex all whose vertex links are \mathbf{k} -homology spheres. A pure d -dimensional simplicial complex Δ is said to be a (\mathbf{k} -)homology d -manifold with boundary if it satisfies

- (i) for all $\emptyset \neq F \in \Delta$, $\beta_i(\text{lk}_{\Delta}(F))$ vanish for $i \neq d - \#F$ and is equal to 0 or 1 for $i = d - \#F$.
- (ii) the boundary $\partial\Delta = \{F \in \Delta : \beta_i(\text{lk}_{\Delta}(F)) = 0\} \cup \{\emptyset\}$ of Δ is a \mathbf{k} -homology $(d - 1)$ -manifold without boundary.

Triangulations of topological manifolds are examples of homology manifolds. Also, condition (ii) can be omitted if we replace \mathbf{k} by \mathbb{Z} (see [Mi]).

We say that a homology d -manifold Δ with boundary is *r -stacked* if it has no interior faces (namely, faces which are not in $\partial\Delta$) of dimension $\leq d - r - 1$. Also, a homology manifold without boundary is said to be *r -stacked* if it is the boundary of an r -stacked homology manifold with boundary. We prove the following properties for r -stacked homology manifolds.

- (a) *Enumerative criterion:* We give a simple criterion for the r -stackedness in terms of h -vectors and Betti numbers for homology manifolds with boundary (Theorem 3.1). Also, we give a similar result for $(r - 1)$ -stacked homology $(d - 1)$ -manifolds without boundary with $r \leq \frac{d}{2}$ when all the vertex links are polytopal (Corollary 5.5). In particular, these results prove that r -stackedness depends only on face numbers and Betti numbers for these manifolds.
- (b) *Vanishing of Betti numbers and missing faces:* We show that if a homology manifold (with or without boundary) is r -stacked, then it has zero Betti numbers and no missing faces in certain dimensions (Corollary 3.2 and Theorem 4.4).
- (c) *Uniqueness of stacked manifolds:* For an $(r - 1)$ -stacked $(d - 1)$ -manifold Δ without boundary, it is shown that if $r \leq \frac{d}{2}$ then there is a unique $(r - 1)$ -stacked homology manifold Σ such that $\partial\Sigma = \Delta$ (Theorem 4.2).
- (d) *Local criterion:* For $r < \frac{d}{2}$, we show that a homology $(d - 1)$ -manifold without boundary is $(r - 1)$ -stacked if and only if all its vertex links are $(r - 1)$ -stacked (Theorem 4.6).
- (e) *The \tilde{g} -vector — a new necessary condition for face vectors:* Motivated by a recent conjecture given by Bagchi and Datta, we define the \tilde{g} -vector of a simplicial complex Δ , and show that it is an M -vector if Δ is an $(r - 1)$ -stacked homology $(d - 1)$ -manifolds without boundary when $r \leq \frac{d}{2}$. Moreover, regardless of stackedness, we show that the same result holds for connected orientable rational homology manifolds all whose vertex links are polytopal (Theorem 5.4).

Most of the results listed above are natural extensions of known results for triangulated balls and spheres. However their proofs are not straightforward and we believe that these properties are useful in the study of face numbers of triangulated manifolds. Indeed, the results about the \tilde{g} -vector prove a refinement of [BD2, Conjecture 1.6] for orientable homology manifolds all whose vertex links are polytopal.

About (c) and (d), the same results were proved independently by Bagchi and Datta [BD3, Theorem 2.19] with essentially the same proof. Their results also prove vanishing of missing faces in (b).

This paper is organized as follows. In Section 2, we recall basic properties of h' - and h'' -vectors which play an important role in the study of face numbers of homology manifolds. In Section 3, we study r -stacked homology manifolds with boundary. In Sections 4 and 5, we study r -stacked homology manifolds without boundary and consider the \tilde{g} -vector. Some of the proofs are omitted from this extended abstract, for space limit, and can be found in the full version of this paper, at math arXiv:1209.0868.

2 h' - and h'' -vectors

In this section, we recall h' - and h'' -vectors and their algebraic meanings. We first recall some basics on simplicial complexes. A simplicial complex Δ on the vertex set V is a collection of subsets of V satisfying that $F \in \Delta$ and $G \subset F$ imply $G \in \Delta$. Elements of Δ are called *faces* of Δ and subsets of V which are not faces of Δ are called *non-faces* of Δ . The maximal faces of Δ (with respect to inclusion) are called the *facets* of Δ and the minimal non-faces of Δ are called the *missing faces* of Δ . The dimension of a face (or a missing face) F is $\#F - 1$, where $\#X$ denotes the cardinality of a finite set X , and a face (or a missing face) of dimension k is called a k -*face* (or a *missing k -face*). Also, the dimension of a simplicial complex is the maximum dimension of its faces. For a simplicial complex Δ of dimension $d - 1$, let $f_k = f_k(\Delta)$ be the number of k -faces of Δ for $k = -1, 0, \dots, d - 1$, where $f_{-1} = 1$. The vector $f(\Delta) = (f_{-1}, f_0, \dots, f_{d-1})$ is called the f -vector of Δ . Also, the h -vector $h(\Delta) = (h_0(\Delta), h_1(\Delta), \dots, h_d(\Delta))$ of Δ is defined by the relation

$$\sum_{i=0}^d h_i(\Delta)t^i = \sum_{i=0}^d f_{i-1}(\Delta)t^i(1-t)^{d-i}.$$

Now we define h' - and h'' -vectors. For a simplicial complex Δ of dimension $d - 1$, its h' -vector $h'(\Delta) = (h'_0(\Delta), \dots, h'_d(\Delta))$ and its h'' -vector $h''(\Delta) = (h''_0(\Delta), \dots, h''_d(\Delta))$ are defined by

$$h'_i(\Delta) = h_i(\Delta) - \binom{d}{i} \sum_{k=1}^{i-1} (-1)^{i-k} \beta_{k-1}(\Delta)$$

for $i = 0, 1, \dots, d$, and by

$$h''_i(\Delta) = h_i(\Delta) - \binom{d}{i} \sum_{k=1}^i (-1)^{i-k} \beta_{k-1}(\Delta) = h'_i(\Delta) - \binom{d}{i} \beta_{i-1}(\Delta)$$

for $i = 0, 1, \dots, d - 1$ and $h''_d(\Delta) = h'_d(\Delta)$. Note that

$$h''_d(\Delta) = h'_d(\Delta) = \sum_{\ell=0}^d (-1)^{\ell-d} f_{\ell-1} - \sum_{k=0}^{d-1} (-1)^{d-k} \beta_{k-1}(\Delta) = \beta_{d-1}(\Delta).$$

If one knows the Betti numbers of Δ , then knowing $h(\Delta)$ is equivalent to knowing $h'(\Delta)$ (or $h''(\Delta)$).

h' - and h'' -vectors have nice algebraic meanings in terms of Stanley–Reisner rings. Let $S = \mathbf{k}[x_1, \dots, x_n]$ be a polynomial ring over a field \mathbf{k} with $\deg x_i = 1$ for all i . For a simplicial complex Δ on $[n] = \{1, 2, \dots, n\}$, the *Stanley–Reisner ring* of Δ is the quotient ring

$$\mathbf{k}[\Delta] = S/I_\Delta$$

where $I_\Delta = (x_{i_1} \cdots x_{i_k} : \{i_1, \dots, i_k\} \notin \Delta)$. If Δ has dimension $d - 1$ and \mathbf{k} is infinite, there is a sequence $\Theta = \theta_1, \dots, \theta_d \in S_1$ of linear forms such that $\dim_{\mathbf{k}}(S/(I_\Delta + (\Theta))) < \infty$. This sequence Θ is called a *linear system of parameters* (l.s.o.p. for short) of $\mathbf{k}[\Delta]$. In the rest of this paper, we always assume that \mathbf{k} is infinite.

A simplicial complex Δ of dimension $d - 1$ is said to be *Cohen–Macaulay* (over \mathbf{k}) if, for all $F \in \Delta$, $\tilde{H}_i(\text{lk}_\Delta(F); \mathbf{k})$ vanishes for $i \neq d - 1 - \#F$. Note that any Cohen–Macaulay simplicial complex is pure. A pure simplicial complex is said to be *Buchsbaum* (over \mathbf{k}) if all its vertex links are Cohen–Macaulay. Homology manifolds are examples of Buchsbaum simplicial complexes.

Let $\mathfrak{m} = (x_1, \dots, x_n)$ be the graded maximal ideal of S . For a graded S -module N , let $F_N(t) = \sum_{i \in \mathbb{Z}} (\dim_{\mathbf{k}} N_i) t^i$ be the *Hilbert Series* of N , where N_i is the graded component of N of degree i , and let $\text{Soc}(N) = \{f \in N : \mathfrak{m}f = 0\}$ be the *socle* of N . The following results shown in [Sc, p. 137] and [NS1, Theorem 3.5] give algebraic meanings of h' - and h'' -vectors.

Lemma 2.1 *Let Δ be a Buchsbaum simplicial complex of dimension $d - 1$, $\Theta = \theta_1, \dots, \theta_d$ an l.s.o.p. of $\mathbf{k}[\Delta]$ and $R = S/(I_\Delta + (\Theta))$. Then*

$$(i) \text{ (Schenzel) } F_R(t) = h'_0(\Delta) + h'_1(\Delta)t + \cdots + h'_d(\Delta)t^d.$$

$$(ii) \text{ (Novik–Swartz) } \dim_{\mathbf{k}}(\text{Soc}(R))_i \geq \binom{d}{i} \beta_{i-1}(\Delta) \text{ for all } i. \text{ In particular, there is an ideal } N \subset \text{Soc}(R) \text{ such that } F_{R/N}(t) = h''_0(\Delta) + h''_1(\Delta)t + \cdots + h''_d(\Delta)t^d.$$

In the rest of this section, we study the relation between the vanishing of h'' -numbers and missing faces. For a homogeneous ideal $I \subset S$, let $\mu_k(I)$ be the number of elements of degree k in a minimal generating set of I , namely, $\mu_k(I) = \dim_{\mathbf{k}}(I/\mathfrak{m}I)_k$. Since missing faces of Δ correspond to the minimal generators of I_Δ , $\mu_k(I_\Delta)$ is equal to the number of missing $(k - 1)$ -faces of Δ .

Lemma 2.2 *Let $I \subset S$ be a homogeneous ideal, $w \in S_1$ a linear form and $k \geq 2$ an integer. If the multiplication $\times w : (S/I)_{k-1} \rightarrow (S/I)_k$ is injective then $\mu_k(I) = \mu_k(I + (w))$.*

Proof: It is clear that $\mu_k(I) \geq \mu_k(I + (w))$ for $k \geq 1$ even without injectivity assumption. We show $\mu_k(I) \leq \mu_k(I + (w))$. Let $\sigma_1, \dots, \sigma_t \in I$ be elements of degree k which are linearly independent in $I/\mathfrak{m}I$. What we must prove is that they are also linearly independent in $(I + (w))/\mathfrak{m}(I + (w))$.

Let $\tau = \lambda_1 \sigma_1 + \cdots + \lambda_t \sigma_t \in \mathfrak{m}(I + (w))$, where $\lambda_1, \dots, \lambda_t \in \mathbf{k}$. We claim $\tau \in \mathfrak{m}I$. Indeed, if $\tau \notin \mathfrak{m}I$ then there are $\rho' \in \mathfrak{m}I$ and $\rho'' \notin I$ such that $\tau = \rho' + w\rho''$, which implies ρ'' is in the kernel of the multiplication map $\times w : (S/I)_{k-1} \rightarrow (S/I)_k$, contradicting the assumption. \square

Lemma 2.3 *For a homogeneous ideal $I \subset S$, if $(S/I)_j = 0$ for some $j \geq 0$ then $\mu_k(I) = 0$ for $k \geq j + 1$.*

Proof: Since $(S/I)_j = 0$, we have $I_k = \mathfrak{m}_k$ for $k \geq j$. Thus $\mu_k(I) = \mu_k(\mathfrak{m}) = 0$ for $k \geq j + 1$. \square

The following statement appears in [Sc, Corollary 2.5 and Theorem 4.3].

Lemma 2.4 (Schenzel) *Let Δ be a Buchsbaum simplicial complex of dimension $d - 1$, $R = \mathbf{k}[\Delta]$, $\Theta = \theta_1, \dots, \theta_d$ an l.s.o.p. of $\mathbf{k}[\Delta]$, and let $\mathcal{K}(i)$ be the kernel of*

$$\times \theta_i : R/(\theta_1, \dots, \theta_{i-1})R \rightarrow R/(\theta_1, \dots, \theta_{i-1})R.$$

Then $\dim_{\mathbf{k}} \mathcal{K}(i)_j = \binom{i-1}{j} \beta_{j-1}(\Delta)$ for all i and j .

Proposition 2.5 *Let Δ be a Buchsbaum simplicial complex of dimension $d - 1$. If $h_r''(\Delta) = 0$ then*

- (i) $\beta_k(\Delta) = 0$ for $k \geq r$.
- (ii) Δ has no missing faces of dimension $\geq r + 1$.

Proof: Let Θ be an l.s.o.p. of $\mathbf{k}[\Delta]$. Since $h_r''(\Delta) = 0$, by Lemma 2.1(ii) all elements in $S/(I_\Delta + (\Theta))$ of degree r are contained in the socle of $S/(I_\Delta + (\Theta))$. This fact implies

$$S/(I_\Delta + (\Theta))_k = 0 \quad \text{for all } k \geq r + 1. \quad (1)$$

Then since $\dim_{\mathbf{k}}(S/(I_\Delta + (\Theta)))_k \geq \binom{d}{k}\beta_{k-1}(\Delta)$ by Lemma 2.1(ii), we have $\beta_k(\Delta) = 0$ for $k \geq r$, proving (i). Moreover, this fact and Lemmas 2.2 and 2.4 show $\mu_k(I_\Delta) = \mu_k(I_\Delta + (\Theta))$ for $k \geq r + 1$. Since $S/(I_\Delta + (\Theta))_{r+1} = 0$ by (1), the statement (ii) follows from Lemma 2.3. \square

3 Stacked manifolds with boundary

In this section, we study r -stacked manifolds with boundary. Recall that a homology d -manifold with boundary is said to be r -stacked if it has no interior faces of dimension $\leq d - r - 1$ and that a homology manifold without boundary is said to be r -stacked if it is the boundary of an r -stacked homology manifold with boundary. For a simplicial complex Δ of dimension $d - 1$, let

$$g_i(\Delta) = h_i(\Delta) - h_{i-1}(\Delta)$$

for $i = 0, 1, \dots, d$.

Enumerative criterion

It is known that a homology ball Δ is $(r - 1)$ -stacked if and only if $h_r(\Delta) = 0$. See [Mc, Proposition 2.4]. We first extend this property for stacked manifolds.

Let Δ be a homology $(d - 1)$ -manifold with boundary. Then the Dehn–Sommerville relations for homology manifolds with boundary [Gr, Corollary 2.2] say

$$g_i(\partial\Delta) = h_i(\Delta) - h_{d-i}(\Delta) + \binom{d}{i}(-1)^{d-1-i}\tilde{\chi}(\Delta), \quad (2)$$

where $\tilde{\chi}(\Delta) = \sum_{k=-1}^{d-1}(-1)^k f_k(\Delta)$ is the reduced Euler characteristic. By substituting $h_{d-i}(\Delta) = h_{d-i}''(\Delta) + \binom{d}{i} \sum_{k=1}^{d-i}(-1)^{d-i-k}\beta_{k-1}(\Delta)$ and $\tilde{\chi}(\Delta) = \sum_{k=0}^{d-1}(-1)^k \beta_k(\Delta)$ in (2), we obtain

$$g_i(\partial\Delta) = h_i(\Delta) - h_{d-i}''(\Delta) + \binom{d}{i} \sum_{k=d-i}^{d-1}(-1)^{d-1-i-k}\beta_k(\Delta). \quad (3)$$

Theorem 3.1 *Let $1 \leq r \leq d$ and let Δ be a homology $(d - 1)$ -manifold with boundary. Then Δ is $(r - 1)$ -stacked if and only if $h_r''(\Delta) = 0$.*

Proof: We first prove that Δ is $(r - 1)$ -stacked if and only if $g_i(\partial\Delta) = h_i(\Delta)$ for all $i \leq d - r$. Indeed, it is clear that Δ is $(r - 1)$ -stacked if and only if $f_i(\partial\Delta) = f_i(\Delta)$ for all $i \leq d - r - 1$. Consider the equations

$$\sum_{i=0}^d f_{i-1}(\Delta)t^i = \sum_{i=0}^d h_i(\Delta)t^i(t+1)^{d-i} \quad (4)$$

and

$$\begin{aligned} \sum_{i=0}^{d-1} f_{i-1}(\partial\Delta)t^i &= \sum_{i=0}^{d-1} h_i(\partial\Delta)t^i(t+1)^{d-i-1} \\ &= \sum_{i=0}^{d-1} h_i(\partial\Delta)\{t^i(t+1)^{d-i} - t^{i+1}(t+1)^{d-(i+1)}\}. \end{aligned} \quad (5)$$

By comparing the coefficients of the polynomials in (4) and (5), we conclude that $f_i(\partial\Delta) = f_i(\Delta)$ for $i \leq d - r - 1$ if and only if $h_i(\partial\Delta) - h_{i-1}(\partial\Delta) = h_i(\Delta)$ for all $i \leq d - r$.

We first prove the ‘if’ part. Suppose $h_r''(\Delta) = 0$. Then we have $h_k''(\Delta) = 0$ for all $k \geq r$, as $h''(\Delta)$ is an M -sequence by Lemma 2.1(ii). Also, $\beta_r(\Delta) = \cdots = \beta_{d-1}(\Delta) = 0$ by Proposition 2.5. Then the Dehn–Sommerville relation (3) shows

$$g_i(\partial\Delta) = h_i(\Delta)$$

for all $i \leq d - r$, as desired.

Next, we prove the ‘only if’ part. Suppose $g_i(\partial\Delta) = h_i(\Delta)$ for all $i \leq d - r$. The Dehn–Sommerville relations (3) imply

$$h''_{d-i}(\Delta) = -\binom{d}{i}\beta_{d-i}(\Delta) + \binom{d}{i} \sum_{k=d-i+1}^{d-1} (-1)^{d-1-i-k}\beta_k(\Delta) \quad (6)$$

for all $i \leq d - r$. We show by induction on i that $\beta_{d-i}(\Delta) = 0$ and $h''_{d-i}(\Delta) = 0$ for $i \leq d - r$: The claim is clear for $i = 0$ by (6). For $i > 0$, by induction the second summand on the right-hand side of (6) vanish. Thus $h''_{d-i}(\Delta) = -\binom{d}{i}\beta_{d-i}(\Delta)$. Since h'' -vectors and Betti numbers are non-negative we have $h''_{d-i}(\Delta) = \beta_{d-i}(\Delta) = 0$. \square

Vanishing of missing faces

If Δ is an $(r - 1)$ -stacked triangulated ball then Δ is Cohen–Macaulay and $h_r(\Delta) = 0$. These facts and Lemmas 2.2 and 2.3 say that Δ has no missing faces of dimension $\geq r$ (another proof of this fact was given in [BD3, Lemma 2.10]). Proposition 2.5 and Theorem 3.1 prove an analogue of this fact for manifolds.

Corollary 3.2 *Let Δ be an $(r - 1)$ -stacked homology manifold with boundary. Then*

- (i) $\beta_k(\Delta) = 0$ for $k \geq r$.
- (ii) Δ has no missing k -faces of dimension $\geq r + 1$.

Finally, we give a few known examples of stacked manifolds.

Example 3.3 (Kühnel–Lassmann construction [Kü, KüL]) *Let $K_{d,n}$ be the simplicial complex on $[n]$ generated by the facets*

$$\{\{i, i + 1, \dots, i + d - 1\} : i = 1, 2, \dots, n\},$$

where $i + k$ means $i + k - n$ if $i + k > n$. If $n \geq 2d - 1$ then $K_{d,n}$ is a homology manifold whose boundary triangulates either $S^1 \times S^{d-3}$ or a non-orientable S^{d-3} -bundle over S^1 depending on the parity of d [KüL]. Since the interior faces of $K_{d,n}$ are those containing one of $\{i, i + 1, \dots, i + d - 2\}$ for $i = 1, 2, \dots, n$, the simplicial complex $K_{d,n}$ is 1-stacked and has the h'' -vector $(1, n - d, 0, \dots, 0)$.

Example 3.4 (Klee–Novik construction [KN]) Let $X = \{x_1, \dots, x_d\}$ and $Y = \{y_1, \dots, y_d\}$ be disjoint sets. For integers $0 \leq i \leq d - 2$, let $B_{d,i}$ be the simplicial complex on the vertex set $X \cup Y$ generated by the facets

$$\{\{z_1, \dots, z_d\} : z_i \in \{x_i, y_i\} \text{ and } \#\{k : \{z_k, z_{k+1}\} \not\subset X \text{ and } \{z_k, z_{k+1}\} \not\subset Y\} \leq i\}.$$

The simplicial complex $B_{d,i}$ is a combinatorial manifold whose boundary triangulates $S^i \times S^{d-i-2}$ and its h'' -vector is given by $h''_k(B_{d,i}) = \binom{d}{k}$ for $k \leq i$ and $h''_{i+1}(B_{d,i}) = 0$ [KN, Proposition 5.1]. In particular, these triangulated manifolds are i -stacked by Theorem 3.1.

Remark 3.5 If Δ is an $(r - 1)$ -stacked triangulated ball then Δ has no missing r -faces. However, an $(r - 1)$ -stacked homology manifold with boundary could have missing r -faces. Indeed, the simplicial complex $K_{4,7}$ in Example 3.3 is 1-stacked but has a missing face $\{1, 4, 7\}$.

4 Stacked manifolds without boundary

In Sections 4 and 5, we study $(r - 1)$ -stacked $(d - 1)$ -manifolds without boundary with $r \leq \frac{d}{2}$. In this section, we study these manifolds from combinatorial viewpoints.

Uniqueness of stacked manifolds

A homology d -manifold Δ with boundary is said to be a (\mathbf{k}) -homology d -ball if $\tilde{H}_k(\Delta; \mathbf{k}) = 0$ for all k and $\partial\Delta$ is a (\mathbf{k}) -homology $(d - 1)$ -sphere. For a simplicial complex Δ on $[n]$, let

$$\Delta(r) = \{F \subset [n] : \text{skel}_r(F) \subset \Delta\},$$

where $\text{skel}_r(F) = \{G \subset F : \#G \leq r + 1\}$ is the r -skeleton of F . This simplicial complex can be defined algebraically. For a homogeneous ideal $I \subset S$, let $I_{\leq k}$ be the ideal generated by all elements in I of degree $\leq k$. Then it is easy to see that $(I_\Delta)_{\leq r+1} = I_{\Delta(r)}$.

For an $(r - 1)$ -stacked homology $(d - 1)$ -sphere Δ , it was shown by McMullen [Mc, Theorem 3.3] (for polytopes) and by Bagchi and Datta [BD1, Proposition 2.10] (for triangulated spheres) that an $(r - 1)$ -stacked homology d -ball Σ satisfying $\partial\Sigma = \Delta$ is unique. Moreover, the following result was shown in [BD1, Corollary 3.6] (for polytopes) and in [MN, Lemma 2.1 and Theorem 2.3] (for homology spheres) by a different approach.

Lemma 4.1 Let $1 \leq r \leq \frac{d+1}{2}$ and Δ an $(r - 1)$ -stacked homology $(d - 1)$ -sphere. If Σ is an $(r - 1)$ -stacked homology d -ball with $\partial\Sigma = \Delta$ then $\Sigma = \Delta(r - 1)$.

Proof: Observe that Σ has no missing faces of dimension $\geq r$ (see the discussion before Corollary 3.2). Then we have $I_\Sigma = (I_\Sigma)_{\leq r}$. Since Σ and Δ have the same $(d - r)$ -skeleton and $r - 1 \leq d - r$, we have $(I_\Sigma)_{\leq r} = (I_\Delta)_{\leq r}$. Hence

$$I_\Sigma = (I_\Sigma)_{\leq r} = (I_\Delta)_{\leq r} = I_{\Delta(r-1)},$$

which implies $\Sigma = \Delta(r - 1)$. □

The following is an extension of Lemma 4.1.

Theorem 4.2 Let $1 \leq r \leq \frac{d}{2}$ and Δ an $(r - 1)$ -stacked homology $(d - 1)$ -manifold without boundary. If Σ is an $(r - 1)$ -stacked homology d -manifold with $\partial\Sigma = \Delta$ then $\Sigma = \Delta(r)$.

Proof: Since Σ is $(r - 1)$ -stacked, by Corollary 3.2(ii), Σ has no missing faces of dimension $\geq r + 1$, namely, $I_\Sigma = (I_\Sigma)_{\leq r+1}$. Then the statement follows in the same way as in the proof of Lemma 4.1. \square

Remark 4.3 We cannot replace $\Delta(r)$ by $\Delta(r - 1)$ in Theorem 4.2 by the same reason as in Remark 3.5. Similarly, the statement fails when $r = \frac{d+1}{2}$.

Vanishing of missing faces

It was shown by Kalai [Ka2, Proposition 3.6] and Nagel [Na, Corollary 4.6] that if Δ is an $(r - 1)$ -stacked homology $(d - 1)$ -sphere and $r \leq \frac{d}{2}$ then Δ has no missing k -faces for $r \leq k \leq d - r$ (they write statements only for polytopes but this fact for homology spheres follows from Nagel's proof since an $(r - 1)$ -stacked homology $(d - 1)$ -sphere with $r \leq \frac{d}{2}$ has the weak Lefschetz property and satisfies $h_{r-1} = h_r$). This fact can be generalized as follows.

Theorem 4.4 Let $1 \leq r < \frac{d}{2}$ and let Δ be an $(r - 1)$ -stacked homology $(d - 1)$ -manifold without boundary. Then

- (i) $\beta_k(\Delta) = 0$ for $r \leq k \leq d - 1 - r$.
- (ii) Δ has no missing k -faces with $r + 1 \leq k \leq d - r$.

Proof: Let Σ be an $(r - 1)$ -stacked homology d -manifold with $\partial\Sigma = \Delta$. Since Σ and Δ have the same $(d - r)$ -skeleton, we have $\beta_i(\Delta) = \beta_i(\Sigma)$ for $i < d - r$ and $\mu_j(I_\Delta) = \mu_j(I_\Sigma)$ for $j \leq d - r + 1$. Then the statement follows from Corollary 3.2. \square

Remark 4.5 Lemma 4.1 and the above proof give another proof for the fact that if $r \leq \frac{d}{2}$ and if Δ is an $(r - 1)$ -stacked homology $(d - 1)$ -sphere then Δ has no missing k -faces for $r \leq k \leq d - r$.

Local criterion

Next, we discuss a local criterion of stackedness. We say that a homology d -manifold without boundary is *locally r -stacked* if all its vertex links are r -stacked. It is clear from the definition that if a homology manifold Δ is r -stacked then it is locally r -stacked. It was shown by Kalai [Ka2, Proposition 3.5] that if $r < \frac{d}{2}$ then the converse holds for the boundary of a simplicial d -polytope. This property can be extended as follows:

Theorem 4.6 Let $1 \leq r < \frac{d}{2}$. Then a homology $(d - 1)$ -manifold without boundary is $(r - 1)$ -stacked if and only if it is locally $(r - 1)$ -stacked.

Proof: The 'only if' part is obvious. The proof of the 'if' part is similar to that of [Mc, Theorem 5.3], however, for space limit, it is omitted. \square

Remark 4.7 Theorem 4.6 fails for $r = \frac{d}{2}$. Indeed, the join Δ of boundaries of two r -simplices is a $(2r - 1)$ -sphere which is not $(r - 1)$ -stacked but it is locally $(r - 1)$ -stacked. Indeed, Δ is not $(r - 1)$ -stacked since $\Delta(r - 1)$ is the power set of $[n]$. Also, Δ is locally $(r - 1)$ -stacked since, for every vertex v of Δ , $\text{lk}_\Delta(v)$ is the boundary of the join of an $(r - 1)$ -simplex and the boundary of an r -simplex.

Remark 4.8 Theorems 4.2 (for $r < \frac{d}{2}$), 4.4(ii) and 4.6 were also proved independently by Bagchi and Datta [BD3, Theorem 2.19] with essentially the same method.

5 New necessary condition for face numbers of manifolds

McMullen and Walkup [MW] conjectured that, for the boundary complex Δ of a simplicial d -polytope, one has $h_{r-1}(\Delta) \leq h_r(\Delta)$ for $r \leq \frac{d}{2}$ and if equality holds for some r then Δ is $(r-1)$ -stacked. This conjecture is called the generalized lower bound conjecture (GLBC for short). The first part of the GLBC was solved by Stanley [St1] in his proof of the necessity of the g -theorem and the second part of the GLBC was recently proved in [MN]. Recall that a connected homology d -manifold Δ without boundary is said to be *orientable* if $\beta_d(\Delta) = 1$. Motivated by the GLBC, Bagchi and Datta [BD2, Conjecture 1.6] suggested the following conjecture.

Conjecture 5.1 (GLBC for triangulated manifolds) *Let Δ be a connected triangulated $(d-1)$ -manifold without boundary. Then*

$$(i) \quad h_r(\Delta) \geq h_{r-1}(\Delta) + \binom{d+1}{r} \sum_{j=1}^r (-1)^{r-j} \beta_{j-1}(\Delta) \text{ for } r = 1, 2, \dots, \lfloor \frac{d}{2} \rfloor.$$

(ii) *if an equality holds for some $r < \frac{d}{2}$ in (i) then Δ is locally $(r-1)$ -stacked.*

Concerning part (i) of the conjecture, a similar conjecture was given by Swartz [Sw]. Moreover, it was proved by Novik and Swartz that (i) holds for all homology manifolds all whose vertex links satisfy certain algebraic property called the weak Lefschetz property. See [NS3, p. 270, Inequality (9)]. Also, the conjecture is known to be true for orientable manifolds when $r = 2$ [NS1, Theorem 5.2].

Conjecture 5.1 suggests us to study the following invariant of simplicial complexes, which we call the \tilde{g} -vector. For a simplicial complex Δ of dimension $d-1$, let

$$\tilde{g}_r(\Delta) = h_r(\Delta) - h_{r-1}(\Delta) - \binom{d+1}{r} \sum_{j=1}^r (-1)^{r-j} \beta_{j-1}(\Delta)$$

for $r = 0, 1, 2, \dots, \lfloor \frac{d}{2} \rfloor$, where $\tilde{g}_0(\Delta) = 1$, and let $\tilde{g}(\Delta) = (\tilde{g}_0(\Delta), \tilde{g}_1(\Delta), \dots, \tilde{g}_{\lfloor \frac{d}{2} \rfloor}(\Delta))$. Then Conjecture 5.1(i) asks if $\tilde{g}_k(\Delta) \geq 0$ for all k when Δ is a connected triangulated manifold without boundary. For an $(r-1)$ -stacked homology $(d-1)$ -manifold with $r \leq \frac{d}{2}$, its \tilde{g} -vector has the following simple but interesting form.

Proposition 5.2 *Let $1 \leq r \leq \frac{d}{2}$ and let Δ be an $(r-1)$ -stacked homology d -manifold with boundary. Then $\tilde{g}_i(\partial\Delta) = h''_i(\Delta)$ for $i \leq \frac{d}{2}$.*

Proof: Since Δ and $\partial\Delta$ have the same $\lfloor \frac{d}{2} \rfloor$ -skeleton, $\beta_{k-1}(\Delta) = \beta_{k-1}(\partial\Delta)$ for $k \leq \frac{d}{2}$, and, as shown in the proof of Theorem 3.1,

$$g_i(\partial\Delta) = h_i(\Delta)$$

for $i \leq \frac{d}{2}$. Subtracting $\binom{d+1}{i} \sum_{j=1}^i (-1)^{i-j} \beta_{j-1}(\Delta)$ from the above equation, we obtain the desired equation. \square

Recall that a vector $h = (h_0, h_1, \dots, h_t) \in \mathbb{Z}^{t+1}$ is said to be an M -vector if there is a standard graded \mathbf{k} -algebra A such that $h_k = \dim_{\mathbf{k}} A_k$ for $k = 0, 1, \dots, t$. Lemma 2.1(ii) shows that, in Proposition 5.2, $\tilde{g}(\partial\Delta)$ is not only a non-negative vector but also an M -vector. It is natural to ask if $\tilde{g}(\partial\Delta)$ is an M -vector for *any* homology manifold without boundary. In this section, we prove that this property as

well as Conjecture 5.1 hold for orientable homology manifolds all whose links satisfy a certain algebraic condition described below.

We say that a homology $(d-1)$ -sphere Δ on $[n]$ has the *weak Lefschetz property* (WLP for short) if there is an l.s.o.p. Θ of $\mathbf{k}[\Delta] = S/I_\Delta$ and a linear form $w \in S_1$ such that the multiplication

$$\times w : (S/(I_\Delta + (\Theta)))_{i-1} \rightarrow (S/(I_\Delta + (\Theta)))_i \quad (7)$$

is injective for $i \leq \frac{d+1}{2}$ and is surjective for $i \geq \frac{d+1}{2}$. Note that it is known that the boundary complex of a simplicial polytope has the WLP over the rationals.

The following result is due to Swartz [Sw, Theorem 4.26]

Lemma 5.3 (Swartz) *Let Δ be a connected orientable homology $(d-1)$ -manifold without boundary on the vertex set $[n]$. Suppose that all the vertex links of Δ have the WLP. Then there is an l.s.o.p. Θ of $\mathbf{k}[\Delta]$ and a linear form w such that the multiplication map*

$$\times w : (S/(I_\Delta + (\Theta)))_{i-1} \rightarrow (S/(I_\Delta + (\Theta)))_i$$

is surjective for all $i \geq \frac{d}{2} + 1$.

The main result of this section is the following.

Theorem 5.4 *With the same assumptions and notation as in Lemma 5.3, let $R = S/(I_\Delta + (\Theta))$ and $R' = R/wR$. Then*

- (i) *there is an ideal $J \subset R'$ such that $\dim_{\mathbf{k}}(R'/J)_i = \tilde{g}_i(\Delta)$ for $i \leq \frac{d}{2}$. In particular, $\tilde{g}(\Delta)$ is an M -vector.*
- (ii) *if $\tilde{g}_r(\Delta) = 0$ for some $r < \frac{d}{2}$ then Δ is locally $(r-1)$ -stacked.*

Theorem 5.4(i) extends the result of Novik and Swartz [NS3] who proved the non-negativity of \tilde{g} -vectors for homology manifolds all whose vertex links have the WLP, and Theorem 5.4(ii) proves that Conjecture 5.1(ii) holds for these manifolds. In particular, Conjecture 5.1 holds for any rational orientable homology manifold all whose vertex links are polytopal, namely, are the boundary complexes of simplicial polytopes. It was conjectured that any homology sphere has the WLP. Thus, if this conjecture is true then Conjecture 5.1 holds for all orientable homology manifolds.

The proof of Theorem 5.4, for space limit, is omitted.

The local criterion for stackedness and Theorem 5.4 imply the following criterion for stackedness.

Corollary 5.5 *Let $r < \frac{d}{2}$ and let Δ be a connected orientable homology $(d-1)$ -manifold without boundary. If all the vertex links of Δ have the WLP then Δ is $(r-1)$ -stacked if and only if $\tilde{g}_r(\Delta) = 0$.*

Proof: The ‘if’ part follows from Theorems 4.6 and 5.4. The ‘only if’ part follows from Theorem 3.1 and Proposition 5.2. \square

We end this paper by a few questions.

Conjecture 5.6 *With the same assumptions and notation as in Theorem 5.4, $\dim_{\mathbf{k}}(\text{Soc } R')_r \geq \binom{d+1}{r} \beta_{r-1}(\Delta)$ for $r \leq \frac{d}{2}$.*

If the conjecture is true, it will give a necessary condition for h -vectors of triangulated manifolds stronger than Theorem 5.4(i). Indeed, Conjecture 5.6 implies Theorem 5.4(i) since [NS2, Theorem 3.2] implies

$$\dim_{\mathbf{k}} R'_r = \dim_{\mathbf{k}} R_r - \dim_{\mathbf{k}}(R/\text{Soc}(R))_{r-1} = h'_r - h''_{r-1} = \tilde{g}_r + \binom{d+1}{r} \beta_{r-1}$$

for $i \leq \frac{d}{2}$. For $(r-1)$ -stacked homology $(d-1)$ -manifolds without boundary with $r \leq \frac{d}{2}$, the conjecture follows from Lemma 2.1(ii) by taking (Θ, w) for Δ to be a general l.s.o.p. of $\mathbf{k}[\Sigma]$, where Σ is the $(r-1)$ -stacked homology manifold with $\partial\Sigma = \Delta$. The conjecture also holds for triangulations of the product of spheres (under the WLP assumption) since the ideal J in Theorem 5.4 is concentrated in a single degree in this case.

Question 5.7 *Is it true that if Δ is a homology $(2k-1)$ -manifold without boundary such that $\tilde{g}_k(\Delta) = 0$ then Δ is $(k-1)$ -stacked?*

A similar question was raised by Novik–Swartz [NS1, Problem 5.3] when $k = 2$. However, we do not have an answer even for this case.

Acknowledgements

We would like to thank Isabella Novik for helpful comments on an earlier version of the full paper version of this extended abstract, as well as the referees.

References

- [Ba1] D.W. Barnette, The minimum number of vertices of a simple polytope, *Israel J. Math.* **10** (1971), 121–125.
- [Ba2] D.W. Barnette, A proof of the lower bound conjecture for convex polytopes, *Pac. J. Math.* **46** (1973) 349–354.
- [BD1] B. Bagchi and B. Datta, On stellated spheres, shellable balls, lower bounds and a combinatorial criterion for tightness, arXiv:1102.0856.
- [BD2] B. Bagchi and B. Datta, On stellated spheres and a tightness criterion for combinatorial manifolds, arXiv:1207.5599.
- [BD3] B. Bagchi and B. Datta, On k -stellated and k -stacked spheres, arXiv:1208.1389.
- [BL] L. Billera and C.W. Lee, A proof of the sufficiency of McMullen’s conditions for f -vectors of simplicial convex polytopes, *J. Combin. Theory Ser. A* **31** (1981), 237–255.
- [BH] W. Bruns and J. Herzog, Cohen–Macaulay rings, Revised Edition, Cambridge University Press, Cambridge, 1996.
- [Gr] H.-G. Gräbe, Generalized Dehn–Sommerville equations and an upper bound theorem, *Beitrage Algebra Geom.* **25** (1987), 47–60.

- [Ka1] G. Kalai, Many triangulated spheres, *Discrete Comput. Geom.* **3** (1988), 1–14.
- [Ka2] G. Kalai, Some aspects of the combinatorial theory of convex polytopes, in: T. Bisztriczky (ed.) et al., *Polytopes: Abstract, Convex, and Computational*, NATO ASI Ser., Ser. C, Math. Phys. Sci. 440, 1994, pp. 205–229.
- [KIL] P. Kleinschmidt and C.W. Lee, On k -stacked polytopes, *Discrete Math.* **48** (1984), 125–127.
- [Kü] W. Kühnel, Higher-dimensional analogues of Czászár’s torus, *Results. Math.* **9** (1986), 95–106.
- [KüL] W. Kühnel and G. Lassmann, Permuted difference cycles and triangulated sphere bundles, *Discrete Math.* **162** (1996), 215–227.
- [KN] S. Klee and I. Novik, Centrally symmetric manifolds with few vertices, *Adv. Math.* **229** (2012), 487–500.
- [Mc] P. McMullen, Triangulations of simplicial polytopes, *Beiträge Algebra Geom.* **45** (2004), 37–46.
- [MW] P. McMullen and D.W. Walkup, A generalized lower-bound conjecture for simplicial polytopes, *Mathematika* **18** (1971), 264–273.
- [Mi] W. J. R. Mitchel, Defining the boundary of a homology manifold, *Proc. Amer. Math. Soc.*, **110** (1990), 509–513.
- [MN] S. Murai and E. Nevo, On the generalized lower bound conjecture for polytopes and spheres, to appear in *Acta Math.*, arXiv:1203.1720.
- [Na] U. Nagel, Empty simplices of polytopes and graded Betti numbers, *Discrete Comput. Geom.*, **39** (2008) 389–410.
- [No] I. Novik, Upper bound theorems for homology manifolds, *Israel J. Math.* **108** (1998), 45–82.
- [NS1] I. Novik and E. Swartz, Socles of Buchsbaum modules, complexes and posets, *Adv. Math.* **222** (2009), 2059–2084.
- [NS2] I. Novik and E. Swartz, Gorenstein rings through face rings of manifolds, *Compos. Math.* **145** (2009), 993–1000.
- [NS3] I. Novik and E. Swartz, Applications of Klee’s Dehn-Sommerville relations, *Discrete Comput. Geom.* **42** (2009), 261–276.
- [Sc] P. Schenzel, On the number of faces of simplicial complexes and the purity of Frobenius, *Math. Z.* **178** (1981), 125–142.
- [St1] R.P. Stanley, The number of faces of a simplicial convex polytope, *Adv. in Math.* **35** (1980), 236–238.
- [St2] R.P. Stanley, *Combinatorics and commutative algebra*, Second edition, Progr. Math., vol. 41, Birkhäuser, Boston, 1996.
- [Sw] E. Swartz, Face enumeration: from spheres to manifolds, *J. Eur. Math. Soc.* **11** (2009), 449–485.