

A uniform model for Kirillov–Reshetikhin crystals

Cristian Lenart, Satoshi Naito, Daisuke Sagaki, Anne Schilling, Mark Shimozono

► **To cite this version:**

Cristian Lenart, Satoshi Naito, Daisuke Sagaki, Anne Schilling, Mark Shimozono. A uniform model for Kirillov–Reshetikhin crystals. Alain Goupil and Gilles Schaeffer. 25th International Conference on Formal Power Series and Algebraic Combinatorics (FPSAC 2013), 2013, Paris, France. Discrete Mathematics and Theoretical Computer Science, AS, pp.25-36, 2013, DMTCS Proceedings. <hal-01229719>

HAL Id: hal-01229719

<https://hal.inria.fr/hal-01229719>

Submitted on 17 Nov 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A uniform model for Kirillov–Reshetikhin crystals

Cristian Lenart^{1†} Satoshi Naito^{2‡} Daisuke Sagaki^{3§}
 Anne Schilling^{4¶} Mark Shimozono^{5||}

¹*Department of Mathematics and Statistics, State University of New York at Albany, Albany, NY 12222, U.S.A.*

²*Department of Mathematics, Tokyo Institute of Technology, 2-12-1 Oh-Okayama, Meguro-ku, Tokyo 152-8551, Japan*

³*Institute of Mathematics, University of Tsukuba, Tsukuba, Ibaraki 305-8571, Japan*

⁴*Department of Mathematics, University of California, One Shields Avenue, Davis, CA 95616-8633, U.S.A.*

⁵*Department of Mathematics, Virginia Polytechnic Institute and State University, Blacksburg, VA 24061-0123, U.S.A.*

Abstract. We present a uniform construction of tensor products of one-column Kirillov–Reshetikhin (KR) crystals in all untwisted affine types, which uses a generalization of the Lakshmibai–Seshadri paths (in the theory of the Littelmann path model). This generalization is based on the graph on parabolic cosets of a Weyl group known as the parabolic quantum Bruhat graph. A related model is the so-called quantum alcove model. The proof is based on two lifts of the parabolic quantum Bruhat graph: to the Bruhat order on the affine Weyl group and to Littelmann’s poset on level-zero weights. Our construction leads to a simple calculation of the energy function. It also implies the equality between a Macdonald polynomial specialized at $t = 0$ and the graded character of a tensor product of KR modules.

Résumé. Nous présentons une construction uniforme pour les produits tensoriels des cristaux de Kirillov–Reshetikhin de type colonne, pour tous les types affines symétriques, qui utilise une généralisation des chemins de Lakshmibai–Seshadri (dans la théorie des chemins de Littelmann). Cette généralisation est basée sur un graphe sur les classes parabolique d’un groupe de Weyl appelé le graphe de Bruhat parabolique quantique. Un modèle lié est le modèle des alcôves quantique. La preuve est basée sur deux relèvements du graphe de Bruhat parabolique quantique: dans l’ordre de Bruhat affine et dans un ensemble ordonné des poids de niveau zero. Notre construction donne une formule simple pour la fonction d’énergie. Elle donne aussi l’égalité d’un polynôme de Macdonald spécialisé à $t = 0$ avec le caractère gradué d’un produit tensoriel des modules de Kirillov–Reshetikhin.

Keywords: Parabolic quantum Bruhat graph, Kirillov–Reshetikhin crystals, energy function, Lakshmibai–Seshadri paths, Macdonald polynomials.

[†]Partially supported by the Research in Pairs program of Mathematisches Forschungsinstitut Oberwolfach and the NSF grant DMS-1101264.

[‡]Partially supported by the Research in Pairs program of Mathematisches Forschungsinstitut Oberwolfach and Grant-in-Aid for Scientific Research (C), No. 24540010, Japan.

[§]Partially supported by Grant-in-Aid for Young Scientists (B) No.23740003, Japan.

[¶]Partially supported by the Research in Pairs program of Mathematisches Forschungsinstitut Oberwolfach and the NSF grants DMS-1001256 and OCI-1147247.

^{||}Partially supported by the Research in Pairs program of Mathematisches Forschungsinstitut Oberwolfach and the NSF grant DMS-1200804.

1 Introduction

Our goal in this series of papers (see [LNSSS1, LNSSS2]) is to obtain a uniform construction of tensor products of one-column Kirillov–Reshetikhin (KR) crystals. As a consequence we shall prove the equality $P_\lambda(q) = X_\lambda(q)$, where $P_\lambda(q)$ is the Macdonald polynomial $P_\lambda(q, t)$ specialized at $t = 0$ and $X_\lambda(q)$ is the graded character of a simple Lie algebra coming from tensor products of KR modules. Both the Macdonald polynomials and KR modules are of arbitrary untwisted affine type. The index λ is a dominant weight for the simple Lie subalgebra obtained by removing the affine node. Macdonald polynomials and characters of KR modules have been studied extensively in connection with various fields such as statistical mechanics and integrable systems, representation theory of Coxeter groups and Lie algebras (and their quantized analogues given by Hecke algebras and quantized universal enveloping algebras), geometry of singularities of Schubert varieties, and combinatorics.

Our point of departure is a theorem of Ion [Ion], which asserts that the nonsymmetric Macdonald polynomials at $t = 0$ are characters of Demazure submodules of highest weight modules over affine algebras. This holds for the Langlands duals of untwisted affine root systems (and type $A_{2n}^{(2)}$ in the case of nonsymmetric Koornwinder polynomials). Our results apply to the untwisted affine root systems. The overlapping cases are the simply-laced affine root systems $A_n^{(1)}$, $D_n^{(1)}$ and $E_{6,7,8}^{(1)}$.

It is known [FL, FSS, KMOU, KMOTU, ST, Na] that certain affine Demazure characters (including those for the simply-laced affine root systems) can be expressed in terms of KR crystals, which motivates the relation between P and X . For types $A_n^{(1)}$ and $C_n^{(1)}$, the equality $P = X$ was achieved in [Le2, LeS] by establishing a combinatorial formula for the Macdonald polynomials at $t = 0$ from the Ram–Yip formula [RY], and by using explicit models for the one-column KR crystals [FOS]. It should be noted that, in types $A_n^{(1)}$ and $C_n^{(1)}$, the one-column KR modules are irreducible when restricted to the canonical simple Lie subalgebra, while in general this is not the case. For the cases considered by Ion [Ion], the corresponding KR crystals are perfect. This is not necessarily true for the untwisted affine root systems considered in this work, especially for the untwisted non-simply-laced affine root systems.

In this work we provide a type-free approach to the equality $P = X$ for untwisted affine root systems. Lenart’s specialization [Le2] of the Ram–Yip formula for Macdonald polynomials uses the quantum alcove model [LeL1], whose objects are paths in the quantum Bruhat graph (QBG), which was defined and studied in [BFP] in relation to the quantum cohomology of the flag variety. On the other hand, Naito and Sagaki [NS1, NS2, NS4, NS5] gave models for tensor products of KR crystals of one-column type in terms of projections of level-zero Lakshmibai–Seshadri (LS) paths to the classical weight lattice. Hence we need to establish a bijection between the quantum alcove model and projected level-zero LS paths.

In analogy with [BFP] and inspired by the quantum Schubert calculus of homogeneous spaces [Mi, P] we define the parabolic quantum Bruhat graph (PQBG), which is a directed graph structure on parabolic quotients of the Weyl group with respect to a parabolic subgroup. We construct two lifts of the PQBG. The *first lift* is from the PQBG to the Bruhat order of the affine Weyl group. This is a parabolic analogue of the lift of the QBG to the affine Bruhat order [LS], which is the combinatorial structure underlying Peterson’s theorem [P]; the latter equates the Gromov–Witten invariants of finite-dimensional homogeneous spaces with the Pontryagin homology structure constants of Schubert varieties in the affine Grassmannian. We obtain Diamond Lemmas for the PQBG via projection of the standard Diamond Lemmas for the affine Weyl group. We find a *second lift* of the PQBG into a poset of Littelmann [Li] for level-zero weights and characterize its local structure (such as cover relations) in terms of the PQBG. Littelmann’s poset was defined in connection with LS paths for arbitrary (not necessarily dominant) weights, but the local

structure was not previously known. The weight poset precisely controls the combinatorics of the level-zero LS paths and therefore their classical projections, which we formulate directly as quantum LS paths. Finally, we describe a bijection between the quantum alcove model and the quantum LS paths.

The paper is organized as follows. In Section 2 we prepare the background and define the PQBG. Section 3 is reserved for the two lifts of the PQBG and the statement of the Diamond Lemmas. In Section 4 we describe KR crystals in terms of the quantum LS paths and the quantum alcove model in [LeL1]; we also give simple combinatorial formulas for the energy function. Finally, we conclude in Section 5 with the results on (nonsymmetric) Macdonald polynomials at $t = 0$.

Acknowledgments

We would like to thank Thomas Lam for helpful discussions during FPSAC 2012, and Daniel Orr for his discussions about Ion’s work [Ion]. We used SAGE [Sa] and SAGE-COMBINAT [Sa-comb] to discover properties about the level-zero weight poset and to obtain some of the pictures in [LNSSS1].

2 Background

2.1 Untwisted affine root datum

Let $I_{\text{af}} = I \sqcup \{0\}$ (resp. I) be the Dynkin node set of an untwisted affine algebra \mathfrak{g}_{af} (resp. its canonical subalgebra \mathfrak{g}), W_{af} (resp. W) the affine (resp. finite) Weyl group with simple reflections r_i for $i \in I_{\text{af}}$ (resp. $i \in I$), and $X_{\text{af}} = \mathbb{Z}\delta \oplus \bigoplus_{i \in I_{\text{af}}} \mathbb{Z}\Lambda_i$ (resp. $X = \bigoplus_{i \in I} \mathbb{Z}\omega_i$) the affine (resp. finite) weight lattice. Let $\{\alpha_i \mid i \in I_{\text{af}}\}$ be the simple roots, $\Phi^{\text{af}} = W_{\text{af}}\{\alpha_i \mid i \in I_{\text{af}}\}$ (resp. $\Phi = W\{\alpha_i \mid i \in I\}$) the set of affine real roots (resp. roots), and $\Phi^{\text{af}+} = \Phi^{\text{af}} \cap \bigoplus_{i \in I_{\text{af}}} \mathbb{Z}_{\geq 0}\alpha_i$ (resp. $\Phi^+ = \Phi \cap \bigoplus_{i \in I} \mathbb{Z}_{\geq 0}\alpha_i$) the set of positive affine real (resp. positive) roots. Furthermore, $\Phi^{\text{af}-} = -\Phi^{\text{af}+}$ (resp. $\Phi^- = -\Phi^+$) are the negative affine real (resp. negative) roots. Let $X_{\text{af}}^{\vee} = \text{Hom}_{\mathbb{Z}}(X_{\text{af}}, \mathbb{Z})$ be the dual lattice, $\langle \cdot, \cdot \rangle : X_{\text{af}}^{\vee} \times X_{\text{af}} \rightarrow \mathbb{Z}$ the evaluation pairing, and $\{d\} \cup \{\alpha_i^{\vee} \mid i \in I_{\text{af}}\}$ the dual basis of X_{af}^{\vee} . The natural projection $\text{cl} : X_{\text{af}} \rightarrow X$ has kernel $\mathbb{Z}\Lambda_0 \oplus \mathbb{Z}\delta$ and sends $\Lambda_i \mapsto \omega_i$ for $i \in I$.

The affine Weyl group W_{af} acts on X_{af} and X_{af}^{\vee} by

$$r_i\lambda = \lambda - \langle \alpha_i^{\vee}, \lambda \rangle \alpha_i \quad \text{and} \quad r_i\mu = \mu - \langle \mu, \alpha_i \rangle \alpha_i^{\vee},$$

for $i \in I_{\text{af}}$, $\lambda \in X_{\text{af}}$, and $\mu \in X_{\text{af}}^{\vee}$. For $\beta \in \Phi^{\text{af}}$, let $w \in W_{\text{af}}$ and $i \in I_{\text{af}}$ be such that $\beta = w\alpha_i$. Define the associated reflection $r_{\beta} \in W_{\text{af}}$ and associated coroot $\beta^{\vee} \in X_{\text{af}}^{\vee}$ by $r_{\beta} = wr_iw^{-1}$ and $\beta^{\vee} = w\alpha_i^{\vee}$.

The null root is the unique element $\delta \in \bigoplus_{i \in I_{\text{af}}} \mathbb{Z}_{>0}\alpha_i$ which generates the rank 1 sublattice $\{\lambda \in X_{\text{af}} \mid \langle \alpha_i^{\vee}, \lambda \rangle = 0 \text{ for all } i \in I_{\text{af}}\}$. We have $\delta = \alpha_0 + \theta$, where θ is the highest root for \mathfrak{g} . The canonical central element is the unique element $c \in \bigoplus_{i \in I_{\text{af}}} \mathbb{Z}_{>0}\alpha_i^{\vee}$ which generates the rank 1 sublattice $\{\mu \in X_{\text{af}}^{\vee} \mid \langle \mu, \alpha_i \rangle = 0 \text{ for all } i \in I_{\text{af}}\}$. The level of a weight $\lambda \in X_{\text{af}}$ is defined by $\text{level}(\lambda) = \langle c, \lambda \rangle$. Let $X_{\text{af}}^0 \subset X_{\text{af}}$ be the sublattice of level-zero elements.

We denote by $\ell(w)$ for $w \in W_{\text{af}}$ (resp. W) the length of w and by \leq the Bruhat cover. The element $t_{\mu} \in W_{\text{af}}$ is the translation by the element μ in the coroot lattice $Q^{\vee} = \bigoplus_{i \in I} \mathbb{Z}\alpha_i^{\vee}$.

2.2 Affinization of a weight stabilizer

Let $\lambda \in X$ be a dominant weight, which is fixed throughout the remainder of Sections 2 and 3. Let W_J be the stabilizer of λ in W . It is a parabolic subgroup, being generated by r_i for $i \in J$, where

$J = \{i \in I \mid \langle \alpha_i^\vee, \lambda \rangle = 0\}$. Let $Q_J^\vee = \bigoplus_{i \in J} \mathbb{Z}\alpha_i^\vee$ be the associated coroot lattice, W^J the set of minimum-length coset representatives in W/W_J , $\Phi_J \supset \Phi_J^+$ the set of roots and positive roots respectively, and $\rho_J = (1/2) \sum_{\alpha \in \Phi_J^+} \alpha$ (if $J = \emptyset$, then ρ_J is denoted by ρ). Define

$$(W_J)_{\text{af}} = W_J \times Q_J^\vee = \{wt_\mu \in W_{\text{af}} \mid w \in W_J, \mu \in Q_J^\vee\}, \quad (1)$$

$$\Phi_J^{\text{af}+} = \{\beta \in \Phi^{\text{af}+} \mid \text{cl}(\beta) \in \Phi_J\} = \Phi_J^+ \cup (\mathbb{Z}_{>0}\delta + \Phi_J), \quad (2)$$

$$(W^J)_{\text{af}} = \{x \in W_{\text{af}} \mid x \cdot \beta > 0 \text{ for all } \beta \in \Phi_J^{\text{af}+}\}. \quad (3)$$

By [LS, Lemma 10.5] [P], any $w \in W_{\text{af}}$ factors uniquely as $w = w_1 w_2$, where $w_1 \in (W^J)_{\text{af}}$ and $w_2 \in (W_J)_{\text{af}}$. Therefore, we can define $\pi_J : W_{\text{af}} \rightarrow (W^J)_{\text{af}}$ by $w \mapsto w_1$. We say that $\mu \in Q^\vee$ is *J-adjusted* if $\pi_J(t_\mu) = z_\mu t_\mu$ with $z_\mu \in W$. Say that $\mu \in Q^\vee$ is *J-superantidominant* if μ is antidominant (i.e., $\langle \mu, \alpha \rangle \leq 0$ for all $\alpha \in \Phi^+$), and $\langle \mu, \alpha \rangle \ll 0$ for $\alpha \in \Phi^+ \setminus \Phi_J^+$.

2.3 The parabolic quantum Bruhat graph

The *parabolic quantum Bruhat graph* $\text{QB}(W^J)$ is a directed graph with vertex set W^J , whose directed edges have the form $w \xrightarrow{\alpha} [wr_\alpha]$ for $w \in W^J$ and $\alpha \in \Phi^+ \setminus \Phi_J^+$. Here we denote by $[v]$ the minimum-length representative in the coset vW_J for $v \in W$. There are two kinds of edges:

1. (Bruhat edge) $w \prec wr_\alpha$. (One may deduce that $wr_\alpha \in W^J$.)
2. (Quantum edge) $\ell([wr_\alpha]) = \ell(w) + 1 - \langle \alpha^\vee, 2\rho - 2\rho_J \rangle$.

If $J = \emptyset$, then we recover the quantum Bruhat graph $\text{QB}(W)$ defined in [BFP].

3 Two lifts of the parabolic quantum Bruhat graph

3.1 Lifting $\text{QB}(W^J)$ to W_{af}

We construct a parabolic analogue of the lift of $\text{QB}(W)$ to W_{af} given in [LS].

Let $\Omega_J^\infty \subset W_{\text{af}}$ be the subset of elements of the form $w\pi_J(t_\mu)$ with $w \in W^J$ and $\mu \in Q^\vee$ *J-superantidominant* and *J-adjusted*. We have $\Omega_J^\infty \subset (W^J)_{\text{af}} \cap W_{\text{af}}^-$, where W_{af}^- is the set of minimum-length coset representatives in W_{af}/W . Impose the Bruhat covers in Ω_J^∞ whenever the connecting root has classical part in $\Phi \setminus \Phi_J$. Then Ω_J^∞ is a subposet of the Bruhat poset W_{af} .

Proposition 3.1 *Every edge in $\text{QB}(W^J)$ lifts to a downward Bruhat cover in Ω_J^∞ , and every cover in Ω_J^∞ projects to an edge in $\text{QB}(W^J)$. More precisely:*

1. For any edge $[wr_\alpha] \xleftarrow{\alpha} w$ in $\text{QB}(W^J)$, and $\mu \in Q^\vee$ that is *J-superantidominant* and *J-adjusted* with $\pi_J(t_\mu) = zt_\mu$, there is a covering relation $y \prec x$ in Ω_J^∞ where

$$x = wzt_\mu, \quad y = xr_{\tilde{\alpha}} = wr_\alpha t_{\chi\alpha^\vee} zt_\mu, \quad \tilde{\alpha} = z^{-1}\alpha + (\chi + \langle \mu, z^{-1}\alpha \rangle)\delta \in \Phi^{\text{af}-},$$

and χ is 0 or 1 according as the arrow in $\text{QB}(W^J)$ is of Bruhat or quantum type respectively.

2. Suppose $y \triangleleft x$ is an arbitrary covering relation in Ω_J^∞ . Then we can write $x = wzt_\mu$ with $w \in W^J$, $z = z_\mu \in W_J$, and $\mu \in Q^\vee$ J -superantidominant and J -adjusted, as well as $y = xr_\gamma$ with $\gamma = z^{-1}\alpha + n\delta \in \Phi^{\text{af}}$, $\alpha \in \Phi^+ \setminus \Phi_J^+$, and $n \in \mathbb{Z}$. With the notation $\chi := n - \langle \mu, z^{-1}\alpha \rangle$, we have

$$\chi \in \{0, 1\}, \quad \gamma = z^{-1}\alpha + (\chi + \langle \mu, z^{-1}\alpha \rangle)\delta \in \Phi^{\text{af}^-};$$


furthermore, there is an edge $wr_\alpha z \xleftarrow{z^{-1}\alpha} wz$ in $\text{QB}(W)$ and an edge $[wr_\alpha] \xleftarrow{\alpha} w$ in $\text{QB}(W^J)$, where both edges are of Bruhat type if $\chi = 0$ and of quantum type if $\chi = 1$.

3.2 The Diamond Lemmas


In the following, a dotted (resp. plain) edge represents a quantum (resp. Bruhat) edge in $\text{QB}(W^J)$, whereas a dashed edge can be of both types. Given $w \in W^J$ and $\gamma \in \Phi^+$, define $z \in W_J$ by $r_\theta w = [r_\theta w]z$. We now state the Diamond Lemmas for $\text{QB}(W^J)$. They are proved based on the lift of $\text{QB}(W^J)$ to W_{af} in Proposition 3.1 and the fact that such a lemma holds for any Coxeter group [BB].

Lemma 3.2 *Let $\alpha \in \Phi$ be a simple root, $\gamma \in \Phi^+ \setminus \Phi_J^+$, and $w \in W^J$. Then we have the following cases, in each of which the bottom two edges imply the top two edges, and vice versa.*

1. In the left diagram we assume $\gamma \neq w^{-1}(\alpha)$. In both cases we have $r_\alpha [wr_\gamma] = [r_\alpha wr_\gamma]$.



2. Here z is defined as above. We assume $\gamma \neq -w^{-1}(\theta)$ whenever both of the hypothesized edges are quantum ones. In the left diagram, the dashed edge is a quantum (resp. a Bruhat) edge depending on $\langle \gamma^\vee, w^{-1}(\theta) \rangle$ being nonzero (resp. zero). In the right diagram, the dashed edge is a Bruhat (resp. a quantum) edge depending on $\langle \gamma^\vee, w^{-1}(\theta) \rangle$ being nonzero (resp. zero).



3.3 Lifting $\text{QB}(W^J)$ to the level-zero weight poset

In [Li], Littelmann introduced a poset related to LS paths for arbitrary (not necessarily dominant) integral weights. Littelmann did not give a precise local description of it. We consider this poset for level-zero weights and characterize its cover relations in terms of the PQBG.

Let $\lambda \in X$ be a fixed dominant weight (cf. Section 2.2 and the notation thereof, e.g., W_J is the stabilizer of λ). We view X as a sublattice of X_{af}^0 . Let $X_{\text{af}}^0(\lambda)$ be the orbit $W_{\text{af}}\lambda$.

Definition 3.3 (Level-zero weight poset [Li]) *A poset structure is defined on $X_{\text{af}}^0(\lambda)$ as the transitive closure of the relation*

$$\mu < r_\beta(\mu) \iff \langle \beta^\vee, \mu \rangle > 0,$$

where $\beta \in \Phi^{\text{af}+}$. This poset is called the level-zero weight poset for λ .

The cover $\mu < \nu = r_\beta(\mu)$ of $X_{\text{af}}^0(\lambda)$ is labeled by the root $\beta \in \Phi^{\text{af}+}$. The projection map cl restricts to the map $\text{cl} : X_{\text{af}}^0(\lambda) \rightarrow W\lambda$. We identify $W\lambda \simeq W/W_J \simeq W^J$, and consider $\text{QB}(W^J)$. Our main result is the construction of a lift of $\text{QB}(W^J)$ to $X_{\text{af}}^0(\lambda)$. The proof is based on Lemma 3.2.

Theorem 3.4 *Let $\mu \in X_{\text{af}}^0(\lambda)$ and $w := \text{cl}(\mu) \in W^J$. If $\mu < \nu$ is a cover in $X_{\text{af}}^0(\lambda)$, then its label β is in Φ^+ or $\delta - \Phi^+$. Moreover, $w \rightarrow \text{cl}(\nu)$ is an up (respectively down) edge in $\text{QB}(W^J)$ labeled by $w^{-1}(\beta) \in \Phi^+ \setminus \Phi_J^+$ (respectively $w^{-1}(\beta - \delta)$), depending on $\beta \in \Phi^+$ (respectively $\beta \in \delta - \Phi^+$). Conversely, if $w \xrightarrow{\gamma} wr_\gamma = w'$ (respectively $w \xrightarrow{\gamma} [wr_\gamma] = w'$) in $\text{QB}(W^J)$ for $\gamma \in \Phi^+ \setminus \Phi_J^+$, then there exists a cover $\mu < \nu$ in $X_{\text{af}}^0(\lambda)$ labeled by $w(\gamma)$ (respectively $\delta + w(\gamma)$) with $\text{cl}(\nu) = w'$.*

4 Models for KR crystals and the energy function

4.1 Quantum LS paths

Throughout this section, we fix a dominant integral weight $\lambda \in X$, and set $J := \{i \in I \mid \langle \alpha_i^\vee, \lambda \rangle = 0\}$.

Definition 4.1 *Let $x, y \in W^J$, and let $\sigma \in \mathbb{Q}$ be such that $0 < \sigma < 1$. A directed σ -path from y to x is, by definition, a directed path*

$$x = w_0 \xrightarrow{\gamma^1} w_1 \xrightarrow{\gamma^2} w_2 \xrightarrow{\gamma^3} \dots \xrightarrow{\gamma^n} w_n = y$$

from y to x in $\text{QB}(W^J)$ such that $\sigma \langle \gamma_k^\vee, \lambda \rangle \in \mathbb{Z}$ for all $1 \leq k \leq n$.

A quantum LS path of shape λ is a pair $\eta = (\underline{x}; \underline{\sigma})$ of a sequence $\underline{x} : x_1, x_2, \dots, x_s$ of elements in W^J with $x_u \neq x_{u+1}$ for $1 \leq u \leq s-1$ and a sequence $\underline{\sigma} : 0 = \sigma_0 < \sigma_1 < \dots < \sigma_s = 1$ of rational numbers such that there exists a directed σ_u -path from x_{u+1} to x_u for each $1 \leq u \leq s-1$. Denote by $\text{QLS}(\lambda)$ the set of quantum LS paths of shape λ . We identify an element $\eta = (x_1, x_2, \dots, x_s; \sigma_0, \sigma_1, \dots, \sigma_s) \in \text{QLS}(\lambda)$ with the following piecewise linear, continuous map $\eta : [0, 1] \rightarrow \mathbb{R} \otimes_{\mathbb{Z}} X$:

$$\eta(t) = \sum_{u'=1}^{u-1} (\sigma_{u'} - \sigma_{u'-1}) x_{u'} \cdot \lambda + (t - \sigma_{u-1}) x_u \cdot \lambda \quad \text{for } \sigma_{u-1} \leq t \leq \sigma_u, 1 \leq u \leq s,$$

and set $\text{wt}(\eta) =: \eta(1)$. Following [Li], we define the root operators e_i and f_i for $i \in I_{\text{af}} = I \sqcup \{0\}$ as follows. For $\eta \in \text{QLS}(\lambda)$ and $i \in I_{\text{af}}$, we set

$$H(t) = H_i^\eta(t) := \langle \alpha_i^\vee, \eta(t) \rangle \quad \text{for } t \in [0, 1],$$

$$m = m_i^\eta := \min\{H_i^\eta(t) \mid t \in [0, 1]\};$$

in fact, $m \in \mathbb{Z}_{\leq 0}$. If $m = 0$, then $e_i \eta := \mathbf{0}$. If $m \leq -1$, then we define $e_i \eta$ by:

$$(e_i \eta)(t) = \begin{cases} \eta(t) & \text{if } 0 \leq t \leq t_0, \\ r_{i, m+1}(\eta(t)) = \eta(t_0) + s_i(\eta(t) - \eta(t_0)) & \text{if } t_0 \leq t \leq t_1, \\ r_{i, m+1} r_{i, m}(\eta(t)) = \eta(t) + \tilde{\alpha}_i & \text{if } t_1 \leq t \leq 1, \end{cases}$$

where

$$t_1 := \min\{t \in [0, 1] \mid H(t) = m\},$$

$$t_0 := \max\{t \in [0, t_1] \mid H(t) = m + 1\},$$

$r_{i, n}$ is the reflection with respect to the hyperplane $H_{i, n} := \{\mu \in \mathbb{R} \otimes_{\mathbb{Z}} X \mid \langle \alpha_i^\vee, \mu \rangle = n\}$ for each $n \in \mathbb{Z}$, and

$$\tilde{\alpha}_i := \begin{cases} \alpha_i & \text{if } i \neq 0, \\ -\theta & \text{if } i = 0, \end{cases} \quad s_i := \begin{cases} r_i & \text{if } i \neq 0, \\ r_\theta & \text{if } i = 0. \end{cases}$$




Fig. 1: Root operator e_i .

The definition of $f_i \eta$ is similar. The following theorem is one of our main results.

Theorem 4.2 (1) The set $\text{QLS}(\lambda)$ together with crystal operators e_i, f_i for $i \in I_{\text{af}}$ and weight function wt , becomes a regular crystal with weight lattice X .

(2) For each $i \in I$, the crystal $\text{QLS}(\omega_i)$ is isomorphic to the crystal basis of $W(\omega_i)$, the fundamental representation of level zero, introduced by Kashiwara [Ka].

(3) Let $\mathbf{i} = (i_1, i_2, \dots, i_p)$ be an arbitrary sequence of elements of I , and set $\lambda_{\mathbf{i}} := \omega_{i_1} + \omega_{i_2} + \dots + \omega_{i_p}$. There exists a crystal isomorphism $\Psi_{\mathbf{i}} : \text{QLS}(\lambda_{\mathbf{i}}) \xrightarrow{\sim} \text{QLS}(\omega_{i_1}) \otimes \text{QLS}(\omega_{i_2}) \otimes \dots \otimes \text{QLS}(\omega_{i_p})$.

Remark 4.3 It is known that the fundamental representation $W(\omega_i)$ (of level zero) is isomorphic to the KR module $W_1^{(i)}$ in the sense of [HKOTT, §2.3] (but for the explicit form of the Drinfeld polynomials of $W(\omega_i)$, see [N, Remark 3.3]), and that it has a global crystal basis (see [Ka, Theorem 5.17]). Furthermore,

the crystal basis of $W(\omega_i) \cong W_1^{(i)}$ is unique, up to a nonzero constant multiple (see also [NS3, Lemma 1.5.3]); we call it a (one-column) KR crystal. By the theorem above, the crystal $\text{QLS}(\lambda)$ is a model for the corresponding tensor product of KR crystals.

4.2 Sketch of the proof of Theorem 4.2

First, let us recall the definition of LS paths of shape λ from [Li].

Definition 4.4 For $\mu, \nu \in X_{\text{af}}^0(\lambda)$ with $\mu > \nu$ (see Definition 3.3) and a rational number $0 < \sigma < 1$, a σ -chain for (μ, ν) is, by definition, a sequence $\mu = \xi_0 \succ \xi_1 \succ \cdots \succ \xi_n = \nu$ of covers in $X_{\text{af}}^0(\lambda)$ such that $\sigma \langle \gamma_k^\vee, \xi_{k-1} \rangle \in \mathbb{Z}$ for all $k = 1, 2, \dots, n$, where γ_k is the label for $\xi_{k-1} \succ \xi_k$.

Definition 4.5 An LS path of shape λ is, by definition, a pair $(\underline{\nu}; \underline{\sigma})$ of a sequence $\underline{\nu} : \nu_1 > \nu_2 > \cdots > \nu_s$ of elements in $X_{\text{af}}^0(\lambda)$ and a sequence $\underline{\sigma} : 0 = \sigma_0 < \sigma_1 < \cdots < \sigma_s = 1$ of rational numbers such that there exists a σ_u -chain for (ν_u, ν_{u+1}) for each $u = 1, 2, \dots, s-1$.

We denote by $\mathbb{B}(\lambda)$ the set of all LS paths of shape λ . We identify an element

$$\pi = (\nu_1, \nu_2, \dots, \nu_s; \sigma_0, \sigma_1, \dots, \sigma_s) \in \mathbb{B}(\lambda)$$

with the following piecewise linear, continuous map $\eta : [0, 1] \rightarrow \mathbb{R} \otimes_{\mathbb{Z}} X_{\text{af}}$:

$$\pi(t) = \sum_{u'=1}^{u-1} (\sigma_{u'} - \sigma_{u'-1}) \nu_{u'} + (t - \sigma_{u-1}) \nu_u \quad \text{for } \sigma_{u-1} \leq t \leq \sigma_u, \quad 1 \leq u \leq s.$$

Let $\mathbb{B}(\lambda)_{\text{cl}} := \{\text{cl}(\pi) \mid \pi \in \mathbb{B}(\lambda)\}$, where $\text{cl}(\pi)$ is the piecewise linear, continuous map $[0, 1] \rightarrow \mathbb{R} \otimes_{\mathbb{Z}} X$ defined by $(\text{cl}(\pi))(t) := \text{cl}(\pi(t))$ for $t \in [0, 1]$. For $\eta \in \mathbb{B}(\lambda)_{\text{cl}}$ and $i \in I_{\text{af}}$, we define $e_i \eta$ and $f_i \eta$ in exactly the same way as above. Then it is known from [NS1, NS2] that the same statement as in Theorem 4.2 holds for $\mathbb{B}(\lambda)_{\text{cl}}$. Thus, Theorem 4.2 follows immediately from the following proposition.

Proposition 4.6 *The affine crystals $\text{QLS}(\lambda)$ and $\mathbb{B}(\lambda)_{\text{cl}}$ are isomorphic.*

This proposition is a consequence of Theorem 3.4. Let us show that if $\eta \in \text{QLS}(\lambda)$, then $\eta \in \mathbb{B}(\lambda)_{\text{cl}}$. Here, for simplicity, we assume that $\eta = (x, y; 0, \sigma, 1) \in \text{QLS}(\lambda)$, and $x = w_0 \xrightarrow{\gamma^1} w_1 \xrightarrow{\gamma^2} w_2 = y$ is a directed σ -path from y to x . Take an arbitrary $\mu \in X_{\text{af}}^0(\lambda)$ such that $\text{cl}(\mu) = y$. By applying Theorem 3.4 to $w_1 \xrightarrow{\gamma^2} w_2 = y$, we obtain a cover $\nu_1 \succ \mu$ for some $\nu_1 \in X_{\text{af}}^0(\lambda)$ with $\text{cl}(\nu_1) = w_1$. Then, by applying Theorem 3.4 to $x = w_0 \xrightarrow{\gamma^1} w_1$, we obtain a cover $\nu \succ \nu_1$ for some $\nu \in X_{\text{af}}^0(\lambda)$ with $\text{cl}(\nu) = x$. Thus we get a sequence $\nu \succ \nu_1 \succ \mu$ of covers in $X_{\text{af}}^0(\lambda)$. It can be easily seen that this is a σ -chain for (ν, μ) , which implies that $\pi = (\nu, \mu; 0, \sigma, 1) \in \mathbb{B}(\lambda)$. Therefore, $\eta = \text{cl}(\pi) \in \mathbb{B}(\lambda)_{\text{cl}}$. The reverse inclusion can be shown similarly.

4.3 Description of the energy function in terms of quantum LS paths

Recall the notation in Theorem 4.2 (2); for simplicity, we set $\lambda := \lambda_i$. In [NS5], Naito and Sagaki introduced a degree function $\text{Deg}_\lambda : \mathbb{B}(\lambda)_{\text{cl}} = \text{QLS}(\lambda) \rightarrow \mathbb{Z}_{\leq 0}$, and proved that Deg_λ is identical to the energy function [HKOTT, HKOTY] on the tensor product $\mathbb{B}(\omega_{i_1})_{\text{cl}} \otimes \mathbb{B}(\omega_{i_2})_{\text{cl}} \otimes \cdots \otimes \mathbb{B}(\omega_{i_p})_{\text{cl}} = \text{QLS}(\omega_{i_1}) \otimes \text{QLS}(\omega_{i_2}) \otimes \cdots \otimes \text{QLS}(\omega_{i_p})$ (which is isomorphic to the corresponding tensor product of

KR crystals; see Remark 4.3) via the isomorphism Ψ_i . The function $\text{Deg}_\lambda : \mathbb{B}(\lambda)_{\text{cl}} \rightarrow \mathbb{Z}_{\leq 0}$ is described in terms of $\text{QB}(W^J)$ as follows. For $x, y \in W^J$ let

$$\mathbf{d} : x = w_0 \xleftarrow{\beta_1} w_1 \xleftarrow{\beta_2} \cdots \xleftarrow{\beta_n} w_n = y$$

be a shortest directed path from y to x , and define

$$\text{wt}(\mathbf{d}) := \sum_{\substack{1 \leq k \leq n \text{ such that} \\ w_{k-1} \xleftarrow{\beta_k} w_k \text{ is a down arrow}} \beta_k^\vee.$$

The value $\langle \text{wt}(\mathbf{d}), \lambda \rangle$ does not depend on the choice of a shortest directed path \mathbf{d} from y to x , and

Theorem 4.7 *Let $\eta = (x_1, x_2, \dots, x_s; \sigma_0, \sigma_1, \dots, \sigma_s) \in \text{QLS}(\lambda) = \mathbb{B}(\lambda)_{\text{cl}}$. Then,*

$$\text{Deg}(\eta) = - \sum_{u=1}^{s-1} (1 - \sigma_u) \langle \lambda, \text{wt}(\mathbf{d}_u) \rangle, \quad (6)$$

where \mathbf{d}_u is a shortest directed path from x_{u+1} to x_u .

4.4 The quantum alcove model

The quantum alcove model is a generalization of the alcove model of the first author and Postnikov [LP1, LP2], which, in turn, is a discrete counterpart of the Littelmann path model [Li]. For the affine Weyl group terminology below, we refer to [H]. Fix a dominant weight λ . We say that two alcoves are adjacent if they are distinct and have a common wall. Given a pair of adjacent alcoves A and B , we write $A \xrightarrow{\beta} B$ for $\beta \in \Phi$ if the common wall is orthogonal to β and β points in the direction from A to B .

Definition 4.8 [LP1] *The sequence of roots $(\beta_1, \beta_2, \dots, \beta_m)$ is called a λ -chain if*

$$A_0 = A_\circ \xrightarrow{-\beta_1} A_1 \xrightarrow{-\beta_2} \cdots \xrightarrow{-\beta_m} A_m = A_\circ - \lambda$$

is a shortest sequence of alcoves from the fundamental alcove A_\circ to its translation by $-\lambda$.

The *lex λ -chain* Γ_{lex} is a particular λ -chain defined in [LP2, Section 4]. Given an arbitrary λ -chain $\Gamma = (\beta_1, \beta_2, \dots, \beta_m)$, let $r_i = r_{\beta_i}$ and define the *level sequence* (l_1, \dots, l_m) of Γ by $l_i = |\{j \geq i \mid \beta_j = \beta_i\}|$.

Definition 4.9 [LeL1] *A (possibly empty) finite subset $J = \{j_1 < j_2 < \cdots < j_s\}$ of $\{1, \dots, m\}$ is a Γ -admissible subset if we have the following path in $\text{QB}(W)$:*

$$1 \xrightarrow{\beta_{j_1}} r_{j_1} \xrightarrow{\beta_{j_2}} r_{j_1} r_{j_2} \xrightarrow{\beta_{j_3}} \cdots \xrightarrow{\beta_{j_s}} r_{j_1} r_{j_2} \cdots r_{j_s} = \kappa(J). \quad (7)$$

Let $\mathcal{A}^\Gamma(\lambda)$ be the collection of Γ -admissible subsets. Define the level of $J \in \mathcal{A}^\Gamma(\lambda)$ by $\text{level}(J) = \sum_{j \in J^-} l_j$, where $J^- \subseteq J$ corresponds to the down steps in Bruhat order in (7).

Theorem 4.10 *There is an isomorphism of graded classical crystals $\Xi : \mathcal{A}^{\Gamma_{\text{lex}}}(\lambda) \rightarrow \text{QLS}(\lambda)$.*

The map Ξ is the following forgetful map. Given the path (7), based on the structure of Γ_{lex} we select a subpath, and project its elements under $W \rightarrow W/W_\lambda$ where W_λ is the stabilizer of λ in W , thereby obtaining a quantum LS path. The inverse map is more subtle, and is based on the so-called *tilted Bruhat theorem* of [LNSS1]; this is a QBG analogue of the minimum-length Deodhar lift [De] $W/W_\lambda \rightarrow W$.

An affine crystal structure was defined on $\mathcal{A}^{\Gamma_{\text{lex}}}(\lambda)$ in [LeL1]. We show that Ξ is an affine crystal isomorphism, up to removing some f_0 -arrows from $\text{QLS}(\lambda)$. The remaining arrows are called *Demazure arrows* in [ST], as they correspond to the arrows of a certain affine Demazure crystal, cf. [FSS]. Moreover the above bijection sends the level statistic to the Deg statistic of (6). Thus, we have proved that $\mathcal{A}^{\Gamma_{\text{lex}}}(\lambda)$ is also a model for KR crystals.

In [LeL2], we show that all the affine crystals $\mathcal{A}^\Gamma(\lambda)$, for various Γ , are isomorphic. Making particular choices in classical types, we can translate the level statistic into a so-called *charge statistic* on sequences of the corresponding *Kashiwara–Nakashima columns* [KN]. In type A , we recover the classical Lascoux–Schützenberger charge [LSc]. Type C was worked out in [Le2, LeS], while type B is considered in [BL].

5 Macdonald polynomials

The symmetric Macdonald polynomials $P_\lambda(x; q, t)$ [M] are a remarkable family of orthogonal polynomials associated to any affine root system (where λ is a dominant weight for the canonical finite root system), which depend on parameters q, t . They generalize the corresponding irreducible characters, which are recovered upon setting $q = t = 0$. For untwisted affine root systems the Ram–Yip formula [RY] expresses $P_\lambda(x; q, t)$ in terms of all subsequences of any λ -chain Γ (cf. Definition 4.8). In [Le2], it was shown that the Ram–Yip formula takes the following simple form for $t = 0$:

$$P_\lambda(x; q, 0) = \sum_{J \in \mathcal{A}^\Gamma(\lambda)} q^{\text{level}(J)} x^{\text{wt}(J)}, \quad (8)$$

where $\text{wt}(J)$ is a weight associated with J . By using the results in Section 4.4, we can write the right-hand side of (8) as a sum over the corresponding tensor product of KR crystals. It follows that $P_\lambda(x; q, 0) = X_\lambda(q)$. Furthermore, we can express the nonsymmetric Macdonald polynomial $E_{w\lambda}(x; q, 0)$, for $w \in W$, in a similar way to (8), by summing over those $J \in \mathcal{A}^\Gamma(\lambda)$ with $\kappa(J) \leq w$ (in Bruhat order), where $\kappa(J)$ was defined in (7). This formula can be derived both by induction, based on Demazure operators, and from the Ram–Yip formula (in this case, for nonsymmetric Macdonald polynomials); however, the latter derivation is more involved than the one of (8), as it uses the transformation on admissible subsets defined in [Le1, Section 5.1].

References

- [BB] A. Björner and F. Brenti. *Combinatorics of Coxeter groups*. Graduate Texts in Mathematics Vol. 231, Springer, New York, 2005.
- [BFP] F. Brenti, S. Fomin, A. Postnikov. Mixed Bruhat operators and Yang–Baxter equations for Weyl groups. *Int. Math. Res. Not.* 1999, no. 8, 419–441.
- [BL] C. Briggs, C. Lenart. A charge statistic in type B . in preparation.
- [De] V. Deodhar. A splitting criterion for the Bruhat orderings on Coxeter groups. *Comm. Algebra*, 15:1889–1894, 1987.

- [FL] G. Fourier, P. Littelmann. Tensor product structure of affine Demazure modules and limit constructions. *Nagoya Math. J.* 182:171–198, 2006.
- [FOS] G. Fourier, M. Okado, A. Schilling. Kirillov–Reshetikhin crystals for nonexceptional types. *Adv. Math.*, 222:1080–1116, 2009.
- [FSS] G. Fourier, A. Schilling, M. Shimozono. Demazure structure inside Kirillov–Reshetikhin crystals. *J. Algebra*, 309:386–404, 2007.
- [HKOTT] G. Hatayama, A. Kuniba, M. Okado, T. Takagi, Z. Tsuboi. Paths, crystals and fermionic formulae, in “MathPhys Odyssey 2001, Integrable Models and Beyond” (M. Kashiwara and T. Miwa, Eds.), *Prog. Math. Phys.* Vol. 23, pp. 205–272, Birkhäuser, Boston, 2002.
- [HKOTY] G. Hatayama, A. Kuniba, M. Okado, T. Takagi, Y. Yamada. Remarks on fermionic formula, in “Recent Developments in Quantum Affine Algebras and Related Topics” (N. Jing and K.C. Misra, Eds.), *Contemp. Math.* Vol. 248, pp. 243–291, Amer. Math. Soc., Providence, RI, 1999.
- [H] J. E. Humphreys. *Reflection groups and Coxeter groups*. Cambridge Studies in Advanced Mathematics Vol. 29, Cambridge: Cambridge University Press, 1990.
- [Ion] B. Ion. Nonsymmetric Macdonald polynomials and Demazure characters. *Duke Math. J.*, 116:299–318, 2003.
- [Ka] M. Kashiwara. On level-zero representations of quantized affine algebras, *Duke Math. J.* 112:117–175, 2002.
- [KN] M. Kashiwara, T. Nakashima. Crystal graphs for representations of the q -analogue of classical Lie algebras. *J. Algebra*, 165:295–345, 1994.
- [KMOU] A. Kuniba, K. .C. Misra, M. Okado, J. Uchiyama. Demazure modules and perfect crystals. *Comm. Math. Phys.* 192:555–567, 1998.
- [KMOTU] A. Kuniba, K. .C. Misra, M. Okado, T. Takagi, J. Uchiyama. Crystals for Demazure modules of classical affine Lie algebras. *J. Algebra* 208:185–215, 1998.
- [LS] T. Lam, M. Shimozono. Quantum cohomology of G/P and homology of affine Grassmannian. *Acta Math.*, 204:49–90, 2010.
- [LSc] A. Lascoux, M.-P. Schützenberger. Sur une conjecture de H. O. Foulkes. *C. R. Acad. Sci. Paris Sér. I Math.*, 288:95–98, 1979.
- [Le1] C. Lenart. On the combinatorics of crystal graphs, I. Lusztig’s involution. *Adv. Math.*, 211:204–243, 2007.
- [Le2] C. Lenart. From Macdonald polynomials to a charge statistic beyond type A . *J. Combin. Theory Ser. A*, 119:683–712, 2012.
- [LeL1] C. Lenart, A. Lubovsky. A generalization of the alcove model and its applications. 24th International Conference on Formal Power Series and Algebraic Combinatorics (FPSAC 2012), 875–886, *Discrete Math. Theor. Comput. Sci. Proc.*, AR, Assoc. Discrete Math. Theor. Comput. Sci., Nancy, 2012.
- [LeL2] C. Lenart, A. Lubovsky. A uniform realization of the combinatorial R -matrix. in preparation.
- [LNSSS1] C. Lenart, S. Naito, D. Sagaki, A. Schilling, M. Shimozono. A uniform model for Kirillov–Reshetikhin crystals I: Lifting the parabolic quantum Bruhat graph, preprint arXiv:1211.2042

- [LNSSS2] C. Lenart, S. Naito, D. Sagaki, A. Schilling, M. Shimozono. A uniform model for Kirillov–Reshetikhin crystals II. in preparation.
- [LP1] C. Lenart, A. Postnikov. Affine Weyl groups in K -theory and representation theory. *Int. Math. Res. Not.* no. 12, 1–65, 2007. Art. ID rnm038.
- [LP2] C. Lenart, A. Postnikov. A combinatorial model for crystals of Kac-Moody algebras. *Trans. Amer. Math. Soc.*, 360:4349–4381, 2008.
- [LeS] C. Lenart, A. Schilling. Crystal energy via the charge in types A and C . *Math. Zeitschrift*, to appear, doi:10.1007/s00209-012-1011-2.
- [Li] P. Littelmann. Paths and root operators in representation theory. *Ann. of Math.*, (2) 142:499–525, 1995.
- [M] I. G. Macdonald. *Affine Hecke Algebras and Orthogonal Polynomials*. Cambridge Tracts in Mathematics. Cambridge University Press, Cambridge, 2003.
- [Mi] L. C. Mihalcea. Positivity in equivariant quantum Schubert calculus. *Amer. J. Math.*, 128:787–803, 2006.
- [NS1] S. Naito, D. Sagaki. Path model for a level-zero extremal weight module over a quantum affine algebra. *Int. Math. Res. Not.* 1731–1754, 2003.
- [NS2] S. Naito, D. Sagaki. Path model for a level-zero extremal weight module over a quantum affine algebra. II. *Adv. Math.*, 200:102–124, 2006.
- [NS3] S. Naito, D. Sagaki. Construction of perfect crystals conjecturally corresponding to Kirillov–Reshetikhin modules over twisted quantum affine algebras. *Commun. Math. Phys.* 263:749–787, 2006.
- [NS4] S. Naito, D. Sagaki. Crystal structure on the set of Lakshmibai–Seshadri paths of an arbitrary level-zero shape. *Proc. Lond. Math. Soc.*, 96:582–622, 2008.
- [NS5] S. Naito, D. Sagaki. Lakshmibai–Seshadri paths of level-zero shape and one-dimensional sums associated to level-zero fundamental representations. *Compos. Math.*, 144:1525–1556, 2008.
- [N] H. Nakajima. Extremal weight modules of quantum affine algebras, in “Representation Theory of Algebraic Groups and Quantum Groups” (T. Shoji et al., Eds.), *Adv. Stud. Pure Math.* Vol. 40, pp. 343–369, Math. Soc. Japan, 2004.
- [Na] K. Naoi. Weyl modules, Demazure modules and finite crystals for non-simply laced type. *Adv. Math.*, 229:875–934, 2012.
- [P] D. Peterson. Quantum cohomology of G/P . Lecture notes, Massachusetts Institute of Technology, Cambridge, MA, Spring 1997.
- [RY] A. Ram, M. Yip. A combinatorial formula for Macdonald polynomials. *Adv. Math.*, 226:309–331, 2011.
- [Sa] W. A. Stein and others. Sage Mathematics Software (Version 5.4). *The Sage Development Team*, 2012. <http://www.sagemath.org>.
- [Sa-comb] The Sage-Combinat community. Sage-Combinat: enhancing Sage as a toolbox for computer exploration in algebraic combinatorics, 2008-2012. <http://combinat.sagemath.org>.
- [ST] A. Schilling, P. Tingley. Demazure crystals, Kirillov–Reshetikhin crystals, and the energy function. *The Electronic Journal of Combinatorics*, 19:P2, 2012.