

# Improvement of the assembly of heterozygous genomes of non-model organisms, a case study of the genomes of two *Spodoptera frugiperda* host strains

Anaïs Gouin, Anthony Bretaudeau, K Labadie, Jean-Marc Aury, Emmanuelle D'Alençon, Claire Lemaitre, Fabrice Legeai

## ► To cite this version:

Anaïs Gouin, Anthony Bretaudeau, K Labadie, Jean-Marc Aury, Emmanuelle D'Alençon, et al.. Improvement of the assembly of heterozygous genomes of non-model organisms, a case study of the genomes of two *Spodoptera frugiperda* host strains. *Arthropod Genomics* 2015, Jun 2015, Manhattan (Kansas), United States. 2015. <hal-01240443>

**HAL Id: hal-01240443**

**<https://hal.inria.fr/hal-01240443>**

Submitted on 9 Dec 2015

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Improvement of the assembly of heterozygous genomes of non-model organisms, a case study of the genomes of two *Spodoptera frugiperda* host strains

Anaïs GOUIN<sup>1</sup>, Anthony BRETAUDEAU<sup>2</sup>, Karine Labadie<sup>3</sup>, Jean-Marc Aury<sup>3</sup>, Emmanuelle d'Alençon<sup>4</sup>, Claire LEMAITRE<sup>1</sup> and Fabrice LEGEA1<sup>1,2</sup>



<sup>1</sup>INRIA/IRISA/GenScale, Rennes, France

<sup>2</sup>INRA, Institut de Génétique, Environnement et Protection des Plantes (IGEPP), Rennes, France


<sup>3</sup>CEA Genoscope, Evry, France

<sup>4</sup>INRA, Diversité, Génomes et Interaction Micro-organismes (DGIMI), Montpellier France




**Motivation** : Some heterozygous regions have a significant divergence between the two haplotypes and the assembly process can lead to the construction of two different contigs, instead of one consensus sequence.


**Objective** : Set up a strategy to detect and correct false duplications in already-built assemblies.




## RE-ASSEMBLY



## RE-ANNOTATION



## APPLICATION : *Spodoptera frugiperda* genomes



### Assembly stats

	Initial assembly <i>Allpaths</i>	<i>Platanus</i> assembly	Corrected assembly
Total size (Mb)	526.0	470.1	437.9
Nb. scaffolds	48 272	41 633	41 577
N50 (bp)	39 593	75 578	52 781
Total Gap length	13.6 Mb	56.1 Mb	11.4 Mb
BUSCO* stats	No hit	14	15
	Single hit	1497	1885
	Multi hit	782	393

### Annotation stats

Previous release : 25,041 genes  
3 746 genes to remap

	# genes	% success
Case A	34	0
Case B	747	45.4%
Case C	643	100.0%
Case D/E	2322	86.3%

**New release : 21,578 genes**

**Comparison of 2 *Spodoptera* strains:** the genome of the *Spodoptera frugiperda* "rice variant" has been sequenced and assembled with *Platanus*, and annotated with the help of *Maker*. The comparison of the two draft sequences leads to the identification of about 16,000 pairs of orthologous genes, and thousands of segmental rearrangements (deletions, duplications or inversions).

