



HAL
open science

Dyshomeostasis, obesity, addiction and chronic stress

David F Marks

► **To cite this version:**

David F Marks. Dyshomeostasis, obesity, addiction and chronic stress. *Health Psychology Open*, 2016, 2 (2). hal-01274337

HAL Id: hal-01274337

<https://inria.hal.science/hal-01274337>

Submitted on 16 Feb 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

Dyshomeostasis, obesity, addiction and chronic stress

David F Marks*

*editorjhp@gmail.com; 2 impasse Enclos Reboul, 13200 Arles, France.

ABSTRACT

When eating control is overridden by hedonic reward, a condition of obesity dyshomeostasis occurs. Appetitive hedonic reward is a natural response to an obesogenic environment containing endemic stress and easily accessible high-energy palatable foods and beverages. Obesity dyshomeostasis is mediated by the prefrontal cortex, amygdala and HPA axis. Ghrelin provides the perfect signalling for feeding dyshomeostasis, affect control and hedonic reward.

Dyshomeostasis plays a central role in obesity causation, the addictions, chronic conditions and in persons with diverse bodies. Prevention and treatment efforts that target sources of dyshomeostasis provide ways of reducing adiposity, ameliorating addiction, and raising the quality of life in people suffering chronic stress.

KEY WORDS

Dyshomeostasis, obesity, addiction, chronic stress, hedonic reward, ghrelin, Circle of Discontent

Homeostasis is omnipresent in nature and all living things. It occurs within individual organisms, in social milieu, and in the environment. At the biochemical, physiological, psychological and social levels, smooth functioning of healthy organisms relies upon the successful operation of homeostasis. However, wherever there is homeostasis, there is the potential of dyshomeostasis. When homeostasis is disrupted, the well-being of an individual, family or population is placed in jeopardy. Recently, the principle of dyshomeostasis was applied to the explanation of obesity (Marks, 2015).

The key scientific dilemma is to understand how obesity can happen in the first place, and on the global scale that exists at the present time. There has been a theoretical vacuum concerning obesity that defies logic and imagination. A phenomenon that is so pervasive cannot lie beyond explanation in science. The explanation, I believe, is a relatively simple but neglected one: obesity is a form of dyshomeostasis. In this paper, I elucidate preliminary hypotheses concerning the neurobiological basis of obesity dyshomeostasis and discuss issues raised by commentators (Annunziato & Grossman, 2016; Brindal & Wittert, 2016; DiClemente & Delahanty, 2016; Markey et al., 2016; Pelletier et al., 2016; Piko & Brassai, 2016; Roosen & Mills, 2016; Rosenbaum & White, 2016).

A THEORETICAL VACUUM

The accepted explanation of overweight and obesity has been the Energy Balance Theory (EBT) in which weight gain is a consequence of energy expenditure being less than energy intake. This mechanistic approach led to the modern obsession with calorie counting and dieting (Marks et al., 2015). It is true that short-term weight loss can be achieved by any calorie-reducing diet but, in the long-term, studies show that calorie-counting is not associated with significant weight loss. One reason for this outcome is that all calories are not equal (Feinman and Fine, 2004). If you eat an equal number of calories of protein, fat and carbohydrates, the metabolic processes are different and calories from fat are more likely to end up on your waist as fewer calories are burned by the thermic effect of eating. The quality and type of foods one consumes influences diverse pathways related to weight homeostasis, such as satiety, hunger, brain reward, glucose-insulin responses, hepatic de novo lipogenesis, adipocyte function, metabolic expenditure, and the microbiome (Mozaffarian, 2016). All calories are not equal: some foods impair pathways of weight homeostasis and others promote the integrity of weight regulation. In sum, the EBT is an oversimplified, descriptive approach that has promoted victim-blaming and stigmatization, which has done little to reduce the prevalence of obesity (Marks et al., 2015). One might even say, it increased it.

Associated with the EBT is the view that obesity and overweight are the consequences of

inactivity. This belief has been responsible for much disillusionment among people striving for significant weight-loss. A 100-kg man needs to run about 20 km. each week to reach a weight of 85 kg. However, this outcome would take approximately 5 years using exercise alone. That would mean running 5000 km., one-eighth of the circumference of the planet, over 5 years to lose 15 kg (Marks et al., 2015). It is perhaps unsurprising that systematic reviewers have concluded that adding physical activity to a dietary intervention for obese individuals has a marginal, if any, effect on average weight loss (Loveman et al., 2011; Swedish Council on Health Technology Assessment, 2013).

The inability of EBT to yield effective long-term interventions for obesity treatment or prevention, suggests to this author that the energy balance approach is theoretically bankrupt. It is a purely descriptive theory of energy transfers in and out of the body but it fails to tell us predictively why any one individual will develop obesity rather than another. The energy balance theory falls short, not only for lack of explanatory power, but because it has done actual harm by stigmatisation of overweight individuals who have been blamed for being both 'greedy' and 'lazy'. The EBT is no longer considered helpful to the complete understanding of obesity and should be retired.

Enter homeostasis theory. The theory of the Circle of Discontent (COD) proposes that homeostatic control of eating can become disrupted under the conditions of modern living in which large sectors of the population are exposed to chronic stress and negative affect while simultaneously being offered supplies of low-cost fatty and sugary foods. Under such oppressive conditions, palliation of stress and negative affect is facilitated by the hedonic eating of high-energy, high-fat or high-sugar foods and beverages, indubitably the main cause of obesity. Over a protracted period of time, obesity dyshomeostasis (OD) has a highly deleterious impact on human physical and mental health and is associated with metabolic syndrome, insulin resistance/diabetes, cardiovascular disease, cancers, fatty liver disease, polycystic ovarian disease, depression and many other conditions that are not easily reversed, or are irreversible.

THE NEUROBIOLOGICAL BASIS OF OBESITY

The first place to look for an explanation of obesity dyshomeostasis is neurobiology. In the obese person, something has gone wrong within the psychoneuroendocrinal system. Clearly, the mechanisms responsible for feeding control have been disrupted. But what is the nature of the disruption? And why one person rather than another?

Obesity occurs as a consequence of a disruption to the homeostatic mechanisms that regulate the control of eating. When addressing the spectrum of clinical conditions that are the concern of

medical and clinical science, the idea of homeostasis imbalance is quite ancient. Since classical theories of Hippocrates and Galen, the history of clinical medicine has been associated with the fundamental principle of dyshomeostasis. What is perhaps surprising is that dyshomeostasis has not previously been cited as the cause of obesity. In the following, the case for dyshomeostasis in obesity will be elaborated. Interesting parallels will become apparent between eating and other forms of consumption which rely on similar neurobiological mechanisms: the addictions to nicotine, alcohol, and illicit drugs, and also behavioural addictions. The following sections discuss known biological mechanisms that relate to the psychological and social issues included in the “Circle of Discontent”. In doing so, the opportunity to lend “a truly integrative biopsychosocial lens” is embraced as suggested by Rosenbaum and White (2016).

Dyshomeostasis in human feeding

In an environment that promotes widespread body dissatisfaction, angst and depression, homeostatic feedback loops are producing excessive consumption of unhealthy processed foods that over a protracted period causes obesity in large numbers of vulnerable people. Multiple clinical studies in different areas of medicine demonstrate the primary role of homeostasis in healthy functioning and the consequences of dyshomeostasis. Homeostasis can be overloaded or over-ridden with too strong a flow of inputs or outputs that disrupt its normal functioning: “The homeostatic behaviour of inflow controllers breaks down when there are large uncontrolled inflows, whereas outflow controllers lose their homeostatic behaviour in the presence of large uncontrolled outflows” (Drengstig et al., 2012). Homeostasis can be disrupted anywhere and perturbations will inevitably occur in normal functioning (Richards, 1960).

There are many examples of dyshomeostasis in clinical medicine. Well-known to psychologists, Hans Selye reported that a persistent environmental stressor (e.g., temperature extremes), together with an associated homeostatic hormonal response, leads to tissue injury that he termed a 'disease of adaptation' (Selye, 1946). Intestinal homeostasis breaks down in inflammatory bowel disease (Maloy & Powrie, 2011) and in the microbial ecology of dental plaque causing dental disease (Marsh, 1994). This form of dyshomeostasis can result from local infection and inflammation and give rise to complications that affect the nervous and endocrine systems (Maynard et al., 2012). An altered balance between the two major enteric bacterial phyla, the Bacteroidetes and the Firmicutes, has been associated with clinical conditions. Within the microbiota of the gut, obesity has been associated with a decreased presence of bacteroidetes and an increased presence of actinobacteria (Turnbaugh & Gordon, 2009; Ley, 2010). Kamalov, Bhattacharya and Weber (2010) proposed a dyshomeostasis theory of congestive heart failure. Craddock et al. (2012) suggested a zinc dyshomeostasis hypothesis of Alzheimer's Disease.

Homeostatic regulation within the endocrinal and central nervous systems has been associated

with feeding control. Cortical areas conveying sensory and behavioral influences on feeding provide inputs to the Nucleus Accumbens (NAc) and the Lateral Hypothalamic Area (LHA) is the site of homeostatic and circadian influences (Saper et al., 2002). Hormones such as leptin circulate in proportion to body fat mass, enter the brain, and act on neurocircuits that govern food intake (Morton et al., 2004). Through direct and indirect actions, it is hypothesized that leptin diminishes the perception of food reward while enhancing the response to satiety signals generated during food consumption that inhibit feeding and lead to meal termination.

Another important hormone is ghrelin which is the only mammalian peptide hormone able to increase food intake. Interestingly, ghrelin also responds to emotional arousal and stress (Labarthe, Fiquet, Hassouna, Zizzari et al., 2015; Müller et al., 2015). During chronic stress, increased ghrelin secretion induces emotional eating by acting at the level of the hedonic/reward system. As ghrelin has anxiolytic action in response to stress, this adaptative response may contribute to control excessive anxiety and prevent depression (LaBarthe et al., 2015). In obesity, studies have shown a reduced ability to mobilize ghrelin in response to stress or central ghrelin resistance at the level of the hedonic/reward system which may explain the inability to cope with anxiety and increased susceptibility to depression (Figure 1). Reciprocally, studies have shown that people with depression have increased susceptibility to obesity and eating disorders (Marks, 2015).

INSERT FIGURE 1 ABOUT HERE

In addition to leptin and ghrelin, other lipid messengers that modulate feeding by sending messages from the gut to the brain have been identified. For example, oleoylethanolamine has been associated with control of the reward value of food in the brain (Lo Verme et al., 2005; Tellez et al., 2013). Mice fed a high-fat diet had abnormally low levels of oleoylethanolamine in their intestines and did not release as much dopamine compared to mice on low-fat diets. Thus, alterations in gastrointestinal physiology induced by excess dietary fat may be one factor responsible for excessive eating in the obese (Tellez et al., 2013).

Obesity Dishomeostasis Theory holds that obesity is caused by the imposition of the hedonic reward system, designed for the amelioration of chronic stress, anxiety and depression, overriding homeostasis. Within OD theory, the Circle of Discontent (Marks, 2015) is closely paralleled by the model of the hedonic/reward response to ghrelin (LaBarthe et al., 2014; Figure 1).¹ The LaBarthe et al. model contains some redundant features and duplication of constructs that can be avoided. In Figure 1, “Chronic stress” and “Anxiety/Depression” are merged into the single construct, “Negative affect”. Similarly, in the context of obesity, “Hedonic reward/response” and “Emotional

eating” are also operationally a single process. With these amendments, it can be seen that the simple diamond structure of the Circle of Discontent emerges from the LaBarthe model (Figure 2). The model of LaBerthe et al. (2015) provides the COD with a framework inside neurochemistry.

INSERT FIGURE 2 ABOUT HERE

Convergence of Obesity Dyshomeostasis Theory with Neurobiology

Traditionally, the control of feeding has been associated with the hypothalamus (Kelley, Baldo, Pratt & Will, 2005). Circulating factors in the blood modulate the activity of energy sensing neurons in the arcuate nucleus, that modulate food-directed behaviours via their activation of outputs from lateral hypothalamic regions to thalamo-cortical systems, central autonomic effectors, and motor pattern generators. There is a convergence of inputs from amygdala, prefrontal cortex, and nucleus accumbens shell that allow direct modulation of feeding behaviours based upon cognitive and affective signaling. These avenues of influence on feeding control provide the entry point for the Circle of Discontent. When the environmental conditions are: (A) obesogenic, due to the ready availability of highly palatable high-energy foods; (B) stressful, due to the presence of stigma, depression and anxiety; and (C) engendering body dissatisfaction, due to the omnipresent socio-cultural thin-ideal, we have all the ingredients necessary for obesity formation. According to OD theory, the cognitive and affective processes of the Circle of Discontent over-ride the neurobiological processes that regulate feeding and energy-homeostasis breaks down.

The amygdala, prefrontal cortex and nucleus accumbens participate in the regulation of both affect and feeding. The amygdala consists of a group of nuclei involved in emotional learning and expression, a key element of the neural basis of emotion. Damage to the amygdala may lead to an increased threshold for emotional perception and expression, impairments in emotional learning, deficits in the perception of facially expressed emotion, and impaired memory for emotional events (Cardinal, Parkinson, Hall and Everitt, 2002).

Among younger adults, it has been found that the ability to wilfully regulate negative affect, enabling effective responses to stressful experiences, engages regions of prefrontal cortex (PFC) and the amygdala. Urry et al. (2006) tested whether PFC and amygdala responses during emotion regulation predict the diurnal pattern of salivary cortisol secretion. They also tested whether PFC and amygdala regions are involved in emotion regulation in older (62- to 64-year-old) individuals. They measured brain activity using functional magnetic resonance imaging as participants regulated (by intentionally increasing or decreasing) their affective responses or attended to negative picture stimuli. Urry et al. also collected saliva samples for 1 week at home

for cortisol assay. Increasing negative affect resulted in ventral lateral, dorsolateral, and dorsomedial regions of PFC and amygdala activation. The predicted link between brain function in the PFC and amygdala occurred while reducing negative affect in the laboratory and diurnal regulation of endocrine activity in the home environment (Urry et al., 2006). The authors concluded that functional coupling between the PFC and the amygdala enables effective regulation of negative emotion and the activity of PFC–amygdala circuitry during regulation of negative affect predicts longer-term regulation of endocrine activity that may be important for health and well-being.

In the obesity dyshomeostasis theory, negative affect causes increased feeding. This causal relationship is made possible by the fact that the system that regulates negative affect, the hypothalamo-pituitary-adrenal axis, also regulates feeding and thus each process influences the other in eliciting increased consumption (Maniam and Morris, 2012). Negative affect induces increased comfort food intake and body weight gain in humans (Dallman et al. 2003). In rats chronic stress produces decreases corticotropin-releasing factor (CRF) mRNA in the hypothalamus. Depressed people who overeat have decreased cerebrospinal CRF, catecholamine concentrations, and hypothalamo–pituitary–adrenal activity. In line with the Circle of Discontent theory, it has been proposed that people eat comfort food in an attempt to reduce the activity of the chronic stress-response network with its attendant anxiety (Dallman et al., 2003; Dallman et al., 2004).

Obesity is associated with neuroendocrine disturbances, in which the HPA axis plays a central role. The HPA axis is stimulated by negative affect which is associated with discrete, periodical elevations of cortisol (Björntorp and Rosmond, 2000). Prolonged HPA axis stimulation is followed by a continuous degradation of the mechanisms controlling eating and affect. The net effects of neuroendocrine-endocrine perturbations in the HPA axis is insulin resistance and accumulation of body fat. These are effects of cortisol combined with diminished secretion of growth and sex hormone secretions. The outcome of these changes is hypothalamic arousal and Metabolic Syndrome. The feedback regulation of the HPA axis has a key position in this chain of events with control being mediated by glucocorticoid receptors (Björntorp and Rosmond, 2000).

The impact of negative affect, whether in the form of anxiety, depression or stress, is modulated by the prefrontal cortex that appraises, evaluates, interprets and monitors the self and the outside world, including responses to the current appearance of the body. A person's body dissatisfaction is at once a cognitive and affective product based on cognitive appraisal and autochthonous perception of the body's attributes and one's feelings about these. In response, the HPA axis produces glucocorticoids that regulate the homeostasis of consumption.

In addition to the mediation of stress responses by the HPA axis, recent studies have observed that there is an alternative system for mediation of stress responses in circulating ghrelin, a peptide hormone, acting upon the amygdala (Meyer et al., 2013). We return to the role of ghrelin later in this article.

Based on the above review of evidence, a provisional description of the neurobiological substrate of the Circle of Discontent is summarised in Figure 3. The model shows feedback loops between the pre-frontal cortex, the amygdala, the HPA axis and visceral adiposity as mediators of body dissatisfaction, negative affect, eating behaviour, and obesity respectively.

INSERT FIGURE 3 ABOUT HERE

The evidence from neurobiology suggests that the homeostasis of eating can be overridden by the hedonic reward system acting to relieve stress through the excessive consumption of palatable foods (Figure 4). Furthermore, eating is controlled by a complex neural network including the mesocorticolimbic pathway, which consists of the ventral tegmental area, nucleus accumbens, amygdala, hippocampus and prefrontal cortex. These regions are the neural substrates of mood, pleasure, desire, experience of self, body satisfaction and self recognition, and have a significant influence on eating patterns and can generate excessive eating. The hedonic system overrides and disrupts homeostatic control when there is chronic negative affect and accessibility to palatable energy-dense foods. In obese people, eating to excess is produced by a Circle of Discontent, a difficult-to-control self-medication of hedonic reward to assuage stress, anxiety and depression which is paralleled by, but not identical to, the use of nicotine, alcohol and drugs among addicted users.

INSERT FIGURE 4 ABOUT HERE

ENTERING AND EXITING THE CIRCLE OF DISCONTENT

A key question concerns entry and exit to the Circle of Discontent (DiClemente and Delahanty, 2016). Who enters the Circle for the first time, who stays and who leaves, and is it a revolving door? What are the prospects, once inside the Circle, of making positive change?

Important work on the issue of change in health-related behaviours has been carried out by the DiClemente group (DiClemente, 2003; DiClemente, Delahanty and Havas, 2015). If the theory is to have true explanatory value, these issues need to be addressed by the homeostasis theory. As stated by DiClemente and Delahanty (2016): "The challenge is to understand how early problems in attachment may influence some to overeating or anorexia, others to sociopathy and drug abuse,

others to depression or anxiety, and still others to being successful professionals. It depends on how the experiences, environment, knowledge, and opportunities filter the early experiences and influence movement forward in the process of change for these different outcomes.”

The Homeostasis Theory of Obesity describes two main systems, the Circle of Discontent (COD) and the Motivation and Energy Mobilization (MEM) system (Figure 5). In the Circle of Discontent, the levels of body dissatisfaction, negative affect and high-energy consumption run out of control. A key connection in the Circle is between chronic stress and comfort eating (Dallman et al., 2003). In the MEM system, lowered motivation causes changes in restraint, dietary intake and activity, which cause reductions to subjective well-being, mobility, and positive affect. The whole complex establishes unhealthy eating habits, low activity levels, negative affect, overweight and obesity.

It is commonly assumed in Psychology that obesity is caused primarily by 'lifestyle changes'. This assumption, however, is not borne out by the evidence. Many drivers push a person towards entry of the Circle. We can use the analogy of a lottery in which people are allocated tickets. For the most part, tickets are allocated at different critical period in the life cycle starting from the moment of conception. The tickets carry percentage points according to their importance as obesity determinants. At any time, a person's BMI is linearly related to the total number of 'obesity points' they have been allotted. A schematic plan of obesity determinants is shown in Table 2.

INSERT TABLE 2 ABOUT HERE

The prenatal period and adolescence present critical periods for the development of obesity that persists into adulthood (Dietz, 1997). Genetic predisposition, epigenetic factors and prenatal maternal stress, including problems with the partner (Sominsky and Spencer, 2014; Talge, Neal and Glover, 2007) all have an influence. Socioeconomic disadvantage in the form of poverty causes chronic life stress at all stages from infancy and adolescence through to adulthood. People living with low levels of income suffer social oppression, chronic stress and multiple episodes of negative affect, restraint and hedonic reward eating of fatty and sugary foods leading to obesity (Drewnowski and Specter, 2004; McLaren, 2007; Monteiro et al., 2004). Early life stress including general parenting, parent childhood abuse and attachment style influence appetite, feeding behaviour and metabolism throughout life (Felliti, 1993; Felliti et al., 1998; Puig et al., 2013; Sleddens et al., 2011; Sominsky and Spencer, 2014).

The field of epigenetics and obesity is relatively new but early steps are being made in identifying biomarkers for obesity. Findings suggest that several epigenetic marks are modifiable, not only by changing the exposure *in utero*, but also by lifestyle changes in adult life, so there is the potential for interventions to reform unfavourable epigenomic profiles (Van Dijk et al., 2015).

Genetic and neurobiological factors help to explain why many people develop obesity while others head towards other consumption-related conditions such as alcoholism, nicotine or drug addiction. Once again, ghrelin helps to tell the story. Ghrelin levels in children with Prader Willi Syndrome are 3- to 4-times higher compared with BMI-matched obese controls (Haqq et al., 2003). Ghrelin shows wide differences between obese and normal weight adults (Tschop, Weyer, Tataranni, Devanarayan et al., 2001) and among adolescents with different kinds of disorders such as anorexia nervosa and obesity. Baseline ghrelin concentrations increased and decreased, respectively, to a mixed meal in anorexic and obese female adolescents (Stock et al., 2005). Low plasma ghrelin has been independently associated with type 2 diabetes, insulin concentration, insulin resistance, and elevated BP (Pöykkö, Kellokoski, Hörkkö, Kauma et al., 2003). Ghrelin levels were also found to be significantly higher in female alcohol-dependent patients compared to controls, but not in male alcoholics (Wurst, Graf, Ehrental, Klein et al., 2007). A review of the burgeoning literature on psychoendocrinological correlates of different COD pathways across gender and population groups will be published elsewhere.

INSERT FIGURE 5 ABOUT HERE

Once inside the Circle, is there any escape? As we have seen in Table 2, the majority of available 'tickets' for the obesity 'lottery' are allocated by the time a person reaches adulthood. Obesity is 90% determined before early adulthood with only limited scope for change. The obesity die has been cast. If we allow for the possibility that around half of epigenetic influences on obesity might be reversible, and a further 10% for potentially reversible lifestyle determinants, we conclude that 80-90% of obesity determinants are irreversible by treatment.

The Circle of Discontent is a vicious, self-sustaining one. Exit options are few. Breaking out of the vicious circle requires momentarily strong motivation and a transformational change in eating habits, lifestyle and philosophy of living. Obesity is a persistent condition that is intractable to treatment. A modest average weight loss of 2-4 kilograms may be achievable by committed adherence to a structured diet regime (Dansinger et al., 2005) but diet systems generally do not provide the keys to a cure (Foster et al., 2003; Katz, 2002). Psychological therapies lead to disappointing outcomes, with cognitive behaviour therapy producing the highest weight loss of a few kilograms (Shaw et al., 2005). Drug treatments involve safety issues and also yield relatively low levels of weight loss. Weight loss relative to placebo ranges from 3% for orlistat and lorcaserin to 9% for phentermine plus topiramate-ER at 1 year (Yanovski & Yanovski, 2014). The only effective treatment for long term weight loss in patients with clinical obesity is surgery, which is costly and inaccessible for the majority of patients (Buchwald et al., 2004).

Within current knowledge, the determinants of obesity are practically irreversible; the illness is persistent and it almost untreatable. To give any other impression to patients would be unethical and misleading. A person inside the Circle of Discontent is likely to remain inside. The most likely exit point will be an early death. To continue to offer treatments that are known to be minimally effective and, quite possibly, harmful to physical or mental health is unethical. Until new, fully authenticated endocrinological treatments become available, all necessary resources should be directed towards prevention.

THE REACH OF PSYCHOLOGICAL HOMEOSTASIS

Looking beyond the topic of obesity, evidence of dyshomeostasis occurs in several fields of health psychology and behavioural medicine. In general, these fields suffer from a theoretical vacuum. I discuss here two specific areas where dyshomeostasis is a prominent feature, addictions and body diversity.

Addictions

In obesity, it has been argued above that hedonistic reward plays a significant disrupting function in weight homeostasis. Many authors have pointed out that food and drug rewards share some common neural substrates, with opioid receptors playing a role in both feeding and reward (Kelley et al., 2005). Saper et al. (2002) stated: “endogenous opioid systems regulate the hedonic value of food intake independently from the ongoing metabolic needs of the individual. Furthermore, food deprivation, which enhances the hedonic response to food, also increases the motivational value of non-food rewards, such as psychostimulants ...intracranial self-stimulation ... and heroin intake”. This perspective places food and addictive drugs such as nicotine and heroin into a similar category. However, while there are definitely similarities, reviews of the comparison and distinctions between mechanisms of food reward and drug addiction also indicate major differences between the two types of consumption (DiLeone, Taylor and Picciotto, 2012). While eating is necessary for survival, and susceptible to selection pressures during evolution, drug addiction begins as a voluntary choice, and is seen as having “piggybacked” onto pre-evolved reward pathways, engaging a subset of the circuits required for feeding (Figure 6).

INSERT FIGURE 6 ABOUT HERE

The Circle of Discontent (COD) has relevance to a variety of conditions that are marked by compulsivity such as the addictions to tobacco, alcohol, illicit drugs, and behaviours such as gambling and internet gaming. These habits/addictions involve compulsion and loss of control which can be costly to the individuals concerned in both health and monetary terms; all have been associated with chronic stress and negative affect in the form of anger, anxiety or depression

(Breslau et al., 1993; Helzer, and Pryzbeck, 1988; Patton et al., 1996, 1998; Swendsen et al., 1998). Diverse consumption patterns across different population groups prove that “no size that fits all” but the causal mechanisms remains essentially the same.

Excessive consumption is a hedonic strategy to increase reward and reinforces habitual behaviour by reducing negative affect and dissatisfaction. Alcohol, drugs, gambling, gaming, shopping, internet use, TV viewing, sports, fitness training, running, swimming, tanning, and sex are all activities that have been said to be addictive or habit forming by one authority or another. It will suffice here to consider tobacco addiction.

Smoking a cigarette is a homeostatic behaviour that corrects the imbalance of the dopaminergic reward system at the biochemical and physiological levels and reduces dissatisfaction and negative affect. The different kinds of homeostasis complement each other to stabilize physiological and psychological well-being. There are many examples of unhealthy habits being strengthened by hedonic reward and palliation of negative affect in Circles of Discontent.

Nicotine addiction is the result of neurochemical changes to the brain. Long-term tobacco use results in physical dependence and a compulsion to use tobacco. The cigarette is the most efficient and rapid method for delivering nicotine to the brain. Nicotine from cigarette smoke is quickly absorbed in the lungs and then rapidly passes into the brain where it binds to specialized nicotinic acetylcholine receptors (nAChRs). Stimulation of nAChRs by nicotine results in the release of a variety of neurotransmitters in the brain of which dopamine is the most important because it produces pleasure. In an addicted smoker, nicotine therefore produces pleasure, arousal, and mood modulation. However, the effects of a single cigarette are short-lived and the smoker requires frequent top-ups of nicotine to maintain a state of cognitive and affective stability. For the addicted smoker, smoking is a homeostatic process which maintains the required level of nicotine in the brain (Prochaska and Benowitz, 2016).

With chronic nicotine addiction, tolerance develops so that more nicotine is required to deliver the same neurochemical effect. Nicotine is needed to maintain normal brain functioning and stopping smoking, or leaving a longer interval between smokes, is associated with withdrawal symptoms of irritability, anxiety, poor concentration, hunger, weight gain and problems getting along with others. Nicotine addiction is therefore a 'two-edged sword' that is sustained both by positive effects of pleasure and arousal and by the avoidance of the unpleasant effects of nicotine withdrawal. Conditioning sustains tobacco use through the reinforced association between smoking and 'triggers' in the form of specific behaviors such as drinking coffee or alcohol, talking on the phone,

driving a car, and/or completing a meal. Sensorimotor trigger factors associated with the act of smoking, e.g., the smell, taste, and feel of the cigarette smoke become cues for smoking and maintain tobacco use (Sulzberger and Marks, 1977).

In the formation of tobacco addiction the novice inhales tobacco smoke which in the early stages gives a toxic and unpleasant sensation in the mouth and throat. However, with each successive inhalation, the unpleasant sensations in the throat and mouth are replaced by feelings of satisfaction as the smoker strengthens the habit. The feelings of satisfaction grow stronger as the habit is reinforced by the sensation of pleasure and the reduction of negative affect. As the habit strength increases and the addiction is established, the smoker feels withdrawal symptoms that increase in intensity the longer he/she waits before lighting the next cigarette. Symptoms of addiction appear within days or weeks after occasional smoking first begins (Russell, 1990).

The smoker is effectively using cigarette smoking as a form of mood control, as self-medication, titrating the dose to match momentary mood. Smokers are able to regulate smoke and nicotine intake on a puff-by-puff basis, an aspect of smoking control that is acquired early in the tobacco-dependence process (Collins et al., 2010). For this reason, smokers report that cigarettes help relieve their feelings of stress (Figure 7).

INSERT FIGURE 7 ABOUT HERE

Contrary to the subjective experience of smokers, smokers' stress levels are higher than those of nonsmokers, and adolescent smokers report increasing levels of stress as they develop regular patterns of smoking (Parrott, 1999). Nicotine consumption rapidly increases heart rate, and blood pressure (Rose, 2001).

Nicotine addiction exacerbates stress yet yields the delusory impression to smokers that it is stress-reducing. Thus, the alleged "relaxation effect" of smoking is a consequence of reversing the tension and irritability that develop during nicotine depletion between cigarettes. Addicted smokers need nicotine to feel normal (Parrott, 1999). Unpleasant withdrawal symptoms are often associated with increases in urges and intentions to take drugs. In addition, addicted individuals rate coping with negative affect as the prepotent motive for drug use (Baker et al., 2004). The act of smoking cessation leads to reduced stress.

One potential mechanism of nicotine addiction is increased dopamine transmission, which gives a feeling of pleasure or satisfaction. The increase in dopamine activity from nicotine results in pleasant feelings of satisfaction for the smoker, but the subsequent decrease in dopamine leaves the smoker craving for more cigarettes (Gamberino & Gold, 1999; Arias-Carrión et al., 2010).

Negative affect influences how much an individual tends to consume, whether it is food, smoking, alcohol, other drugs, or behaviours and how intensely one craves and, ultimately, whether an abstinent individual will return to harmful consumption (Holgate and Bartlett, 2015). Chronic consumption of alcohol alters the normal function of the affect system causing an increased susceptibility to stress (Adinoff et al., 1998). This increases the likelihood of progression as it produces a cycle of degeneration where exposure to stress leads to escalations in consumption, further reducing the ability to cope with stress and shortening the length of intervals between periods of abstinence.

Many individuals in the population have multiple addictions (Anthony, Warner and Kessler, 1994; Lorains, Cowlishaw and Thomas, 2011). In such individuals multiple Circles of Discontent operate in complementary fashion. Figure 8 illustrates a model of a person who is addicted to nicotine, ethanol, cocaine, and gambling. The four concurrent addictions each has its own homeostatic system and Circle of Discontent. The same person could quite possibly have other addictions as well (e.g. to caffeine, other drugs and the internet) and the already complex diagram would need to be extended to include these. The different addictions have associative connections and any one of the behaviours may act as a trigger for one or more of the others. The brain areas mediating appetitive drug seeking and addictive behaviours may differ between addictions, but include at least some of the areas shown in Figure 5. The four addictions reinforce one other and, after prolonged exposure, the addictions become closed off from external influence and compulsive in nature (Vanderschuren and Everitt, 2004; Volkow and Fowler, 2000). The total system becomes self-sustaining with all of the addictions under the control of a single hedonic reward system designed to palliate negative affect by repeated appetitive behaviours. As previously mentioned, the peptide ghrelin activates reward systems, and its receptor (GHS-R1A) appears to be required for alcohol, cocaine, amphetamine and nicotine induced reward (Jerlhag and Engel, 2010). The hedonic reward system, under the influence of ghrelin, overrides the normal functioning of homeostasis, maintaining the Circle of Discontent and placing the individual at significant long-term risk.

INSERT FIGURE 8 ABOUT HERE

Diverse Bodies

In discussing obesity stigmatisation, Roosen and Mills (2016) suggest the need for a cultural shift “*not only to reduce thin valorization, but also to promote social acceptance of diverse bodies, including bodies that are traditionally understood as unattractive, unhealthy, and unproductive (i.e., disabled and/or obese).*” Roosen and Mills (2016) suggest that this cultural shift is already underway, valorising a “fit” ideal body instead of a thin or muscular one (Musolino et al., 2015).

In line with this perspective, homeostasis and dyshomeostasis are evident in a diverse array of life circumstances and conditions (see Table 1). Behavioural homeostasis occurs in a variety of ways including coping strategies, compensatory actions, life identity projects and an infinite array of sophisticated adaptations to illness, injury and life events. Of key significance to stigmatisation is the plain visibility of obesity, gigantism, dwarfism and, in many instances, disfigurement. The degree of stigmatisation may be influenced in part by the perceived self-responsibility for the condition. Gigantism, dwarfism and disfigurement are genetic and unavoidable. Obesity is often viewed as controllable, changeable and a matter of personal choice. The social perception that obese people can *choose* to lose weight if they want to, but fail to do so, could explain the relatively strong stigmatisation of obese people in modern society (Backstrom, 2012).

INSERT TABLE 1 ABOUT HERE

MOTIVATION

Pelletier, Guertin, Pope and Rocchi (2016) focus their review on motivation constructs from Self-Determination Theory (Deci and Ryan, 1985). They suggest that the Circle of Discontent “does not explain why some people exposed to the same conditions (e.g., abundance of unhealthy foods, negative life events) do not gain weight and become obese” and that it emphasizes strategies that do not consider individuals as active agents of their own behaviours. I clarify here the motivational aspects of my theory.

There can be no question that motivation plays a pivotal role in human behaviour change and in the aetiology of obesity. As previously stated: “It remains necessary to explain *how* or *why* overweight or obesity can develop in a susceptible individual, and *why some individuals develop it and not others*” (Marks, 2015). The Homeostasis Theory of health (HTO) holds that human health is regulated at all times by *multiple systems of homeostasis* that are operating in parallel and in cascades all directed towards stability of function. All of the many thousands of homeostatic systems are interconnected and complementary in maintaining the stability of the human

organism. I refer the reader to Figure 8. In my earlier article, I concentrated on one only of the many homeostatic systems, the *Circle of Discontent (COD)*, a feedback loop that includes Physical Health, Life Satisfaction, Affect and Consumption.

Of equal importance to the COD is the *Motivation and Energy Mobilization (MEM) system*. The MEM system incorporates Motivation, Restraint, Diet, Physical Health, Activity, Subjective Well-Being, Mobility and Affect. As the diagram of Figure 8 shows, the MEM and COD systems are both equally involved in regulating Physical Health and Affect, but only the MEM system includes individual motivation. Without any doubt, the MEM system is of great significance in the maintenance of healthy habits and behaviours and, when things go wrong, in the generation of overweight and obesity.

FIGURE 8 ABOUT HERE

It is helpful to consider the regulatory styles of SDT that are differentiated along an alleged continuum that ranges from non-self-determined styles (i.e. amotivation, external regulation, and introjection) to self-determined ones (i.e. identification, integration, and intrinsic motivation). As suggested by Pelletier et al. (2016) there are parallels between SDT concepts concerning motivational style and the HTO. The Circle of Discontent (COD) is a perfect fit to the profile of 'Controlled Motivation' ¹ within SDT.

Studies by Pelletier and colleagues have provided interesting evidence concerning the style of motivation most likely to be associated with unhealthy eating, depressive symptoms and increased BMI, namely 'Controlled Motivation' (Pelletier, Dion, Slovenic-D'Angelo & Reid, 2004). Pelletier et al. (2004) revealed a response pattern that is consistent with the COD, namely unsuccessful regulation of eating, a concern with quantity but not quality of food eaten, bulimic and depressive symptomatology, low self-esteem and low life satisfaction, and increased BMI, all significantly associated with controlled regulation (Table IV, Pelletier et al, 2004). On the other hand, autonomous regulation was found to be significantly correlated with a concern for quality rather than quantity of food eaten, with successful regulation of eating, healthy eating behaviors, high self-esteem and high life satisfaction. One could not wish for a more positive confirmation of the theory, although I was unaware of it until Pelletier et al. drew my attention to it.

In this light, the two profiles of Controlled Motivation and Autonomous Motivation represent the opposite ends of the homeostasis continuum. Autonomous Motivation brings satisfactory internalized control of eating behaviours, relatively high life satisfaction and positive affect, a state

of positive homeostasis. Controlled Motivation, on the other hand, is a component of homeostatic imbalance in which the individual fails to enjoy, or internalize, the desired goals of eating behaviour (Ryan & Deci, 2006). The Circle of Discontent is perfectly represented by the 'Controlled Regulator', a person whose eating habits are running out of control and whose life satisfaction, and affect levels have deteriorated. In SDT, Controlled Regulation occurs in three forms:

- 1) The Introjected Regulator, not wanting to be ashamed of how they look and eat, feeling that they must absolutely be thin, feeling they would be humiliated if they were not in control of their eating behaviors.
- 2) The External Regulator, other people close to them insist that they do things a certain way, other people close to them will be upset if they don't eat well, people around them will nag them to do it, or it is expected of them.
- 3) The Amotivated Regulator, the worst case scenario, feeling helpless and hopeless, not really knowing what to do, having the impression that they are wasting their time trying to regulate their eating behaviors, not seeing how their efforts can ever lead to eating healthily or helping to improve their health.

In SDT, motivation is King, with a commanding role in need satisfaction for autonomy, competence and relatedness (Ryan and Deci, 2000). In the HTO, motivation is more a Courtier than a King, but a key player, nevertheless, in the MEM system. From the perspective of HTO, the role of motivation in real behaviour change should be evaluated on the basis of the hard-won findings of systematic reviews and meta-analyses. Ng et al.'s (2012) meta-analysis of self-determination theory studies in health care found only low correlations: between autonomous self-regulation and mental and physical health of .06 and .11 respectively; between controlled regulation and mental and physical health of -.19 and .09 respectively; and between amotivation and mental and physical health of -.05 and -.15 respectively. These findings suggests that motivational style controls, at most, 3-4% of variance in mental and physical health.

These modest empirical associations between self-determination theory constructs and health outcomes may, in part, be explained by methodological problems concerning the scoring of self-determination motivation. The validity of the assumed self-determination continuum, forming the basis of the measures employed, has not been supported by robust state-of-the-art psychometric analyses. In a Rasch analysis of the continuum concept, Chemolli and Gagne (2014) found strong evidence of a multidimensional factor structure rather than evidence of a continuum. This significant issue places a serious limitation on the use of self-determination theory in prevention of obesity. Until these methodological issues are solved, the status of SDT remains uncertain and unclear. Unless psychological theories and interventions can be cashed out in objective benefits to health outcomes, they tend to lead only to false hopes and disappointment.

A RCT with SDT-based exercise motivation variables evaluated a behavioral weight control intervention on 3-yr weight change (Silva et al., 2009, 2011). The 1-yr SDT-based intervention was immediately followed up and then again 2-years later with 221 female participants. The intervention group attended 30 sessions, targeted at increasing PA and energy expenditure, adopting a diet consistent with a moderate energy deficit, and integrating exercise and eating patterns that would support weight maintenance. The control group received 29-sessions of general health education on the basis of several educational courses covering various topics, e.g., preventive nutrition, stress management, self-care, and effective communication skills.

Treatment had significant effects on 1- and 2-yr autonomous regulations, 2-yr physical activity, and 3-yr weight change. Average weight loss at 12 months was -7.29% vs. -1.74% in the control group, but the intervention effect tapered off over time showing only -3.9% vs. -1.9% in the control at 36 months. The intervention produced a 2.0% greater average weight loss at 36 months than the control condition. Autonomous style motivation correlated -.31 with 3-year weight change, explaining only 10% of the variance in weight change.

Unfortunately, the abstract, theoretical importance of motivation in SDT has not yet been established in the form of concrete health outcomes. The role of individual motivation appears to be a quite modest one, one process within a complex system, as explicated in the HTO.

UPSTREAM VERSUS DOWNSTREAM INTERVENTION

To have any significant impact on the obesity epidemic, effective interventions must be delivered. Any long-term strategy to curtail the obesity epidemic needs to be based on effectiveness and cost-effectiveness. In this regard, upstream interventions (primary prevention) have been shown to be more effective and more cost-effective than downstream (secondary prevention) ones. A recent economic analysis of the obesity epidemic concluded: *“Education and personal responsibility are critical elements of any program to reduce obesity, but not sufficient on their own. Additional interventions are needed that rely less on conscious choices by individuals and more on changes to the environment and societal norms”* (Dobbs et al., 2014). There are one-billion-plus sufferers alive today. The infrastructure required for individual-level psychological interventions on a universal basis for these one billion people far outstrips available resources. To make any real impact on the obesity epidemic, it is essential to combine prevention efforts with individuals with upstream policies to change the context that is currently promoting the spread of obesity at all levels of society.

Pelletier et al. (2016) argue that “*Environmental changes ... may be slow to implement, can be very expensive, and could be stalled by industries with competing interests*”. However, environmental changes in the form of regulations on sugar or advertizing can generate significant revenue. Both the sugar-sweetened beverage excise tax and the elimination of the tax subsidy for advertising unhealthy food to children would lead to substantial yearly tax revenues (\$12.5 billion and \$80 million, respectively; see Gortmaker et al., 2015 b). The analyses of Gortmaker et al. (2015 a, b) have shown that *the cost effectiveness of these preventive interventions is greater than that obtained from published clinical interventions to treat obesity*. Individual approaches using social cognition models have been tried and tested for many years and outcomes have been disappointing (Marks, Murray, Evans & Estacio, 2015). Lehnert et al. (2012) reviewed long-term economic findings (at least 40 years) for 41 obesity prevention interventions. Interventions were grouped according to their method of delivery, setting and risk factors targeted into behavioural ($n = 21$), community ($n = 12$) and environmental interventions ($n = 8$). Interventions that modified a target population’s environment, i.e. fiscal and regulatory measures, reported the most favourable cost-effectiveness. There can be little doubt that obesity prevention requires the use of cost-effective interventions at all levels of society.

For the one billion-plus individuals living today with obesity these words will not be very welcome. But it is better to face the truth than live in a dreamworld with impossible hopes and expectations. For the vast majority of obese people alive today, there will be no significant reversal. Current treatments are disappointingly weak, expensive, and, often, have unwanted side effects, especially drugs and surgery (Marks et al., 2015). The only way forward that makes sense is prevention - to prevent new cases, as many as possible. The emphasis should be placed on upstream approaches, preventing new floods of cases before they arrive at the point of no return.

SPIRITUALITY HOMEOSTASIS?

Piko and Brassai (2015) make a case for *spiritual balance* as a form of homeostasis. They contend, correctly I believe, that existential attitudes are closely related to “identity formation, moral development, value-related attitudes, personal goals, and lifestyle choices”. Having meaning in life encourages engagement in health-promoting behaviors and the avoidance of health-risking behaviors, such as obesity and eating disorders. Along with physical, cultural, psychosocial and economic needs, a definition of health can also include spiritual needs, not simply the absence of illness (Marks et al., 2015, p. 5).

Piko and Brassai (2016) discuss the meaning-making model of Park (2010), which proposes that people’s perceptions may contribute to content/discontent with life, body and the world. Park (2013) states: “According to the Meaning Making Model, the degree to which one perceives one’s illness as discrepant from one’s global beliefs, such as those regarding identity (e.g., I live a

healthy life style) and health (e.g., living a healthy lifestyle protects people from illness), and global goals (e.g., desire to live a long time with robust health) determines the extent to which the illness is distressing” (p. 43). The Meaning Making Model of Park (2013) assumes that a discrepancy between global beliefs and identity produces distress. In some cases, those beliefs are spiritual in nature. However, primary sources of research on spirituality are not all supportive of the model proposed by Park (2013).

The central role of meaning and purpose in life was previously advocated by Viktor E Frankl (1959) and, later, in the Salutogenic Theory of Aaron Antonovsky (1979, 1987). Neither Frankl's (1952) study nor Antonovsky's theory of salutogenesis is discussed by Park (2010, 2013). We must never forget what Frankl said about the prisoners living in concentration camps: *“Every man was controlled by one thought only: to keep himself alive for the family waiting for him at home, and to save his friends”* (Frankl, 1959). In describing the inmates' dream life, he stated: *“What did the prisoner dream about most frequently? Of bread, cake, cigarettes, and nice warm baths. The lack of having these simple desires satisfied led him to seek wish-fulfilment in dreams.”* In another place, Frankl describes his ultimate realisation, that it is love that satisfies a person's needs for meaning: *“A thought transfixed me: for the first time in my life I saw the truth as it is set into song by so many poets, proclaimed as the final wisdom by so many thinkers. The truth —that love is the ultimate and the highest goal to which man can aspire. Then I grasped the meaning of the greatest secret that human poetry and human thought and belief have to impart: The salvation of man is through love and in love. I understood how a man who has nothing left in this world still may know bliss, be it only for a brief moment, in the contemplation of his beloved...”* *“Set me like a seal upon thy heart, love is as strong as death.”*” No mention is made in Frankl's account of the search for meaning to find spirituality. Frankl (1952) asserted what he called “the will to meaning”: man's search for meaning as the primary motivation in his life.

The HTO is a particular case of the General Theory of Well-Being, which posits causal reciprocal relationships between subjective well-being and life satisfaction (Marks, 2015; Marks, Murray, Evans & Estacio, 2015). Empirical studies suggest the existence of a strong and stable relationship between meaning in life and subjective well-being (Zika & Chamberlain, 1992). People who experience their lives as meaningful tend to be more optimistic and self-actualized (Compton et al., 1996), experience more self-esteem (Steger et al., 2006) and positive affect (King et al., 2006), as well as suffering less depression and anxiety (Steger et al., 2006) and less suicidal ideation (Harlow et al., 1986). The Salutogenic Theory of Antonovsky emphasized the relationship between meaning, purpose in life and positive health outcomes (Eriksson & Lindström, 2006).

For many people, spiritual experience is a source of great meaning to their lives. However spiritual beliefs and experiences are far from universal. To quote one statistic, in the region of 500-750

million people worldwide have no religious or spiritual beliefs and are living as declared atheists (Zuckerman, 2009). In homeostasis the organism actively strives to reduce the discrepancy between an optimum level of a quantity or quality and its current state. While many people certainly strive for meaning, and may feel that they lead 'empty lives', there is no evidence of an optimum level or a homeostatic mechanism for spirituality.

ISSUES REQUIRING MORE RESEARCH

Homeostasis theory proposes that weight gain is fostered by a COD consisting of body dissatisfaction, negative affect and overconsumption. Drawing on this framework, Annunziato and Grossman (2016) describe research in two domains, victim-blaming and devalorizing the thin-ideal. They suggest that university-based clinical health psychologists are uniquely positioned to implement large-scale approaches that have shown promise in addressing core issues in the HTO. Annunziato and Grossman cite examples of research that include a “Social and Emotional Learning” curriculum in Sweden that demonstrated decreases in victimization (Kimber et al., 2008) and the “Body Project” which produced reductions in eating disorders (Stice et al., 2013), in thin-ideal internalization, body image dissatisfaction and negative affect in female students (Stice et al., 2011), and an internet-based programme that demonstrated large weight gain prevention effects (Stice et al., 2014). Annunziato and Grossman (2016) propose more extensive use of both systemic and individual interventions with adolescents and young adults in school settings. For example, large-scale programs in secondary schools and universities could be designed to bring about culture change.

A school-based programme is described by Tran et al. (2014) based in the Canadian province of Alberta. Fung et al. (2012) demonstrated the feasibility and effectiveness of a school-based program in preventing childhood obesity, the Alberta Project Promoting active Living and healthy Eating in Schools (APPLE Schools). The intervention involved a full-time School Health Facilitator in each of 10 schools for implementing healthy eating and active living policies, practices and strategies while engaging stakeholders, including parents, staff and the community. The Facilitators contributed to the schools' health curriculum, and organized activities such as cooking clubs and healthy breakfast, lunch and snack programs, after school physical activity programs, walk-to-school days, community gardens, weekend events and circulated newsletters. By 2010 the students' eating habits and physical activity levels at APPLE Schools had improved significantly while obesity prevalence had declined relative to their peers attending other Albertan schools (Figure 9). Other comprehensive school-based programs have achieved similarly positive results (Greening et al., 2011; Khambalia, 2012; Verstraeten et al., 2012; Veugelers & Fitzgerald, 2005). Ideally, education about and training of healthy eating habits and regular physical activity will become a part of every school curriculum universally.

INSERT FIGURE 9 ABOUT HERE

Markey et al. (2016) referred to the role of social relationships in eating patterns and romantic partners who appear to be especially important and an understudied factor in eating behaviors, body image, and obesity risk. In line with Markey et al.'s (2016) ideas, the influence of the quality of marital relationships in appetite regulation was investigated in a double-blind randomized crossover trial (Jaremka et al., 2015). Both members in 43 couples ate a standardized meal at the beginning of two visits. Observational recordings of marital conflict was employed to assess marital distress. Ghrelin and leptin were sampled premeal and postmeal at 2, 4, and 7 hr. People in more distressed marriages were found to have higher postmeal ghrelin and a poorer quality diet than those in less distressed marriages, but only among participants with lower BMI. Ghrelin and diet quality therefore may be the links between marital distress and its negative health effects (Jaremka et al., 2015).

Children growing up in a disharmonious environment, whether the disharmony is caused by socioeconomic disadvantage or other factors, are exposed to parental frustrations, relationship discord, a lack of support and cohesion, negative belief systems, unmet emotional needs and general insecurity. These stressful experiences increase the risk of psychological and emotional distress, including low self-esteem and self-worth, negative emotions, negative self-belief, powerlessness, depression, anxiety, insecurity and a heightened sensitivity to stress (Hemmingsson, 2014).

Brindal and Wittert (2016) suggest consideration of allostasis, coping style and habituation in addition to the COD model. They argue that the incorporation of these elements into the Homeostatic Theory of Obesity may help to “expand its explanatory power and associated avenues of intervention”. Furthermore they suggest that a meaningful approach to the obesity epidemic and associated chronic disease will require “policy and regulation as well as targeted behavioural strategies aiming to reduce allostatic load”. However, in this author's opinion, the concept of allostasis does not add anything new to the COD model, which is founded on the concept of homeostasis described by Cannon (1932). The concepts of 'allostasis' and 'allostatic load' appear to be based on a misunderstanding of the original concept of homeostasis, which covers all of the functions that the proponents wish to attribute to allostasis (Day, 2005). In addition, the construct of allostasis does not help us to better define stress. I concur with Day (2005), who provided a helpful précis of the 'allostasis theory' as follows: McEwen and Wingfield (2003) wrote: “(the term) stress will be used to describe events that are threatening to the

individual and which elicit physiological and behavioural responses as part of allostasis *in addition to that imposed by the normal life cycle*' (my italics). They propose, in effect, that stress is just one type of challenge that can activate ... allostatic (or, as I prefer, homeostatic) responses.

Accordingly, we can summarise their position as follows: life is a series of challenges; some are part of the normal life cycle; some can be described as stressors; all of these challenges must be met, i.e. homeostasis must be maintained; the process of maintaining homeostasis (a process they would refer to as allostasis) involves wear and tear (which they refer to as allostatic load) which can impact adversely on health. This re-statement of McEwen and Wingfield's thesis may seem banal but reading it with the bracketed words eliminated will demonstrate that understanding their thesis does not require the adoption of allostasis terminology. The critical question that remains then is this: does the concept of allostasis help us to better define stress? I suggest that the answer is 'no' " (Day, 2005, p. 1198).

CONCLUSIONS

Homeostasis is an omnipresent process that has been neglected in the theoretical Psychology. Homeostasis is the primary process for the maintenance of healthy organisms. The breakdown of homeostasis causes disorders including obesity, the addictions, chronic conditions including stress in persons with diverse bodies. All such conditions entail the self-reinforcing activity of a vicious Circle of Discontent. Hedonic reward overrides weight homeostasis to produce obesity dyshomeostasis. A preliminary model suggests that obesity dyshomeostasis is mediated by the prefrontal cortex, amygdala and HPA axis with signalling by the peptide hormone ghrelin which simultaneously controls feeding, affect and hedonic reward. The totality of evidence within current knowledge suggests that obesity is a persistent, intractable condition. Prevention and treatment efforts targeting sources of dyshomeostasis provide ways of reducing adiposity, ameliorating addiction, and raising the quality of life in people suffering chronic stress.

REFERENCES

- Adinoff, B.; Iranmanesh, A.; Veldhuis, J.; Fisher, L. Disturbances of the stress response: The role of the hpa axis during alcohol withdrawal and abstinence. *Alcohol Health Res. World* **1998**, *22*, 67–72.
- Anderberg R H, Anefors C, Bergquist F, Nissbrandt H and Skibicka K P (2014) Dopamine signaling in the amygdala, increased by food ingestion and GLP-1, regulates feeding behavior *Physiology and Behavior* *136*: 135-144.
- Anthony J C, Warner L A and Kessler R C (1994) Comparative epidemiology of dependence on tobacco, alcohol, controlled substances, and inhalants: basic findings from the National Comorbidity Survey. *Experimental and clinical psychopharmacology* *2*: 244.
- Antonovsky, A (1979) *Health, Stress and Coping*. San Francisco: Jossey-Bass.
- Antonovsky A (1987) *Unraveling The Mystery of Health - How People Manage Stress and Stay Well*, San Francisco: Jossey-Bass.
- Arias-Carrión O, Stamelou M, Murillo-Rodríguez E, Menéndez-Gonzalez M, et al. (2010) Dopaminergic reward system: a short integrative review. *International Archives of Medicine* *3*: 24.
- Backstrom L (2012) From the freak show to the living room: Cultural representations of dwarfism and obesity. *Sociological Forum* *27*:682-707.
- Baker T B, Piper M E, McCarthy D E, Majeskie M R et al. (2004) Addiction motivation reformulated: an affective processing model of negative reinforcement. *Psychological Review* *111*: 33.
- Bjork S, Jonsson B, Westphal O and Levin, J E. (1989) Quality of life of adults with growth hormone deficiency: a controlled study. *Acta Paediatrica* *78*(s356): 55-59.
- Björntorp P and Rosmond R (2000) Neuroendocrine abnormalities in visceral obesity. *International journal of obesity and related metabolic disorders: journal of the International Association for the Study of Obesity*, *24*, S80-5.
- Boden J M and Fergusson DM (2011) Alcohol and depression. *Addiction* *106*: 906-914.
- Brindal E and Wittert G (2016) The weight balancing act and allostasis: Commentary on the Homeostasis Theory of Obesity. *Health Psychology Open* (this issue).
- Brown CG and Stewart, SH (2008) Exploring women's use of alcohol as self-medication for depression among women receiving community-based treatment for alcohol problems. *Journal of Prevention and Intervention in the Community* *35*:33-47.
- Buchwald H, Avidor Y, Braunwald E, Jensen MD et al. (2004) Bariatric surgery: a systematic review and meta-analysis. *JAMA* *292*:1724-37.
- Cannon W B (1932) *The wisdom of the body*. New York: Norton.
- Cardinal R N, Parkinson J A, Hall J and Everitt B J (2002) Emotion and motivation: the role of the amygdala, ventral striatum, and prefrontal cortex. *Neuroscience and Biobehavioral Reviews*, *26*: 321-352.

Collins CC et al. (2010). Puffing behavior during the smoking of a single cigarette in tobacco-dependent adolescents *Nicotine and Tobacco Research* 12:164-167.

Craddock T J A, Tuszynski J A, Chopra D, Casey N et al. (2012). The zinc dyshomeostasis hypothesis of Alzheimer's Disease. *PLoS ONE* 7(3), e33552.

Crum R M, La Flair L, Storr CL, Green K M. et al. (2013) Reports of drinking to self-medicate anxiety symptoms: longitudinal assessment for subgroups of individuals with alcohol dependence. *Depression and Anxiety* 30:174-183.

Dallman M F, Pecoraro N, Akana S F, La Fleur S E et al. (2003) Chronic stress and obesity: a new view of "comfort food". *Proceedings of the National Academy of Sciences* 100: 11696-11701.

Dallman M F, la Fleur S E, Pecoraro N C, Gomez F et al. (2004) Minireview: glucocorticoids—food intake, abdominal obesity, and wealthy nations in 2004. *Endocrinology* 145: 2633-2638.

Dansinger M L, Gleason J A, Griffith J L, Selker H P et al. (2005). Comparison of the Atkins, Ornish, Weight Watchers, and Zone diets for weight loss and heart disease risk reduction: a randomized trial. *JAMA*, 293: 43-53.

Dietz WH (1994) Critical periods in childhood for the development of obesity. *The American Journal of Clinical Nutrition* 59:955-9.

Day T A (2005) Defining stress as a prelude to mapping its neurocircuitry: no help from allostasis. *Progress in Neuro-Psychopharmacology and Biological Psychiatry* 29:1195-1200.

DiClemente CC (2003) *Addiction and Change: How Addictions Develop and Addicted People Recover*. New York: Guilford Press.

DiClemente CC, Delahanty JC, Havas SW and Van Orden OR. (2015). Understanding self-reported staging of dietary behavior in low-income women. *Journal of Health Psychology* 20: 741-53.

DiClemente C and Delahanty J (2016) Homeostasis and change. *Health Psychology Open* (this issue).

Dijkers M. (1997) Quality of life after spinal cord injury: a meta analysis of the effects of disablement components. *Spinal Cord* 35:12.

DiLeone R J, Taylor J R and Picciotto M R (2012) The drive to eat: comparisons and distinctions between mechanisms of food reward and drug addiction. *Nature Neuroscience* 15(10): 1330-1335.

Drengstig T, Jolma I W, Ni X Y, Thorsen K, et al. (2012) A basic set of homeostatic controller motifs. *Biophysical Journal* 103: 2000-2010.

Drewnowski A and Specter S E (2004) Poverty and obesity: the role of energy density and energy costs. *The American Journal of Clinical Nutrition*, 79: 6-16.

Elliott T R and Frank R G (1996). Depression following spinal cord injury. *Archives of Physical Medicine and Rehabilitation* 77: 816-823.

Feinman R D and Fine E J (2004) "A calorie is a calorie" violates the second law of thermodynamics. *Nutrition Journal* 3: 10-186.

Felitti V J (1993). Childhood sexual abuse, depression, and family dysfunction in adult obese

patients: a case control study. *Southern Medical Journal*, 86: 732-736.

Felitti V J, Anda R F, Nordenberg D, Williamson D F et al. (1998). Relationship of childhood abuse and household dysfunction to many of the leading causes of death in adults: The Adverse Childhood Experiences (ACE) Study. *American Journal of Preventive Medicine*, 14: 245-258.

Foster G D, Wyatt H R, Hill J O, McGuckin et al. (2003). A randomized trial of a low-carbohydrate diet for obesity. *New England Journal of Medicine*, 348: 2082-2090.

Frankl V E (1959). *Ein Psycholog erlebt das Konzentrationslager. [English] Man's Search For Meaning: An Introduction To Logotherapy*. Boston: Beacon Books.

Fung C, Kuhle S, Lu C, Purcell M et al. (2012) From “best practice” to “next practice”: the effectiveness of school-based health promotion in improving healthy eating and physical activity and preventing childhood obesity. *International Journal of Behavioral Nutrition and Physical Activity* 9: 27.

Gallagher P and MacLachlan M (1999). Psychological adjustment and coping in adults with prosthetic limbs. *Behavioral Medicine* 25: 117-124.

Gamberino WC and Gold MS (1999) Neurobiology of tobacco smoking and other addictive disorders. *The Psychiatric Clinics of North America*, 22:301-312.

Gillman MW (2008) The first months of life: a critical period for development of obesity. *The American Journal of Clinical Nutrition* 87:1587-9.

Goff D C, Henderson D C and Amico E (1992) Cigarette smoking in schizophrenia: relationship to psychopathology and medication side effects. *American Journal of Psychiatry* 149:1189-94.

Gordon-Larsen P, Adair LS, Nelson MC, Popkin BM (2004) Five-year obesity incidence in the transition period between adolescence and adulthood: the National Longitudinal Study of Adolescent Health. *The American Journal of Clinical Nutrition* 80(3):569-75.

Gortmaker S L et al. (2015 a). Cost effectiveness of childhood obesity interventions. *American Journal of Preventive Medicine* 49:102 – 111.

Gortmaker S L et al. (2015 b). Three interventions that reduce childhood obesity are projected to save more than they cost to implement. *Health Affairs* 34:1932-1939.

Greening L, Harrell KT, Low AK and Fielder CE (2011) Efficacy of a school-based childhood obesity intervention program in a rural southern community: TEAM Mississippi Project. *Obesity (Silver Spring)* 19: 1213–1219.

Grogan S (2006). Body image and health contemporary perspectives. *Journal of Health Psychology* 11:523-530.

Guo SS, Wu W, Chumlea WC and Roche AF (2002) Predicting overweight and obesity in adulthood from body mass index values in childhood and adolescence. *The American Journal of Clinical Nutrition* 76:653-8.

Heijnders M and Van Der Meij S (2006). The fight against stigma: an overview of stigma-reduction

strategies and interventions. *Psychology, Health & Medicine* 11:353-363.

Hemmingsson E (2014) A new model of the role of psychological and emotional distress in promoting obesity: conceptual review with implications for treatment and prevention. *Obesity Reviews*, 15: 769-779.

Horgan O and MacLachlan M (2004) Psychosocial adjustment to lower-limb amputation: a review. *Disability and rehabilitation*, 26: 837-850.

Hurxthal I M (1961) Pituitary gigantism in a child five years of age: effect of x-radiation, estrogen therapy and self-imposed starvation diet during an eleven year period. *Journal of Clinical Endocrinology and Metabolism* 21:343-353.

Hyman S E (2009) How adversity gets under the skin. *Nature neuroscience*, 12: 241-243.

Kamalov G, Bhattacharya S K and Weber K T (2010) Congestive heart failure: where homeostasis begets dyshomeostasis. *Journal of Cardiovascular Pharmacology* 56: 320–328.

Jaremka LM, Belury MA, Andridge RR, Lindgren ME et al. (2015) Novel links between troubled marriages and appetite regulation marital distress, ghrelin, and diet quality. *Clinical Psychological Science* 29:2167702615593714.

Jerlhag E and Engel J A (2011) Ghrelin receptor antagonism attenuates nicotine-induced locomotor stimulation, accumbal dopamine release and conditioned place preference in mice. *Drug and Alcohol Dependence*, 117: 126-131.

Katz D L (2002) Pandemic obesity and the contagion of nutritional nonsense. *Public Health Reviews*, 31: 33-44.

Kelley A E, Baldo B A, Pratt W E and Will M J (2005) Corticostriatal-hypothalamic circuitry and food motivation: integration of energy, action and reward. *Physiology and Behavior* 86: 773-795.

Kennedy P, Lude P and Taylor N (2006) Quality of life, social participation, appraisals and coping post spinal cord injury: a review of four community samples. *Spinal cord* 44: 95-105.

Khambalia AZ, Dickinson S, Hardy LL, Gill T et al. (2012) A synthesis of existing systematic reviews and meta-analyses of school-based behavioural interventions for controlling and preventing obesity. *Obesity Reviews* 13:214–233.

Khantzian E J (2014) The self-medication hypothesis of addictive disorders: focus on heroin and cocaine dependence. *American Journal of Psychiatry* 142:1259-1264.

Kuo S F, Chuang W Y, Ng S, Chen C H et al. (2013) Pituitary gigantism presenting with depressive mood disorder and diabetic ketoacidosis in an Asian adolescent. *Journal of Pediatric Endocrinology and Metabolism* 26:945-948.

Labarthe A, Fiquet O, Hassouna R, Zizzari P et al. (2015). Ghrelin-derived peptides: a link between appetite/reward, GH axis, and psychiatric disorders?. *Neurological and Psychiatric Disorders in Endocrine Diseases*, 71.

Lamis D A, Malone P S, Langhinrichsen-Rohling J and Ellis T E (2015) Body investment, depression, and alcohol use as risk factors for suicide proneness in college students. *Crisis*.

- Lange K, Thamocharan S, Racine M, Hirko, C. et al. (2014) The relationship between weight and smoking in a national sample of adolescents: Role of gender. *Journal of Health Psychology* p.1359105313517275.
- Lehnert T, Sonntag D, Konnopka A, Riedel-Heller S et al. (2012) The long-term cost-effectiveness of obesity prevention interventions: systematic literature review. *Obesity Reviews* 13: 537-553.
- Ley R E (2010) Obesity and the human microbiome. *Current Opinion in Gastroenterology* 26: 5–11.
- Li S, Zhao JH, Luan J, Luben RN, Rodwell SA, et al. (2010) Cumulative effects and predictive value of common obesity-susceptibility variants identified by genome-wide association studies. *American Journal of Clinical Nutrition* 91: 184–190.
- Lorains F K, Cowlshaw S and Thomas S A (2011) Prevalence of comorbid disorders in problem and pathological gambling: Systematic review and meta-analysis of population surveys. *Addiction* 106: 490-498.
- Lo Verme J, Gaetani S, Fu J, Oveisi F et al. (2005) Regulation of food intake by oleoylethanolamide. *Cellular and Molecular Life Sciences* 62: 708-716.
- Loveman E, Frampton G K, Shepherd J, Picot J et al. (2011) The clinical effectiveness and cost-effectiveness of long-term weight management schemes for adults: a systematic review.
- McLaren L (2007) Socioeconomic status and obesity. *Epidemiologic reviews*, 29: 29-48.
- McEwen BS and Wingfield JC (2003) The concept of allostasis in biology and biomedicine. *Hormones and Behavior* 43:2–15.
- Maes HH, Neale MC and Eaves LJ (1997) Genetic and environmental factors in relative body weight and human adiposity. *Behavioral Genetics* 27:325–51.
- Magno da Costa Maranduba C et al. (2015). Intestinal microbiota as modulators of the immune system and neuroimmune system: impact on the host health and homeostasis. *Journal of Immunology Research* Article ID 931574.
- Maloy K J, and Powrie F (2011) Intestinal homeostasis and its breakdown in inflammatory bowel disease. *Nature*, 474(7351): 298-306.
- Maniam J and Morris M J (2012) The link between stress and feeding behaviour. *Neuropharmacology* 63: 97-110.
- Markey C N, August K J, Bailey L C, Markey P M et al. (2016) The pivotal role of psychology in a comprehensive theory of obesity. *Health Psychology Open* (this issue).
- Marks D F (2015) Homeostasis theory of obesity. *Health Psychology Open* January-June 2015 2: 2055102915590692, first published on June 29, 2015 doi:10.1177/2055102915590692
- Marsh P D (1994) Microbial ecology of dental plaque and its significance in health and disease.

Advances in Dental Research 8: 263-271.

Maynard L, Elson, C O, Hatton R D and Weaver, C T (2012) Reciprocal interactions of the intestinal microbiota and immune system. *Nature* 489 (7415): 231–241.

Meyer R M, Burgos-Robles A, Liu E, Correia S S et al. (2014) A ghrelin–growth hormone axis drives stress-induced vulnerability to enhanced fear. *Molecular Psychiatry* 19:1284-1294.

Monteiro C A, Moura E C, Conde W L and Popkin B M (2004). Socioeconomic status and obesity in adult populations of developing countries: a review. *Bulletin of the World Health Organization* 82: 940-946.

Morton G J, Cummings D E, Baskin D G, Barsh, G S et al. (2006). Central nervous system control of food intake and body weight. *Nature* 443(7109):289-295.

Mozaffarian D (2016). Dietary and policy priorities for cardiovascular disease, diabetes, and obesity: a comprehensive review. *Circulation*, CIRCULATIONAHA-115.

Müller T D, Nogueiras R, Andermann M L, Andrews Z B et al. (2015) Ghrelin. *Molecular Metabolism* 4: 437-460.

Neumark-Sztainer D and Eisenberg M E (2014). Body image concerns, muscle-enhancing behaviors, and eating disorders in males. *JAMA* 312:2156-2157.

Ng J, Ntoumanis N, Thøgersen-Ntoumani, E C, Deci et al. (2012) Self-Determination Theory applied to health contexts : A meta-analysis. *Perspectives on Psychological Science* 7:325-340.

Obuchowski K, Zienkiewicz H and Graczykowska-Koczorowska A (1970) Psychological studies in pituitary dwarfism. *Polish Medical Journal*, 9:1229-1235.

Park C L (2010) Making sense of the meaning literature: an integrative review of meaning making and its effects on adjustment to stressful life events. *Psychological Bulletin* 136:257-301.

Park C L (2013) The Meaning Making Model: A framework for understanding meaning, spirituality, and stress-related growth in health psychology. *The European Health Psychologist* 2: 40–47.

Parrott A C (1999) Does cigarette smoking cause stress?. *American Psychologist* 54: 817.

Pelletier L G, Dion S C , Slovenic-D'Angelo M and Reid R (2004) Why do you regulate what you eat? Relationship between forms of regulation, eating behaviors, sustained dietary behavior change, and psychological adjustment. *Motivation & Emotion* 28: 245- 277.

Pöykkö S M, Kellokoski E, Hörkkö S, Kauma, H et al. (2003). Low plasma ghrelin is associated with insulin resistance, hypertension, and the prevalence of type 2 diabetes. *Diabetes*, 52: 2546-2553.

Potthoff AD, Ellison G and Nelson L (1983) Ethanol intake increases during continuous administration of amphetamine and nicotine, but not several other drugs. *Pharmacology Biochemistry and Behavior* 18: 489-493.

Prochaska JJ and Benowitz NL (2016) The past, present, and future of nicotine addiction therapy. *Annual Review of Medicine* 67:4.1–4.20.

- Puig J, Englund M M, Simpson J A and Collins W A (2013). Predicting adult physical illness from infant attachment: A prospective longitudinal study. *Health Psychology* 32: 409.
- Reilly JJ, Armstrong J, Dorosty AR, Emmett PM et al. (2005) Early life risk factors for obesity in childhood: cohort study. *BMJ* 330(7504):1357.
- Remes L, Isoaho R, Vahlberg T, Viitanen M et al. (2010). Quality of life three years after major lower extremity amputation due to peripheral arterial disease. *Aging clinical and experimental research* 22: 395-405.
- Richards DW (1960) Homeostasis: its dislocations and perturbations. *Perspectives in Biology and Medicine* 3:238–251.
- Rose J E, Behm F M and Westman E C (2001) Acute effects of nicotine and mecamylamine on tobacco withdrawal symptoms, cigarette reward and ad lib smoking. *Pharmacology, Biochemistry and Behavior* 68: 187-97.
- Rosenbaum D and White K (2016) Understanding the complexity of biopsychosocial factors in the public health epidemic of overweight and obesity. *Health Psychology Open* (this issue).
- Rumsey N and Harcourt D (2004) Body image and disfigurement: issues and interventions. *Body Image* 1: 83-97.
- Russell M A (1990) The nicotine addiction trap: A 40 year sentence for four cigarettes. *British Journal of Addiction* 85: 293-300.
- Saper C B, Chou T C and Elmquist J K (2002) The need to feed: homeostatic and hedonic control of eating. *Neuron* 36: 199-211.
- Selye H (1946) The general adaptation syndrome and the diseases of adaptation. *Journal of Clinical Endocrinology and Metabolism* 6:117-230.
- Silva M N et al. (2011). Exercise autonomous motivation predicts 3-yr weight loss in women. *Medicine & Science in Sports and Exercise* 43: 728–737.
- Shaw K, O'Rourke P, Del Mar C, Kenardy J (2005) Psychological interventions for overweight or obesity. *Cochrane Database Systematic Reviews* 18:2(2).
- Sleddens S F, Gerards S M, Thijs C, De Vries N K et al. (2011). General parenting, childhood overweight and obesity-inducing behaviors: a review. *International Journal of Pediatric Obesity* 6(sup3):e12-27.
- Sominsky L and Spencer S J (2014) Eating behavior and stress: a pathway to obesity. *Frontiers in Psychology* 5: 1-8.
- Stock S, Leichner P, Wong A C, Ghatel M A et al. (2005) Ghrelin, peptide YY, glucose-dependent insulinotropic polypeptide, and hunger responses to a mixed meal in anorexic, obese, and control female adolescents. *The Journal of Clinical Endocrinology & Metabolism* 90: 2161-2168.

Sulzberger P and Marks D (1977) *The Isis Smoking Cessation Programme*. Dunedin: Isis Research Centre.

Swedish Council on Health Technology Assessment (2013) *Dietary treatment for obesity*. Stockholm:SBU.

Talge N M, Neal C and Glover V (2007) Antenatal maternal stress and long-term effects on child neurodevelopment: how and why? *Journal of Child Psychology and Psychiatry*, 48: 245-261.

Taveras EM, Rifas-Shiman SL, Belfort MB, Kleinman KP (2009) Weight status in the first 6 months of life and obesity at 3 years of age. *Pediatrics* 123:1177-83.

Taylor G, McNeill A, Girling A, Farley, A et al. (2014) Change in mental health after smoking cessation: systematic review and meta-analysis. *BMJ*, 348: 1151.

Thompson A and Kent G (2001) Adjusting to disfigurement: processes involved in dealing with being visibly different. *Clinical Psychology Review* 21: 663-682.

Tran BX, Ohinmaa A, Kuhle S, Johnson JA et al. (2014). Life course impact of school-based promotion of healthy eating and active living to prevent childhood obesity. *PLoS ONE* 9(7): e102242.

Tschop M, Weyer C, Tataranni PA, Devanarayan V et al. (2001) Circulating ghrelin levels are decreased in human obesity. *Diabetes* 50:707–709.

Tsutsumi A, Izutsu T, Islam M A, Amed J U et al. (2004) Depressive status of leprosy patients in Bangladesh: association with self-perception of stigma. *Leprosy review* 75: 57-66.

Turnbaugh P J and Gordon, J I (2009). The core gut microbiome, energy balance and obesity. *Journal of Physiology* 587: 4153–4158.

Urry H L, Van Reekum C M, Johnstone T, Kalin N H et al. (2006) Amygdala and ventromedial prefrontal cortex are inversely coupled during regulation of negative affect and predict the diurnal pattern of cortisol secretion among older adults. *The Journal of Neuroscience* 26: 4415-4425.

Vanderschuren L J and Everitt B J (2004) Drug seeking becomes compulsive after prolonged cocaine self-administration. *Science*, 305: 1017-1019.

Van Dijk SJ, Molloy PL, Varinli H, Morrison JL et al. (2015) Epigenetics and human obesity. *International Journal of Obesity* 39:85-97.

Van Vugt D A (2010). Brain imaging studies of appetite in the context of obesity and the menstrual cycle. *Human reproduction update* 16: 276-292.

Verstraeten R, Roberfroid D, Lachat C, Leroy JL et al. (2012) Effectiveness of preventive school based obesity interventions in low- and middle-income countries: a systematic review. *American Journal of Clinical Nutrition* 96: 415–438.

Veugelers PJ and Fitzgerald A L (2005) Effectiveness of school programs in preventing childhood obesity: a multilevel comparison. *American Journal of Public Health* 95: 432–435.

- Volkow N D and Fowler J S (2000) Addiction, a disease of compulsion and drive: involvement of the orbitofrontal cortex. *Cerebral Cortex*, 10: 318-325.
- Yanovski SZ and Yanovski JA (2014) Long-term drug treatment for obesity: a systematic and clinical review. *JAMA* 311:74-86.
- Weiss R D, Griffin M L and Mirin S M (1992). Drug abuse as self-medication for depression: An empirical study. *American Journal of Drug and Alcohol Abuse* 18:121-129.
- Whitehead E M, Shalet S M, Davies D, Enoch B A et al. (1982) Pituitary gigantism: a disabling condition. *Clinical Endocrinology* 17: 271-277.
- Woodhouse L J, Mukherjee A, Shalet S M and Ezzat S. (2006). The influence of growth hormone status on physical impairments, functional limitations, and health-related quality of life in adults. *Endocrine Reviews* 27: 287-317.
- Wurst F M, Graf I, Ehrental H D, Klein, S et al. (2007) Gender differences for ghrelin levels in alcohol-dependent patients and differences between alcoholics and healthy controls. *Alcoholism: Clinical and Experimental Research*, 31: 2006-2011.
- Zuckerman P (2009) Atheism, secularity, and well-being: How the findings of social science counter negative stereotypes and assumptions. *Sociology Compass* 3/6: 949-971.

TABLE 1: The Circle of Discontent for Diverse Conditions. For each issue the table lists publications providing evidence of causal links to negative affect (column 2), to dissatisfaction (column 3), and evidence of homeostasis in the form of self-medication, compensatory actions, coping or psychological adaptations (column 4).

Issue	Evidence of Causal Link to Depression	Evidence of Causal Link to Dissatisfaction	Evidence of Homeostasis
Disfigurement	Thompson & Kent (2001).	Rumsey & Harcourt (2004). Grogan (2006).	Rumsey & Harcourt (2004). Camouflaging.
Gigantism	Whitehead (1982). Kuo et al. (2013).	Woodhouse et al. (2006).	Hurxthal (1961). Self-imposed starvation diet in a young girl.
Dwarfism	Obuchowski (1970).	Bjork (1989).	Backstrom (2012). Positive identity advocacy.
Leprosy	Tsutsumi et al. (2004)	Tsutsumi et al. (2004)	Heijnders & Van Der Meij (2006). Stigma reduction/ Empowerment
Amputation	Horgan & MacLachlan (2004)	Remes et al. (2010)	Gallagher & MacLachlan (1999). Wearing a prosthetic limb.
Spinal cord injury	Elliott & Frank (1996)	Dijkers (1997). Kennedy, Lude & Taylor (2006)	Kennedy, Lude & Taylor (2006) Social integration.

Table 2: Determination of obesity at the main entry points to the Circle of Discontent.

Individual obesity risk accumulates over time with exposure to different periods of obesity risk. Percentage points can be allocated to each critical period to reflect its weighting as a determinant of obesity over an individual's lifetime. By the end of adolescence, the pathway to obesity has been 90% determined. Only the remaining 10% of the variance can be controlled by individual lifestyle components (diet, exercise).

Period	Evidence	Estimated percentage points	Accumulated percentage
Genetic predisposition	Maes, Neale and Eaves (1997)	50	50
Epigenesis	Van Dijk et al. (2015)	15 (?)	65
First few months of life	Reilly et al. (2005) Taveras et al. (2009)	15 (?)	80
Adolescence	Guo et al. (2002) Gordon-Larsen et al. (2004)	10	90
Adult lifestyle	Multiple	10	100

FIGURE 1 Model of hedonic/reward response to ghrelin after chronic stress in relation to anxiety and depression. During chronic stress, increased ghrelin secretion induces emotional eating as hedonic reward. Ghrelin has anxiolytic actions in response to stress; this adaptive response helps to control excessive anxiety and prevent depression. In obesity, a lower ability to mobilize ghrelin in response to stress or central ghrelin resistance at the level of the hedonic/reward system may explain the inability to cope with anxiety and increased susceptibility to depression. Reciprocally, people suffering from depressed show increased susceptibility to obesity or eating disorders (due to an altered hedonic/reward response). Elevated ghrelin may also contribute to alcohol/drug craving as higher ghrelin levels correlate with greater alcohol craving. Reproduced from LaBarthe et al. (2014).




FIGURE 2 The potential role of ghrelin in obesity dishomeostasis and the hedonic reward system in the amelioration of negative affect, chronic stress, anxiety and depression. The Circle of Discontent Theory is paralleled by the model of hedonic/reward response to ghrelin after chronic stress proposed by LaBarthe et al. (2014) (as previously shown in Figure 1). The original version of the model by LaBarthe et al. has been synced with the COD theory. Green arrows represent increased levels of ghrelin while red arrows represent decreased ghrelin activity, as shown in Figure 1. The part of the Circle of Discontent shown in light gray was not included in the LaBarthe et al. Model. The amended LaBarthe model is a perfect fit to the OD theory and gives the theory a neurochemical basis.




FIGURE 3 Model of the neurobiological basis of the Circle of Discontent. Disruption of homeostasis in feedback loops between self-consciousness/cognition mediated by the pre-frontal cortex, negative affect mediated by ghrelin signalling in amygdala, eating behaviour mediated by the HPA axis and increased visceral adiposity leading to increased ghrelin secretion. Ghrelin is a significant part of the signalling system throughout the systems of hedonic reward and homeostatic control of appetitive behaviour.




FIGURE 4 Interaction of homeostatic and hedonic control of food intake. Hormones from adipose tissue, the gastrointestinal tract and, in females, the ovary influence food intake through hypothalamic and extra-hypothalamic sites of action. Activation of the corticolimbic pathway, which includes the ventral tegmental area (VTA), nucleus accumbens (NAC), prefrontal cortex (PFC), hippocampus (Hippo) and amygdala (Amyg) are associated with the hedonic pleasure of eating and is influenced by hunger level, food palatability, experience and mood. From Van Vugt (2010), reproduced by permission.




FIGURE 5

Obesity Dyshomeostasis: Causing overweight and obesity, feedback loops required for equilibrium become disrupted by the hedonic reward system. In one system, the Circle of Discontent, the levels of body dissatisfaction, negative affect and high-energy consumption run out of control. Another system provides homeostasis for Motivation and Energy Mobilization (MEM). In the MEM system, lowered motivation causes reduced restraint, increased dietary intake and lowered activity, which cause reductions to subjective well-being, mobility, and positive affect. The whole complex establishes unhealthy eating habits, low activity levels, negative affect, overweight and obesity. Early lifetime stress which interferes with attachment can influence appetite, feeding behaviour and metabolism throughout life. People living with low levels of income will also be at high risk of negative affect, hedonic reward eating of fatty and sugary foods and, ultimately, obesity. Reproduced from Marks (2015).




FIGURE 6 Areas of the brain mediating food intake and drug seeking. Areas that are most critical for food intake are depicted in lighter shades and those areas most critical for drug reward and seeking are depicted in darker shades. Most areas have some influence on both food and drug intake, and the spectrum represents this overlap. The hypothalamus is critical for food intake and is modulated by the darker shaded areas. The VTA and NAc are critical for drug seeking and are modulated by many other brain areas. Inputs from cortex and amygdala provide control over both food- and drug-related behaviors. Arc, arcuate nucleus; LH, lateral hypothalamus; MHb, medial habenula. Red lines, inhibitory connections; dashed lines, indirect projections. Thicker lines indicate stronger connections. From DiLeone, Taylor and Picciotto (2012), reproduced with permission.




FIGURE 7

The Circle of Discontent in Addiction: Homeostatic reduction of negative affect and low satisfaction induce increased consumption, which increases habit strength through the positive reinforcement by hedonic reward and negative reinforcement by palliation of negative affect.




FIGURE 8 Multiple Circles of Discontent: A person addicted to nicotine, ethanol, cocaine, and gambling. This diagram shows a model of four concurrent addictions, each having a homeostatic system and Circle of Discontent. The green arrows indicate facilitation; the red arrows indicate temporary palliation of negative affect. The grey dashed lines indicate associative connections between habits where one acts as a trigger for the other. The brain areas mediating appetitive drug seeking and addictive behaviours may differ between addictions, but will include the areas shown in the previous Figure 5. The four addictions reinforce one other and the four homeostatic systems become closed off from external influence. The hedonic reward system serves to palliate negative affect by appetitive behaviours and controls all four of the addictions in an identical manner. The hedonic reward system overrides homeostasis, ultimately putting the individual at significant risk of different catastrophes including life-limiting conditions, mental illness, crime and bankruptcy. Not all pathways are included to avoid cluttering the diagram. The same person could well have several other addictions, (e.g. caffeine, licit or illicit drugs, gaming, sex, the internet) and the model could be extended to include them.





FIGURE 9 Life course projections of avoided health care costs for Canada and the province of Alberta (in million dollars) in light of a school-based obesity prevention programme (Reproduced from Tran et al., 2014, Figure 6).



Endnotes

1 A similar model, recently published by Hemmingsson (2014), discusses emotional distress in causing obesity: "... inner disturbances eventually cause a psycho-emotional overload, triggering a cascade of weight gain-inducing effects including maladaptive coping strategies such as eating to suppress negative emotions, chronic stress, appetite up-regulation, low-grade inflammation and possibly reduced basal metabolism. Over time, this causes obesity, circular causality and further weight gain " (Hemmingsson, 2014, p. 770).

2 In Self-Determination Theory, the term for non-autonomous motivation is 'Controlled Motivation'. Perhaps a more apposite term would be '*Uncontrolled Motivation*'.

Declaration of conflicting interests The author declares no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Acknowledgements

I warmly thank the commentators on the Homeostasis Theory of Obesity for their insights concerning the development of the theory: Rachel Annunziato, Kristin August, Lindzee Bailey, Laszlo Brassai, Emily Brindal, Janine Delahanty, Carlo DiClemente, Stephanie Grossman, Camille Guertin, Charlotte Markey, Patrick Markey, Jennifer Mills, Christopher Nave, Luc Pelletier, Bettina Piko, Paige Pope, Meredith Rocchi, Kaley Roosen, Diane Rosenbaum, Kamila White and Gary Wittert.