

# A Combining Forecasting Modeling and Its Application

Dengmei Qiu, Wenjie Wang

► **To cite this version:**

Dengmei Qiu, Wenjie Wang. A Combining Forecasting Modeling and Its Application. Kecheng Liu; Stephen R. Gulliver; Weizi Li; Changrui Yu. 15th International Conference on Informatics and Semiotics in Organisations (ICISO), May 2014, Shanghai, China. Springer, IFIP Advances in Information and Communication Technology, AICT-426, pp.246-252, 2014, Service Science and Knowledge Innovation. <10.1007/978-3-642-55355-4\_25>. <hal-01350930>

**HAL Id: hal-01350930**

**<https://hal.inria.fr/hal-01350930>**

Submitted on 2 Aug 2016

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



# A Combining Forecasting Modeling and Its Application

Dengmei Qiu, WenJie Wang\*

Glorious Sun School of Business and Management, Donghua University  
Shanghai, 200051, China  
dmqiu@dhu.edu.cn, wenjiew@dhu.edu.cn

**Abstract.** The supply chain coordination has abstracted more and more attention from industries and academics. This paper studies a Bayesian combination forecasting model to integrate multiple forecasting resources and coordinate forecasting process among partners in retail supply chain. The simulation results based on the retail sales data show the effectiveness of this Bayesian combination forecasting model to coordinate the collaborative forecasting process. This Bayesian combination forecasting model can improve demand forecasting accuracy of supply chain.

**Keywords:** Combination forecasting, Bayesian model, Supply chain

## 1. Introduction

With the development of information and network technology, various innovative supply chain solutions are created. Collaborative planning, forecasting and replenishment (CPFR), which is an retailing supply chain coordination innovation based on the network technology, has been adopted and implemented by many world-renowned retailers and manufacturers, such as Wal-Mart, Proctor & Gamble, etc.. CPFR concerns the collaboration where two or more parties in the supply chain jointly plan a number of promotional activities and work out synchronized forecasts, on the basis of which the production and replenishment processes are determined [1]. The first CPFR project was piloted by Wal-Mart with its suppliers in 1995. The results of two-year project showed that CPFR could simultaneously reduce inventory levels and increase sales for both retailers and suppliers. Since its original application was initiated, CPFR has had many successful applications in North America, Europe and China [2].

The collaborative forecasting plays an important part in CPFR implementation procedure. We will briefly review the CPFR concept and its implementation process at the beginning of this paper. And then, the collaborative forecasting process which is the core part of CPFR will be mainly discussed. As the basics phase of the implementation of CPFR, the collaborative forecasting process is the cornerstone to the success of CPFR projects. The collaborative forecasting process of CPFR requires a solid forecasting approach to synthesis information and knowledge from multiply parties in the supply chain. The combination forecasting method can combines

forecasting models from different parties to smooth coordination in the supply chain and reduce forecasting discrepancies. Thus, considering the multiple forms of forecasting resources in the retail supply chain, the Bayesian combination forecasting method is applied for CPFR collaborative forecasting modeling with improved forecasting accuracy and supply chain collaboration performance in this paper.

Combining forecasts is a well-established procedure for improving forecasting accuracy which takes advantage of the availability of both multiple information and computing resources for data intensive forecasting [3]. Since Bates-Granger first proposed the combination forecasting method in 1969 [4], many kinds of combining methods have been developed [3]. Zeng et al [5] studied the combination forecasting model with error correction and changing weight coefficient. Hoogerheide et al [6] compared several Bayesian combination schemes in terms of forecast accuracy and economic gains. Bayesian combination methods use the distributional properties of the individual forecasts to construct the combination. The demand forecasting in retail supply chain is impacted by many factors such as product promotion or social development trend. And, subjective forecasting based on the expert experiments is often used in retail market forecasting. So, the Bayesian combination forecasting model is considered as the collaborative forecasting approach in retail supply chain coordination.

In the first part of this paper, the CPFR retail supply chain coordination and collaborative forecasting process are discussed briefly. In the second part of the paper, a Bayesian combination forecasting method is modeled to coordinate forecasting process in retail supply chain. Finally, the simulation for this model is completed based on the Carrefour sales data. The simulation results showed the effective of this Bayesian combination forecasting model in retail supply chain collaboration process.

## 2. Bayesian Combination Forecasting Modeling for Collaborative Forecasting Process

The combination method proposed by Bates and Branger in 1969 is normally called as optimal linear combination model or B-G method. The forecasting results  $f_{c1}, f_{c2}, \dots, f_{cn}$  are supposed as random variables with the covariance matrix  $\Sigma$  in this model. Based on the minimizing the variance criteria (MV), the optimal forecasting results can be calculated as the following formula (1)

$$f_c = \sum_{k=1}^n W_k f_k = W^T \cdot f \quad (1)$$

Here, weighting vector  $W^T = (1_n^T \Sigma^{-1} 1_n)^{-1} 1_n^T \Sigma^{-1} = (W_1, W_2, \dots, W_n)$ ,  $1_n^T = (1, 1, \dots, 1)$ .

The Bayesian combination forecasting model is developed based on the B-G method and uses the distributional properties of the individual forecasts to construct the combination. Supposed the Y is presenting the samples of actual demand. The forecasting results obtained from the different parties in the supply chain which are

forecasted with  $m$  kinds of individual forecasting methods are presented by  $f_1, f_2, \dots, f_m$  ( $j=1, 2, \dots, m$ ).  $i=1, 2, \dots, n$  present different forecasting time periods. The Bayesian combination forecasting method makes use of the Bayesian rule to decide the optimal combination ways and weights of individual forecasting methods in combination model and get the combined forecasting results in different time periods  $\hat{f}_{ic}$  ( $i=1, 2, \dots, n$ ), which can be fully approximated to actual demand values.

Set  $Z_i^T = (f_1, f_2, \dots, f_m)$ ,  $i=1, 2, \dots, n$ , and then the joint probability density function of  $m$  individual forecasting samples on independent time  $z_1, z_2, \dots, z_n$  can be calculated as follows.

$$f(z_1, z_2, \dots, z_n | Y; \varphi) = \prod_{i=1}^n f(z_i | Y_i; \varphi) \quad (2)$$

Here  $\varphi$  is parameter vector, and  $Y^T = (y_1, y_2, \dots, y_n)$  is the vector of the actual demand samples on the different forecasting time periods.

According to the Bayesian rule, the probability density function that  $Y$  is specified as a vector value  $y$  is as follows:

$$\begin{aligned} f(y | z_1, z_2, \dots, z_n) &= \frac{\prod_{i=1}^n f(z_i | y_i) f(y_i)}{\prod_{i=1}^n \int_{A_i} f(z_i | y_i) f(y_i) dy_i} \\ &= \prod_{i=1}^n \frac{f(f_{i1}, f_{i2}, \dots, f_{im} | y_i) f(y_i)}{\int_{A_i} f(f_{i1}, f_{i2}, \dots, f_{im} | y_i) f(y_i) dy_i} \end{aligned} \quad (3)$$

Here  $f(y_i)$  ( $i=1, 2, \dots, n$ ) is the prior probability distribution of  $y_i$ , which presents the prior estimation or preference of decision maker to  $y_i$ .  $A_i$  is the definition set of  $y_i$ .

If only one forecasting time period is considered as the general case and prior distribution  $f(y)$  is uniform distribution form, that is  $f(y) \propto 1$ , the formula (2) can be simplified as the following.

$$f(y | f_1, f_2, \dots, f_m) = \frac{f(f_1, f_2, \dots, f_m | y)}{\int_A f(f_1, f_2, \dots, f_m | y) dy} \quad (4)$$

The forecasting error distribution of individual forecasting results  $(f_1, f_2, \dots, f_m)$  is chosen as normal distribution or logarithm normal distribution in most case. In this paper, more general distribution of forecasting error is introduced through *Box-Cox* conversion.

$$Z^{(\lambda)} = \begin{cases} \frac{Z^\lambda - 1}{\lambda} & \lambda \neq 0 \\ \ln Z & \lambda = 0 \end{cases} \quad (5)$$

Here,  $\lambda$  is conversion parameter

Supposed  $\Sigma$  is the covariance matrix of  $(f_1, f_2, \dots, f_m)$  and  $W_i$  presents the weights of individual forecasting  $f_i$ . Then, weights can be calculated as following formula based on the minimum error variance criteria:

$$W^T = \left( \mathbf{1}_m^T \Sigma^{-1} \mathbf{1}_m \right)^{-1} \mathbf{1}_m^T \Sigma^{-1} = (W_1, W_2, \dots, W_m) \quad (6)$$

And, nonlinear combination forecasting formula can be obtained through Bayesian analysis as follows, which will be used to calculate the optimal combination forecasting results approximated to the actual values  $Y$ .

$$\hat{f}_c = \begin{cases} \prod_{i=1}^m f_i^{w_i} \times e^{\frac{s^2}{2}} & \lambda = 0 \\ \left[ 1 + \sum_{i=1}^m \lambda W_i f_i^{(\lambda)} \right]^{\frac{1}{\lambda}} = \left[ \sum_{i=1}^m W_i f_i^\lambda \right]^{\frac{1}{\lambda}} & \lambda \neq 0 \end{cases} \quad (7)$$

Here,  $\frac{1}{S^2} = \mathbf{1}_m^T \Sigma^{-1} \mathbf{1}_m = \frac{1}{|\Sigma|} \mathbf{1}_m^T \Sigma^{-1} \mathbf{1}_m$ , and  $\sum_{i=1}^m W_i = 1$

And, the optimal conversion parameter  $\lambda^*$  can be calculated as the following formula.

$$\lambda^* = \arg \min_{\lambda} \sum_{j=1}^t \left[ Y_j - \left( \sum_{i=1}^m W_i f_{ij}^\lambda \right)^{\frac{1}{\lambda}} \right]^2 \quad (8)$$

Here,  $f_{ij}$  present the forecasting result on  $j$  time period by  $i$  individual forecasting methods. The weights  $W_i$  can be calculated by formula (4). The covariance matrix  $\Sigma$  can be determined by prior value or estimated by past proximate samples values.

### 3. Simulation

The simulation of the Bayesian combination forecasting model will be based on the sales data of one kind of biscuit product in Carrefour China in this paper. The detailed sales data of this biscuit product in 39 weeks in Carrefour China is showed in table 1. During the simulation process, the sales data from week 1 to week 28 are used to estimate parameters in Bayesian combination forecasting model creation. The sales data from week 29 to week 39 are used to compare with the forecasting results obtained from combination forecasting methods.

**Table 1** The Sales Data of Biscuit in Carrefour Supermarket

Week	1	2	3	4	5	6	7	8	9	10	11	12	13
Demand	64	78	62	38	59	13	99	82	105	56	56	88	93
Week	14	15	16	17	18	19	20	21	22	23	24	25	26
Demand	100	110	135	124	105	86	117	172	185	169	192	195	156
Week	27	28	29	30	31	32	33	34	35	36	37	38	39
Demand	136	132	183	84	92	119	137	107	80	218	167	156	170

Based on the Carrefour biscuit sale data and statics analysis, the Bayesian forecasting model can be created following three main steps, which includes proper individual forecasting methods selection, combination ways determination and optimal parameter estimation. The characteristics of the individual forecasts combining in the model has substantial implications on the overall forecasting performance of model, and thus it is very important to make a rigorous analysis on the individual forecast errors. The first step of combination modeling is to compare and select proper individual forecasting methods for combination.

**Table 2** Forecasting Results of Five Individual Forecasting Methods

Week	Actual Demand	F1	F2	F3	F4	F5
29	183	141.3333	138.9865	179.2469	136.3747	125.3344
30	84	150.3333	165.3946	184.8024	173.2965	159.2878
31	92	133	116.5578	190.4484	112.6916	68.3652
32	119	119.6667	101.8231	196.185	93.30308	111.0044
33	137	98.3333	112.1293	202.0121	115.4397	106.5959
34	107	116	127.0517	207.9297	135.7434	117.6981
35	80	121	115.0207	213.9379	118.0299	93.1682
36	218	108	94.0083	220.0367	90.26195	82.5842
37	167	135	168.4033	226.2259	186.0273	177.1399
38	156	155	167.5613	232.5058	183.4052	115.0543
39	170	180.3333	160.6245	238.8761	162.1665	150.9677

In general, different parties in the retail supply chain may use the different patterns of individual forecasting. So, the different patterns of individual forecasting methods, which include the simple moving average, the exponential smoothing, the trend extrapolation method, ARIMA (autoregressive integrated moving average) method and artificial neural network method, are applied to combine model to forecast Carrefour biscuit demand from week 29 to week 39. Through comparison study of forecasting results of each individual forecasting method, the best parameters of each individual forecasting method are estimated. The forecasting results of five different individual forecasting methods are indicated in Table 2. The F1 row data indicated the best result forecasted by the simple moving average method when moving period  $N$  equal to 3. The F2 row data indicated the best result forecasted by the exponential smoothing method when smoothing coefficient  $a$  equal to 0.6. The F3 row data indicated the best result forecasted by the two polynomial regression method ( $n=2$ ). The F4 row data indicated the best result forecasted by the ARIMA method when parameter  $d$  equal to 1. The F5 row data indicated the best result forecasted by the artificial neural network method when there are three neurons and two hidden layers in the neural network.

After the five individual forecasting methods selected, the second step of combination modeling is to determine proper combination ways of these individual methods. With the Matlab simulation tool, the forecasting results are calculated using different combination ways of these individual forecasting methods. As it is widely accepted that only "good" forecasts should be included in a combination, strong differences in forecast error variances between the individual forecasts are not to be expected [7]. Four individual methods (F1, F2, F4, F5) which have good forecasting performance is decided to be combined into Bayesian combination model.

The optimal conversion parameter  $\lambda^*$  can be calculated according to the formula (8). The sum of square of forecasting error is minimized when  $\lambda^* = 6.9419$ . The optimal conversion parameter  $\lambda^*$  could be simplified as  $\lambda^*=7$  in the Bayesian combination modeling. The forecasting error between forecasting result and actual demand are used as the evaluation standards of forecasting methods performance. There are many kinds of measure indexes of forecasting errors [8]. In this paper, four main measure indexes of forecasting error, which are the squares sum error (SSE), the mean square error (MSE), the mean absolute percentage error (MAPE) and the mean square percentage error (MSPE), are applied to comprehensively evaluate the forecasting accuracy of Bayesian combination forecasting model.

**Table 3** The Comparison of Bayesian Combination Models Forecasting Accuracy

Combination	Forecasting	SSE	MSE	MAPE	MSPE
	Simple Average Method	2.64E+04	2.40E+03	0.3589	0.3725
	Optimal Linear Method	2.52E+04	2.29E+03	0.2734	0.3416
	Bayesian Combination $\lambda^*=7$	1.92E+03	1.74E+02	0.0803	0.0561

The Bayesian combination models with different parameters  $\lambda$  are compared with the simple average method and optimal linear methods in the simulation based

on Carrefour biscuit sale data. The forecasting errors of different combination forecasting methods are showed in the table 3. It can be found that the four measure indexes of forecasting error of optimal Bayesian combination method when  $\lambda^* = 7$  are lower than those of simple combination methods. So, the optimal Bayesian combination forecasting model performs better than other combination models in retail collaborative forecasting process. This simulation research proved that the optimal Bayesian combination forecasting method is an effective approach to integrate and coordinate the forecasting process among partners in retail supply chain. Bayesian combination forecasting method can highly improve the demand forecasting accuracy of collaborative forecasting activity in the retail supply chain.

#### 4. Conclusion

The collaborative planning, forecasting and replenishment framework provides the practical roadmap for retail supply chain coordination. The collaborative forecasting is taken as the core part in CPFR solution implementation. A Bayesian combination forecasting method, which can combine individual forecasting methods from different parties in the retail supply chain, is modeled for CPFR collaborative forecasting process. The simulation results showed that forecasting discrepancies are reduced and collaborative forecasting accuracy is improved after integrating forecasting process with the optimal Bayesian combination forecasting model. It is turned out that the Bayesian combination forecasting method is an effective means for collaborative forecasting process in retail supply chain. The further research on collaborative forecasting methodology for supply chain coordination will be extended into different statistic features of product demand situation in the future.

**Acknowledgments.** This research was supported by a grant from the Shanghai Science Foundation Council (12ZR1400900) and the Chinese National Science Foundation Council (71172174).

#### References

1. Liu, X., Sun, Y., 2012. Information Integration of CPFR in Inbound Logistics of Automotive Manufactures Based on Internet of Things. *Journal of Computers*, 7(2):349-355.
2. Danese, P., 2007. Designing CPFR collaborations: insights from seven case studies, *International Journal of Operations & Production Management*, 27(2): 181 - 204
3. Bunn, D.W., 1989. Forecasting with more than one model. *Journal of Forecasting*, 8(3), 161-166.
4. Bates, J.M., Greanger, C. W. J, 1969. Combination of Forecasts, *Operational Research Quarterly*, 20(4): 451-468
5. Zeng, Y., Tang, X., Zheng, W., Combination forecasting based on Stein-rule



estimation and error correction, *Journal of Management Sciences in China*, 2001, 4 (6):39-47.

6. Hoogerheide, L., Kleijn, R., Ravazzolo, F., Van Dijk, H.K., Verbeek, M., Forecast Accuracy and Economic Gains from Bayesian Model Averaging Using Time-Varying Weights, *Journal of Forecasting*, 2010, 29(1-2): 251-269,
7. Menezes, L.M. de, Bunn, D. W., Taylor, J. W. 2000. Review of guidelines for the use of combined forecasts. *European Journal of Operational Research*, 120(1):190-204
8. Zhang, C., Huang, L., Zhao, Z., 2013. Research on combination forecast of port cargo throughput based on time series and causality analysis. *Journal of Industrial Engineering and Management*, 6(1):124-134