



The inapproximability for the $(0, 1)$ -additive number

Arash Ahadi, Ali Dehghan

► **To cite this version:**

Arash Ahadi, Ali Dehghan. The inapproximability for the $(0, 1)$ -additive number. Discrete Mathematics and Theoretical Computer Science, DMTCS, 2016, Vol. 17 no. 3 (3), pp.217-226. <hal-01352839>

HAL Id: hal-01352839

<https://hal.inria.fr/hal-01352839>

Submitted on 16 Aug 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The inapproximability for the $(0, 1)$ -additive number

Arash Ahadi¹Ali Dehghan^{2*}¹ Sharif University of Technology, Tehran, Iran² Carleton University, Ottawa, Ontario, Canadareceived 27th Apr. 2013, revised 15th Apr. 2016, accepted 8th June 2016.

An *additive labeling* of a graph G is a function $\ell : V(G) \rightarrow \mathbb{N}$, such that for every two adjacent vertices v and u of G , $\sum_{w \sim v} \ell(w) \neq \sum_{w \sim u} \ell(w)$ ($x \sim y$ means that x is joined to y). The *additive number* of G , denoted by $\eta(G)$, is the minimum number k such that G has an additive labeling $\ell : V(G) \rightarrow \mathbb{N}_k$. The *additive choosability* of a graph G , denoted by $\eta_\ell(G)$, is the smallest number k such that G has an additive labeling for any assignment of lists of size k to the vertices of G , such that the label of each vertex belongs to its own list.

Seamone in his PhD thesis conjectured that for every graph G , $\eta(G) = \eta_\ell(G)$. We give a negative answer to this conjecture and we show that for every k there is a graph G such that $\eta_\ell(G) - \eta(G) \geq k$.

A $(0, 1)$ -*additive labeling* of a graph G is a function $\ell : V(G) \rightarrow \{0, 1\}$, such that for every two adjacent vertices v and u of G , $\sum_{w \sim v} \ell(w) \neq \sum_{w \sim u} \ell(w)$. A graph may lack any $(0, 1)$ -additive labeling. We show that it is **NP**-complete to decide whether a $(0, 1)$ -additive labeling exists for some families of graphs such as perfect graphs and planar triangle-free graphs. For a graph G with some $(0, 1)$ -additive labelings, the $(0, 1)$ -additive number of G is defined as $\sigma_1(G) = \min_{\ell \in \Gamma} \sum_{v \in V(G)} \ell(v)$ where Γ is the set of $(0, 1)$ -additive labelings of G . We prove that given a planar graph that admits a $(0, 1)$ -additive labeling, for all $\varepsilon > 0$, approximating the $(0, 1)$ -additive number within $n^{1-\varepsilon}$ is **NP**-hard.

Keywords: Additive labeling, additive number, lucky number, $(0, 1)$ -additive labeling, $(0, 1)$ -additive number, Computational complexity.

1 Introduction

Throughout the paper we denote $\{1, 2, \dots, k\}$ by \mathbb{N}_k . An *additive labeling* of a graph G , which was introduced by Czerwiński et al. [11], is a function $\ell : V(G) \rightarrow \mathbb{N}$, such that for every two adjacent vertices v and u of G , $\sum_{w \sim v} \ell(w) \neq \sum_{w \sim u} \ell(w)$ ($x \sim y$ means that x is joined to y). The *additive number* of G , denoted by $\eta(G)$, is the minimum number k such that G has an additive labeling $\ell : V(G) \rightarrow \mathbb{N}_k$. Initially, additive labeling is called a lucky labeling of G . The following important conjecture was proposed by Czerwiński et al. [11].

*Corresponding author.

E-mail Addresses: arash_ahadi@mehr.sharif.edu (Arash Ahadi) alidehghan@sce.carleton.ca (Ali Dehghan).

1365–8050 © 2016 Discrete Mathematics and Theoretical Computer Science (DMTCS), Nancy, France

Conjecture 1 [Additive Coloring Conjecture [11]] For every graph G , $\eta(G) \leq \chi(G)$.

Czerwiński et al. also, considered the list version of above problem [11]. The *additive choosability* of a graph G , denoted by $\eta_\ell(G)$, is the smallest number k such that G has an additive labeling from any assignment of lists of size k to the vertices of G . Idem above, about list-coloring proved that if T is a tree, then $\eta_\ell(T) \leq 2$, and if G is a bipartite planar graph, then $\eta_\ell(G) \leq 3$ (for more information about the recent results see [9]). Seamone in his Ph.D dissertation posed the following conjecture about the relationship between additive number and additive choosability [23].

Conjecture 2 [Additive List Coloring Conjecture [23]] For every graph G , $\eta(G) = \eta_\ell(G)$.

For a given connected graph G with at least two vertices, if no two adjacent vertices have a same degree, then $\eta(G) = 1$ and $\eta_\ell(G) > 1$. We show that not only there exists a counterexample for the above equality but also the difference between $\eta(G)$ and $\eta_\ell(G)$ can be arbitrary large.

Theorem 1 For every k there is a graph G such that $\eta(G) \leq k \leq \eta_\ell(G)/2$.

Chartrand et al. introduced another version of additive labeling and called it sigma coloring [10]. For a graph G , let $c : V(G) \rightarrow \mathbb{N}$ be a vertex labeling of G . If for every two adjacent vertices v and u of G , $\sum_{w \sim v} c(w) \neq \sum_{w \sim u} c(w)$, then c is called a *sigma coloring* of G . The minimum number of labels required in a sigma coloring is called the *sigma chromatic number* of G and is denoted by $\sigma(G)$. Chartrand et al. proved that, for every graph G , $\sigma(G) \leq \chi(G)$ [10]. Note that the only difference between additive labeling and sigma coloring is the objective function, but the feasible labelings are the same.

Additive labeling and sigma coloring have been studied extensively by several authors, for instance see [3, 4, 6, 8, 10, 11, 13, 21, 22]. It is proved, in [3] that it is **NP**-complete to determine whether a given graph G has $\eta(G) = k$ for any $k \geq 2$. Also, it was shown that, it is **NP**-complete to decide for a given planar 3-colorable graph G , whether $\eta(G) = 2$ [3]. Furthermore, it was proved that, it is **NP**-complete to decide for a given 3-regular graph G , whether $\eta(G) = 2$ [13].

The edge version of additive labeling was introduced by Karoński, Łuczak and Thomason [18]. They introduced an edge-labeling which is additive vertex-coloring that means for every edge uv , the sum of labels of the edges incident to u is different from the sum of labels of the edges incident to v [18]. It is conjectured that three integer labels \mathbb{N}_3 are sufficient for every connected graph, except K_2 [18]. Currently the best bound is 5 [17]. This labeling has been studied extensively by several authors, for instance see [1, 2, 5, 14, 19, 20].

A *clique* in a graph $G = (V, E)$ is a subset of its vertices such that every two vertices in the subset are connected by an edge. The *clique number* $\omega(G)$ of a graph G is the number of vertices in a maximum clique in G . There is no direct relationship between the additive number and the clique number of graphs. For any natural number ω there exists a graph G , such that $\omega(G) = \omega$ and $\eta(G) = 1$. To see this for given number ω , consider a graph G with the set of vertices $V(G) = \{v_i | i \in \mathbb{N}_\omega\} \cup \{u_{i,j} | i, j \in \mathbb{N}_\omega, j < i\}$ and the set of edges $E(G) = \{v_i v_j | i \neq j\} \cup \{v_i u_{i,j} | i, j \in \mathbb{N}_\omega, j < i\}$.

Theorem 2 We have the following:

- (i) For every graph G , $\eta(G) \geq \frac{\omega}{n-\omega+1}$.
- (ii) If G is a regular graph and $\omega > \frac{n+4}{3}$, then $\eta(G) \geq 3$.

A $(0, 1)$ -additive labeling of a graph G is a function $\ell : V(G) \rightarrow \{0, 1\}$, such that for every two adjacent vertices v and u of G , $\sum_{w \sim v} \ell(w) \neq \sum_{w \sim u} \ell(w)$. A graph may lack any $(0, 1)$ -additive labeling. It was

proved that, it is **NP**-complete to decide for a given 3-regular graph G , whether $\eta(G) = 2$ [13]. So, it is **NP**-complete to decide whether a (0, 1)-additive labeling exists for a given 3-regular graph G . In this paper, we study the computational complexity of (0, 1)-additive labeling for perfect graphs and planar graphs.

A graph G is called *perfect* if $\omega(H) = \chi(H)$ for every induced subgraph H of G . Here, we show that it is **NP**-complete to decide whether a (0, 1)-additive labeling exists for perfect graphs.

Theorem 3 *The following problem is NP-complete: Given a perfect graph G , does G have any (0, 1)-additive labeling?*

Next, we show that it is **NP**-complete to decide whether a (0, 1)-additive labeling exists for planar triangle-free graphs.

Theorem 4 *It is NP-complete to determine whether a given planar triangle-free graph G has a (0, 1)-additive labeling.*

For a graph G with some (0, 1)-additive labelings, the (0, 1)-additive number of G is defined as $\sigma_1(G) = \min_{\ell \in \Gamma} \sum_{v \in V(G)} \ell(v)$ where Γ is the set of (0, 1)-additive labelings of G . For a given graph G with a (0, 1)-additive labeling ℓ the function $f(v) = 1 + \sum_{w \sim v} \ell(w)$ is a proper vertex coloring, so we have the following trivial lower bound for $\sigma_1(G)$.

$$\chi(G) - 1 \leq \sigma_1(G).$$

We prove that given a planar graph that admits a (0, 1)-additive labeling, for all $\varepsilon > 0$, approximating the (0, 1)-additive number within $n^{1-\varepsilon}$ is **NP**-hard.

Theorem 5 *If $\mathbf{P} \neq \mathbf{NP}$, then for any constant $\varepsilon > 0$, there is no polynomial-time $n^{1-\varepsilon}$ -approximation algorithm for finding $\sigma_1(G)$ for a given planar graph with at least one (0, 1)-additive labeling.*

For $v \in V(G)$ we denote by $N(v)$ the set of neighbors of v in G . Also, for every $v \in V(G)$, the degree of v is denoted by $d(v)$. We follow [16, 24] for terminology and notation not defined here, and we consider finite undirected simple graphs $G = (V, E)$.

2 Counterexample

Proof of Theorem 1: For every k we construct a graph G such that $\eta_\ell(G) - \eta(G) \geq k$. For every α , $\alpha \in \mathbb{N}_{2k-1}$ consider a copy of complete graph $K_{2k}^{(\alpha)}$, with the vertices $\{x_\beta^\alpha : \beta \in \mathbb{N}_k\} \cup \{y_\beta^\alpha : \beta \in \mathbb{N}_k\}$. Next, consider an isolated vertex t and join every vertex y_β^α to t . Call the resulting graph G . First, note that in every additive labeling ℓ of G , for every (i, j) , where $i < j$ and $i, j \in \mathbb{N}_k$ we have $\sum_{z \in N(x_i^1)} \ell(z) \neq \sum_{z \in N(x_j^1)} \ell(z)$, thus $\ell(x_i^1) \neq \ell(x_j^1)$ (because all the neighbors of x_i^1 and x_j^1 are common except x_i^1 as a neighbor of x_j^1 , and vice versa). Therefore $\ell(x_1^1), \ell(x_2^1), \dots, \ell(x_k^1)$ are k distinct numbers, that means $\eta(G) \geq k$. Define (for every α and β):

$$\begin{aligned} \ell : V(G) &\rightarrow \mathbb{N}_k, \\ \ell(x_\beta^\alpha) &= \ell(y_\beta^\alpha) = \beta, \\ \ell(t) &= k. \end{aligned}$$

It is easy to see that ℓ is an additive labeling for G . Next, we show that $\eta_\ell(G) > 2k - 1$. Consider the following lists for the vertices of G (for every α and β).

$$\begin{aligned} L(x_\beta^\alpha) &= \mathbb{N}_{2k-1}, \\ L(y_\beta^\alpha) &= \{i + \alpha : i \in \mathbb{N}_{2k-1}\}, \\ L(t) &= \mathbb{N}_{2k-1}. \end{aligned}$$

To the contrary suppose that $\eta_\ell(G) \leq 2k - 1$ and let ℓ be an additive labeling from the above lists.

Suppose that $\ell(t) = r$. Consider the complete graph $K_{2k}^{(r)}$, for every β we have:

$$\begin{aligned} L(x_\beta^r) &= \mathbb{N}_{2k-1}, \\ L(y_\beta^r) &= \{i + r : i \in \mathbb{N}_{2k-1}\}. \end{aligned}$$

Now, consider the following partition for $\mathbb{N}_{2k-1} \cup \{i + r : i \in \mathbb{N}_{2k-1}\}$,

$$\{1 + r, 1\}, \{2 + r, 2\}, \dots, \{2k - 1 + r, 2k - 1\}.$$

By Pigeonhole Principle, there are indices i, n and m such that $\ell(x_m^r), \ell(y_n^r) \in \{i + r, i\}$, so $\ell(x_m^r) = i$ and $\ell(y_n^r) = i + r$. Therefore, $\sum_{z \in N(x_m^r)} \ell(z) = \sum_{z \in N(y_n^r)} \ell(z)$. This is a contradiction, so $\eta_\ell(G) \geq 2k$. \square

3 Lower bounds

Proof of Theorem 2: (i) Let $\ell : V(G) \rightarrow \mathbb{N}_k$ be an additive labeling of G and suppose that $T = \{v_i | i \in \mathbb{N}_w\}$ is a maximum clique in G . For each vertex $v \in T$, define the function Y_v .

$$Y_v \stackrel{\text{def}}{=} \sum_{\substack{x \in V(G) \setminus T \\ x \sim v}} l(x) - l(v).$$

For every two adjacent vertices v and u in T , we have:

$$\begin{aligned} \sum_{x \sim v} l(x) &\neq \sum_{x \sim u} l(x), \\ \sum_{\substack{x \notin T \\ x \sim v}} l(x) + \sum_{\substack{x \in T \\ x \neq v}} l(x) &\neq \sum_{\substack{x \notin T \\ x \sim u}} l(x) + \sum_{\substack{x \in T \\ x \neq u}} l(x), \\ \sum_{\substack{x \notin T \\ x \sim v}} l(x) + l(v) &\neq \sum_{\substack{x \notin T \\ x \sim u}} l(x) + l(u), \\ Y_v &\neq Y_u. \end{aligned}$$

Thus, Y_{v_1}, \dots, Y_{v_w} are distinct numbers. On the other hand, for each vertex $v \in T$, the image of the function Y_v is $[-k, k(n-w)-1]$. So $w \leq k(n-w+1)$, therefore $k \geq \frac{w}{n-w+1}$ and the proof is completed.

(ii) Let G be a regular graph, obviously $\eta(G) \geq 2$. To the contrary suppose that $\eta(G) = 2$. Let T be a maximum clique in G and $c : V(G) \rightarrow \{1, 2\}$ be an additive labeling of G . Define:

$$\begin{aligned} X_1 &= c^{-1}(1) \cap T, & X_2 &= c^{-1}(2) \cap T, \\ Y_1 &= c^{-1}(1) \setminus T, & Y_2 &= c^{-1}(2) \setminus T. \end{aligned}$$

Suppose that $X_1 = \{v_1, \dots, v_k\}$ and $X_2 = \{v_{k+1}, \dots, v_w\}$. For each $i \in \mathbb{N}_w$, denote the number of neighbors of v_i , in Y_1 by d_i . Since c is an additive labeling of the regular graph, every two adjacent vertices have different numbers of neighbors in $c^{-1}(1)$. Therefore $d_1, \dots, d_k, 1 + d_{k+1}, \dots, 1 + d_w$ are distinct numbers. Since for each $i \in \mathbb{N}_w$, $0 \leq d_i \leq |Y_1|$, we have $|Y_1| \geq w - 2$. Similarly, $|Y_2| \geq w - 2$, so

$$n = |T| + |Y_1| + |Y_2| \geq 3\omega - 4.$$

This is a contradiction. So the proof is completed. □

4 List Coloring Problem

Proof of Theorem 3: Let G be a graph and let L be a function which assigns to each vertex v of G a set $L(v)$ of positive integers, called the list of v . A proper vertex coloring $c : V(G) \rightarrow \mathbb{N}$ such that $f(v) \in L(v)$ for all $v \in V$ is called a *list coloring* of G with respect to L , or an L -coloring, and we say that G is L -colorable.

Next, for a given graph G and a list $L(v)$ for every vertex v , we construct a graph H_G such that H_G has a (0, 1)-additive labeling if and only if G is L -colorable.

Define $W = \bigcup_{v \in V(G)} L(v)$ and let f be a bijective function from the set W to the set $\mathbb{N}_{|W|+1} \setminus \{1\}$. For every vertex $v \in V(G)$, let $L_f(v) = \{f(i) | i \in L(v)\}$. The graph G is L -colorable if and only if G is L_f -colorable. Now, we construct H_G from G and L_f .

Construction of H_G .

We use three auxiliary graphs $T(w)$, $I(j)$ and $G(v, L_f(v), s)$. The gadgets $I(j)$ and $T(w)$ are shown in Figure 1. Consider a vertex v and a copy of auxiliary graph $T(w)$. Join the vertex v to $T(w)$. Next, for every $j \in (\mathbb{N}_s \setminus \{1\}) \setminus L_f(v)$ consider a copy of $I(j)$ and join the vertex v to the vertex u_j . Finally, put s isolated vertices and join each of them to the vertex v . Call the resulting graph $G(v, L_f(v), s)$. Now, for every vertex $v \in V(G)$ put a copy of $G(v, L_f(v), |W| + 1)$ and for every edge vv' in the graph G join the vertex $v \in V(G(v, L_f(v), |W| + 1))$ to the vertex $v' \in V(G(v', L_f(v'), |W| + 1))$. Call the resulting graph H_G .

For a family \mathcal{F} of graphs, define: $\mathcal{F}' \stackrel{\text{def}}{=} \{H_G | G \in \mathcal{F}\}$. We show that if \mathcal{F} is a family of graphs such that *list coloring problem* is NP-complete for that family. Then, the following problem is NP-complete: "Given a graph $H_G \in \mathcal{F}'$, does H_G have a (0, 1)-additive labeling?"




Fig. 1: The auxiliary graphs $I(j)$ and $T(w)$.

First consider the following facts.

Fact 1 Let G be a graph with a $(0, 1)$ -additive labeling ℓ and assume that it has the auxiliary graph $T(w)$ as a subgraph, $\ell(v) = 0$, $\ell(w) = 1$ and $\sum_{x \in N(w)} \ell(x) = 1$.

Proof of Fact 1. By attention to the two triangles $x_1x_2x_3$ and $y_1y_2y_3$, $\ell(w) = 1$ and $\ell(y_4) = 1$. Also $\ell(x_1) \neq \ell(x_2)$, without loss of generality suppose that $\ell(x_1) = 1$ and $\ell(x_2) = 0$. Therefore, $\ell(x_3) = 0$, thus $\sum_{x \in N(w)} \ell(x) = 1 + \ell(v)$. Since $\sum_{x \in N(x_3)} \ell(x) = 2$, therefore $\sum_{x \in N(w)} \ell(x) = 1$, consequently $\ell(v) = 0$. ♠

Fact 2 Let G be a graph with a $(0, 1)$ -additive labeling ℓ and assume that it has the auxiliary graph $I(j)$ as a subgraph, $\sum_{x \in N(u_j)} \ell(x) \geq j$.

Proof of Fact 2. By Fact 1, $\ell(w) = 1$, by using a similar argument $\ell(z_1) = \dots = \ell(z_{j-1}) = 1$. So $\sum_{x \in N(u_j)} \ell(x) \geq j$. ♠

Fact 3 Let ℓ be a $(0, 1)$ -additive labeling for $G(v, L_f(v), |W| + 1)$, $\sum_{x \in N(v)} \ell(x) \in L_f(v)$.

Proof of Fact 3. By Fact 1 and Fact 2 it is clear.

First, suppose that the graph H_G has a $(0, 1)$ -additive labeling ℓ , define $c : V(G) \rightarrow \mathbb{N}$, $c(v) = \sum_{x \in N(v)} \ell(x)$. The function c is a proper vertex coloring and for every vertex v , by Fact 3, $c(v) \in L_f(v)$. Next, suppose that the graph G is L_f -colorable, then it is clear that the graph H_G has a $(0, 1)$ -additive labeling.

The list coloring problem is **NP**-complete for perfect graphs and planar graphs (see [7]). Obviously if G is a planar graph, then H_G is a planar graph. Also, if G is a perfect graph, then it is easy to see that the graph H_G is a perfect graph. This completes the proof. \square

5 Planar graphs

Proof of Theorem 4: Let Φ be a 3-SAT formula with the set of clauses C and the set of variables X . Let $G(\Phi)$ be a graph with the vertices $C \cup X \cup (\neg X)$, where $\neg X = \{\neg x : x \in X\}$, such that for each clause $c = y \vee z \vee w$, c is adjacent to y , z and w , also every $x \in X$ is adjacent to $\neg x$. Φ is called planar 3-SAT(type 2) formula if $G(\Phi)$ is a planar graph. It was shown that the problem of satisfiability of planar 3-SAT(type 2) is **NP**-complete [15]. In order to prove our theorem, we reduce the following problem to our problem.

Problem: *Planar 3-SAT(type 2)*.

INPUT: A planar 3-SAT(type 2) formula Φ .

QUESTION: Is there a truth assignment for Φ that satisfies all the clauses?

Consider an instance of planar 3-SAT(type 2) with the set of variables X and the set of clauses C . We transform this into a graph $G'(\Phi)$ such that $G'(\Phi)$ has a $(0, 1)$ -additive labeling, if and only if Φ is satisfiable. The graph $G'(\Phi)$ has a copy of $B(x)$ for each variable x and a copy of $A(c)$ for each clause c . The gadgets $B(x)$ and $A(c)$ are shown in Figure 2. Also, for every $c \in C$, $x \in X$, the edge $w_c^1 x$ is added if c contains the literal x . Furthermore, for every $c \in C$, $\neg x \in \neg X$, the edge $w_c^1 \neg x$ is added if c contains the literal $\neg x$. Call the resulting graph $G'(\Phi)$. Clearly the graph $G'(\Phi)$ is triangle-free and planar.




Fig. 2: The two auxiliary graphs $A(c)$ and $B(x)$.

Fact 4 Let ℓ be a (0, 1)-additive labeling for the graph $G'(\Phi)$, for each clause $c = a \vee b \vee d$, $\ell(a) + \ell(b) + \ell(d) \geq 1$.

Proof of Fact 4. To the contrary suppose that there exists a clause $c = a \vee b \vee d$, such that $\ell(a) + \ell(b) + \ell(d) = 0$, then $\sum_{t \in N(w_c^1)} \ell(t) = \ell(w_c^2) + \ell(w_c^3)$. Notice that in that case, ℓ restricted to the odd cycle $w_c^1 w_c^2 w_c^4 w_c^5 w_c^3$, is a (0,1)-additive labeling, but an odd cycle does not have any (0, 1)-additive labeling, this is a contradiction. ♠

Fact 5 Let $G'(\Phi)$ be a graph with a (0, 1)-additive labeling ℓ , for each variable x , $\ell(x) + \ell(\neg x) \leq 1$.

Proof of Fact 5. To the contrary, suppose that there is a variable x , such that $\ell(x) + \ell(\neg x) = 2$. Consider the auxiliary graph $B(x)$. Because of the odd cycle $y_x^1 y_x^2 y_x^4 y_x^5 y_x^3$, $\ell(y_x^6) = 1$. Now two cases for $\ell(y_x^5)$ can be considered.

Case 1. $\ell(y_x^5) = 1$. Thus $\sum_{t \in N(y_x^6)} \ell(t) = 3$, therefore $\sum_{t \in N(y_x^5)} \ell(t) \in \{1, 2\}$.

- If $\sum_{t \in N(y_x^5)} \ell(t) = 1$, then $\ell(y_x^3) = \ell(y_x^4) = 0$. Thus, $\ell(y_x^1) + \ell(y_x^2) = 1$; without loss of generality suppose that $\ell(y_x^1) = 1$ and $\ell(y_x^2) = 0$, in this case $\sum_{t \in N(y_x^2)} \ell(t) = \sum_{t \in N(y_x^4)} \ell(t)$, but this is a contradiction.

- If $\sum_{t \in N(y_x^5)} \ell(t) = 2$. Suppose that $\ell(y_x^3) = 1$, $\ell(y_x^4) = 0$. Four subcases for $\ell(y_x^1), \ell(y_x^2)$ can be considered, each of them produces a contradiction.

Case 2. $\ell(y_x^5) = 0$. Thus $\sum_{t \in N(y_x^6)} \ell(t) = 2$, therefore $\sum_{t \in N(y_x^5)} \ell(t) \in \{1, 3\}$.

- If $\sum_{t \in N(y_x^5)} \ell(t) = 1$, then $\ell(y_x^3) = \ell(y_x^4) = 0$. Therefore, $\ell(y_x^1) + \ell(y_x^2) = 1$. With no loss of generality suppose that $\ell(y_x^1) = 1$, $\ell(y_x^2) = 0$, therefore $\sum_{t \in N(y_x^3)} \ell(t) = \sum_{t \in N(y_x^5)} \ell(t)$, but this is a contradiction.

- If $\sum_{t \in N(y_x^5)} \ell(t) = 3$, then $\ell(y_x^3) + \ell(y_x^4) = 2$. Thus $\ell(y_x^1) + \ell(y_x^2) = 1$. Suppose that $\ell(y_x^1) = 1$, $\ell(y_x^2) = 0$, therefore $\sum_{t \in N(y_x^1)} \ell(t) = \sum_{t \in N(y_x^3)} \ell(t)$, this is a contradiction. ♠

First, suppose that Φ is satisfiable with the satisfying assignment $\Gamma : X \rightarrow \{true, false\}$. We present a (0, 1)-additive labeling ℓ for $G'(\Phi)$. For every variable x if $\Gamma(x) = true$, then put $\ell(x) = 1$, otherwise put $\ell(\neg x) = 1$. Also put $\ell(z_1) = \dots = \ell(z_{10}) = \ell(y_x^1) = \ell(y_x^3) = \ell(y_x^4) = \ell(y_x^5) = \ell(y_x^6) = 1$. Moreover, for every clause c , put $\ell(w_c^1) = \ell(w_c^2) = \ell(w_c^3) = \ell(w_c^5) = 1$. It is easy to extend this labeling to a (0, 1)-additive labeling for the graph $G'(\Phi)$. Next, suppose that the graph $G'(\Phi)$ has a (0, 1)-additive

labeling ℓ . For each variable x , by Fact 5, $\ell(x) + \ell(\neg x) \leq 1$. If $\ell(x) = 1$, put $\Gamma(x) = \text{true}$, if $\ell(\neg x) = 1$, then put $\Gamma(x) = \text{false}$ and otherwise put $\Gamma(x) = \text{true}$. By Fact 4, Γ is a satisfying assignment for Φ . \square

6 Inapproximability

Proof of Theorem 5: Let $\varepsilon > 0$ and k be a sufficiently large number. It was shown that 3-colorability of 4-regular planar graphs is NP-complete [12]. We reduce this problem to our problem. In other words, for a given 4-regular planar graph G with k vertices, we construct a planar graph G^* with $7k + 10k^{\lceil \frac{3}{\varepsilon} \rceil + 2}$ vertices, such that if $\chi(G) \leq 3$, then $\sigma_1(G^*) \leq 5k$, otherwise $\sigma_1(G^*) > 5k^{\lceil \frac{3}{\varepsilon} \rceil + 1}$, therefore there is no θ -approximation algorithm for determining $\sigma_1(G^*)$ for planar graphs, where:

$$\begin{aligned} \theta = \frac{\text{Approximate Answer}}{OPT} &> \frac{5k^{\lceil \frac{3}{\varepsilon} \rceil + 1}}{5k} \\ &= k^{\lceil \frac{3}{\varepsilon} \rceil} \\ &= (k^{\lceil \frac{3}{\varepsilon} \rceil + 3})^{\frac{\lceil \frac{3}{\varepsilon} \rceil}{\lceil \frac{3}{\varepsilon} \rceil + 3}} \\ &\geq (7k + 10k^{\lceil \frac{3}{\varepsilon} \rceil + 2})^{\frac{\lceil \frac{3}{\varepsilon} \rceil}{\lceil \frac{3}{\varepsilon} \rceil + 3}} \\ &\geq |V(G^*)|^{\frac{\lceil \frac{3}{\varepsilon} \rceil}{\lceil \frac{3}{\varepsilon} \rceil + 3}} \\ &\geq |V(G^*)|^{1-\varepsilon} \end{aligned}$$

In order to construct the graph G^* , we use the auxiliary graph $D(v)$ which is shown in Figure 3. Using simple local replacements, for every vertex v of the graph G , put a copy of $D(v)$, and for every edge vu of the graph G , join the vertex v of $D(v)$ to the vertex u of $D(u)$. Call the resulting graph G^* . First, suppose that G is not 3-colorable and let ℓ be a $(0, 1)$ -additive labeling for G^* . By the structure of $D(v)$ we have $\ell(v) = 1$ and $\ell(p_3) = 0$, so $\sum_{x \in N(v)} \ell(x) = 4 + \ell(p_4) + \ell(p_5) + \ell(p_6)$. Since G is not 3-colorable, there exists a vertex v such that $\sum_{x \in N(v)} \ell(x) = 4$, therefore in the subgraph $D(v)$, $\ell(p_4) + \ell(p_5) + \ell(p_6) = 0$, so $\ell(p_5) = 0$. Consequently for every i , $1 \leq i \leq d$, in the subgraph $D(v)$, $\ell(v_i) + \ell(v'_i) \geq 1$. So $\sigma_1(G^*) > 5k^{\lceil \frac{3}{\varepsilon} \rceil + 1}$. Next, suppose that $\chi(G) \leq 3$. So G has a proper vertex coloring $c : V(G) \rightarrow \{1, 2, 3\}$. For every vertex v of G , if $c(v) = 1$ put $\ell(p_4) = \ell(p_6) = 0$ and $\ell(p_5) = 1$, else if $c(v) = 2$ let $\ell(p_4) = 0$ and $\ell(p_5) = \ell(p_6) = 1$ and if $c(v) = 3$ let $\ell(p_4) = \ell(p_5) = \ell(p_6) = 1$. It is easy to extend ℓ to a $(0, 1)$ -additive labeling for the graph G^* such that $\sigma_1(G^*) \leq 5k$. \square

7 Concluding remarks

In this paper we study the computational complexity of $(0, 1)$ -additive labeling of graphs. A $(0, 1)$ -additive labeling of a graph G is a function $\ell : V(G) \rightarrow \{0, 1\}$, such that for every two adjacent vertices v and u of G , $\sum_{w \sim v} \ell(w) \neq \sum_{w \sim u} \ell(w)$. For future work, someone can consider another version of this problem that we call proper total dominating set. A *proper total dominating set* of a graph $G = (V, E)$, is




Fig. 3: The auxiliary graph $D(v)$. This graph has $7 + 10k^{\lceil \frac{3}{\varepsilon} \rceil + 1}$ vertices, where $d = 5k^{\lceil \frac{3}{\varepsilon} \rceil + 1}$.

a subset D of V such that every vertex has a neighbor in D (all vertices in the graph including the vertices in the dominating set have at least one neighbor in the dominating set) and every two adjacent vertices have a different number of neighbors in D (note that in a (0,1)-additive labeling every vertex does not need to have a neighbor labeled 1).

In this work, we proved that for every k there is a graph G such that $\eta(G) \leq k \leq \eta_\ell(G)/2$. What can we say about the difference in bipartite graphs?

Acknowledgements

The authors wish to thank Saieed Akbari who drew their attention to Additive List Coloring Conjecture.

References

- [1] L. Addario-Berry, R. E. L. Aldred, K. Dalal, and B. A. Reed. Vertex colouring edge partitions. *J. Combin. Theory Ser. B*, 94(2):237–244, 2005.
- [2] L. Addario-Berry, K. Dalal, and B. A. Reed. Degree constrained subgraphs. *Discrete Appl. Math.*, 156(7):1168–1174, 2008.
- [3] A. Ahadi, A. Dehghan, M. Kazemi, and E. Mollaahmadi. Computation of lucky number of planar graphs is NP-hard. *Inform. Process. Lett.*, 112:109–112, 2012.
- [4] S. Akbari, M. Ghanbari, R. Manaviyat, and S. Zare. On the lucky choice number of graphs. *Graphs Combin.*, 29(2):157–163, 2013.
- [5] T. Bartnicki, J. Grytczuk, and S. Niwczyk. Weight choosability of graphs. *J. Graph Theory*, 60(3):242–256, 2009.
- [6] Tomasz Bartnicki, Bartłomiej Bosek, Sebastian Czerwiński, Jarosław Grytczuk, Grzegorz Matecki, and Wiktor Żelazny. Additive coloring of planar graphs. *Graphs and Combinatorics*, 30(5):1087–1098, 2014.
- [7] F. Bonomo, G. Duřan, and J. Marengo. Exploring the complexity boundary between coloring and list-coloring. *Electronic Notes in Discrete Mathematics*, 25:41–47, 2006.
- [8] M. Borowiecki, J. Grytczuk, and M. Piłśniak. Coloring chip configurations on graphs and digraphs. *Inform. Process. Lett.*, 112(1–2):1–4, 2012.

- [9] Axel Brandt, Jennifer Diemunsch, and Sogol Jahanbekam. Lucky choice number of planar graphs with given girth. *Submitted*, 2015.
- [10] Gary Chartrand, Futaba Okamoto, and Ping Zhang. The sigma chromatic number of a graph. *Graphs Combin.*, 26(6):755–773, 2010.
- [11] Sebastian Czerwiński, Jarosław Grytczuk, and Wiktor Żelazny. Lucky labelings of graphs. *Inform. Process. Lett.*, 109(18):1078–1081, 2009.
- [12] David P. Dailey. Uniqueness of colorability and colorability of planar 4-regular graphs are NP-complete. *Discrete Math.*, 30(3):289–293, 1980.
- [13] A. Dehghan, M.-R. Sadeghi, and A. Ahadi. The complexity of the sigma chromatic number of cubic graphs. *Submitted*.
- [14] Ali Dehghan, Mohammad-Reza Sadeghi, and Arash Ahadi. Algorithmic complexity of proper labeling problems. *Theoretical Computer Science*, 495:25–36, 2013.
- [15] Ding-Zhu Du, Ker-K Ko, and J. Wang. *Introduction to Computational Complexity*. Higher Education Press, 2002.
- [16] M. R. Garey and D. S. Johnson. Computers and intractability: A guide to the theory of NP-completeness. *Bull. Amer. Math. Soc. (N.S.)*, 3(2):898–904, 1980.
- [17] Maciej Kalkowski, Michał Karoński, and Florian Pfender. Vertex-coloring edge-weightings: towards the 1-2-3-conjecture. *J. Combin. Theory Ser. B*, 100(3):347–349, 2010.
- [18] Michał Karoński, Tomasz Łuczak, and Andrew Thomason. Edge weights and vertex colours. *J. Combin. Theory Ser. B*, 91(1):151–157, 2004.
- [19] M. Khatirinejad, R. Naserasr, M. Newman, B. Seamone, and B Stevens. Digraphs are 2-weight choosable. *Electron. J. Combin.*, 18(1):Paper 21,4, 2011.
- [20] M. Khatirinejad, R. Naserasr, M. Newman, B. Seamone, and B. Stevens. Vertex-colouring edge-weightings with two edge weights. *Discrete Math. Theor. Comput. Sci.*, 14(1):1–20, 2012.
- [21] Michał Lasoń. A generalization of combinatorial Nullstellensatz. *Electron. J. Combin.*, 17(1):Note 32, 6, 2010.
- [22] Javier Marenco, Marcelo Mydlarz, and Daniel Severín. Topological additive numbering of directed acyclic graphs. *Information Processing Letters*, 115(2):199–202, 2015.
- [23] B. Seamone. Derived colourings of graphs. PhD thesis, Carleton University, 2012.
- [24] Douglas B. West. *Introduction to graph theory*. Prentice Hall Inc., Upper Saddle River, NJ, 1996.