

Study on Dynamic Variation Regularities of Regional Groundwater Depth Based on Complexity Diagnosis

Dong Liu, Miao Yu, Nan Sun, Ying Qi

► **To cite this version:**

Dong Liu, Miao Yu, Nan Sun, Ying Qi. Study on Dynamic Variation Regularities of Regional Groundwater Depth Based on Complexity Diagnosis. 5th Computer and Computing Technologies in Agriculture (CCTA), Oct 2011, Beijing, China. pp.72-83, 10.1007/978-3-642-27278-3_9 . hal-01360966

HAL Id: hal-01360966

<https://hal.inria.fr/hal-01360966>

Submitted on 6 Sep 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Study on Dynamic Variation Regularities of Regional Groundwater Depth Based on Complexity Diagnosis¹

Dong Liu, Miao Yu, Nan Sun, Ying Qi

(School of Water Conservancy & Civil Engineering, Northeast Agricultural University, Harbin Heilongjiang 150030, China)

Abstract: In the course of analyzing and forecasting regional groundwater depth, its variation process complexity was ignored in the past, so that it was difficult to realize scientific management of groundwater resources. Aiming at the aforesaid defect, taking Jiansanjiang branch bureau as research platform, this paper diagnosed the complexity of each monthly groundwater depth series through using complexity measurement methods which include wavelet theory, fractal theory, Approximate Entropy and so on. The comprehensive measurement analysis results show that the complexity order of monthly groundwater depth series in the eighth district of Farm Nongjiang is highest in four farms of northern region. On that basis, the most complexity groundwater depth series combination stochastic forecasting model based on wavelet transform in northern region of Jiansanjiang branch bureau was set up. The model forecasting results show that local annual mean groundwater depth will be in a state of fluctuant lifting, and have considerable gap in comparison to groundwater ideal depth in the future. So, local government should reinforce the scientific management of groundwater. The research achievements break through the traditional study paradigm of hydrological forecasting, provide a research mode for studying regional hydrologic process, and provide scientific evidence for sustainable utilizing regional groundwater resources.

Key words: northern region of Jiansanjiang branch bureau; groundwater depth; complexity diagnosis; wavelet transform; combination stochastic forecasting model

1 Introduction

Groundwater is an important basic strategy resource to support the economic sustainable development of the regional society. Groundwater resources is a complex system^[1], containing system factor such as water level, water quality, water quantity and water temperature etc., among which, groundwater level is the most sensitive and active one. Under conditions of unreasonable exploitation of groundwater, excessive use of fertilizers and pesticides and some nature factors, fall of groundwater level, water quality deterioration and intensification of agricultural non-point source pollution etc. serious problem are induced, making the complexity of groundwater more and more clear which is general concerned by all circles. But the complexity of changing process was ignored by the domestic scholars when they studied the

About the Author: Dong Liu (1972-) Male (Han nationality), from Heilongjiang Anda, professor, doctor, main research direction is agriculture and water resources system analysis and optimal using. E-mail: liu72dong@126.com.

groundwater depth dynamic change regularity of the groundwater, resulting it imposible to fully excavate information of groundwater buried deeply change process and lack of practicality. Under this background, it is necessary to analyze predict the regional groundwater buried deeply series dynamic change to achieve the sustainable use of local groundwater, premised the complexity diagnoses.

Generally speaking, the higher complexity of the hydrological time series, the lower predictability of it^[2]. So, to increase the prediction accuracy of the higher hydrological time series complexity, appropriate method should be chosen. Traditional prediction method of groundwater groundwater depth contains time series analysis^[3], gray theory^[4], artificial neural networks^[5], multiple linear regression, seasonal decomposition model^[6], self-memory model^[7], etc., some of them base on linear theory, considered questions simply, some has ineffective extension, some has complex theories, these all resulting lower prediction accuracy or difficult to use. Wavelet analysis, developed recently, can demonstrate the fine structure series fully and extract change regularity easily of the hydrological time. The authors study the case of north district of Jiansanjiang branch bureau, to measure the complexity of the groundwater groundwater depth series to found the random predicting model of complex groundwater depth series of north district of Jiansanjiang branch bureau.


2 Research area and methods

2.1 Research area

Heilongjiang Land Reclamation Jiansanjiang branch bureau is rice production base of China, locating at the junction of two cities and two countries of Fujin, Tongjiang, Raohe in Sanjiang Plain with the agricultural acreage of 682 thousand hm²^[8] of total area of 12.3 thousand km²^[9]. The north district of Jiansanjiang branch bureau includes farms of Qinglongshan, Qindeli, Nongjian and Yalu River with total land area of 2900.8km², and in the year of 2009, the agricultural acreage is 147 hm². The district has provincial nature reserve of Qindeli sturgeon fish, Qinglonghe and Yalu River branch level nature reserve, mainly planted crops of rice, maize, soybean and wheat with higher mechanization and commodity grain. From 1920s, the district tied rice planting, and the acreage increased year by year, by the year of 2009, it was 105 thousand, taking up 71.4% of the total agricultural acreage. Because of the lack of control projects, agriculture irrigation is mainly used groundwater, resulting continuous decline of the groundwater level. Meanwhile, under the influence of precipitation, evaporation and hydrological conditions, the complexity of the groundwater dynamic change become more and more clear of the district. Therefore, the authors try to analysis and predict the complex groundwater depth series dynamic change of the north district of Jiansanjiang branch bureau under the premise of complexity diagnose, to realize the sustainable use of the local groundwater resource.

2.2 Data Sources

From Heilongjiang Land Reclamation Jiansanjiang branch bureau Water Authority collected the monthly groundwater level series monitoring data including 15 farms practice of Jiansanjiang branch bureau from 1997 to 2007($n=132$). Draw the groundwater level series changing curves of every areas in fig. 1.As can be seen from fig. 1, depth of the groundwater monitoring sites in Jiansanjiang branch bureau showed a gradual growth trend overall.



a. 1st team of Farm 859; b.69th team of Farm Qixing; c.22nd team of Farm Qianjin; d.24th team of Farm Chuanye; e.5th team of Farm Yalu River; f.28th team of Farm Hongwei; g.8th district of Farm Nongjiang; h.17th team of Farm Qinglongshan; i.36th team of Farm Qindeli; j.main field of Farm Qianfeng; k.6th district of Farm Honghe; l.5th district of Farm Erdaohe; m.11th team of Farm Daxing; n.31st team of Farm Shengli; o.12th team of Farm Qianshao

Fig. 1 Monthly groundwater level series changing curves of Jiansanjiang branch bureau (1997~2007)

2.3 Diagnose Measurements of Hydrological Series complexity

Frequently used diagnose measurements of hydrological series contain information cost function based on wavelet transform(abbr. *WT-ICF*), approximate Entropy(abbr. *ApEn*), Lempel-Ziv Complexity(abbr. *LZC*), Rescaled Range Analysis, (abbr. *R/S*), discrete wavelet transform(abbr. *DWT*), continue wavelet transform(abbr. *CWT*), etc., specific references as [10] and [11], not repeated here.

2.4 Modeling Principles of Hydrological Series Random Model Based on Wavelet Transform

2.4.1 Fast Wavelet Transform Method

Computational complexity is big for information redundancy^[12] of the obtained wavelet transform coefficient when the hydrological time series are wavelet transformed. So, fast wavelet transform (abbr. *FWT*) is usually used in practical application to calculate the wavelet transform coefficient. The authors use *A Trous* method^[10] which is simple, fast and small computational complexity.

To wavelet transform the hydrological series $f(t) (t=1, 2, \dots, N)$, letting $C^0(t) = f(t)$, the factorization and reconstruction process of *A Trous* method is as following:

$$\begin{cases} C^j(t) = \sum_{k=-\infty}^{+\infty} h(k)C^{j-1}(t+2^j k) \\ W^j(t) = C^{j-1}(t) - C^j(t) \end{cases} \quad (1)$$

$$C^0(t) = C^J(t) + \sum_{j=1}^J W^j(t) \quad (2)$$

Among which, $C^j(t)$, $W^j(t)$ are the scale coefficient (background signal) and wavelet coefficient (detail signal) under j scale; $h(k)$ is discrete lowpass filter, generally be chosen the compactly supported symmetric third-order *B-spline* one, as $h(k)=(1/16, 1/4, 3/8, 1/4, 1/16)$. $\{W^1(t), W^2(t), \dots, W^j(t), C^j(t)\}$ is called the wavelet transform series under J scale.

2.4.2 Modeling Procedures

Wavelet transform method (e.g. *A Trous* Method) is used to transform the hydrological time series $f(t)$ to obtain wavelet transform series $\{W^1(t), W^2(t), \dots, W^j(t), C^j(t)\}$ under some J scale firstly, then recognize the main components (random or certain components) of the wavelet transform series and found suitable mathematic models of each wavelet transform series, finally, to get the random predicting model^[10] of the studying hydrological time series $f(t)$ by the method of wavelet transform and reconstruction.

3 Results and Analyses

3.1 Complexity measures of groundwater level series

Estimated the complexity of each monthly groundwater depth series through using six kinds of complexity measurement methods mentioned above in Jiansanjiajiang

branch bureau. The complexity diagnosis results of northern region of Jiansanjiang branch bureau shown in table 1.

Table 1. Calculated Results of the complexity index about monthly groundwater depth series in the northern region of Jiansanjiang branch bureau

Long view well location	WT-ICF		ApEn		LZC		R/S		D				comprehensive complexity index	complexity sort
	(0.16)		(0.16)		(0.16)		(0.12)		Wavelet estimation					
									DWT		CWT			
	score	sort	score	sort	score	sort	score	sort	score	sort	score	sort		
	t	e	t	e	t	e	t	e	t	e	t	e		
17 th team of Farm Qinglongshan	⑧	8	⑧	8	③	13	①	15	⑩	6	⑥	10	9.80	③
36 th team of Farm Qindeli	⑤	11	⑩	6	②	14	⑨	7	⑬	3	⑮	1	6.52	⑩
8 th district of Farm Nongjiang	⑦	9	⑫	4	⑦	9	⑤	11	②	14	①	15	10.68	②
5 th team of Farm Yalu River	④	12	⑨	7	⑤	11	④	12	⑤	11	⑩	6	9.44	④

Note: ①The digital in parentheses is the weight of groundwater depth sequence complexity measure;
 ②D is fractal dimension.

3.2 Synthetical complexity measures of groundwater level series

In the above complexity measurement methods, Range Analysis is sensitive to the sequence length which is belongs to biased estimation and poor stability^[13], while the method Continuous Wavelet Transform Fractal Theory has good stability^[10], and others' be placed in the middle. In order to give full play to the advantages of various complexity measurement methods, according to the above analysis, determined the weight of that six complexity measures w_i ($i=1,2,\dots,6$) (shown in table 1). Assigned the corresponding score $s_i = 15 \sim 1$ to the sort result (①~⑮) of each monthly groundwater depth sequence complexity in Jiansanjiang branch bureau. Then get the groundwater depth sequence comprehensive complexity calculation formula:

$$C_{j_j} = \sum_{i=1}^6 s_i w_i \quad (3)$$

C_{j_j} is the comprehensive complexity index of which in the sort of j in groundwater depth sequence, $j=1, 2, \dots, 15$.

Calculate the comprehensive complexity index of groundwater depth sequence in northern region of Jiansanjiang branch bureau followed calculation formula (3), and the result seen in table 1. From table 1 we known that the synthetical complexity sort of northern region of Jiansanjiang branch bureau monthly groundwater level series is

the eighth district of Farm Nongjiang >the 17th team of Farm Qinglongshan>the fifth district of Farm Yalu River>the 36th team of Farm Qindeli>the 12th team of Farm Qianshao.

3.3 The model of complexity groundwater depth series combination stochastic forecasting model which is based on wavelet transform in northern region of Jiansanjiang branch bureau.

By the calculation analysis shows that the complexity of the eight district of Farm Nongjiang is the highest in the sort of five farms in northern region. Therefore, chose the monthly groundwater depth series of the eighth district of Farm Nongjiang as a representative, describe northern region of Jiansanjiang branch bureau groundwater depth dynamic change general characteristics based on analyzes dynamic change regulation above. Set up groundwater depth series combination stochastic forecasting model based on wavelet transform according to the monthly groundwater depth data surveied in the eighth district of Farm Nongjiang from 1997 to 2006. Reserved the data of 2007 as verify results.

(1) Wavelet decomposition and reconstruction of monthly groundwater depth series

Adopted *A Trous* algorithm mentioned above and boundary extension method which include 35 percent symmetric extension and the same proportion of linear extension and 30 percent orthogonal polynomials fitting extension to decompose the monthly groundwater depth series $G(t)$ ($t = 1 \sim 120$) of the eighth district of Farm Nongjiang from 1997 to 2006 through comparative analysis. The scale number $p=2$ here .Obtained wavelet decomposition sequence $\{W^1(t), W^2(t), C^2(t)\}$, shown in fig. 2 (a),(b),(c). Then we can got reconstruction sequences by superposed every wavelet decomposition sequence. Shown in fig. 2(d). As can be seen from fig. 2(d), reconstruction complete agreement with the monthly groundwater depth series change precess results in fig. 1. So, used *A Trous* algorithm to decompose monthly groundwater depth data series in the eighth district of Farm Nongjiang is feasible.




Fig. 2 Wavelet decomposition and reconstruction of monthly groundwater depth series in the eighth district of Farm Nongjiang

(2) Component identification about wavelet transform sequence

Draw the self-relevant chart and variance spectral density fig. ^[14] of the wavelet transform sequence. Seen fig. 3 and fig. 4.




Fig. 3 The self-relevant charts of each wavelet transform series




Fig. 4 The variance spectral density of each wavelet transform series

It can be seen from fig. 2, fig.3, fig. 4 that the wavelet decomposition sequence has a periodical for 12 months, and it has changing regularity powerful. So $W^1(t)$ which belongs to deterministic component can represent periodic term of $G(t)$. Although the wavelet transform sequence $W^2(t)$ also has a certain of periodicity that inferior to $W^1(t)$, it mainly reflect randomness. So we used $W^2(t)$ of which belongs to random component stand for randomness of $G(t)$. The sequence $C^2(t)$ not has periodic changing regularity but clear trend. For this reason, $C^2(t)$ which belongs to deterministic component^[14] can be approximated as trend term of $G(t)$.

(3) Cross-correlation analysis about wavelet transform sequence

It can be seen each cross-correlation analysis curve^[14] about wavelet transform sequence from fig. 5. In this fig. $r_k(W^1, W^2)$, $r_k(W^1, C^2)$, and $r_k(W^2, C^2)$ showed the cross correlation coefficients among sequences $W^1(t)$, $W^2(t)$, and $C^2(t)$ respectively. In addition, as the complex characteristics of original sequence is obvious, data sequences $W^1(t)$ and $W^2(t)$ are non stationary sequence with periodic or approximate periodic components, both part of the information contained overlap lead to many points in the cross correlation coefficient $r_k(W^1, W^2)$ beyond 95% allows limited scope. So, Need to smooth with sequence $W^1(t)$ and $W^2(t)$.




Fig. 5 The cross-correlation charts of each wavelet transform series




Fig. 6 The cross-correlation charts of each wavelet transform series after smooth processing

(4) Smoothly with the wavelet transform $W^1(t)$ and $W^2(t)$

Take seasonal difference^[15] to the series $W^1(t)$ who has 12 months (1 year) cycle of change rule obviously, will get the wavelet transform difference sequence $W^{1*}(t)$. For the sequences $W^2(t)$ also has 12 months (1 year) cycle of change rule obviously, and has nonstationarity remarkable. So, deal it with logarithmic transformation^[16] method first of all, obtain the logarithmic transformation series $W^{2*}(t)$. Then, take seasonal difference to $W^{2*}(t)$ to get the wavelet logarithmic transformation difference sequences $W^{2**}(t)$.

(5) Cross-correlation analysis about wavelet transform sequence after smooth processing

Draw the cross-correlation curve about $W^{1*}(t)$, $W^{2**}(t)$ and $C^2(t)$ in fig. 6. It can be seen from the fig. that after smooth processing, the cross correlation coefficient are almost followed in 95% allows limited scope, and close to zero. Therefore, the mutual correlation of sequence is lesser. We can approximate think that they are independent of each other. The above cross-correlation analysis result showed that the components of $W^{1*}(t)$, $W^{2**}(t)$ and $C^2(t)$ nearly single, more simple than $G(t)$. At the same time series $W^{1*}(t)$ and $W^{2**}(t)$ can be reduced to the wavelet sequences $W^1(t)$ and $W^2(t)$ after smooth processing. So analysis and deal with $G(t)$ is means to manage $W^{1*}(t)$, $W^{2**}(t)$ and $C^2(t)$

(6) Set up the combination stochastic model

From those analysis mentioned above, we known that sequence $W^{1*}(t)$ is got from periodic term $W^1(t)$ of original sequence after seasonal difference, and sequence $C^2(t)$ which belongs to deterministic component is approximate trend term of original sequence. At the same time, sequence $W^{2**}(t)$ is got from stochastic $W^2(t)$ of original sequence after logarithmic transformation and seasonal difference, it belongs to stochastic component and it is approximate random component of original sequence. Therefore, set up auto-regression model^[14] of $W^{1*}(t)$ and $C^2(t)$ (cut out random variables ε_t) and auto-regression model of $W^{2**}(t)$ by using point of view of stochastic hydrology. Reduction and superposition each series' mathematical model then will get the groundwater depth series combination stochastic forecasting model which is based on wavelet transform in the eighth district of Farm Nongjiang

$$\begin{cases} \hat{W}^{1*}(t) = 0.7216W_{t-1}^{1*} - 0.2698W_{t-12}^{1*} \\ \hat{W}^1(t) = \hat{W}^{1*}(t) + W^1(t-12) \\ \hat{W}^{2**}(t) = 0.0595 + 0.8300(W_{t-1}^{2**} - 0.0595) + 0.6934(W_{t-2}^{2**} - 0.0595) \\ \quad - 0.8232(W_{t-3}^{2**} - 0.0595) + 0.3840(W_{t-5}^{2**} - 0.0595) + \varepsilon_t \\ \hat{W}^{2*}(t) = \hat{W}^{2**}(t) + W^{2*}(t-12) \\ \hat{W}^2(t) = 2 - e^{\hat{W}^{2*}(t)} \\ \hat{C}^2(t) = 8.5940 + 0.8889(C_{t-1}^2 - 8.5940) \\ \quad + 0.2438(C_{t-2}^2 - 8.5940) - 0.2247(C_{t-3}^2 - 8.5940) \\ \hat{G}(t) = \hat{W}^1(t) + \hat{W}^2(t) + \hat{C}^2(t) \end{cases} \quad (4)$$

Type with "" said the corresponding sequence of predictive value

(7) Model fitting

Adopted the most complexity groundwater depth series combination stochastic forecasting model which is based on wavelet transform to fit the monthly groundwater depth series in eighth district of Farm Nongjiang from 1997 to 2006. Shown in fig. 7. As can be seen from fig. 7, the model fitting effect is good.




Fig.7 The fitting curve of monthly groundwater depth series combination stochastic forecasting model in the eighth district of Farm Nongjiang (1997~2006)




Fig.8 The forecasting curve of monthly groundwater depth series combination stochastic forecasting model in the eighth district of Farm Nongjian (2008~2011)

(8) Model Accuracy Test

Used the data of most complexity groundwater depth series combination stochastic forecasting model which is based on wavelet transform to test the fitting effect, and used the left monthly modeling groundwater depth measurement data of 2007 which is not adopted in the model to test trial reported result. The specific results in Table 2.

Table 2 Accuracy test result of combination stochastic forecasting model based on wavelet transform

Fitting Effect index				Simulated forecast index E3 (%)
C	p	E1 (%)	E2	
0.3208	0.9444	14.13	0.9259	83.33

According to relevant literature we can see that posterior error ratio C, fitting accuracy rate E2 and trial reported effect indexes E3 were reached I level standard, minimum error frequency p is II level, and the relative mean square error E1 reached III level standard^[17]. Therefore, the model based on wavelet transform mentioned above have good fitting effect and highly forecast precision, can be used to predict future groundwater depth in the eighth district of Farm Nongjiang.

(9) Groundwater depth prediction

Now adopted the most complexity groundwater depth series combination stochastic forecasting model which is based on wavelet transform to predict monthly groundwater level series of the eighth district of Farm Nongjiang from 2008 to 2011. The prediction data and prediction curve of groundwater table can be seen respectively in table 3 and fig. 8. From table 3 and fig. 8 shown that the groundwater level of the eighth district of Farm Nongjiang will fluctuate lift, if still using the past groundwater exploitation mode. The inter-annual luffing of groundwater level from 2008 to 2011 were -1.28 m, 0.52 m, 0.08 m and -0.12 m respectively in the fluctuation change down-up-up-down state. And the groundwater level maintained at about 10m. This result have considerable gap in comparison to groundwater ideal depth which is about 4m to 6m in the future^[18]. Therefore, the eighth district of Farm

Nongjiang should increase groundwater management efforts, to protect local limited of groundwater resources.

Table 3.The prediction data of monthly groundwater depth in the eighth district of Farm Nongjiang (2008~2011) (m)

Month Year	1	2	3	4	5	6	7	8	9	10	11	12	Average
2008	7.4118	7.9645	7.8243	8.4863	11.9801	12.0779	12.5336	11.1655	9.9586	9.1503	8.1838	7.9454	9.56
2009	7.3383	8.0817	8.1816	9.0356	12.6974	12.7893	13.2663	11.8944	10.6546	9.8109	8.7365	8.4515	10.08
2010	7.7792	8.4760	8.5244	9.3032	12.9034	12.9148	13.3161	11.8589	10.5339	9.6153	8.4798	8.1595	10.16
2011	7.4709	8.1710	8.2395	9.0524	12.6953	12.7523	13.2019	11.7968	10.5272	9.6653	8.5838	8.3112	10.04

4. Conclusions

It is more obvious, driven in a variety of factors, the complexity of characteristics during the course of the regional groundwater table changed has became. So, it has broken the traditional hydrological forecasting patterns and has more practical significance for the regional groundwater resource management that analysis of groundwater depth development trend based on identify regional groundwater depth sequence complex. This paper diagnosed the complexity of 15 monthly groundwater level series through many kinds of data mining methods in Heilongjiang Agricultural Reclamation Jiansanjiang branch bureau. The complexity diagnosis results show that the complexity order of monthly groundwater depth series in the eighth district of Farm Nongjiang is highest in five farms of northern region. On that basis, the most complexity groundwater depth series combination stochastic forecasting model based on wavelet transform in northern region of Jiansanjiang branch bureau in the eighth district of Farm Nongjiang was set up. This model concepts are clear, structured and high reliability. The model forecasting results show that local annual mean groundwater depth will be keep 10 m, and have considerable gap in comparison to groundwater ideal depth in the future. So, local government should reinforce the scientific management of groundwater. Need to point out is that The stability of the complexity measure results by certain effect as the groundwater depth monitoring data insufficient length in Jiansanjiang branch bureau. In the future, with the monitoring data accumulation, we should update this paper research.

Acknowledgments.

Thanks to National Natural Science Foundation of China (No.41071053) 、 China Postdoctoral Science Foundation (No.20080440832) 、 China Postdoctoral Scientific Fund special funding (No.201003410) 、 Research Fund for the Doctoral Program of Higher Education of China (No.20102325120009) 、 Natural Science Foundation of Heilongjiang Province, China (No.C201026) 、 Science and Technology Foundation of Education

Department of Heilongjiang Province (No.11541024) 、 Dr Start-up Fund Research of Northeast Agricultural University (No.2009RC37)

References

- [1]Guoli Wang, Guohua Liang. Application of Multi-period, Multi-level and Multi-objective Fuzzy Optimization Model in Groundwater System Management Decision-making. [J] Journal of China Hydrological, 2009,29 (6) : 24-27.
- [2] Tao Peng, Xiaohong Chen, Chengbin Zhuang. Analysis on complexity of monthly runoff series based on sample entropy in the Dongjiang river [J]. Ecology and Environmental Science, 2009,18 (4) : 1379-1382.
- [3] Dong Liu, Yong-sheng Ma, Qiang Fu. Study on Dynamic Variation Regularities of Groundwater in Area of Well Irrigation Paddy in Sanjiang Plain [J].Journal of Irrigation and Drainage ,2006, 25 (5) : 42-46.
- [4] Wei-tao Zhang, Jun-yan Sun, Cai-yun Wu. Research on the application of groundwater dynamic change regulation based on the gray-theory in pingdu [J] Shandong Water Resources, 2009, (3~4) : 45-48.
- [5] Banerjee P, Prasad R K, Singh V S. Forecasting of groundwater level in hard rock region using artificial neural network[J].Environmental Geology, 2009, 58 (6) : 1239-1246.
- [6] Qiang Xu, Longcang Shu, Dan Yang, et al. Model on Groundwater Level Dynamic Statistical in Pinggu Plain of Beijing [J]. Water Resources and Power, 2009,27 (5) : 58-61.
- [7] Rongfeng Li, Bing Shen, Jinkai Zhang. Self-memory model for predicting groundwater depth series with periodical fluctuation [J]. Transaction of The Chinese Society of Agricultural Engineering, 2005, 21 (7) : 34-37.
- [8] Qing Zhao. Research on the Trend of Underground Water Change in Jiansanjiang Area Based on Gray Prediction [J]. Journal of Water Resources and Water Engineering, 2009,20 (5) : 128-130,134.
- [9] Lei Guo, Ke-Ming Ma, Yi Zhang. Landscape assessment on wetland degradation during thirty years in Jiansanjiang region of Sanjiang Plain,Northeast China [J]. Acta Ecologica Sinica,2009,29 (6) : 3126-3135.
- [10] Wen-sheng Wang, Jing Ding, Yue-qing Li. Hydrological wavelet analysis [M].Beijing: Chemical Industry Press, 2005.
- [11] Chun-sheng Tong, Qiang Huang, Han Liu, et al. Study on runoff series complexity based on approximate entropy [J]. Journal of Northwest Sci-Tech University of Agriculture and Forestry, 2005, 33 (6) : 121-126.
- [12] Su-ying Ding. Laplacian Pyramid image data compression based on wavelet transformation [J]. Journal of Weifang University, 2009, 9 (4) : 34-36.
- [13] Rakhshandehroo G R, Shaghaghian M R, Keshavarzi A R, et al. Temporal variation of velocity components in a turbulent open channel flow: Identification of fractal dimensions[J].Applied Mathematical Modelling, 2009, 33(10): 3815-3824.
- [14] Jing Ding, Quan-shou Liu. Stochastic hydrology [M].Beijing: China Water Power Press, 1997.
- [15] Yan-hua Yu, Jun Wang, Jun-de Song. Research on traffic prediction and result analysis of using multiplicative seasonal autoregressive integrated moving average [J]. Computer Engineering and Application, 2009,45 (20) : 99-102.
- [16] Dao-gui Chen, Nai-lian Hu, Guo-qing Li. Stationarity for the non-normal distribution of regionalized variables [J]. Journal of University of Science and Technology Beijing, 2009,31 (4) : 412-417.

- [17] Nan-xiang Chen. Evaluating the Precision of Forecasting Model for Groundwater Dynamic [J]. GEOTECHNICAL INVESTIGATION AND SURVEYING, 1999, (3) : 35-38.
- [18] Bin Yuan, Shao-min Du, Wei-qing Yang, et al. Status Overdraft of Groundwater on Sanjiang Plain(2) [J]. Journal of Heilongjiang Hydraulic Engineering College, 2008,35 (1) : 18-21.