

## Rapid estimation of 3D ventricular activation from electroanatomic mapping

Simone Pezzuto, Peter Kalavsky, Mark Potse, François Regoli, Maria Luce Caputo, Giulio Conte, Tiziano Moccetti, Frits W. Prinzen, Angelo Auricchio, Rolf Krause

► **To cite this version:**

Simone Pezzuto, Peter Kalavsky, Mark Potse, François Regoli, Maria Luce Caputo, et al.. Rapid estimation of 3D ventricular activation from electroanatomic mapping . Cardiostim EHRA Europe, Jun 2016, Nice, France. Cardiostim Europe, <<http://www.cardiostim.com/>>. <hal-01386922>

**HAL Id: hal-01386922**

**<https://hal.inria.fr/hal-01386922>**

Submitted on 9 Nov 2016

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## **TITLE**

Accurate estimation of 3D ventricular activation from electroanatomic mapping

## **AUTHORS**

Pezzuto S., Kalavsky P., Potse M., Regoli F., Caputo M.L., Conte G., Moccetti T., Prinzen F.W., Auricchio A., Krause R.

## **AFFILIATION**

- Center for Computational Medicine in Cardiology, Institute of Computational Science, Università della Svizzera italiana, Via Giuseppe Buffi 13, 6904 Lugano


- Fondazione Cardiocentro Ticino, Via Tesserete 48, 6900 Lugano

## **ABSTRACT**

**Purpose** Heart failure patients are often affected by a ventricular conduction disorder. Accurate characterization of the tridimensional (3D) activation sequence of the ventricles is fundamental for an effective personalized therapy. Electroanatomical mapping provides the activation times at the endocardium, with limited coverage due to anatomical constraints. Our aim is to develop a robust tool to generate a complete 3D, biventricular activation map from a limited number of endocardial measurements.

**Method** In 6 candidates to cardiac resynchronization therapy a high density endocardial activation map of the right and left ventricle was obtained. High resolution propagating activation was simulated in patient-tailored anatomical heart models using the anisotropic eikonal equation and realistic fiber orientation. Subsequently, the mean square error between the measured and simulated activation times was minimized by varying early activation site(s) and local anisotropic conduction velocities.

**Results.** A close correlation was achieved between the simulated and measured activation time for a single patient (Panel A) while detailed 3D activation maps were achieved (panel B and C). In all cases, the optimal parameters were identified within few minutes and the full activation sequence was computed in less than 2 seconds. A 0.9 to 0.95 Pearson correlation between measured and simulated activation times was obtained in all cases. The mathematical formulation is flexible in terms of the number of activation sites and a possible delay between them, in order to mimic the presence of a partially working Purkinje network.



**Conclusion** Our methodology is able to generate in a clinically affordable computational time (few minutes) reliable, patient-specific, fully volumetric activation map of both ventricles using a local workstation or a laptop.