



HAL
open science

On the Interval of Boolean Strong Partial Clones Containing Only Projections as Total Operations

Miguel Couceiro, Lucien Haddad, Victor Lagerkvist, Biman Roy

► **To cite this version:**

Miguel Couceiro, Lucien Haddad, Victor Lagerkvist, Biman Roy. On the Interval of Boolean Strong Partial Clones Containing Only Projections as Total Operations. 47th IEEE International Symposium on Multiple-Valued Logic (ISMVL), May 2017, Novi Sad, Serbia. hal-01504012

HAL Id: hal-01504012

<https://hal.inria.fr/hal-01504012>

Submitted on 8 Apr 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

On the Interval of Boolean Strong Partial Clones Containing Only Projections as Total Operations

Miguel Couceiro*, Lucien Haddad†, Victor Lagerkvist‡, Biman Roy§

*LORIA (CNRS - Inria Nancy Grand Est - Université de Lorraine) B.P. 239 54506 Vandœuvre-les-Nancy Cedex, France
Email: miguel.couceiro@inria.fr

† Department of Mathematics & Computer Science, Royal Military College of Canada, Kingston, Ontario, Canada
Email: haddad-l@rmc.ca

‡ Institut für Algebra, TU Dresden, Dresden, Germany
Email: victor.lagerkvist@tu-dresden.de

§ Department of Computer and Information Science, Linköpings Universitet, Sweden,
Email: biman.roy@liu.se

Abstract—A *strong partial clone* is a set of partial operations closed under composition and containing all partial projections. Let \mathcal{X} be the set of all Boolean strong partial clones whose total operations are the projections. This set is of practical interest since it induces a partial order on the complexity of NP-complete constraint satisfaction problems. In this paper we study \mathcal{X} from the algebraic point of view, and prove that there exists two intervals in \mathcal{X} , corresponding to natural constraint satisfaction problems, such that one is at least countably infinite and the other has the cardinality of the continuum.

I. INTRODUCTION

A k -ary *polymorphism* of a relation R is a homomorphism from the k th power of R to R . It is well-known that the set of all polymorphisms of a set of relations Γ form a *clone*, that is a set of operations closed under composition and containing all projections. The notion of a clone easily generalizes to partial operations, and it is known that the set of all *partial polymorphisms* of a set of relations Γ , $\text{pPol}(\Gamma)$, form a *strong partial clone*, which is a set of partial operations closed under composition and containing all partial projections. Clone theory is not only a well-studied topic in universal algebra, but has many practical applications in computational complexity. One such example is the *constraint satisfaction problem* over a set of relations Γ ($\text{CSP}(\Gamma)$), which can be viewed as the problem of deciding whether an existentially quantified conjunctive formula over Γ admits at least one model. For example, the NP-complete problem 1-in-3-SAT can be seen as a CSP problem over the ternary relation $R_{1/3} = \{(0, 0, 1), (0, 1, 0), (1, 0, 0)\}$. It has been proven that the polymorphisms of Γ determine the complexity of the CSP problem over Γ up to polynomial-time many-one reductions [5], while the partial polymorphisms of Γ can be used to study the complexity of the CSP problem over Γ with respect to stronger notions of reductions [6], [14]. More specifically, Jonsson et al. [6] proved that if $\text{pPol}(\Gamma) \subseteq \text{pPol}(\Delta)$ then $\text{CSP}(\Delta)$ is solvable in $O(c^n)$ time whenever $\text{CSP}(\Gamma)$ is solvable in $O(c^n)$ time (if Γ and Δ are both finite sets of relations). Hence, the partial polymorphisms of finite sets of relations induce a partial order on the complexity of NP-complete CSPs. Moreover, if Γ is a Boolean set of relations such that each operation in $\text{Pol}(\Gamma)$ is idempotent, then $\text{CSP}(\Gamma)$ is NP-complete if and only if $\text{Pol}(\Gamma)$ consists only of projections [11], [13]. Put together, this implies

that the set of strong partial clones containing only projections as total operations is a particularly interesting object of study, due to its relationship with NP-complete CSP problems. With these observation Jonsson et al. [6] then proved that there exists a relation $R_{1/3}^{\neq 01}$ such that (1) the CSP problem over $R_{1/3}^{\neq 01}$ is NP-complete but (2) there does not exist any NP-complete Boolean CSP problem with a strictly lower worst-case time-complexity. It is worth noting that the relation $R_{1/3}^{\neq 01}$ is also interesting from a purely algebraic point of view since it is known that $\text{pPol}(R_{1/3}^{\neq 01})$ is the largest set of Boolean partial operations which does not contain any total operations (except the projections) [6], [14]. Jonsson et al. [6] then conjectured that the set of strong partial clones between $\text{pPol}(R_{1/3})$ and $\text{pPol}(R_{1/3}^{\neq 01})$ had a particularly simple structure consisting of only five elements. This conjecture turned out to be incorrect, and Lagerkvist & Roy [8] proved that the cardinality of this set is at least countably infinite.

In this paper we continue the investigation of $\text{pPol}(R_{1/3}^{\neq 01})$, with a particular focus on the strong partial clones between $\text{pPol}(R_{1/3})$ and $\text{pPol}(R_{1/3}^{\neq 01})$. After having introduced the basic notions in Section II, we begin (in Section III-B) by proving that there exists at least countably many strong partial clones between $\text{pPol}(R_{1/3})$ and $\text{pPol}(R_{1/3}^{01})$, where $R_{1/3}^{01} = R_{1/3} \times \{(0)\} \times \{(1)\}$. To prove this, we find a class of relations σ^k with the property that $\text{pPol}(R_{1/3}) \subset \text{pPol}(\{\sigma^k, R_{1/3}^{01}\}) \subset \text{pPol}(R_{1/3}^{01})$ and $\text{pPol}(\{\sigma^k, R_{1/3}^{01}\}) \subset \text{pPol}(\{\sigma^{k+1}, R_{1/3}^{01}\})$, for each $k \geq 4$. In Section III-C we study the strong partial clones between $\text{pPol}(R_{1/3}^{01})$ and $\text{pPol}(R_{1/3}^{\neq 01})$, and prove that this set is of continuum cardinality, and therefore strengthen the results from Lagerkvist & Roy [8]. Hence, even for extremely simple relations such as $R_{1/3}$, trying to fully determine the strong partial clones between $\text{pPol}(R_{1/3})$ and $\text{pPol}(R_{1/3}^{\neq 01})$ appears to be an insurmountable task.

II. PRELIMINARIES

In this section we introduce the basic terminology that will be used throughout the paper.

A. Operations and Relations

Let X be a finite and non-empty set of values. Without loss of generality we may assume that $X \subset \mathbb{N}$. In the particular case

when $|X| = 2$ we assume that $X = \{0, 1\}$ and denote this set by \mathbb{B} . A k -ary operation over X , $k \geq 1$, is a function $X^k \rightarrow X$. We write OP_X for the set consisting of all operations over the set X . We will sometimes denote Boolean operations by their defining logical formulas, and for example write \bar{x} for the operation $1 - x$, and we use $x \oplus y$ to denote addition modulo 2. A k -ary partial operation over X is a map $D \rightarrow X$ where $D \subseteq X^k$, and we write PAR_X for the set of all partial operations over X . The set D is called the *domain* of the partial operation f and we let $\text{dom}(f) = D$. For both total and partial operations we let $\text{ar}(f)$ denote the arity of f . If f and g are two k -ary partial operations g is said to be a *suboperation* of f if $\text{dom}(g) \subseteq \text{dom}(f)$ and $f(x) = g(x)$ for all $x \in \text{dom}(g)$. We write $f|_{D'}$ for the suboperation obtained by restricting f to a set $D' \subseteq \text{dom}(f)$.

A k -ary relation over X is a subset of X^k . The set of all relations over X is denoted by Rel_X . For $R_1, R_2 \in \text{Rel}_X$ of arity n and m , we let $R_1 \times R_2 = \{(x_1, \dots, x_{n+m}) \mid (x_1, \dots, x_n) \in R_1, (x_{n+1}, \dots, x_{n+m}) \in R_2\}$ denote their Cartesian product. Given a k -ary tuple $t = (x_1, \dots, x_k) \in X^k$ we let $t[i] = x_i$ denote the i th component of t , and we let $\text{Proj}_{i_1, \dots, i_{k'}}(t) = (x_{i_1}, \dots, x_{i_{k'}})$, $1 \leq k' \leq k$, denote the projection of t on the coordinates $i_1, \dots, i_{k'} \in \{1, \dots, k\}$. Similarly, if R is a k -ary relation we let $\text{Proj}_{i_1, \dots, i_{k'}}(R) = \{\text{Proj}_{i_1, \dots, i_{k'}}(t) \mid t \in R\}$. As a notational shorthand we let $\vec{0}^k$ and $\vec{1}^k$ denote the k -ary tuples $(0, \dots, 0)$ and $(1, \dots, 1)$.

It will be convenient to view a sequence of tuples t_1, \dots, t_n from a k -ary relation R as a $k \times n$ matrix where the i th column consists of the tuple t_i . We let

$$\text{Rows}(t_1, \dots, t_n) = ((t_1[1], \dots, t_n[1]), \dots, (t_1[k], \dots, t_n[k]))$$

denote the tuple consisting of all rows of this matrix. When the exact ordering is not important we instead write $\text{RowSet}(t_1, \dots, t_n) = \{(t_1[1], \dots, t_n[1]), \dots, (t_1[k], \dots, t_n[k])\}$ to denote the set consisting of all rows. For example, if $t_1 = (0, 0, 0)$, $t_2 = (0, 1, 0)$, and $t_3 = (1, 1, 0)$ then

$$\text{Rows}(t_1, t_2, t_3) = ((0, 0, 1), (0, 1, 1), (0, 0, 1))$$

while

$$\text{RowSet}(t_1, t_2, t_3) = \{(0, 0, 1), (0, 1, 1)\}.$$

We will sometimes represent a relation as the set of models of a first-order formula, and if Γ is a set of relations we use the notation $R(x_1, \dots, x_k) \equiv \varphi(x_1, \dots, x_k)$, where $\varphi(x_1, \dots, x_k)$ is a first-order formula over Γ with the free variables x_1, \dots, x_k , to define the relation

$$R = \{(f(x_1), \dots, f(x_k)) \mid f \text{ is a model of } \varphi(x_1, \dots, x_k)\}.$$

B. Clones

Let Π_X be the set of all projections over $X \subseteq \mathbb{N}$, i.e. all operations π_i^n of the form $\pi_i^n(x_1, \dots, x_i, \dots, x_n) = x_i$. A *clone* over X is a set of operations $\mathcal{C} \subseteq \text{OP}_X$ such that (1) $\mathcal{C} \supseteq \Pi_X$ and (2) \mathcal{C} is closed under functional composition. More formally, the latter condition means that if $f, g_1, \dots, g_m \in \mathcal{C}$, where f has arity m and the functions g_1, \dots, g_m all have the same arity n , then the composition $f \circ (g_1, \dots, g_m)(x_1, \dots, x_n) = f(g_1(x_1, \dots, x_n), \dots, g_m(x_1, \dots, x_n))$ for all $x_1, \dots, x_n \in$

X , is included in \mathcal{C} . For a set of total operations F we let $[F]$ be the smallest clone containing F , and call F a *basis* of $[F]$. A clone \mathcal{C} is said to be *finitely generated* if there exists a finite set F such that $[F] = \mathcal{C}$, and is said to be *infinitely generated* otherwise. Every Boolean clone is known to be finitely generated, and the lattice of Boolean clones, *Post's lattice*, is countably infinite [11].

C. Strong Partial Clones

A *partial projection* is a suboperation of a projection. Let Π_X^p be the set of all partial projections over a set X . A set of partial operations \mathcal{C} is a *strong partial clone* if (1) $\mathcal{C} \supseteq \Pi_X^p$ and (2) \mathcal{C} is closed under composition of partial operations. It is well-known that this definition implies that \mathcal{C} is closed under taking suboperations [12]. Composition of partial operations is defined analogously to the total case, i.e., if $f, g_1, \dots, g_m \in \text{PAR}_X$, $\text{ar}(f) = m$, and $\text{ar}(g_1) = \dots = \text{ar}(g_m) = n$, then $f \circ (g_1, \dots, g_m)(x_1, \dots, x_n) = f(g_1(x_1, \dots, x_n), \dots, g_m(x_1, \dots, x_n))$, and the resulting function is defined for every tuple

$$(x_1, \dots, x_n) \in \bigcap_{i=1}^m \text{dom}(g_i)$$

such that

$$(g_1(x_1, \dots, x_n), \dots, g_m(x_1, \dots, x_n)) \in \text{dom}(f).$$

For a set of partial operations F we let $[F]_s$ be the smallest strong partial clone containing F , and similar to the total case the set F is said to be a *basis* of $[F]_s$. A strong partial clone \mathcal{C} is said to be finitely generated if there exists a finite set of partial operations F such that $[F]_s = \mathcal{C}$, and is infinitely generated otherwise.

D. Galois Connections

It is known that clones and strong partial clones admit relational representations. As a shorthand we let

$$f(t_1, \dots, t_n) = (f(t_1[1], \dots, t_n[1]), \dots, f(t_1[k], \dots, t_n[k]))$$

and say that a relation R is *closed* or *invariant* under f if $f(t_1, \dots, t_n) \in R$ for every sequence $t_1, \dots, t_n \in R$. We also say that f is a *polymorphism* of R and let $\text{Pol}(R)$ denote the set of all polymorphisms of the relation R . A relation R with $\text{Pol}(R) = \Pi_X$ is said to be *strongly rigid*. Similarly, for a set of relations Γ we let $\text{Pol}(\Gamma) = \bigcap_{R \in \Gamma} \text{Pol}(R)$, and it is readily verified that f is a polymorphism of Γ if it preserves every relation in Γ . These notions can be generalized to partial operations, and we say that the n -ary partial operation f is a *partial polymorphism* of a k -ary relation R if $f(t_1, \dots, t_n) \in R$ for every sequence $t_1, \dots, t_n \in R$ such that $(t_1[1], \dots, t_n[1]), \dots, (t_1[k], \dots, t_n[k]) \in \text{dom}(f)$. We then let $\text{pPol}(R)$ denote the set of all partial polymorphisms of the relation R , and let $\text{pPol}(\Gamma)$ denote the set of all partial polymorphisms of the set of relations Γ . It is not difficult to verify that sets of the form $\text{Pol}(\Gamma)$ and $\text{pPol}(\Gamma)$ form clones and strong partial clones, respectively. Moreover, if we let $\text{Inv}(F)$ be the set of all invariant relations under the set of (partial) operations F , it is well-known (cf. Chapter 2.9 in Lau [10]) that $\text{Pol}(\cdot)$ and $\text{Inv}(\cdot)$ give rise to a *Galois connection*.

Theorem 1 ([1], [2], [4], [12]): Let Γ and Γ' be two sets of relations. Then $\Gamma \subseteq \text{Inv}(\text{Pol}(\Gamma'))$ if and only if $\text{Pol}(\Gamma') \subseteq \text{Pol}(\Gamma)$.

An analogous result is known for $\text{Inv}(\cdot)$ and $\text{pPol}(\cdot)$, due to Geiger [4] and Romov [12].

Theorem 2 ([4], [12]): Let Γ and Γ' be two sets of relations. Then $\Gamma \subseteq \text{Inv}(\text{pPol}(\Gamma'))$ if and only if $\text{pPol}(\Gamma') \subseteq \text{pPol}(\Gamma)$.

As a shorthand we let $\langle \Gamma \rangle_{\exists} = \text{Inv}(\text{pPol}(\Gamma))$. It is known that $\langle \Gamma \rangle_{\exists}$ is a set of relations closed under *quantifier-free primitive positive definitions*, (qfpp-definitions) over Γ , i.e., logical formulas of the form

$$\varphi(x_1, \dots, x_n) \equiv R_1(\mathbf{x}_1) \wedge \dots \wedge R_m(\mathbf{x}_m)$$

where each $R_i \in \Gamma \cup \{\text{Eq}\}$ and each \mathbf{x}_i is a tuple of variables over x_1, \dots, x_n . Here, Eq denotes the equality relation $\{(x, x) \mid x \in X\}$ over X . Hence, we can prove that $\text{pPol}(\Gamma) \subseteq \text{pPol}(\Gamma')$ by proving that each relation in Γ' admits a qfpp-definition over Γ .

E. Weak Bases

It turned out that the lattice of partial clones on a set with at least two elements is very complex, and significant efforts were made by several authors to study parts of it. The question of describing the general position of the lattice of all total clones within the lattice of partial clones was raised by D. Lau. For example, given a total clone C , determine cardinality of the set of the partial clones on X whose total component is C . For \mathbb{B} , this problem was completely solved in [3] and the corresponding problem for strong partial clones was solved in [15]. We need the following definition:

Definition 3: Let C be a clone over X . We let

$$\text{Int}(C) = \{\text{pPol}(\Gamma) \mid \Gamma \subseteq \text{Rel}_X, \text{Pol}(\Gamma) = C\}.$$

We will sometimes be concerned with even more fine-grained sets of strong partial clones and, given two strong partial clones C_1 and C_2 , write $\text{Int}(C_1, C_2)$ to denote the set

$$\{\text{pPol}(\Gamma) \mid \Gamma \subseteq \text{Rel}_X, C_1 \subseteq \text{pPol}(\Gamma) \subseteq C_2\}.$$

It is worth noting that the smallest element in the set $\text{Int}(C)$ is simply $[C]_s$, the strong partial clone obtained by closing C under suboperations, but it is also known that each such set admits a largest element.

Theorem 4: [14] Let C be a finitely generated clone. Then

$$\bigcup_{\text{pPol}(\Gamma) \in \text{Int}(C)} \text{pPol}(\Gamma) \in \text{Int}(C).$$

It is also known that whenever $\text{Inv}(C)$ is finitely generated, there exists a relation R over X such that

$$\text{pPol}(R) = \bigcup_{\text{pPol}(\Gamma) \in \text{Int}(C)} \text{pPol}(\Gamma).$$

Using the terminology of Schnoor and Schnoor [14] we call a relation satisfying this condition a *weak basis* of $\text{Inv}(C)$. Moreover, for the Boolean domain, all weak bases have been fully described [7]. In this paper we

have a particular interest in the weak basis $R_{1/3}^{\#\#\#01} = \{(0, 0, 1, 1, 1, 0, 0, 1), (0, 1, 0, 1, 0, 1, 0, 1), (1, 0, 0, 0, 1, 1, 0, 1)\}$ of $\text{Rel}_{\mathbb{B}}$. Note that $\text{Proj}_{1,2,3}(R_{1/3}^{\#\#\#01}) = R_{1/3}$, that the third, fourth, and fifth arguments are complement of the three first, and that the two last arguments are constant 0 and 1, respectively. As remarked in Section I, $\text{pPol}(R_{1/3}^{\#\#\#01})$ can be viewed as the set of all Boolean partial operations f such that $[\{f\}]_s \cap \text{OP}_{\mathbb{B}} = \Pi_{\mathbb{B}}$. Hence, it is the largest Boolean strong partial clone whose only total operations are the projections. Somewhat surprisingly, it is known that this strong partial clone is not finitely generated, in contrast to the strong partial clone consisting of all Boolean partial operations (which can be generated by a partial Sheffer operation).

Theorem 5: [9] Let Γ be a finite set of relations over a finite domain such that $\text{Pol}(\Gamma)$ has a basis consisting of unary operations. Then $\text{pPol}(\Gamma)$ is infinitely generated.

Since $\text{Pol}(\{R_{1/3}^{\#\#\#01}\}) = \Pi_{\mathbb{B}}$, it can be generated by a single projection, and so by Theorem 5 we have that $\text{pPol}(R_{1/3}^{\#\#\#01})$ is infinitely generated.

III. THE STRUCTURE BETWEEN $\text{pPol}(R_{1/3})$ AND $\text{pPol}(R_{1/3}^{\#\#\#01})$

Jonsson et al. [6] conjectured that $\text{Int}(\text{pPol}(R_{1/3}), \text{pPol}(R_{1/3}^{\#\#\#01}))$ consisted of $\text{pPol}(R_{1/3})$, $\text{pPol}(R_{1/3}^{01})$, $\text{pPol}(R_{1/3}^{01})$, $\text{pPol}(R_{1/3}^{\#\#\#01})$, and $\text{pPol}(R_{1/3}^{\#\#\#01})$, where $R_{1/3}^{\#\#\#01} = \text{Proj}_{1,2,3,4,5,7,8}(R_{1/3}^{\#\#\#01})$, $R_{1/3}^{01} = \text{Proj}_{1,2,3,4,6,7}(R_{1/3}^{\#\#\#01})$, and $R_{1/3}^{01} = R_{1/3} \times R^0 \times R^1$. This was disproven by Lagerkvist & Roy who proved that $\text{Int}(\text{pPol}(R_{1/3}^{01}), \text{pPol}(R_{1/3}^{\#\#\#01}))$ is at least countably infinite [8]. However, the questions of whether $\text{pPol}(R_{1/3})$ is covered by $\text{pPol}(R_{1/3}^{01})$, and whether $\text{Int}(\text{pPol}(R_{1/3}^{01}), \text{pPol}(R_{1/3}^{\#\#\#01}))$ is in fact of continuum cardinality, were left open. After introducing some simplifying notation in Section III-A, we answer both of these questions in Section III-B and Section III-C. Before reading these sections in greater detail, the reader may first consult Figure 1 for a visualization of the main results.

A. Partial Polymorphisms of Strongly Rigid Relations

In order to describe the partial polymorphisms of strongly rigid relations we make the following definition.

Definition 6: Let R be a relation and f a k -ary partial operation. We let

$$\text{Cover}_R(\text{dom}(f)) = \{ \text{RowSet}(t_1, \dots, t_k) \mid t_1, \dots, t_k \in R, \text{RowSet}(t_1, \dots, t_k) \subseteq \text{dom}(f) \}.$$

This immediately gives rise to the following lemma.

Lemma 7: Let R be a strongly rigid relation and f a partial operation over a finite set X . If $f|_C \in \Pi_X^p$ for every $C \in \text{Cover}_R(\text{dom}(f))$ then $f \in \text{pPol}(R)$.

For $R_{1/3}$ and related relations the following concept can be used to characterize $\text{Cover}_{R_{1/3}}(\text{dom}(f))$. For $k, k' \geq 1$, a set $\{\omega_1, \dots, \omega_{k'}\} \subseteq \mathbb{B}^k$ is an *exact k' -cover* if $\omega_1[i] + \dots + \omega_{k'}[i] = 1$ for every $i \in \{1, \dots, k\}$.




Fig. 1. A visualization of $\text{Int}(\text{pPol}(R_{1/3}), \text{pPol}(R_{1/3}^{\neq 01}))$. A directed arrow from Γ to Δ means that $\text{pPol}(\Gamma) \subset \text{pPol}(\Delta)$. Some inclusions are omitted.

B. $\text{pPol}(R_{1/3})$ and $\text{pPol}(R_{1/3}^{01})$

In this section we study the structure of the set $\text{Int}(\text{pPol}(R_{1/3}), \text{pPol}(R_{1/3}^{01}))$ and prove that it is at least countably infinite. First we define a useful class of relations.

Definition 8: For each $k \geq 4$ let σ^k denote the relation

$$\{(x_1, \dots, x_k, \overline{x_1}, \dots, \overline{x_k}, y_1, \dots, y_{k-3}, \overline{y_1}, \dots, \overline{y_{k-3}}, 0) \mid \sum_{i=1}^k x_i = 1, y_1 + x_1 + x_2 = 1, \dots, y_{k-3} + \sum_{i=1}^{k-2} x_i = 1\}.$$

Hence, if $t_1, \dots, t_{k'} \in \sigma^k$ then $\text{RowSet}(t_1, \dots, t_{k'})$ contains an exact k' -cover, the complement of these tuples, $2(k-3)$ tuples determined by the exact k' -cover, and a constant 0 tuple. For example, the relation σ^5 may be visualized as

$$\begin{pmatrix} 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 \\ 1 & 1 & 1 & 1 & 0 \\ 1 & 1 & 1 & 0 & 1 \\ 1 & 1 & 0 & 1 & 1 \\ 1 & 0 & 1 & 1 & 1 \\ 0 & 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 0 & 0 \\ 1 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 1 & 1 & 1 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}.$$

We first claim that $\text{pPol}(R_{1/3}) \subseteq \text{pPol}(\sigma^4)$. To see this, note that σ^4 can be qfpp-defined as

$$\sigma^4(x_1, x_2, x_3, x_4, x_5, x_6, x_7, x_8, y_1, z_1, C_0) \equiv R_{1/3}(x_1, x_2, y_1) \wedge R_{1/3}(x_3, x_4, z_1) \wedge R_{1/3}(y_1, z_1, C_0) \wedge \bigwedge_{i=1}^4 (x_i, x_{i+4}, C_0).$$

It is then not difficult to see that the partial operation $f(0, 0, 1) = f(0, 1, 0) = f(1, 0, 0) = 0$ preserves σ^4 and $R_{1/3}^{01}$ but not $R_{1/3}$. Since $\text{pPol}(R_{1/3}) \subset \text{pPol}(R_{1/3}^{01})$ it therefore follows that $\text{pPol}(R_{1/3}) \subset \text{pPol}(\{\sigma^4, R_{1/3}^{01}\})$. More generally it also holds that $\text{pPol}(\{\sigma^k, R_{1/3}^{01}\}) \subset \text{pPol}(\{\sigma^{k+1}, R_{1/3}^{01}\})$.

Lemma 9: $\text{pPol}(\{\sigma^k, R_{1/3}^{01}\}) \supset \text{pPol}(\{\sigma^{k-1}, R_{1/3}^{01}\})$ for each $k \geq 5$.

Proof: To prove the inclusion $\text{pPol}(\{\sigma^k, R_{1/3}^{01}\}) \supseteq \text{pPol}(\{\sigma^{k-1}, R_{1/3}^{01}\})$ we show that $\sigma^k \in \langle \{\sigma^{k-1}\} \rangle_{\overline{\mathcal{A}}}$, via the qfpp-definition

$$\sigma^k(x_1, \dots, x_k, x_{k+1}, \dots, x_{2k}, y_1, \dots, y_{k-3}, z_1, \dots, z_{k-3}, C_0) \equiv \sigma^{k-1}(x_1, \dots, x_{k-2}, y_{k-3}, x_{k+1}, \dots, x_{2k-2}, z_{k-3}, y_1, \dots, y_{k-4}, z_1, \dots, z_{k-4}, C_0) \wedge \sigma^{k-1}(x_k, x_{k-1}, \dots, x_3, z_1, x_{2k}, x_{2k-1}, \dots, x_{k+3}, y_1, z_{k-3}, z_{k-4}, \dots, z_1, y_{k-4}, \dots, y_1, C_0).$$

For the proper inclusion, define the $(k-1)$ -ary partial operation f such that $\text{dom}(f) = \text{RowSet}(\sigma^{k-1})$, and let $f(\vec{0}^{k-1}) = 1$ and $f(x) = \pi_1^{k-1}(x)$ otherwise. By definition f does not preserve σ^{k-1} , and it directly follows from Theorem 10 in Lagerkvist & Roy that f preserves $R_{1/3}^{01}$ [8]. Hence, all that remains is to prove that f preserves σ^k , which we do by showing that $\text{Cover}_{\sigma^k}(\text{dom}(f)) = \emptyset$. Assume there exists $t_1, \dots, t_{k-1} \in \sigma^k$ such that $\text{RowSet}(t_1, \dots, t_{k-1}) \subseteq \text{dom}(f)$. Since $\sum_{j=1}^k t_i[j] = 1$ for each $i \in \{1, \dots, k-1\}$ and $k > k-1$, there exists $l \in \{1, \dots, k\}$ such that $t_1[l] = \dots = t_{k-1}[l] = 0$. But due to Definition 8, $t_1[l+k] = \dots = t_{k-1}[l+k] = 1$, implying that $\text{RowSet}(t_1, \dots, t_{k-1}) \not\subseteq \text{Cover}_{\sigma^k}(\text{dom}(f))$ since $\vec{1}^{k-1} \notin \text{dom}(f)$. ■

We have thus proved the following.

Theorem 10: The set $\text{Int}(\text{pPol}(R_{1/3}), \text{pPol}(R_{1/3}^{01}))$ is at least countably infinite.

C. $\text{pPol}(R_{1/3}^{01})$ and $\text{pPol}(R_{1/3}^{\neq 01})$

Following Lagerkvist & Roy [8] we define the following class of relations.

Definition 11: Let $k \geq 5$. The relation α^k is defined as

$$\alpha^k(x_1, \dots, x_k, y_1, \dots, y_{k-3}, z_1, \dots, z_{k-3}, w_1, \dots, w_{k-4}, C_0, C_1) \equiv \exists x_{k+1}, \dots, x_{2k} \cdot \bigwedge_{i=3}^{k-2} R_{1/3}(x_1, x_i, w_{i-2}) \wedge R^1(C_1) \wedge \sigma^k(x_1, \dots, x_k, x_{k+1}, \dots, x_{2k}, y_1, \dots, y_{k-3}, z_1, \dots, z_{k-3}, C_0).$$

For each $k \geq 5$, let $\alpha_1, \dots, \alpha_k$ be an enumeration of the tuples in α^k such that $\alpha_1[k] = 1, \alpha_2[k-1] = 1, \dots, \alpha_k[1] = 1$.

Under this enumeration of tuples we for example see that the matrix corresponding to α^6 can be visualized as:

$$\begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 1 & 1 & 1 & 0 & 0 \\ 1 & 1 & 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 & 1 & 1 \\ 0 & 0 & 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 0 & 1 & 0 \\ 1 & 1 & 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 1 & 1 & 1 & 1 & 1 \end{pmatrix}.$$

Let $\mathcal{V}_k = \text{Rows}(\alpha_1, \dots, \alpha_k)[2(k-2)+2(k-3)]$, and define the k -ary partial operation f^k as $\text{dom}(f^k) = \text{RowSet}(\alpha_1, \dots, \alpha_k)$ and $f^k(x) = \pi_1^k(x)$ for every $x \in \text{dom}(f^k) \setminus \{\mathcal{V}_k\}$ and $f^k(\mathcal{V}_k) = \alpha_1[2(k-2) + 2(k-3)] \oplus 1$. For example, we have that $\mathcal{V}_6 = (1, 1, 0, 1, 1, 0)$ and therefore that $f(\mathcal{V}_6) = 0$. We then have the following result from Lagerkvist & Roy [8].

Theorem 12 (Lagerkvist & Roy [8]): Let $k \geq 5$. Then (1) $\text{pPol}(R_{1/3}^{01}) \subset \text{pPol}(\alpha^k) \subset \text{pPol}(R_{1/3}^{\neq 01})$ and (2) $f^k \notin \text{pPol}(\alpha^k)$ but $f^k \in \text{pPol}(\alpha^{k'})$ for every $5 \leq k < k'$.

We will shortly see that f^k preserves $\alpha^{k'}$ whenever $k' \geq 5$ and $k' \neq k$. First we state the following lemma, whose proof is trivial due to the fact that f^k is defined as a projection on the first argument whenever its arguments are distinct from \mathcal{V}_k .

Lemma 13: Let $k, k' \geq 5$. Then $f_C^k \in \Pi_{\mathbb{B}}^p$ for every $C \in \text{Cover}_{\alpha^{k'}}(\text{dom}(f^k))$ such that $\mathcal{V}_k \notin C$.

Before the proof of the following lemma we make a few observations regarding the domain of the function f^k . Due to Definition 11 this set consists of

- 1) k tuples x_1, \dots, x_k such that $\{x_1, \dots, x_k\}$ forms an exact k -cover,
- 2) $k-3$ tuples y_1, \dots, y_{k-3} such that $\{x_1, \dots, x_i, y_{i-1}\}$, $2 \leq i \leq k-2$, forms an exact k -cover,
- 3) $k-3$ tuples $\overline{y}_1, \dots, \overline{y}_{k-3}$ such that $\{x_k, \dots, x_{i+1}, \overline{y}_{i-1}\}$, $2 \leq i \leq k-2$, forms an exact k -cover,
- 4) $k-4$ tuples z_1, \dots, z_{k-4} such that $\{x_1, x_i, z_{i-2}\}$ forms an exact k -cover for $3 \leq i \leq k-2$, and
- 5) two constant tuples $\overline{0}^k$ and $\overline{1}^k$.

It is also worth remarking that each y_i satisfies the condition that $y_i[j] = 1$ if $1 \leq j \leq k-i+1$ and $y_i[j] = 0$ if $k-i+1 < j \leq k$, and that $z_i[k] = 0$, $z_i[i+2] = 0$, and $z_i[j] = 1$ otherwise. In particular, $\text{dom}(f^k)$ contains the tuple \mathcal{V}_k , which satisfies $\mathcal{V}_k[3] = 0$ and $\mathcal{V}_k[k] = 0$, and $\mathcal{V}_k[i] = 1$ otherwise. We are now ready to prove the main result of this section.

Lemma 14: Let $k \geq 5$. Then $f^k \notin \text{pPol}(\alpha^k)$ but $f^k \in \text{pPol}(\alpha^{k'})$ for every $k' \geq 5$ such that $k' \neq k$.

Proof: Assume first that $k' > k$. Then the result follows from Theorem 12. Hence, in the forthcoming proof, assume that

$k' < k$. We will prove that whenever $C \in \text{Cover}_{\alpha^{k'}}(\text{dom}(f^k))$ then $\mathcal{V}_k \notin C$, which by combining Lemma 7 and Lemma 13 proves that $f^k \in \text{pPol}(\alpha^{k'})$. Hence, for $t_1, \dots, t_k \in \alpha^{k'}$ let

$$(a_1, \dots, a_{k'}, b_1, \dots, b_{k'-3}, c_1, \dots, c_{k'-3}, d_1, \dots, d_{k'-4}, \overline{0}^k, \overline{1}^k)$$

denote the tuples in $\text{Rows}(t_1, \dots, t_k)$. There are now a few cases to consider. However, if $\mathcal{V}_k = a_i$, $1 \leq i \leq k'$, or if $\mathcal{V}_k = b_i$, $\mathcal{V}_k = c_i$ for $1 \leq i \leq k'-3$, then the proof follows the same argument as Lemma 13 in Lagerkvist & Roy [8].

Therefore, assume there exists some $i \in \{1, \dots, k'-4\}$ such that $d_i = \mathcal{V}_k$. This implies that $\{a_1, a_{i+2}, d_i\}$ forms an exact k -cover, and we may without loss of generality assume that $\overline{0}^k \notin \{a_1, a_{i+2}, d_i\}$ as otherwise $\mathcal{V}_k \in \{a_1, a_{i+2}, d_i\}$, which is impossible since $\mathcal{V}_k \notin \text{dom}(f^k)$. Note in particular that this implies that a_1 and a_{i+2} both contain exactly one entry equal to 1. Assume without loss of generality that $a_1[k] = 1$ and that $a_{i+2}[3] = 1$. We will first prove that $\Sigma a_j = 1$ for any $j \in \{1, \dots, k'-2\}$, where $\Sigma a_j = a_j[1] + \dots + a_j[k]$.

First, assume that $a_j = \overline{0}^k$ for some $j \in \{1, \dots, k'-2\}$. The case when $j = 1$ cannot occur since we have already established that $\Sigma a_1 = 1$, and $j = 2$ is impossible since it implies that $b_1 = \overline{a_1} \notin \text{dom}(f)$, due to the fact that $\{a_1, a_2, b_1\}$ must be an exact k -cover. Similarly, we see that $d_{j-2} = \overline{a_1} \notin \text{dom}(f)$ if $\Sigma a_j = 0$ for $j \in \{3, \dots, k'-2\}$, since $\{a_1, a_j, d_{j-2}\}$ must be an exact k -cover.

Second, assume that $\Sigma a_j > 1$ for some $1 \leq j \leq k'-2$. Since the cases when $j = 1$ or $j = i+2$ are impossible we assume that $j \neq 1$ and that $j \neq i+2$. Since $a_1 = (0, 0, \dots, 0, 1)$ and $a_{i-2} = (0, 0, 1, 0, \dots, 0)$, this implies that $a_j = (1, 1, 0, \dots, 0)$, as otherwise $a_j \notin \text{dom}(f^k)$. But then, since $\{a_1, a_j, d_{j-2}\}$ must form an exact k -cover, f^k cannot be defined on d_{j-2} .

Hence, $\Sigma a_j = 1$ for each $1 \leq j \leq k'-2$, but since $k' < k$ this implies that $\Sigma a_{k'-1} > 1$ or that $\Sigma a_{k'} > 1$. We assume that $\Sigma a_{k'-1} > 1$ since the case when $\Sigma a_{k'} > 1$ is entirely symmetric. Due to the assumption that $a_{k'-1} \in \text{dom}(f^k)$, $a_1 = (0, \dots, 0, 1)$ and $a_{i-2} = (0, 0, 1, 0, \dots, 0)$, this in fact implies that $a_{k'-1} = (1, 1, 0, \dots, 0)$. Now, since $\{a_{k'-1}, a_{k'}, c_{k'-2}\}$ must form an exact k -cover, either $a_{k'} = \overline{0}^k$ or $c_{k'-2} = \overline{0}^k$, as otherwise $a_{k'} \notin \text{dom}(f^k)$ or $c_{k'-2} \notin \text{dom}(f^k)$. However, the latter case is impossible since it implies that $a_{k'-1} = \overline{a_{k'}}$, which cannot happen since $\{a_1, \dots, a_{k'}\}$ is an exact k -cover, and since $\Sigma a_j = 1$ for $1 \leq j \leq k'-2$. Hence, $a_{k'} = \overline{0}^k$, and $c_{k'-2} = \overline{a_{k'}}$. Now, since $\Sigma a_j = 1$ for every $1 \leq j \leq k'-2$, it follows that $|\{t_3, \dots, t_k\}| = k-2$, i.e., the tuples t_3, \dots, t_k do not contain any repetitions. However, since $t_1 \notin \{t_3, \dots, t_k\}$, and there cannot exist t_j such that $t_j[k'] = 1$, there are only $k'-2$ tuples to choose from. This is impossible since $k' < k$, and we conclude that $\mathcal{V}_k \notin \text{RowSet}(t_1, \dots, t_k)$. ■

Theorem 15: The set $\text{Int}(\text{pPol}(R_{1/3}^{01}), \text{pPol}(R_{1/3}^{\neq 01}))$ is of continuum cardinality.

Proof: Let $\mathbb{N}_{\geq 5} = \mathbb{N} \setminus \{0, 1, 2, 3, 4\}$. We will prove that there exists an injective function h from $\{X \mid X \subseteq \mathbb{N}_{\geq 5}\}$ to $\text{Int}(\text{pPol}(R_{1/3}^{01}), \text{pPol}(R_{1/3}^{\neq 01}))$. Define $h(X) = \text{pPol}(\{\alpha^i \mid i \in X\})$ for every $X \subseteq \mathbb{N}_{\geq 5}$. We claim that $\text{pPol}(\{\alpha^i \mid i \in X\}) \neq \text{pPol}(\{\alpha^i \mid i \in Y\})$ whenever $X, Y \subseteq \mathbb{N}_{\geq 5}$ and $X \neq Y$. Indeed, if $X \neq Y$ then there exist $i_1 \in X$ or $i_2 \in Y$ such that $i_1 \notin Y$ or $i_2 \notin X$. But then $f_{i_1} \notin \text{pPol}(\{\alpha^i \mid i \in X\})$

and $f_{i_1} \in \text{pPol}(\{\alpha^i \mid i \in Y\})$, or $f_{i_2} \notin \text{pPol}(\{\alpha^i \mid i \in X\})$ and $f_{i_2} \in \text{pPol}(\{\alpha^i \mid i \in Y\})$, due to Lemma 14. ■

IV. CONCLUDING REMARKS

We have studied the set $\text{Int}(\Pi_{\mathbb{B}})$, with a particular focus on describing the strong partial clones between $\text{pPol}(R_{1/3})$ and $\text{pPol}(R_{1/3}^{\neq 01})$. By generalizing the results from Lagerkvist & Roy [8] we have proven that $\text{Int}(\text{pPol}(R_{1/3}), \text{pPol}(R_{1/3}^{01}))$ is at least countably infinite and that $\text{Int}(\text{pPol}(R_{1/3}), \text{pPol}(R_{1/3}^{\neq 01}))$ is of continuum cardinality. These results open up a few distinct directions for future research. First, note that Lemma 9 raises the possibility that $\text{pPol}(\{R_{1/3}^{01}, \sigma^k\})$ is covered by $\text{pPol}(\{R_{1/3}^{01}, \sigma^{k+1}\})$, i.e., that there does not exist any strong partial clone inbetween. If this is indeed the case then the cardinality of $\text{Int}(\text{pPol}(R_{1/3}), \text{pPol}(R_{1/3}^{01}))$ might be countably infinite. Second, it would be interesting to determine the strong partial clones covered by $\text{pPol}(R_{1/3}^{\neq 01})$. Since $\text{pPol}(R_{1/3}^{\neq 01})$ is infinitely generated (by Theorem 5) a good starting point is to determine whether every strong partial subclone of $\text{pPol}(R_{1/3}^{\neq 01})$ is contained in a maximal strong partial subclone of $\text{pPol}(R_{1/3}^{\neq 01})$, or if there exists a countably infinite chain of strong partial clones in $\text{Int}(\Pi_{\mathbb{B}})$ such that the union of this chain equals $\text{pPol}(R_{1/3}^{\neq 01})$.

ACKNOWLEDGEMENTS

We thank the anonymous reviewers for several helpful suggestions on how to improve the paper. The third author has received funding from the DFG-funded project ‘‘Homogene Strukturen, Bedingungserfullungsprobleme, und topologische Klone’’ (Project number 622397).

REFERENCES

[1] V. G. Bodnarchuk, L. A. Kaluzhnin, V. N. Kotov, and B. A. Romov. Galois theory for Post algebras. I. *Cybernetics*, 5:243–252, 1969.

[2] V. G. Bodnarchuk, L. A. Kaluzhnin, V. N. Kotov, and B. A. Romov. Galois theory for Post algebras. II. *Cybernetics*, 5:531–539, 1969.

[3] M. Couceiro, L. Haddad, K. Scholzel, and T. Waldhauser. A solution to a problem of D. Lau: Complete classification of intervals in the lattice of partial boolean clones. *Multiple-Valued Logic and Soft Computing*, 28(1):47–58, 2017.

[4] D. Geiger. Closed systems of functions and predicates. *Pacific Journal of Mathematics*, 27(1):95–100, 1968.

[5] P. Jeavons, D. Cohen, and M. Gyssens. Closure properties of constraints. *Journal of the ACM*, 44(4):527–548, July 1997.

[6] P. Jonsson, V. Lagerkvist, G. Nordh, and B. Zanuttini. Strong partial clones and the time complexity of SAT problems. *Journal of Computer and System Sciences*, 84:52 – 78, 2017.

[7] V. Lagerkvist. Weak bases of Boolean co-clones. *Information Processing Letters*, 114(9):462–468, 2014.

[8] V. Lagerkvist and B. Roy. A Preliminary Investigation of Satisfiability Problems Not Harder than 1-in-3-SAT. In *Proceedings of the 41st International Symposium on Mathematical Foundations of Computer Science (MFCS-2016)*, volume 58, pages 64:1–64:14, 2016.

[9] V. Lagerkvist and M. Wahlstrom. The power of primitive positive definitions with polynomially many variables. *Journal of Logic and Computation*, 2016.

[10] D. Lau. *Function Algebras on Finite Sets: Basic Course on Many-Valued Logic and Clone Theory (Springer Monographs in Mathematics)*. Springer-Verlag New York, Inc., Secaucus, NJ, USA, 2006.

[11] E. Post. The two-valued iterative systems of mathematical logic. *Annals of Mathematical Studies*, 5:1–122, 1941.

[12] B.A. Romov. The algebras of partial functions and their invariants. *Cybernetics*, 17(2):157–167, 1981.

[13] T. Schaefer. The complexity of satisfiability problems. In *Proceedings of the 10th Annual ACM Symposium on Theory Of Computing (STOC-78)*, pages 216–226. ACM Press, 1978.

[14] H. Schnoor and I. Schnoor. Partial polymorphisms and constraint satisfaction problems. In N. Creignou, P. G. Kolaitis, and H. Vollmer, editors, *Complexity of Constraints*, volume 5250 of *Lecture Notes in Computer Science*, pages 229–254. Springer Berlin Heidelberg, 2008.

[15] K. Scholzel. Dichotomy on intervals of strong partial Boolean clones. *Algebra Universalis*, 73(3-4):347–368, 2015.