

CENTROIDAL LOCALIZATION GAME

BARTŁOMIEJ BOSEK, PRZEMYSŁAW GORDINOWICZ, JAROSŁAW GRZYTCZUK, NICOLAS NISSE,
JOANNA SOKÓŁ, AND MAŁGORZATA ŚLESZYŃSKA-NOWAK

ABSTRACT. One important problem in a network G is to locate an (invisible) moving entity by using distance-detectors placed at strategical locations in G . For instance, the famous *metric dimension* of a graph G is the minimum number k of detectors placed in some vertices $\{v_1, \dots, v_k\}$ such that the vector (d_1, \dots, d_k) of the distances $d(v_i, r)$ between the detectors and the entity's location r allows to uniquely determine r whatever be $r \in V(G)$. In a more realistic setting, each device does not get the exact distance to the entity's location. Rather, given locating devices placed in $\{v_1, \dots, v_k\}$, we get only relative distances between the moving entity's location r and the devices (roughly, for every $1 \leq i, j \leq k$, it is provided whether $d(v_i, r) >$, $<$, or $=$ to $d(v_j, r)$). The *centroidal dimension* of a graph G is the minimum number of devices required to locate the entity, in one step, in this setting.

In this paper, we consider the natural generalization of the latter problem, where vertices may be probed sequentially (i.e., in several steps) until the moving entity is located. Roughly, at every turn, a set $\{v_1, \dots, v_k\}$ of vertices are probed and then the relative order of the distances between the vertices v_i and the current location r of the moving entity is given. If it not located, the moving entity may move along one edge. Let $\zeta^*(G)$ be the minimum k such that the entity is eventually located, whatever it does, in the graph G .

We first prove that $\zeta^*(T) \leq 2$ for every tree T and give an upper bound on $\zeta^*(G \square H)$ in cartesian product of graphs G and H . Our main result is that $\zeta^*(G) \leq 3$ for any outerplanar graph G . We then prove that $\zeta^*(G)$ is bounded by the pathwidth of G plus 1 and that the optimization problem of determining $\zeta^*(G)$ is NP-hard in general graphs. Finally, we show that approximating (up to a small constant distance) the location of the robber in the Euclidean plane requires at most two vertices per turns.

1. INTRODUCTION

The problem of locating or capturing an intruder in a graph has been widely studied using many different approaches. One approach is to place detection devices at some vertices of the graph such that these devices precisely determine the position of the intruder at any moment and wherever it is. For instance, this is the approach taken by identifying codes, metric bases and centroidal bases.

Recall that a *dominating set* $D \subseteq V$ of a graph $G = (V, E)$ is a set such that any vertex of V is in the closed neighbourhood of some vertex of D . A vertex u *separates* two vertices v and w if it is in the closed neighbourhood of exactly one of them. A set $D \subseteq V$ separates the vertices of a set X if, for every two vertices $v, w \in X$, there exists $u \in D$ which separates them. A set D is an *identifying code* of a graph G if it is dominating and separates all vertices from $V(G)$ [15]. Similarly, Slater defined the notion of *locating-dominating set*, that is a dominating set D separating vertices of $V(G) \setminus D$ [23, 24] (see [16] and the references

This work was supported by The National Center for Research and Development under the project PBS2/B3/24/2014, by the ANR project Stint (ANR-13-BS02-0007) (N. Nisse) and the associated Inria team AIDyNet (N. Nisse).

therein). Both previous approaches model the situation when the detection devices can detect an intruder at distance at most one from them.

The case when the devices have a longer range of detection has also been considered. For instance, Slater considered the case of an infinite range of detection. Precisely, a *locating set* is a vertex set $L = \{w_1, \dots, w_k\} \subseteq V(G)$ such that, for each vertex $v \in V(G)$, the ordered k -tuple $(d(v, w_1), d(v, w_2), \dots, d(v, w_k))$ of distances between the detectors and the intruder vertex v is unique [22, 13] (i.e., the vector of distances allows to determine uniquely the vertex v). The minimum cardinality of a locating set is called *metric dimension* of a graph, denoted by $\text{MD}(G)$, and a locating set of minimal cardinality is called a *metric basis* of G . For more details on the complexity of computing the metric dimension and on bounds on metric dimension of various graph classes, see for instance [10, 11].

The concept of *centroidal bases* (see eg. [9]) is similar to the one of metric bases. Once again, detecting devices placed at the vertices $C = \{c_1, \dots, c_k\} \subseteq V(G)$ are assumed to have unlimited range, but they do not determine the exact distance from the intruder. Instead, the devices report in an order: if c_i is closer to the intruder than c_j , then c_i reports before c_j and, if they are at the same distance from the intruder, then they report simultaneously. In other words, the received information is an ordered partition of C (ordered by nondecreasing distance from the intruder and ties noted). If, for any intruder position $v \in V(G)$, the received information allows us to uniquely determine v , then C is called a *centroidal locating set*. If C is a centroidal locating set of minimal cardinality, then it is said to be a *centroidal basis* of G , and its cardinality is called the *centroidal dimension* of G , denoted by $\text{CD}(G)$. For instance, it is known that, for any n -node graph G with maximum degree at least 2, then $(1 + o(1)) \frac{\ln n}{\ln \ln n} \leq \text{CD}(G) \leq n - 1$ [9].

All above mentioned models aim at locating the intruder at any moment of the time or, equivalently, at any turn. Another approach consists in locating it in a finite number of turns. This is in the vein of the famous Cops and Robber games where a team of cops must capture a (generally) visible robber by moving alternately in a graph (e.g., see the book [3]).

The *localization game* [19, 4, 14] somehow generalizes the notion of metric dimension of a graph by allowing to probe several sets (of bounded size) instead of only one. Of course, between any two probes, the intruder may move (since otherwise, it would be sufficient to check every vertex one by one). Precisely, the localization game is defined as follows. Let $G = (V, E)$ be a simple undirected graph and let $k \geq 1$ be a fixed integer. Two players, the *Cop-player* and the *Robber-player* (the *robber*), play alternately as follows. In the first turn, the robber chooses a vertex $r \in V$ but keeps it in a secret. Then, at every turn, first the Cop-player picks (or probes) k vertices $B = \{v_1, v_2, \dots, v_k\} \in V^k$ and, in return, gets the vector $D(B) = (d_1, d_2, \dots, d_k)$ where $d_i = d_G(r, v_i)$ is the distance (in G) from r to v_i for every $i = 1, 2, \dots, k$. If the location of the robber is uniquely identified thanks to this information, the game ends with the victory of the Cop-player. Otherwise, the robber may move along one edge. The robber wins if its location is never known. Let the *localization number* of G , denoted by $\zeta(G)$, be the least integer k for which the cops have a winning strategy whatever be the strategy of the robber.

This game restricted to $k = 1$ has been introduced by Seager [19], and studied further in [5, 6, 20]. The parameter $\zeta(G)$ can be seen as the game theoretic variant of $\text{MD}(G)$ and, by the definition, $\zeta(G) \leq \text{MD}(G)$ in any graph G . The localization game with many cops has been introduced recently by the authors of this paper in [4] and, independently, in [14]. For instance, in [14], it was shown that $\zeta(G) \leq \lfloor \frac{(\Delta+1)^2}{4} \rfloor + 1$ for any graph G with maximum

degree Δ . The main result in [4] is that $\zeta(G)$ is unbounded in the class of planar graphs (more precisely, in the class of graphs obtained from a tree by adding a universal vertex). Moreover, computing $\zeta(G)$ is NP-hard [4].

The goal of this paper is to propose and study a “generalization” of the centroidal dimension, in the same way as the localization game somehow extends the notion of metric dimension. This is inspired by localization problems in wireless networks (such as the network of Wi-Fi access points). We are interested in locating a person with a mobile device, who may walk along the network changing his position in time. The strength of the signal from the mobile device is proportional to its distance to particular access points. Unfortunately the signal may be easily disturbed by various factors, hence its strength depends highly on the circumstances. On the other hand, the relative order of the strengths of two signals is expected to be invariant regardless of the circumstances.

The *centroidal localization game* is a turn-by-turn 2-Player game that proceeds as follows. First an invisible robber is placed at some vertex r . Then, at every turn, first the Cop-player probes a set $\{v_1, \dots, v_k\}$ of k vertices. In return, the Cop-player receives, for any $1 \leq i < j \leq k$, the information whether $d(v_i, r) = 0$ or $d(v_i, r) = d(r, v_j)$ or $d(v_i, r) < d(r, v_j)$ or $d(v_i, r) > d(r, v_j)$. If the location of the robber is uniquely identified thanks to this information, the game ends with the victory of the Cop-player. Otherwise, the robber may move along one edge. The robber wins if its location is never known. Note that it is not necessary for the Cop-player to probe the vertex occupied by the robber to win.

The *centroidal localization number* of G , denoted by $\zeta^*(G)$, is the minimum k that ensures the victory for the Cop-player whatever be the strategy of the robber. Note that $\zeta^*(G)$ may be viewed as game-theoretical version of the centroidal dimension, and the inequality $\zeta^*(G) \leq \text{CD}(G)$ obviously holds for every graph G . Note also that $\zeta(G) \leq \zeta^*(G)$ holds by definition. Hence, by [4], $\zeta^*(G)$ is unbounded on planar graphs (even in the class of graphs obtained from a tree by adding a universal vertex), while any upper bound proven in this paper holds for $\zeta(G)$ as well. It is also interesting to note that, some results obtained in the context of the localization game, have been proved without using the exact distances but only their relative order. For this reason, from the proof of [14], it is possible to directly derive that

Corollary 1. *For any graph G with maximum degree Δ there is $\zeta^*(G) \leq \lfloor \frac{(\Delta+1)^2}{4} \rfloor + 1$.*

Our results. As a warm-up, we give easy results on the centroidal localization number (Section 2). Then, we show that $\zeta^*(T) \leq 2$ for any tree T (Section 3) and we provide an upper bound on $\zeta^*(G \square H)$ (where \square denotes the cartesian product) in Section 4. Our main result is that $\zeta^*(G) \leq 3$ for any outerplanar graph G (Section 5), which also gives the best known bound for $\zeta(G)$ in this class of graphs. Then we show that deciding whether $\zeta^*(G) \leq k$ is NP-hard in the class of graphs G with diameter 2 (Section 6). Finally, we show that approximating (up to a small constant distance) the location of the robber in the Euclidean plane requires to probe at most two vertices per turns (Section 7). In the final section (Section 8) we set several open problems for future research.

2. WARM-UP

Let us start with very simple observations.

Proposition 2. *$\zeta^*(G) = 1$ iff G is a graph with at most one edge.*

Proof. First let G be a graph with at most one edge. Then G is a collection of isolated vertices and possibly one pair of vertices connected by the edge $e = \{u, v\}$. Note that if the robber starts game in one of the isolated vertices, then he will remain in the same location throughout the whole game. The strategy for the Cop-player is to first check all isolated vertices one by one. If by doing so she doesn't catch the robber then he is in u or v . Then the Cop-player checks u and either immediately catches the robber, or knows that he is in v .

It is easy to see that if G has more than 1 edge, then the robber should choose one of the edges and move between between it's vertices. the Cop-player can locate the robber only if he checks the exact vertex with the robber, which can always be avoided. \square

Proposition 3. *For any graph G , $\zeta^*(G) = \max\{\zeta^*(C) \mid C \text{ is a connected component of } G\}$.*

Proof. The result is obvious if G has at most one edge. Otherwise, by the above Proposition, we may assume that the Cop-player can probe at least two vertices per turn. We show that, by checking two vertices in each round the Cop-player can determine the component of G containing the robber.

Assume G has more than one component. First note that the robber will never leave the component of the vertex of his first location. Moreover only vertices from this component are at finite distance from him. Say the Cop-player chooses vertices c_1, c_2 , each from different component. Then at most one of c_1, c_2 are at finite distance from the robber. If their distances from the robber are equal, then both are from different component than the robber. Otherwise the component of vertex with smaller distance contains the robber. \square

Proposition 4. *Every bipartite graph G with partition classes of size a and b satisfies $\zeta^*(G) \leq \max(2, \min(a, b))$.*

Proof. As the Cop-player can probe at each round at least 2 vertices, by Proposition 3 we may assume that G is connected. Assume $a \leq b$.

When $a = 1$ the Cop-player may use the following strategy. Check at each round the vertex of least partition class, which forces the robber not to move (otherwise he will be caught immediately). Second probed vertex is chosen from another partition class checking the whole class in a sequence. Eventually exact location of the robber will be checked.

When $a > 1$ at first round the Cop-player probes all a vertices from least partition class. Either the robber is caught or there is some vertex, say v in the set S_1 (we know that the robber is in neighbourhood of v). In further rounds of the game the Cop-player probes $a - 1$ vertices from least partition class except v and step by step one vertex, say u_i , from $N(v)$. Note, that when the robber moves, he can either go to some probed vertex (he is then caught immediately) or to vertex v . But then $d(u_i, v) = 1$, while other distances are at least 2, hence the robber is located at v . Therefore, the robber is forced to stay in a vertex chosen at the first round. Eventually exact location of the robber will be checked. \square

It is easy to note, that for a path P_n for $n \geq 2$ there is $\zeta^*(P_n) = 2$. Indeed, the Cop-player wins sweeping the path from one of its ends probing at each step 2 adjacent vertices. Since paths are precisely the graphs with pathwidth one, this remark can be generalized as follows.

A *path-decompositon* of a graph $G = (V, E)$ is a sequence $\mathcal{X} = (X_1, \dots, X_t)$ of subsets of V , called *bags*, such that, for every edge $\{u, v\} \in E$, there exists a bag containing both u and v , and such that, for every $1 \leq i \leq k \leq j \leq t$, $X_i \cap X_j \subseteq X_k$. The *width* of \mathcal{X} equals $\max_{1 \leq i \leq t} |X_i| - 1$ and the *pathwidth* of G , denoted by $\text{pw}(G)$, is the minimum width of its path-decompositions. Pathwidth and path-decompositions are closely related to some kind of pursuit-evasion games [2].

Proposition 5. *Every graph G satisfies $\zeta^*(G) \leq \text{pw}(G) + 1$.*

Proof. By Proposition 3 we may assume that G is connected (as $\text{pw}(G) = 0$ means that $E(G) = \emptyset$). Let (X_1, \dots, X_t) be an optimal path-decomposition (of width $\text{pw}(G)$) of G and such that, for every $1 < i \leq t$, $|X_{i-1} \setminus X_i| \geq 1$. The strategy for the Cop-player is to probe X_i for $i = 1$ to t step by step. As, for k, j such that $k \leq i \leq j$, the robber cannot move from any vertex $v \in X_j$ to a vertex $u \in X_k$ without being caught then eventually (for $i = t$), he will be located. \square

3. TREES

Following results of Seager [19, 20] one can deduce that for any tree, say T , $\zeta(T)$ is either 1 or 2. More precisely, she proved that one cop is sufficient to locate a robber on any tree when robber is not allowed to move to a vertex just checked by the Cop-player (in the previous round) and that this restriction is necessary for trees that contains a ternary regular tree of height 2 as a subtree. The same bound holds true for $\zeta^*(T)$ which can be easily proven.

Theorem 6. *If T is a tree with at least 3 vertices, then $\zeta^*(T) = 2$.*

Proof. We will describe a simple recursive strategy for the Cop-player (see Figure 1). In the first round she picks a pair of adjacent vertices c_1 and c_2 . Assuming the robber is not




FIGURE 1. Cop's strategy for trees.

immediately caught, he says which of distances d_1 and d_2 from his current position to c_1 , respectively to c_2 , is smaller. As each edge of the tree is a bridge (cut-edge) it splits the tree into 2 subtrees and information about d_1 and d_2 allows the Cop-player to determine the subtree in which the robber is hiding. W.l.o.g., let us assume that $d_2 < d_1$. Then the robber is staying in the subtree containing c_2 , denoted by T_2 , otherwise he is staying in some subtree containing other neighbor of c_1 .

Suppose he is in T_2 . Then in the second round the Cop-player will pick again c_2 and some neighbor c_3 of c_2 in T_2 . After getting a response from the robber she will know if the robber is hiding in a subtree of T_2 containing c_3 or in some subtree of T_2 containing other neighbor of c_2 . The process goes on, decreasing the size of the subtree where the robber can be, until the robber is caught. \square

4. CARTESIAN PRODUCT OF GRAPHS

Another result concerns the Cartesian product of graphs. Recall that the Cartesian product of graphs G and H , is the graph $G \square H$ with vertex set $V(G \square H) = V(G) \times V(H)$ in which (u, v) is adjacent to (u', v') if and only if either $uu' \in E(G)$ and $v = v'$ or $u = u'$ and $vv' \in E(H)$.

Theorem 7. *For any graphs G and H there is*

$$\zeta^*(G \square H) \leq \max\{\Delta(G) + \Delta(H) + 1, \Delta(G) + \zeta^*(H), \zeta^*(G) + \Delta(H)\}.$$

Proof. Let G, H be two graphs. The winning strategy for the Cop-player is divided into 2 phases. During the first phase, probing at most $\Delta(G) + \Delta(H) + 1$ vertices per turn, the Cop-player will chase the robber to know the exact value of at least one coordinates of his position. Suppose w.l.o.g. that known coordinate is the one from the graph G . Then, in the second phase, probing at most $\Delta(G) + \zeta^*(H)$ per turn, the Cop-player will locate the robber using the strategy for graph H while always maintaining its knowledge on the G 's coordinate of the robber.

Details goes as follows. Let us describe the first phase. Suppose the robber is hidden at vertex (r_G, r_H) . Let $(u, v) \in V(G \square H)$. Probing at most $\Delta(G) + \Delta(H) + 1$ vertices, the Cop-player probes each vertex in the set $N_{G \square H}[(u, v)]$. Note that $d_{G \square H}((u, v), (r_G, r_H)) < d_{G \square H}((u_G, v), (r_G, r_H))$ for all $u_G \in N_G(u)$ if and only if $u = r_G$. Symmetrically, $d_{G \square H}((u, v), (r_G, r_H)) < d_{G \square H}((u, v_G), (r_G, r_H))$ for all $v_G \in N_G(v)$ if and only if $v = r_H$. Therefore, the Cop-player can recognize when he has found one of the coordinate of the robber's location and, in this case, the second phase starts. Let us show that the Cop-player eventually achieves this situation. Suppose $u \neq r_G$ and $v \neq r_H$. Comparing distances of vertices (\cdot, v) and (u, \cdot) with (u, v) , the Cop-player may recognize a shortest path to the robber position in both graphs G and H . Suppose $d_G(u_G, r_G) < d_G(u, r_G)$ and $d_H(v_H, r_H) < d_H(v, r_H)$ with (u, v) for some $u_G \in N_G(u)$ and $v_H \in N_H(v)$. Note that

$$d_{G \square H}((u_G, v_H), (r_G, r_H)) = d_{G \square H}((u, v), (r_G, r_H)) - 2.$$

Therefore even after the robber move the Cop-player is able to decrease the distance from the robber by choosing to probe $N_{G \square H}[(u_G, v_H)]$ in the next round. So, eventually, at least one coordinate of the robber position will be known by the Cop-player.

Now, the second phase proceeds as follows. Suppose, w.l.o.g., that known coordinate is the one from the graph G , say $r_G \in V(G)$. Now, the Cop-player uses the winning strategy in the graph H probing $\zeta^*(H)$ vertices at each round, say $(r_G, v_1), (r_G, v_2), \dots, (r_G, v_{\zeta^*(H)})$ and at most $\Delta(G)$ vertices of the form (u, v_1) for $u \in N(r_G)$. When the robber does not move on G coordinate, this is simply a winning strategy. If the robber moved, then exactly one vertex of the form (\cdot, v_1) , say (u, v_1) shows the minimal distance. Hence, during the next round, the Cop-player may continue the winning strategy on H coordinates, while the new coordinate of the robber in G is u . Therefore, the robber will eventually be located. \square

Mimicking the proof we get immediately an analogue for the (metric) localization game.

Corollary 8. *For any graphs G and H there is*

$$\zeta(G \square H) \leq \max\{\Delta(G) + \Delta(H) + 1, \Delta(G) + \zeta(H), \zeta(G) + \Delta(H)\}.$$

5. OUTERPLANAR GRAPHS

In this section we prove that $\zeta(G) \leq \zeta^*(G) \leq 3$ for any outerplanar graph G .

We first recall some basic notions of graphs. For any $S \subseteq V$, let $N(S) = \{v \in V \setminus S \mid \exists u \in S, \{u, v\} \in E\}$. A set S is a separator if $G \setminus S$ has at least two connected components. A set S is a *minimal separator* if there exist two vertices a and b in distinct connected components of $G \setminus S$ and no proper subset of S separates a and b . Let S be a set of vertices of G . Any connected component A of $G \setminus S$ is called a *full component* (with respect to S) if $N(A) = S$. A set S is a minimal separator if and only if it has at least two full components.

Definition 9. *A graph is outerplanar if it has a planar embedding such that every vertex stands on the outer-face.*

Let us recall basic properties of outerplanar graphs. A *minor* of G is any graph that can be obtained from a subgraph H of G by contracting some edges of H . Recall that an outerplanar graph does not admit $K_{2,3}$ nor K_4 as a minor. In particular, this implies that any minimal separator of an outerplanar graph has at most two vertices (since otherwise, there would be a $K_{2,3}$ minor). Moreover, for any minimal separator S of an outerplanar, if $|S| = 2$, then $G \setminus S$ has exactly two full connected components (otherwise there would be a $K_{2,3}$ as minor).

Theorem 10. *Every outerplanar graph G satisfies $\zeta^*(G) \leq 3$. Moreover, if G has n vertices, then there is a cop strategy using 3 probes per turn that takes $O(n^2)$ turns.*

Proof. From now on, let us assume that an outerplanar embedding of G is given (this can be computed in polynomial-time) and fixed. Note that, once the embedding is fixed, it defines a cyclic ordering (that may be clockwise or counter-clockwise) of the neighbours of each vertex.

The algorithm proceeds as follows, gradually reducing the set of vertices where the robber may be hidden. Initially, the Cop-player probes any vertex v and knows that the robber stands at some vertex of $R = V \setminus \{v\}$ (unless the robber is immediately localized). Then, after the robber's move, it can be only at some vertex of $R \cup \{v\}$.

Now, let us assume that we have reached one of the following two possible situations:

Situation 1: There is a vertex v and $R \subseteq V \setminus v$, where R is a union of connected component(s) of $V \setminus \{v\}$, such that the robber stands at some vertex of $R \cup \{v\}$ and this is the turn of the Cop-player (note that we are in such situation after the first turn); or

Situation 2: There are two vertices $\{u, v\}$ and $R \subseteq V \setminus \{u, v\}$, where R is a union of connected component(s) of $V \setminus \{u, v\}$ such that the robber stands at some vertex of $R \cup \{u, v\}$ and this is the turn of the Cop-player. Moreover, there is at most one connected component of R that is full with respect to $\{u, v\}$ and there is a uv -path in $G \setminus R$ (possibly u and v are adjacent).

We show how the algorithm proceeds to reach a new such situation and strictly reducing the size of R , or going from Situation 2 to Situation 1 (i.e., reducing the size of the considered separator of R).

First, let us consider the Situation 1.

- If R is not connected, let X be any connected component of R , the Cop-player checks whether the robber is located in X . For this purpose, let $\{v_1, \dots, v_d\}$ be the neighbors of $v = v_0$ in X in the order they appear in the outer-face (w.l.o.g., clockwise). Note that, by outer-planarity, for any connected component Y of $X \setminus \{v_1, \dots, v_d\}$, $N(Y) \subseteq \{v_i, v_{i+1}\}$ for some $0 \leq i < d$.

The Cop-player sequentially probes $\{v_0, v_i, v_{i+1}\}$ for i from 1 to $d - 1$.

If, for some $1 \leq i < d$, the probe at $\{v_0, v_i, v_{i+1}\}$ indicates that the robber is closer to v_i or v_{i+1} than it is to v , then the robber is necessarily in X and we reach Situation 1 for v and $X \subsetneq R$.

Otherwise, we show that the robber cannot be in X . Indeed, for every $1 \leq i < d$, let Y_i be the union of the connected components Y of X such that $N(Y) \subseteq \{v = v_0, v_1, \dots, v_{i+1}\}$ (in particular, $N(Y_i) \subseteq \{v = v_0, v_1, \dots, v_{i+1}\}$). Note that $Y_{d-1} = X \setminus N(v)$. By induction on i , let us assume that $\{v_0, v_j, v_{j+1}\}$ have been probed sequentially for j from 1 to $i - 1$ and that the robber cannot be in Y_{i-1} . When $\{v_0, v_i, v_{i+1}\}$ is probed, if the robber occupies a connected component Z of $X \setminus N(v)$ such that $N(Z) \subseteq \{v_i, v_{i+1}\}$ (i.e., in $Y_i \setminus Y_{i-1}$), then its distance to v_i or v_{i+1} should be strictly less than its distance to v_0 . Moreover, if the robber was in $X \setminus Y_i$ before the probe, it must still be the case since $\{v_0, v_{i+1}\}$ separates Y_i from $X \setminus Y_i$ (and also from $G \setminus X$). Hence, unless the robber is caught or detected in $Y_i \setminus Y_{i-1}$, it cannot be in Y_i after this probe.

Eventually, either the robber is detected in X and we reach Situation 1 (for v and X as described above) or we can certify that the robber cannot be in X , and we reach the Situation 1 where v keeps its role and $R \setminus X$ plays the role of R .

- Otherwise, if R is connected and v has a unique neighbor w in R (note that this is the case iff $\{v, w\}$ is a cut-edge/bridge), the Cop-player probes $\{v, w\}$. Then, the robber is immediately localized or occupies some vertex in $R \setminus \{w\}$. We are back to Situation 1 where w plays the role of v and $R \setminus \{w\}$ plays the role of R .
- Finally, let us assume that R is connected and $v = v_0$ has $d \geq 2$ neighbors v_1, \dots, v_d in R where these neighbors are ranked in the order (w.l.o.g., clockwise) they appear in the outer-face. Then, the Cop-player probes $\{v = v_0, v_1\}$. Clearly, since R is connected, at least one connected component C of $R \setminus \{v_1\}$ is full with respect to $\{v, v_1\}$ (i.e., $N(C) = \{v, v_1\}$) and because v_1 is the first (in clockwise order) neighbor of v and by outer-planarity, it can be at most one such a full component. Hence, we have reached Situation 2 for the edge $\{v, v_1\}$ and $R \setminus \{v_1\}$.

Now, let us consider Situation 2, i.e., two vertices $\{u, v\}$ have been probed; after which the robber is in some component $R \subseteq V \setminus \{u, v\}$. So, when it is the cop turn, the robber is at some vertex of $R \cup \{u, v\}$. Moreover, at most one connected component F of R is full with respect to $\{u, v\}$ (by definition of Situation 2).

- First, let us assume that there is a connected component X of R such that $N(X) = \{v\}$. Note that during the previous probe of u and v , the distance between v and the robber must have been strictly less than the distance between u and the robber, since otherwise it is already clear that the robber cannot be in X . In particular, it means that the distance between u and the robber was at least two (during the previous probe).

Let $\{v_1, \dots, v_d\}$ be the neighbors of $v = v_0$ in X in the order (w.l.o.g., clockwise) they appear in the outer-face. The Cop-player sequentially probes $\{v, v_i, v_{i+1}\}$ (odd turns) and $\{u, v, v_{i+1}\}$ (even turns) for i from 1 to $d - 1$.

The fact that u and v are probed every two turns prevents the robber to reach a vertex in $G \setminus R$ without being localized. Indeed, at every such even turn, the distance between the robber and v must be strictly less than the distance between the robber and u (otherwise, it becomes sure that the robber is not in X). In particular, this implies that the robber is at distance at least two from u and so cannot cross u in two turns.

The fact that v and v_{i+1} are probed both during the odd and even i th turns allows the proof of first case of Situation 1 to apply (because $\{v, v_{i+1}\}$ separates Y_i from $X \setminus Y_i$, where Y_i is defined as in the first case of Situation 1).

Therefore, as for the first case of Situation 1, the robber must be in X if and only if there is some odd turn $i < d$ such that the distance between v_i or v_{i+1} and the robber's location r is less than the distance between v_0 and r . If the robber is in X , we reach the Situation 1 for v and $X \subsetneq R$ plays the role of R . Otherwise, we reach the Situation 2 for $\{u, v\}$ and $R \setminus X$ plays the role of R .

- The case is symmetric (exchanging u and v) if there is some connected component of R that is adjacent only to u .
- Hence, if neither of the above cases occurs, we may assume that R is connected and that it is full with respect to $\{u, v\}$. Let $P = (v = w_0, w_1, \dots, w_d, w_{d+1} = u)$ be the shortest path among the paths from v to u with internal vertices in R (let us emphasize that we consider the paths with at least one vertex in R , even if u and v may be adjacent). Such a path exists because R is a full component and is unique by both outer-planarity and the existence of a uv -path in $G \setminus R$. Note that, by outer-planarity, for any connected component Y of $R \setminus P$, there is $0 \leq i \leq d$ such that $N(Y) \subseteq \{w_i, w_{i+1}\}$.

If there is no connected component Y of $R \setminus P$ such that $N(Y) = \{w_0, w_1\}$, then the Cop-player probes $\{u, v, w_1\}$ and (unless the robber is located), we reach Situation 2 for $\{w_1, u\}$ and $R \setminus \{w_1\}$. Hence, let L be the (unique) component of $R \setminus P$ that is full with respect to $\{w_0, w_1\}$. Let $R' = R \setminus (L \cup \{w_1\})$.

The Cop-player first probes $\{u, v, w_1\}$. Let us discuss the result of such a probe. Let r be the (unknown) position of the robber.

- if $d(r, u) < d(v, r)$ and $d(u, r) \leq d(w_1, r)$, then r must be in R' and therefore, we reach Situation 2 for $\{w_1, u\}$ and R' .
- if $d(r, v) < d(u, r)$ and $d(v, r) \leq d(w_1, r)$, then r must be in L and therefore, we reach Situation 2 for $\{w_1, v\}$ and L .
- if $d(r, v) = d(u, r)$, it can be checked that w_1 must be strictly closer to r than u and v (by outer-planarity and by uniqueness of P in R) unless the robber is localized. Therefore, if none of the above two cases holds, then w_1 must be strictly closer to r than u and v . In particular (unless the robber is at w_1 and then is located), it implies that the robber is at distance at least 2 from u and v .

Let us then assume that we are in the latter case (w_1 closer to the robber). The aim of the following strategy is to decide whether the robber is in L or not. An important ingredient for the correctness of the strategy is that u, v and w_1 will be probed every two steps (even turns) which will prevent the robber to leave R without being caught

(since u and v separate R from the rest of the graph). Indeed, as described above, each time u, v and w_1 are probed, either the robber is located either in L or R' , or it must be at distance at least 2 from u and v , and therefore, probing u and v every two steps is sufficient to avoid the robber crossing them.

W.l.o.g., let us assume that, when going clockwise along the outer-face (recall that the embedding is fixed), we first meet v , then w_1 and finally u . Let s_1, \dots, s_t be the neighbors (in clockwise order) of w_1 in L . By outer-planarity, for every connected component Y of $L \setminus \{s_1, \dots, s_t\}$, there is $1 \leq i < t$ such that $N(Y) \subseteq \{s_i, s_{i+1}\}$, or $N(Y) = \{v, s_1\}$. Let Y_0 be the (unique if any) component of $L \setminus \{s_1, \dots, s_t\}$ that is full with respect to $\{v, s_1\}$, and for any $1 \leq i \leq t$, let Y_i be the union of the connected components Y of $L \setminus \{s_1, \dots, s_t\}$ such that $N(Y) \subseteq \{v, s_1, \dots, s_i\}$.

The strategy of the Cop-player proceeds as follows. First the Cop-player probes $\{v, s_1, w_1\}$ (odd turn) and then $\{u, v, w_1\}$ (even turn) and, then, for $1 \leq i < t$, the Cop-player probes $\{s_i, s_{i+1}, w_1\}$ (odd turns) and then $\{u, v, w_1\}$ (even turns).

We will show that the robber is in L if and only if either $\min\{d(v, r), d(s_1, r)\} < d(w_1, r)$ (during the first odd turn), or there is an odd turn $1 \leq i < t$ such that $\min\{d(s_i, r), d(s_{i+1}, r)\} < d(w_1, r)$.

As already mentioned, at every even turn, $d(w_1, r) < \min\{d(u, r), d(v, r)\}$ since otherwise, the robber is located either in L or in R' . Hence, at every odd turn, $d(w_1, r) \leq d(v, r) + 1$ (since the robber moves along at most one edge). Moreover, if the robber is in R' , then at every odd turn, $d(v, r) \geq d(w_1, r)$. Indeed, let Q be a shortest path from $r \in R'$ to v . If Q does not pass through w_1 (otherwise it would hold that $d(v, r) > d(w_1, r)$), then it passes through u and $d(v, r) > d(u, r)$. However, let $d^*(x, r)$ denote the distance between r and a vertex x during the previous even turn (after the probe but before the possible move of the robber), we have that $d^*(w_1, r) < d^*(u, r)$ (as mentioned above) and that $d(w_1, r) \leq d^*(w_1, r) + 1$ and $d(u, r) \geq d^*(u, r) - 1$ (since the robber has moved along at most one edge). Altogether $d(v, r) > d(u, r) \geq d^*(u, r) - 1 \geq d^*(w_1, r) \geq d(w_1, r) - 1$.

Now, let us consider the odd turns one by one. When probing $\{v, s_1, w_1\}$, if $d(v, r) < d(w_1, r)$, then the robber is in L by previous paragraph. Otherwise, if $d(s_1, r) < d(w_1, r)$, then the robber must be in $Y_1 \subseteq L$. Reciprocally, if the robber is in $Y_1 \subseteq L$ either $d(s_1, r) < d(w_1, r)$ or $d(v, r) < d(w_1, r)$. Now, let us assume by induction on i that, after the previous odd turn, it was ensured that the robber was not in Y_i and that the Cop-player now probes $\{s_i, s_{i+1}, w_1\}$. Note that $\{s_i, v\}$ separates Y_i from the rest of the graph. After previous odd and even turns, the robber may (moving twice) only have reached Y_i by crossing s_i (since if it crosses v , it is immediately localized). If such a case occurs, the robber must be at a neighbor of s_i in Y_i when $\{s_i, s_{i+1}, w_1\}$ are probed, and therefore $d(r, s_i) < d(r, w_1)$. Now, if the robber is in $Y_{i+1} \setminus Y_i$ when $\{s_i, s_{i+1}, w_1\}$ are probed, then clearly $\min\{d(s_i, r), d(s_{i+1}, r)\} < d(r, w_1)$.

It only remains to prove that, if $\min\{d(s_i, r), d(s_{i+1}, r)\} < d(r, w_1)$, then the robber must be in L . For purpose of contradiction, let us assume that the robber is in R' and $\min\{d(s_i, r), d(s_{i+1}, r)\} < d(r, w_1)$: any shortest path from r to s_i or s_{i+1} must pass through v , and hence $d(v, r) < d(r, w_1)$ which is a contradiction as shown above.

Therefore, after these $2(t+1)$ rounds, either the robber has been localized in L , in which case we are in Situation 2 for $\{w_1, v\}$ and L , or the robber is known not to be in $Y_t = L$ and we reach Situation 2 for $\{w_1, u\}$ and R' .

□

6. COMPLEXITY

In this section, we prove that the centroidal localization game (i.e., computing ζ^*) is NP-hard. At first let us introduce some related properties. A set $L \subseteq V$ of vertices is called a *locating set* if, for every $u, v \in V \setminus L$, $N[u] \cap L \neq N[v] \cap L$. Note that a locating set must “see” almost all vertices. Formally, for any locating set L , $|V \setminus N[L]| \leq 1$. Indeed, otherwise, there would be two vertices u, v such that $N[u] \cap L = N[v] \cap L = \emptyset$.

In [4] it was proven that the localization game (i.e., computing ζ) is NP-hard. At first it was proven the following

Lemma 11. [4] *Computing a minimum locating set is NP-hard in the class of graphs with diameter 2.*

Then, for any n -node graph G with diameter 2, it was constructed a graph G' by adding $n+1$ pairwise non-adjacent vertices x_1, \dots, x_{n+1} , each of them being adjacent to every vertex of $V(G)$. Note that, because G has diameter 2, then G is an isometric subgraph of G' (i.e., distances are preserved). Finally it was proven that $\zeta(G') = k + 1$, where k is the minimum size of a locating set of G .

Since $\zeta(G') \leq \zeta^*(G')$, it only remains to prove that $\zeta^*(G') \leq k + 1$. However, in [4], the proof that $\zeta(G') \leq k + 1$ only relies on the relative order of the distances and not on the exact distances. Therefore, the proof in [4] actually shows that $\zeta^*(G') \leq k + 1$.

It follows that:

Theorem 12. *The centroidal localization game is NP-hard.*

7. EUCLIDEAN PLANE

In this section, we study the centroidal localization game on the infinite graph with a vertex in every point of the Euclidean plane and edges between points at distance at most 1.

In such a graph, the graph distance between a vertex and the robber is the ceiling of their Euclidean distance. We will show that the Cop-player can locate the robber with an error of at most $2\sqrt{2} + \varepsilon$ by probing only two vertices in each round.

First, let us consider the possible three results of a single round. Firstly, the Cop-player may probe the vertex in which the robber stays, in which case the Cop-player wins immediately. Secondly, the Cop-player may probe two vertices that their graph distances from the robber are the same. In this case the Cop-player could conclude that the robber is within a strap of width at most 1 (see Figure 2). Last, the Cop-player may probe two vertices such that their distances to the robber differ. In this case, the Cop-player can draw a line between those two vertices (consisting of all points of the plane with equal distance from the two chosen vertices) and say that the robber must be on one side of the “bounding” line - the side with vertex with smaller distance from the robber (see Figure 3).

Lemma 13. *There exists a strategy allowing the Cop-player to determine a rectangle containing the robber, in a finite number of turns and probing two vertices each time.*

Proof. For the convenience of the proof, let us denote the vertices chosen by the Cop-player as c_1 and c_2 , and their distances from the robber as d_1 and d_2 . Let $\Delta > 1$ be any constant. We will show that, in a finite number of rounds, the Cop-player can find a rectangle, bounding the location of the robber, by first bounding his location by two vertical lines, and then by




FIGURE 2. Location of the robber when distances are the same





FIGURE 3. Location of the robber when distances differ

two horizontal lines.

To bound the location of the robber by two vertical lines the Cop-player will choose c_1 and c_2 from x axis (let c_1 have smaller x coordinate than c_2). Let us denote by l_1 and l_2 the lines bounding location of the robber. The strategy is as follows:

- if both l_1 and l_2 are already set, then move l_1 to the left by 1, and move l_2 to the right by 1, and end the procedure;
- if $d_1 = d_2$ then location of the robber is now bounded by the two vertical lines (the robber is within a strap of width at most 1 (see Figure 2));
- if $d_1 < d_2$ then set l_2 to be the bounding line created by all points of equal distance from both c_1 and c_2 , and in the next round choose $c_1 - (\Delta, 0)$ and $c_2 - (\Delta, 0)$ (see Figure 4);
- if $d_1 > d_2$ then set l_1 to be the bounding line created by all points of equal distance from both c_1 and c_2 , and in the next round choose $c_1 + (\Delta, 0)$ and $c_2 + (\Delta, 0)$;

Note that this procedure will end, since Δ is strictly bigger than 1. Indeed, let us assume for purpose of contradiction that the strategy always consider the case when $d_1 < d_2$ (the case $d_1 > d_2$ is symmetric). Then, at each round, our procedure moves the bounding line l_2 to the right by $\Delta > 1$, while the robber moves by at most 1 to the right. Hence, at some point, the bounding line l_2 must be further to the right than the robber, which means that the procedure has reached the case $d_1 > d_2$ (and now, the second line l_1 will be defined). At the end of the procedure, we move lines l_1 and l_2 by one (first item above) since one of them

FIGURE 4. Strategy of Cop, when $d_1 < d_2$

bounded location of the robber in previous round.

When location of the robber is bounded by two vertical lines l_1 and l_2 , then the Cop-player can repeat the bounding procedure with choosing c_1 and c_2 from y axis and thus bounding the robbers location by two horizontal lines, with an extra operation of moving the two bounding vertical lines by one at each round (l_1 to the left and l_2 to the right). \square

Lemma 14. *For any $\varepsilon > 0$, in finite number of rounds with choosing 2 vertices, the Cop-player may show a point P such that the distance between the robber and P is at most $2\sqrt{2} + \varepsilon$.*

Proof. For the convenience of the proof, let us denote the two vertices chosen by the Cop-player as c_1 and c_2 , and their distances to the robber as d_1 and d_2 . By the previous lemma, we know that, in finite number of rounds, the Cop-player can bound the area where the robber is located by a rectangle A_0 . At every round, let us denote the minimal and maximal x coordinates of the bounding area by x_1 and x_2 , and the minimal and maximal y coordinates of the bounding area by y_1 and y_2 . The strategy for the Cop-player is as follows.

Probe $c_1 = (x_1, 0)$ and $c_2 = (x_2, 0)$. If $d_1 = d_2$, then the robber is in the intersection of the resulting stripe of width at most 1 and the area A_0 . Otherwise the Cop-player knows that the robber is either on the “left” or “right” side of the vertical line with x coordinate equal to $(x_1 + x_2)/2$. Hence, he is in one half of the previous bounding area. Let A_1 denote the new bounding area of the current location of the robber. Then, in the next round, the robber is in the area $A'_1 = \{p \in \mathbb{R}^2 : \exists z \in A_1 \text{ dist}(p, z) \leq 1\}$.

In the next round choose $c_1 = (0, y_1)$ and $c_2 = (0, y_2)$. If $d_1 = d_2$, then the robber is in the intersection of the resulting stripe of height at most 1 and the area A'_1 . Otherwise, the Cop-player knows that the robber is either on the “upper” or “lower” side of a horizontal line with y coordinate equal to $(y_1 + y_2)/2$. Hence, he is in one half of previous bounding area. Let us denote by A_2 the bounding area of current location of the robber. Then, in the next round, the robber is in bounding area $A'_2 = \{p \in \mathbb{R}^2 : \exists z \in A_2 \text{ dist}(p, z) \leq 1\}$.

The strategy simply repeats these two steps. It is easy to see that the maximal width w_n of the bounding area A_n changes after every second round with the recursive formula $w_n = \frac{w_{n-2}+4}{2}$, unless $d_1 = d_2$ and then $w_n \leq 1$. Similarly, the maximal height h_n of the bounding area A_n changes after every second round with the recursive formula $h_n = \frac{h_{n-2}+4}{2}$, unless $d_1 = d_2$ and then $h_n \leq 1$. Note that the limit of the sequence satisfying the recursive formula is 4. Therefore, for any ε , there exists an integer n such that $w_n \leq 4 + \varepsilon$ and $h_n \leq 4 + \varepsilon$. Hence, the bounding area A_n is contained in a rectangle with side lengths at most $4 + \varepsilon$. So, the distance between every point of A_n and the point $P = (\frac{x_2-x_1}{2}, \frac{y_2-y_1}{2})$ is at most $\frac{(4+\varepsilon)\sqrt{2}}{2} \leq 2\sqrt{2} + \varepsilon$. \square

8. FURTHER WORK

There are many interesting questions, yet to be asked, about Centroidal Coding Game. One would be to find bounds on $\zeta^*(G)$ for special classes of graphs (e.g., hypercube, partial cubes, chordal graphs, disk intersection graphs etc.). In particular: is the bound given by Theorem 10 tight? One can notice without any effort that $\zeta(C_4) = \zeta^*(C_4) = 2$, but are there outerplanar graphs which require 3 cops?

Another problem worth to be considered is the game without the *immediate catch* rule. That is, a variant of the game where the robber could go into checked vertex unnoticed. It is easy to see that, in case of the Euclidean plane, the game does not change. For trees, let us note that the Cop-player can win such a game by probing at most $\Delta(T)$ vertices in each round (by checking all neighbors of a vertex we can tell in which subtree the robber is).

Finally, we ask what happens if the Cop-player is given a set of vertices $S \subset V(G)$ such that she can only choose vertices from this set, meaning in each round $C \subset S$. Such approach might be useful for practical applications.

The question of the computational complexity of ζ^* in various graph classes such as bipartite graphs, bounded treewidth graphs... is also of interest. Finally, most of the interesting turn-by-turn two-player games are known to be PSPACE-hard or even EXPTIME-complete. The exact status of the complexity of the centroidal localization game is still open.

REFERENCES

- [1] M. Aigner and M. Fromme, A game of cops and robbers, *Discrete Appl. Math.* 8, pp. 1–12 (1984).
- [2] D. Bienstock, Graph searching, path-width, tree-width and related problems (a survey). DIMACS Ser. Discret. Math. Theor. Comput. Sci. 5, pp. 3349 (1991)
- [3] A. Bonato and R.J. Nowakowski, The game of cops and robbers on graphs, Student Mathematical Library, vol. 61, American Mathematical Society (2011).
- [4] B. Bosek, P. Gordinowicz, J. Grytczuk, N. Nisse, J. Sokół, M. Śleszyńska-Nowak, Localization game on geometric and planar graphs, manuscript (2017).
- [5] A. Brandt, J. Diemunsch, C. Erbes, J. LeGrand and C. Moffatt, Locating a robber on a tree, manuscript (2015).
- [6] J. Carraher, I. Choi, M. Delcourt, L. H. Erickson, D. B. West, Locating a robber on a graph via distance queries, *Theoretical Computer Science* 463, pp. 54–61 (2012).
- [7] I. Charon, O. Hudry and A. Lobstein, Minimizing the Size of an Identifying or Locating-Dominating Code in a Graph is NP-Hard. *Theoretical Computer Science*, Vol. 290/3, pp. 2109-2120 (2003)
- [8] G. Cohen, I. Honkala, A. Lobstein and G. Zémor, On Identifying Codes. *Proceedings of DIMACS Workshop on Codes and Association Schemes '99*, Vol. 56, pp. 97-109 (2001)
- [9] F. Foucaud, R. Klasing, P.J. Slater, Centroidal Bases in Graphs, *Networks* 64(2): 96-108 (2014)
- [10] F. Foucaud, G. B. Mertzios, R. Naserasr, A. Parreau, P. Valicov, Identification, location-domination and metric dimension on interval and permutation graphs. I. *Bounds. Theor. Comput. Sci.* 668: 43-58 (2017)
- [11] F. Foucaud, G. B. Mertzios, R. Naserasr, A. Parreau, P. Valicov, Identification, Location-Domination and Metric Dimension on Interval and Permutation Graphs. II. *Algorithms and Complexity. Algorithmica* 78(3): 914-944 (2017)
- [12] F. V. Fomin and D. M. Thilikos, An annotated bibliography on guaranteed graph searching, *Theor. Computer Science* 399, pp. 236–245 (2008).
- [13] F. Harary, R. A. Melter, On the metric dimension of a graph, *Ars Combin.* 2, pp. 191–195 (1976).
- [14] J. Haslegrave, R.A.B. Johnson and S. Koch, Locating a robber with multiple probes: arXiv:1703.06482 (2017).
- [15] M. G. Karpovsky, K. Chakrabarty and L. B. Levitin. On a new class of codes for identifying vertices in graphs. *IEEE Transactions on Information Theory* 44:599611, 1998.
- [16] A. Lobstein, Watching systems, identifying, locatingdominating and discriminating codes in graphs: A bibliography, Available at: <http://www.infres.enst.fr/lobstein/debutBIBidetlocdom.pdf>.

- [17] R. J. Nowakowski, P. Winkler, Vertex-to-vertex pursuit in a graph, *Discrete Mathematics* 43, pp. 235–239 (1983).
- [18] A. Quillot. Jeux et pointes xes sur les graphes. Th ese de 3 eme cycle, Universit e de Paris VI, pp. 131–145 (1978).
- [19] S. Seager, Locating a robber on a graph, *Discrete Math.* 312, pp. 3265–3269 (2012).
- [20] S. Seager, Locating a backtracking robber on a tree, *Theor. Computer Science* 539, pp. 28–37 (2014).
- [21] P. D. Seymour, R. Thomas, Graph searching and a min–max theorem for tree-width, *J. Combin. Theory Ser. B* 58, pp. 22–33 (1993).
- [22] P. J. Slater, Leaves of trees, in *Proc. Sixth Southeastern Conf. Combin., Graph Theory, Computing, Congressus Numer.* 14, pp. 549–559 (1975).
- [23] P. J. Slater, Domination and location in acyclic graphs. *Networks* 17(1) 5564, 1987.
- [24] P. J. Slater, Dominating and reference sets in a graph. *Journal of Mathematical and Physical Sciences* 22(4) 445455, 1988.

THEORETICAL COMPUTER SCIENCE DEPARTMENT, FACULTY OF MATHEMATICS AND COMPUTER SCIENCE, JAGIELLONIAN UNIVERSITY, 30-348 KRAKÓW, POLAND

E-mail address: bosek@tcs.uj.edu.pl

INSTITUTE OF MATHEMATICS, LODZ UNIVERSITY OF TECHNOLOGY, ŁÓDŹ, POLAND

E-mail address: pgordin@p.lodz.pl

FACULTY OF MATHEMATICS AND INFORMATION SCIENCE, WARSAW UNIVERSITY OF TECHNOLOGY, 00-662 WARSAW, POLAND

E-mail address: j.grytczuk@mini.pw.edu.pl

UNIVERSITÉ CÔTE D’AZUR, INRIA, CNRS, I3S, FRANCE

E-mail address: nicolas.nisse@inria.fr

FACULTY OF MATHEMATICS AND INFORMATION SCIENCE, WARSAW UNIVERSITY OF TECHNOLOGY, 00-662 WARSAW, POLAND

E-mail address: j.sokol@mini.pw.edu.pl

FACULTY OF MATHEMATICS AND INFORMATION SCIENCE, WARSAW UNIVERSITY OF TECHNOLOGY, 00-662 WARSAW, POLAND

E-mail address: m.sleszynska@mini.pw.edu.pl