

## Global stabilization of stochastic nonlinear systems

Mohamed Bensoubaya, Abdelhak Ferfera, Abderrahman Iggidr

► **To cite this version:**

Mohamed Bensoubaya, Abdelhak Ferfera, Abderrahman Iggidr. Global stabilization of stochastic nonlinear systems. 37th IEEE Conference on Decision and Control, Dec 1998, Tampa, Florida, United States. pp.2376–2377, 10.1109/cdc.1998.757728 . hal-01862873

**HAL Id: hal-01862873**

**<https://hal.inria.fr/hal-01862873>**

Submitted on 29 Aug 2018

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Global stabilization of stochastic nonlinear systems

M. Bensoubaya      A. Ferfera      A. Iggidr

CONGE Project

INRIA Lorraine & University of Metz (Dept. of Mathematics, URA-CNRS 399)

I.S.G.M.P. Bat. A, Ile du Saulcy 57045 Metz cedex 01, France

e.mail :{bensouba,ferfera,iggidr}@loria.fr

## Abstract

We consider stochastic nonaffine nonlinear control systems  $dx_t = f(x_t, u) dt + g(x_t, u) d\omega_t$ ,  $\omega$  being a standard Wiener process, for which we give a sufficient condition for global stabilization by a bounded smooth state feedback which is explicitly given. This condition generalizes the well known Jurdjevic-Quinn result for deterministic affine control systems.

## 1 Introduction

In this paper, we investigate the global stabilization problem for stochastic nonlinear control differential equations (written in the sens of Itô) of the form

$$x_t = x_0 + \int_0^t f(x_s, u) ds + \sum_{j=1}^p \int_0^t g_j(x_s, u) d\omega_s^j \quad (1)$$

where  $u = (u^1, \dots, u^m)^T$  is a bounded  $\mathbb{R}^m$ -valued control law,  $f, g_j : \mathbb{R}^n \times \mathbb{R}^m \rightarrow \mathbb{R}^n$  are bounded smooth ( $\mathcal{C}^\infty$ ) functions with  $f(0, 0) = g_j(0, 0) = 0$ ,  $1 \leq j \leq p$  and  $\{\omega_t, t \geq 0\}$  is a standard  $\mathbb{R}^p$ -valued Wiener process defined on an usual probability space  $(\Omega, \mathcal{F}, P)$ . System (1) is said globally stabilizable if there exists a feedback law  $u : \mathbb{R}^n \rightarrow \mathbb{R}^m$  with  $u(0) = 0$  such that the zero solution of the closed-loop system

$$x_t = x_0 + \int_0^t f(x_s, u(x_s)) ds + \sum_{j=1}^p \int_0^t g_j(x_s, u(x_s)) d\omega_s^j$$

is globally asymptotically stable in probability (G.A.S.P.) (see [3]). Our goal is to give a sufficient condition for global stabilization of (1). This condition generalizes the well known Jurdjevic-Quinn [2] result for deterministic affine control systems (see [4] and references therein)

$$\dot{x} = X(x) + \sum_{i=1}^m u^i Y^i(x) \quad (2)$$

In [4], it is proved that if there exists a positive definite and proper smooth function  $V : \mathbb{R}^n \rightarrow \mathbb{R}$  such that:

(i) the Lie-derivative of  $V$  with respect to  $X$  satisfies  $XV(x) \leq 0, \forall x \in \mathbb{R}^n$ ; (ii) the set  $\{x \in \mathbb{R}^n | X^{k+1}V(x) =$

$X^k Y^i(V) = 0, k \in \mathbb{N}, i = 1, \dots, m\}$  is reduced to  $\{0\}$ ; then system (2) is globally stabilizable by means of the feedback law  $u^i(x) = -Y^i V(x)$ . [1] extends the Jurdjevic-Quinn theorem to the particular class of stochastic control affine nonlinear systems

$$x_t = x_0 + \int_0^t [X_0(x_s) + \sum_{i=1}^m u^i Y_0^i(x_s)] ds + \sum_{j=1}^p \int_0^t X_j(x_s) dz_s^j \quad (3)$$

for which the coefficients associated with the noise do not depend on the control. For (3) the associated infinitesimal generator  $\mathcal{L}$  satisfies  $\mathcal{L}V(x) = LV(x) + \sum_{i=1}^m u^i Y_0^i V(x)$ , where  $L$  is the second order differential operator:  $L = X_0 + \frac{1}{2} \sum_{j=1}^p X_j^2$ , and  $V$  is a Lyapunov function. So, the feedback  $u^i(x) = -Y_0^i V(x)$  yields  $\mathcal{L}V(x) \leq 0$  if  $LV(x) \leq 0$ , which allows to establish in [1], under stochastic analogous conditions to (i) and (ii), that this feedback globally stabilizes (3). Contrarily to (3), for (1) where the random parametric excitation depends on the control, the linearity on  $u$  disappears in  $\mathcal{L}V(x)$ , and so, it is not obvious to prove the existence of a state feedback law  $u(x)$  yielding  $\mathcal{L}V(x) \leq 0$  if  $LV(x) \leq 0$ .

## 2 Main result

In order to state our result, we need the following notations. Let  $X_0, Y_0^i, X_j$  and  $Y_j^i, 1 \leq j \leq p, 1 \leq i \leq m$ , be the vector fields defined on  $\mathbb{R}^n$  by  $X_0(x) = f(x, 0), Y_0^i(x) = (\partial f / \partial u^i)(x, 0), X_j(x) = g_j(x, 0)$  and  $Y_j^i(x) = (\partial g_j / \partial u^i)(x, 0)$ . Moreover, denote by  $L$  and  $L_i, 1 \leq i \leq m$ , the second order differential operators associated with (1) defined for any function  $\Psi$  in  $\mathcal{C}^2(\mathbb{R}^n)$  by

$$L\Psi(x) = \langle X_0(x), \nabla\Psi(x) \rangle + \frac{1}{2} \sum_{j=1}^p \text{Tr} (X_j(x) X_j^T(x) (\partial^2 \Psi / \partial x^2)(x)) \quad (4)$$

$$L_i\Psi(x) = \langle Y_0^i(x), \nabla\Psi(x) \rangle + \frac{1}{2} \sum_{j=1}^p \text{Tr} ([X_j(x) Y_j^i{}^T(x) + Y_j^i(x) X_j^T(x)] (\partial^2 \Psi / \partial x^2)(x)) \quad (5)$$

Besides, for  $x \in \mathbb{R}^n$  and  $v, w \in \mathbb{R}^m$  we set

$$\tilde{f}(x, v, w) = \int_0^1 (1-t) (\partial^2 f / \partial u^2)(x, tv) w^2 dt$$

$$\tilde{g}_j(x, v, w) = \int_0^1 (1-t)(\partial^2 g_j / \partial u^2)(x, tv) w^2 dt$$

where

$w^2 = (w, w)$  and  $(\partial^2 f / \partial u^2)(x, tv)$ ,  $(\partial^2 g_j / \partial u^2)(x, tv) \in \mathcal{L}_2(\mathbb{R}^m \times \mathbb{R}^m, \mathbb{R}^n)$  (the space of bilinear applications from  $\mathbb{R}^m \times \mathbb{R}^m$  to  $\mathbb{R}^n$ ) are respectively the second order derivatives of  $f$  and  $g_j$  with respect to  $u$  at  $(x, tv)$ , and we consider the  $n \times n$  matrices

$$\begin{aligned} A_j(x, v, w) &= X_j(x) \tilde{g}_j^T(x, v, w) + \tilde{g}_j(x, v, w) X_j^T(x) \\ &+ \sum_{i=1}^m v^i \left[ Y_j^i(x) \tilde{g}_j^T(x, v, w) + \tilde{g}_j(x, v, w) Y_j^{iT}(x) \right] \\ &+ \sum_{i_1, i_2=1}^m w^{i_1} w^{i_2} Y_j^{i_1}(x) Y_j^{i_2 T}(x) + \tilde{g}_j(x, v, w) \tilde{g}_j^T(x, v, w) \end{aligned}$$

Finally, we assume that there exists  $V(x)$  proper positive definite and smooth such that

**(h1)**  $LV(x) \leq 0$ ,  $\forall x \in \mathbb{R}^n$  ;

**(h2)** The set  $W = \{x \in \mathbb{R}^n \mid L^{k+1}V(x) = L^k L_i V(x) = 0, k \in \mathbb{N}, i = 1, \dots, m\}$  is reduced to  $\{0\}$  ;

and we set

$$\begin{aligned} \varphi(x, v, w) &= \langle \tilde{f}(x, v, w), \nabla V(x) \rangle \\ &+ \frac{1}{2} \sum_{j=1}^p \text{Tr} (A_j(x, v, w) (\partial^2 V / \partial x^2)(x)) \end{aligned} \quad (6)$$

Notice that the real valued function  $\varphi$  is homogeneous of degree 2 with respect to  $w$ . Using the recursive notation  $L^{k+1}\Psi(x) = LL^k\Psi(x)$ ,  $L^0\Psi(x) = \Psi(x)$ , for  $k \in \mathbb{N}$  and  $\Psi \in C^\infty(\mathbb{R}^n)$ , our main result can now be stated as follows.

**Theorem 1** *If the conditions **(h1)** and **(h2)** hold, then for any  $\eta > 0$  and any smooth functions  $K_1(x)$  and  $K_2(x)$  satisfying,  $\forall x \in \mathbb{R}^n$ ,  $K_1(x) + K_2(x) \neq 0$  and*

$$K_1(x) \geq \sup_{\|v\| \leq \eta, \|w\|=1} |\varphi(x, v, w)| \quad (7)$$

$$K_2(x) \geq \|(L_1 V(x), \dots, L_m V(x))\| \quad (8)$$

*the stochastic control system (1) is globally stabilizable by means of the feedback law*

$$u(x) = \frac{-\eta}{\eta K_1(x) + K_2(x)} (L_1 V(x), \dots, L_m V(x))^T \quad (9)$$

*which satisfies  $\|u(x)\| \leq \eta$ ,  $\forall x \in \mathbb{R}^n$ .*

**Proof:** The inequality  $\|u(x)\| \leq \eta$  is immediate. Moreover, the closed-loop system is given by

$$\begin{aligned} x_t &= x_0 + \int_0^t [X_0(x_s) + \sum_{i=1}^p u^i(x_s) Y_0^i(x_s) \\ &+ \tilde{f}(x_s, u(x_s), u(x_s))] ds + \sum_{j=1}^p \int_0^t [X_j(x_s) \\ &+ \sum_{i=1}^p u^i(x_s) Y_j^i(x_s) + \tilde{g}_j(x_s, u(x_s), u(x_s))] d\omega_s^j \end{aligned} \quad (10)$$

The infinitesimal generator  $\mathcal{L}$  associated with (10) satisfies

$$\begin{aligned} \mathcal{L}V(x) &= \langle X_0(x_s) + \sum_{i=1}^p u^i(x_s) Y_0^i(x_s) \\ &+ \tilde{f}(x_s, u(x_s), u(x_s)), \nabla V(x) \rangle + \frac{1}{2} \sum_{j=1}^p \text{Tr} ([X_j(x_s) \\ &+ \sum_{i=1}^m u^i(x_s) Y_j^i(x_s) + \tilde{g}_j(x_s, u(x_s), u(x_s))] [X_j(x_s) \\ &+ \sum_{i=1}^m u^i(x_s) Y_j^i(x_s) + \tilde{g}_j(x_s, u(x_s), u(x_s))]^T \frac{\partial^2 V}{\partial x^2}(x)) \end{aligned}$$

and by a simple computation one gets from (4), (5) and (6)  $\mathcal{L}V(x) = LV(x) + \sum_{i=1}^m u^i(x) L_i V(x) + \varphi(x, u(x), u(x))$ . It follows that  $\mathcal{L}V(x) = LV(x)$  if  $u(x) = 0$  and otherwise, from (9) and the homogeneity property of  $\varphi(x, v, w)$  with respect to  $w$  one gets

$$\mathcal{L}V(x) = LV(x) - \frac{1}{K(x)} \|u(x)\|^2 [1 - K(x) \varphi(x, u, \frac{u}{\|u\|})]$$

where  $K(x) = \eta / (\eta K_1(x) + K_2(x))$ . Besides, by (7), (8) and  $\|u(x)\| \leq \eta$  one has  $1 - K(x) \varphi(x, u(x), u(x)) / \|u(x)\| \geq 0$  and so one gets from **(h1)**  $\mathcal{L}V(x) \leq 0$ ,  $\forall x \in \mathbb{R}^n$ . Besides, according to the stochastic version of LaSalle's invariance principle (see [3]), the stochastic process  $x_t$  converges in probability to the largest invariant set whose support is contained in the locus  $\mathcal{L}V(x_t) = 0$  for all  $t \geq 0$ . Therefore, in order to prove that the zero solution of (10) is G.A.S.P. it must be shown that for any complete solution  $x_t$  of (10) along which  $\mathcal{L}V(x_t) = 0$  for all  $t \geq 0$ , one has necessarily  $x_t \equiv 0$ . Notice that, from (7), (8) and (9), one has if  $u(x) \neq 0$  then  $K_2(x) \neq 0$  and so  $1 - K(x) \varphi(x, u(x), u(x)) / \|u(x)\| \neq 0$  and, from (9), it turns out that  $\mathcal{L}V(x) = 0$  if and only if  $LV(x) = 0$  and  $L_i V(x) = 0$ ,  $i = 1, \dots, m$ . So, for any complete solution  $x_t$  of (10) for which  $\mathcal{L}V(x_t) = 0$  for all  $t \geq 0$ , successive differentiations by means of Itô's formula yield  $L^{k+1}V(x_t) = L^k L_i V(x_t) = 0$ , for  $t \geq 0$ ,  $k \in \mathbb{N}$ , and  $i = 1, \dots, m$ . Hence, by assumption **(h2)**, it follows that  $x_t \equiv 0$  which completes the proof.

## References

- [1] P. Florchinger. A stochastic version of jurdjevic-quinn theorem. *J. Stoch. Anal. Appl.*, 12:473–480, 1994.
- [2] V. Jurdjevic and J.P. Quinn. Controllability and stability. *J. Diff. Equ.*, 28:381–389, 1978.
- [3] H.J. Kushner. Stochastic stability. In R. Curtain, editor, *Stability of Stochastic Dynamical Systems, Lecture Notes in Mathematics*, volume 294, pages 97–124. Springer-Verlag, Berlin, 1972.
- [4] R. Outbib and G. Sallet. Stabilizability of the angular velocity of a rigid body revisited. *Systems & Control Letters*, 18:93–98, 1992.