



HAL
open science

Advancing dynamical cores of oceanic models across all scales

Florian Lemarié, Hans Burchard, Laurent Debreu, Knut Klingbeil, Jacques Sainte-Marie

► **To cite this version:**

Florian Lemarié, Hans Burchard, Laurent Debreu, Knut Klingbeil, Jacques Sainte-Marie. Advancing dynamical cores of oceanic models across all scales. *Bulletin of the American Meteorological Society*, 2019, 100, pp.ES109-ES115. 10.1175/BAMS-D-18-0303.1 . hal-01939057

HAL Id: hal-01939057

<https://inria.hal.science/hal-01939057>

Submitted on 29 Nov 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Advancing dynamical cores of oceanic models across all scales

FLORIAN LEMARIÉ*

Univ. Grenoble Alpes, Inria, CNRS, Grenoble INP, LJK, 38000 Grenoble, France

HANS BURCHARD

Leibniz Institute for Baltic Sea Research Warnemünde, Seestraße 15, 18119 Rostock, Germany

LAURENT DEBREU

Univ. Grenoble Alpes, Inria, CNRS, Grenoble INP, LJK, 38000 Grenoble, France

KNUT KLINGBEIL

Leibniz Institute for Baltic Sea Research Warnemünde, Seestraße 15, 18119 Rostock, Germany

JACQUES SAINTE-MARIE

Inria Paris, 75012 Paris & Sorbonne Université, Lab. Jacques-Louis Lions, 75252 Paris, France

ABSTRACT

First COMMODORE workshop: community for the numerical modeling of the global, regional and coastal ocean

WHAT: 47 participants from 9 countries representing 15 different oceanic numerical models met to review our current understanding of future challenges in the design of oceanic dynamical cores

WHEN: 17-19 September 2018

WHERE: Paris, France

Oceanic numerical models are used to understand and predict a wide range of processes from global paleoclimate scales to short-term prediction in estuaries and shallow coastal areas. One of the overarching challenges, and the main topic of the COMMODORE workshop, is the appropriate design of the dynamical cores given the wide variety of scales of interest and their interactions with atmosphere, sea-ice, biogeochemistry, and even societal processes. The construction of a dynamical core is a very long effort which takes years and decades of research and development and which requires a collaborative mixture of scientific disciplines. This work involves a significant number of fundamental choices, such as which equations to solve, which horizontal and vertical grid arrangement is adequate, which discrete algorithms allows

jointly computational efficiency and sufficient accuracy, etc. Nowadays, a broad range of numerical methods are implemented in models used for realistic ocean simulations, and, owed to the advances in computational power, a meeting point has been reached between global circulation models and regional local models such that there can be mutual benefits of a cross-fertilization between communities. This report outlines an initiative to bring together the world-wide leading researchers actively contributing to the development of oceanic model dynamical cores, such that participants could network together and focus on next challenges irrespective of target applications (regional, coastal, or global). The first community for the numerical modeling of the global, regional and coastal ocean (COMMODORE) workshop (<https://commodore2018.sciencesconf.org/>) has been organized in Paris in September 2018. In total, the participants represented 15 oceanic dynamical cores among the

*Corresponding author address: Florian Lemarié, Univ. Grenoble Alpes, Inria, CNRS, Grenoble INP, LJK, 38000 Grenoble, France.
E-mail: florian.lemarie@inria.fr

most widely used by the research and operational community. The motivations, topics of discussion sessions, and outcomes of the workshop are summarized below.

Context

The ocean model developers community has had the tendency to be split depending on the target applications (global vs coastal) and on the type of horizontal grids (structured vs unstructured) and has been organized around relatively small modeling groups. However, the models have now reached such a high level of complexity that model development goes beyond the expertise of one given group and requires interactions between physicists, mathematicians, and computer scientists. In this context, this workshop aimed at gathering a community of model-oriented researchers to foster more regular exchanges and share expertise on outstanding issues and perspectives. During this first workshop, the emphasis was on reviewing the characteristics and diversity in the formulation of oceanic models used for realistic applications as well as on outlining upcoming challenges.

Evolution of oceanic models

Historically, global and regional ocean models have been based on the hydrostatic primitive equations (e.g. Griffies and Adcroft 2008) discretized on structured grids using a mixture of finite-difference and finite-volume techniques for the discretization in space. The time dimension is usually treated using standard predictor-corrector or two-level approaches (e.g. Lemarié et al. 2015). Those choices have been made because of their good compromise between simplicity, efficiency, and accuracy. In recent years, significant progress has been made for ocean modeling on unstructured grids, either via finite volume (e.g. Chen et al. 2003; Ringler et al. 2010; Danilov et al. 2017), or finite element (e.g. Zhang et al. 2016; Korn 2017; Kärnä et al. 2018) approach. Unstructured grid models have reached an unprecedented level of maturity at least for two reasons. First, the vertical dimension is treated in a structured way compared to earlier initiatives trying to get three-dimensional unstructured meshes working. Second, a better understanding of computational modes and dispersion properties associated to a wide range of possible choice of finite element pairs has been reached (e.g. Le Roux et al. 2007; Le Roux 2012; Eldred and Roux 2018). For example, Korn and Danilov (2017) have recently proposed a specific mimetic approach to control the well-known spurious mode occurring in triangular C-grids (Wolfram and Fringer 2013). Unstructured grid models have been used for coastal applications for many years and they have now reached the application phase for global applications (e.g. Sidorenko et al. 2015; Petersen et al. 2018). A long-standing concern is that computational cost per nominal grid point is generally much

larger than for structured-grid models and this problem is further compounded by the absence of time refinement to locally adjust the time-step to the mesh resolution when explicit time-stepping is used. The test strategy presented in the next section should provide a way to quantify more rationally the difference in terms of computational costs among existing models.

Recent advances also include the development of hybrid (or generalized) vertical coordinate systems based on Arbitrary Lagrangian Eulerian (ALE) methods. The vertical distribution of Eulerian coordinate levels is predefined, whereas Lagrangian coordinate levels freely evolve with the flow. ALE methods combine the advantages of well-defined (i.e. undistorted) meshes and reduced numerical mixing, and also allow adaptation strategies (e.g. Bleck 2002; Burchard and Beckers 2004; White et al. 2009; Leclair and Madec 2011; Petersen et al. 2015). A difficulty in this case is the rezoning (a.k.a. regridding) phase to maintain the integrity of the grid locally and globally.

A tendency in the design of the oceanic dynamical cores is the extension to the non-hydrostatic (NH) equations. The most widely used approach nowadays in oceanic models is based on the incompressible NH system solved using a pressure correction/projection method which requires the solution of a 3D Poisson equation (Lai et al. 2011; Vitousek and Fringer 2013; Voltzinger and Androsov 2016). Recently, Auclair et al. (2018) have proposed the use of the compressible non-hydrostatic ocean equations with the advantage that no global algebraic system needs to be solved to compute NH pressure anomalies but with the disadvantage to permit acoustic modes. In this case a specific numerical procedure is required to maintain acceptable stability of the whole code.

Test strategy and benchmark suite

Given the wide variety of choices that need to be made during the development of dynamical cores and their overall complexity, it is crucial to define evaluation methods to compare the behavior of different models. Such effort has been made over the last decade by the global atmospheric community (Dynamical Core Model Intercomparison Project (DCMIP), Ullrich et al. 2012). In particular, within DCMIP, a collection of test-cases that found broad acceptance in the community have been designed and applied by a large number of modeling groups. The workshop highlighted that in the context of the oceanic community, existing test-cases are scattered in the literature and not always fully documented and reproducible. The few existing examples of such effort (e.g. Ezer et al. 2002; Gerritsen et al. 2008; Ilicak et al. 2012; Soufflet et al. 2016) turned out to provide a valuable feedback on the consequences of model formulations. A good test-case should be easy to configure with analytical data suitable for all horizontal grids and different vertical coordinates and easy

to evaluate while being relevant to test a given component of the dynamical core. The evaluation can be done either via analytical solutions (e.g. Bristeau et al. 2018), numerically converged solutions (provided that all models converge toward the same solution) or more subjectively based on an unambiguous physical understanding of the processes (e.g. Marques et al. 2017). Such benchmark suite is also useful to motivate communication between modeling groups and also to open room for prospective approaches from applied mathematicians to highlight their effectiveness. Existing initiatives aiming at oceanic model intercomparison based on realistic simulations (e.g. Chassignet et al. 2000; Griffies et al. 2009) are generally too complex to clearly associate the observed differences to particular numerical choices. A way to evaluate numerical models in such complex configurations which could gain ground in the next few years is the uncertainty quantification (e.g. Iskandarani et al. 2016). Such approach provides tools to characterize modeling and numerical sensitivities.

Challenges

Throughout the 3 days of discussion, different current and future challenges have been identified. Addressing these challenges requires closer collaboration between the modeling groups.

Multi-resolution strategy: block-structured vs unstructured. On one hand, unstructured grid models have the ability to allow variable-resolution meshes provided an efficient mesh generation tool (e.g. Engwirda 2017), for example to adapt the resolution to follow the local Rossby radius (Hallberg 2013; Sein et al. 2017). On the other hand structured grid models can also locally increase the resolution via nesting techniques (Debreu and Blayo 2008; Warner et al. 2010; Debreu et al. 2012) or quadtree-octree refinement (Popinet and Rickard 2007). One advantage of the nesting approach is to allow the adjustment of the time-step and the physical parameters to the local resolution while unstructured models will need scale-aware parameterizations and a specific procedure for time refinement. Other points to investigate are the impact of variable resolution on propagating waves and the optimal layout to build a multiresolution mesh. A more prospective approach could be the use of an adaptive wavelet method (Kevlahan et al. 2015).

Energy consistency and resolved/unresolved scales coupling. Energy consistency is an important aspect for the proper interaction between resolved and parameterized scales (e.g. Burchard 2002; Bachman et al. 2017). However as soon as a numerical core does not globally conserve energy at a discrete level (e.g. because of monotonicity enforcement, vertical remapping, or some form of upwinding), the identification of energy pathways is difficult and requires an in-depth analysis to close the energy cycle (e.g. Marsaleix et al. 2008; Eden 2016) which

can be rather tedious (if not impossible) when advanced high-order numerics is used. An alternative could be to opt for an energy-conserving space and time discretization (e.g. Korn 2017; Eldred et al. 2018) with specific care (based on explicit numerical dissipation in the dynamical core and/or on parameterization of subgrid processes) to avoid instability issues of the existing approaches (Bell et al. 2017). This consistent coupling between dynamical cores and subgrid processes (known as physics-dynamics coupling) is an increasingly important topic for the building of geophysical models in general (Gross et al. 2018).

Vertical coordinates & spurious numerical mixing. Spurious mixing (especially spurious dianeutral mixing) is a long-standing issue for oceanic dynamical cores (e.g. Griffies et al. 2000). The use of an ALE vertical coordinate is a way to mitigate this issue. Despite the fact that Gibson et al. (2017) suggest that the vertical component of spurious mixing dominates as horizontal resolution increases it should not overshadow that many components of dynamical cores can be a source of numerical mixing (e.g. horizontal advection, time-stepping, the stabilization of the mode-splitting procedure). There is still a need to better understand the implications of different choice of momentum / tracers advection schemes, rezoning and remapping procedures on numerical mixing. Idealized test-cases and efficient diagnostic tools are important to tackle this issue (e.g. Ilicak et al. 2012; Klingbeil et al. 2014). It would also be instructive to keep investigating the improvement of quasi-Eulerian vertical coordinates (e.g. Berntsen 2011). In this context schemes for internal pressure gradient and for isoneutral tracer diffusion should also be considered (e.g. Shao et al. 2018).

Non-hydrostatic pressure contribution. As discussed earlier, many oceanic dynamical cores have the possibility to account for non-hydrostatic effects. A difficulty is now to clearly identify under which conditions relaxing the hydrostatic assumption is necessary and which resolution is required for proper NH modeling. Another challenge is the possibility to locally account for NH effects within a primitive equations model either in the form of a superparameterization (e.g. Campin et al. 2011) or in the form of 2-way nesting between coarse hydrostatic and fine non-hydrostatic meshes (e.g. Blayo and Rousseau 2016). Ultimately, further investigations of the merits and flaws of the incompressible vs compressible NH approaches would be worthwhile, also for global applications (Losch et al. 2004).

Coupling with other Earth-System compartments. Oceanic dynamical cores are often used as a component of larger coupled model systems. Coupling to several other Earth system compartments is common, such as to surface wave, sea-ice, atmospheric, biogeochemical, benthic, and hydrological models. The numerical implementation of such coupling can become an issue (e.g. Lemarié et al.

2015; Beljaars et al. 2017) especially at high coupling frequency and/or spatial resolution. More systematic analysis of the coupling stability and consistency using simplified equation sets and the design of simplified coupled test-cases must be encouraged.

Vanishing layers, wetting and drying and shock-resolving numerics. An accurate treatment of wetting and drying is essential for coastal simulations as well as for climate simulations of under-ice-shelf cavities. At a numerical level this requires the non-negativity of the water height and an adequate volume conserving treatment of dry states (a.k.a. vacuum states). Standard numerical methods used in oceanic dynamical cores do not have the ability to handle vacuum, or equivalently shocks. Instead, approaches based on some pre-defined minimum water depth and specific ad-hoc manipulation of discrete fluxes are often used (e.g. Sec. 5.2 in Klingbeil et al. 2018). However there is a vast literature dedicated to the design of numerical schemes preserving positivity and able to correctly treat vacuum states which furthermore satisfy an entropy-preserving property; i.e. the nonlinear solution is physically relevant even in the presence of discontinuities (e.g. Audusse et al. 2004, 2016). Considering advection-diffusion equations e.g. for tracers (temperature, salinity) there is an obvious benefits in using numerical schemes preserving the maximum principle. There could be an interest in comparing these more advanced approaches with the usual treatment adopted in dynamical cores.

Conclusions

The workshop gave a broad and fresh overview of existing numerical methods used in realistic ocean models as well as some examples of alternatives from the applied maths community. The participants have been enthusiastic and very positive about the possibility to sustain this type of workshop into a biennial workshop series. A collective article is currently in preparation to summarize the challenges and prospects for oceanic numerical cores across all scales. Moreover, a particular effort will be directed toward the formation of an active community of model developers with international collaborations, starting e.g. with the standardization of existing idealized test cases as the basis for model/methods intercomparison studies. The next meeting will be organized either in Fall 2019 or Winter 2020 in Hamburg, Germany.

Acknowledgments. The authors thank Gurvan Madec for reading and checking the content of this summary as well as all the workshop's participants¹. The workshop was hosted by the French National Institute for computer science and applied mathematics (Inria), Paris, France. We gratefully acknowledge the support from Inria, from

the Coastal and Regional Ocean Community Model research group (Gdr Croco), and from the french program Earths Fluid Envelopes (LEFE).

References

- Auclair, F., L. Bordoiso, Y. Dossmann, T. Duhaut, A. Paci, C. Ulses, and C. Nguyen, 2018: A non-hydrostatic non-boussinesq algorithm for free-surface ocean modelling. *Ocean Modell.*, doi:<https://doi.org/10.1016/j.ocemod.2018.07.011>.
- Audusse, E., F. Bouchut, M.-O. Bristeau, R. Klein, and B. Perthame, 2004: A fast and stable well-balanced scheme with hydrostatic reconstruction for Shallow Water flows. *SIAM J. Sci. Comput.*, **25** (6), 2050–2065, doi:[10.1137/S1064827503431090](https://doi.org/10.1137/S1064827503431090).
- Audusse, E., F. Bouchut, M.-O. Bristeau, and J. Sainte-Marie, 2016: Kinetic entropy inequality and hydrostatic reconstruction scheme for the Saint-Venant system. *Math. Comp.*, **85** (302), 2815–2837, doi:[10.1090/mcom/3099](https://doi.org/10.1090/mcom/3099), URL <http://dx.doi.org/10.1090/mcom/3099>.
- Bachman, S. D., B. Fox-Kemper, and B. Pearson, 2017: A scale-aware subgrid model for quasi-geostrophic turbulence. *J. Geophys. Res.*, **122** (2), 1529–1554, doi:[10.1002/2016JC012265](https://doi.org/10.1002/2016JC012265).
- Beljaars, A., E. Dutra, G. Balsamo, and F. Lemarié, 2017: On the numerical stability of surface-atmosphere coupling in weather and climate models. *Geosci. Model Dev.*, **10** (2), 977–989, doi:[10.5194/gmd-10-977-2017](https://doi.org/10.5194/gmd-10-977-2017).
- Bell, M. J., P. S. Peixoto, and J. Thuburn, 2017: Numerical instabilities of vector-invariant momentum equations on rectangular c-grids. *Quart. J. Roy. Meteorol. Soc.*, **143** (702), 563–581, doi:[10.1002/qj.2950](https://doi.org/10.1002/qj.2950).
- Berntsen, J., 2011: A perfectly balanced method for estimating the internal pressure gradients in σ -coordinate ocean models. *Ocean Modell.*, **38** (1), 85 – 95, doi:<https://doi.org/10.1016/j.ocemod.2011.02.006>.
- Blayo, E., and A. Rousseau, 2016: About interface conditions for coupling hydrostatic and nonhydrostatic navier-stokes flows. *Discrete Contin. Dyn. Syst. Ser. S*, **9** (5), 1565–1574, doi:[10.3934/dcdss.2016063](https://doi.org/10.3934/dcdss.2016063), URL <https://hal.inria.fr/hal-01185255>.
- Bleck, R., 2002: An oceanic general circulation model framed in hybrid isopycnic-Cartesian coordinates. *Ocean Modell.*, **4**, 55–88.
- Bristeau, M.-O., B. DI Martino, A. Mangeney, J. Sainte-Marie, and F. Soullé, 2018: Various analytical solutions for the incompressible Euler and Navier-Stokes systems with free surface, URL <https://hal.archives-ouvertes.fr/hal-01831622>, working paper or preprint.
- Burchard, H., 2002: Energy-conserving discretisation of turbulent shear and buoyancy production. *Ocean Modell.*, **4** (3), 347 – 361, doi:[https://doi.org/10.1016/S1463-5003\(02\)00009-4](https://doi.org/10.1016/S1463-5003(02)00009-4).
- Burchard, H., and J.-M. Beckers, 2004: Non-uniform adaptive vertical grids in one-dimensional numerical ocean models. *Ocean Modell.*, **6**, 51–81.
- Campin, J.-M., C. Hill, H. Jones, and J. Marshall, 2011: Superparameterization in ocean modeling: Application to deep convection. *Ocean Modell.*, **36** (1), 90 – 101, doi:<https://doi.org/10.1016/j.ocemod.2010.10.003>.
- Chassignet, E. P., and Coauthors, 2000: DAMÉE-NAB: the base experiments. *Dynam. Atmos. Oceans*, **32** (3-4), 155–183, doi:[10.1016/S0377-0265\(00\)00046-4](https://doi.org/10.1016/S0377-0265(00)00046-4).

¹<https://commodore2018.sciencesconf.org/resource/listeparticipants>

- Chen, C., H. Liu, and R. C. Beardsley, 2003: An unstructured grid, finite-volume, three-dimensional, primitive equations ocean model: Application to coastal ocean and estuaries. *J. Atmos. Oceanic Technol.*, **20** (1), 159–186, doi:10.1175/1520-0426(2003)020<0159:AUGFVT>2.0.CO;2.
- Danilov, S., D. Sidorenko, Q. Wang, and T. Jung, 2017: The finite-volume sea ice–ocean model (fesom2). *Geosci. Model Dev.*, **10** (2), 765–789, doi:10.5194/gmd-10-765-2017.
- Debreu, L., and E. Blayo, 2008: Two-way embedding algorithms: a review. *Ocean Dyn.*, **58** (5-6), 415–428, doi:10.1007/s10236-008-0150-9.
- Debreu, L., P. Marchesiello, P. Penven, and G. Cambon, 2012: Two-way nesting in split-explicit ocean models: algorithms, implementation and validation. *Ocean Modell.*, **49-50**, 1–21.
- Eden, C., 2016: Closing the energy cycle in an ocean model. *Ocean Modell.*, **101**, 30–42, doi:https://doi.org/10.1016/j.ocemod.2016.02.005.
- Eldred, C., T. Dubos, and E. Kritsikis, 2018: A Quasi-Hamiltonian Discretization of the Thermal Shallow Water Equations, URL <https://hal.inria.fr/hal-01847698>, submitted.
- Eldred, C., and D. Y. L. Roux, 2018: Dispersion analysis of compatible galerkin schemes for the 1d shallow water model. *J. Comp. Phys.*, **371**, 779–800, doi:https://doi.org/10.1016/j.jcp.2018.06.007.
- Engwirda, D., 2017: Jigsaw-geo (1.0): locally orthogonal staggered unstructured grid generation for general circulation modelling on the sphere. *Geosci. Model Dev.*, **10** (6), 2117–2140, doi:10.5194/gmd-10-2117-2017.
- Ezer, T., H. Arango, and A. F. Shchepetkin, 2002: Developments in terrain-following ocean models: intercomparisons of numerical aspects. *Ocean Modell.*, **4** (3), 249–267, doi:https://doi.org/10.1016/S1463-5003(02)00003-3.
- Gerritsen, H., E. D. Goede, F. Platzek, J. V. Kester, R. Uittenbogaard, and M. Genseberger, 2008: Validation document Delft3D-FLOW. Tech. Rep. X0356/M3470, WL—Delft Hydraulics. URL https://oss.deltares.nl/c/document_library/get_file?uuid=39169f8f-4ab0-4f7b-9771-c3f7d0ddd61f&groupId=183920.
- Gibson, A. H., A. M. Hogg, A. E. Kiss, C. J. Shakespeare, and A. Adcroft, 2017: Attribution of horizontal and vertical contributions to spurious mixing in an arbitrary lagrangianeulerian ocean model. *Ocean Modell.*, **119**, 45–56, doi:https://doi.org/10.1016/j.ocemod.2017.09.008.
- Griffies, S., and A. Adcroft, 2008: Formulating the Equations of ocean models. *Ocean Modeling in an Eddying Regime*, M. Hecht, and H. Hasumi, Eds., Geophysical Monograph Series, Vol. 177, American Geophysical Union, USA, 350.
- Griffies, S., and Coauthors, 2009: Coordinated ocean-ice reference experiments (cores). *Ocean Modell.*, **26** (1-2), 1–46, doi:10.1016/j.ocemod.2008.08.007.
- Griffies, S. M., R. C. Pacanowski, and R. W. Hallberg, 2000: Spurious diapycnal mixing associated with advection in a z-coordinate ocean model. *Mon. Weather Rev.*, **128** (3), 538–564, doi:10.1175/1520-0493(2000)128<0538:SDMAWA>2.0.CO;2.
- Gross, M., and Coauthors, 2018: Physics-dynamics coupling in weather, climate and earth system models: Challenges and recent progress. *Mon. Weather Rev.*, **0** (0), null, doi:10.1175/MWR-D-17-0345.1.
- Hallberg, R., 2013: Using a resolution function to regulate parameterizations of oceanic mesoscale eddy effects. *Ocean Modell.*, **72**, 92–103, doi:https://doi.org/10.1016/j.ocemod.2013.08.007.
- Ilicak, M., A. J. Adcroft, S. M. Griffies, and R. W. Hallberg, 2012: Spurious dianeutral mixing and the role of momentum closure. *Ocean Modell.*, **45-46**, 37–58.
- Iskandarani, M., S. Wang, A. Srinivasan, W. C. Thacker, J. Winokur, and O. M. Knio, 2016: An overview of uncertainty quantification techniques with application to oceanic and oil-spill simulations. *J. Geophys. Res.*, **121** (4), 2789–2808, doi:10.1002/2015JC011366.
- Kärnä, T., S. C. Kramer, L. Mitchell, D. A. Ham, M. D. Piggott, and A. M. Baptista, 2018: Thetis coastal ocean model: discontinuous galerkin discretization for the three-dimensional hydrostatic equations. *Geosci. Model Dev.*, **2018**, 1–36, doi:10.5194/gmd-2017-292.
- Kevalhan, N. K.-R., T. Dubos, and M. Aechtner, 2015: Adaptive wavelet simulation of global ocean dynamics using a new brinkman volume penalization. *Geosci. Model Dev.*, **8** (12), 3891–3909.
- Klingbeil, K., F. Lemarié, L. Debreu, and H. Burchard, 2018: The numerics of hydrostatic structured-grid coastal ocean models: State of the art and future perspectives. *Ocean Modell.*, **125**, 80–105, doi:https://doi.org/10.1016/j.ocemod.2018.01.007.
- Klingbeil, K., M. Mohammadi-Aragh, U. Gräwe, and H. Burchard, 2014: Quantification of spurious dissipation and mixing discrete variance decay in a finite-volume framework. *Ocean Modell.*, **81**, 49–64, doi:https://doi.org/10.1016/j.ocemod.2014.06.001.
- Korn, P., 2017: Formulation of an unstructured grid model for global ocean dynamics. *J. Comp. Phys.*, **339**, 525–552, doi:https://doi.org/10.1016/j.jcp.2017.03.009.
- Korn, P., and S. Danilov, 2017: Elementary dispersion analysis of some mimetic discretizations on triangular c-grids. *J. Comp. Phys.*, **330**, 156–172, doi:https://doi.org/10.1016/j.jcp.2016.10.059.
- Lai, Z., C. Chen, G. W. Cowles, and R. C. Beardsley, 2011: A nonhydrostatic version of fvcom: 1. validation experiments. *J. Geophys. Res.*, **115** (C11), doi:10.1029/2009JC005525.
- Le Roux, D., 2012: Spurious inertial oscillations in shallow–water models. *J. Comp. Phys.*, **231** (24), 7959–7987, doi:https://doi.org/10.1016/j.jcp.2012.04.052.
- Le Roux, D., V. Rostand, and B. Pouliot, 2007: Analysis of numerically induced oscillations in 2d finite–element shallow–water models part i: Inertia–gravity waves. *SIAM J. Sci. Comput.*, **29** (1), 331–360, doi:10.1137/060650106.
- Leclair, M., and G. Madec, 2011: \tilde{z} -coordinate, an arbitrary lagrangian-eulerian coordinate separating high and low frequency motions. *Ocean Modell.*, **37** (3-4), 139–152, doi:DOI:10.1016/j.ocemod.2011.02.001.
- Lemarié, F., E. Blayo, and L. Debreu, 2015: Analysis of ocean-atmosphere coupling algorithms: Consistency and stability. *Procedia Computer Science*, **51** (Supplement C), 2066–2075, doi:https://doi.org/10.1016/j.procs.2015.05.473, international Conference On Computational Science, ICCS 2015.
- Lemarié, F., L. Debreu, G. Madec, J. Demange, J. Molines, and M. Honorat, 2015: Stability constraints for oceanic numerical models: implications for the formulation of time and space discretizations. *Ocean Modell.*, **92**, 124–148, doi:https://doi.org/10.1016/j.ocemod.2015.06.006.

- Losch, M., A. Adcroft, and J.-M. Campin, 2004: How sensitive are coarse general circulation models to fundamental approximations in the equations of motion? *J. Phys. Oceanogr.*, **34** (1), 306–319, doi:10.1175/1520-0485(2004)034<0306:HSACGC>2.0.CO;2.
- Marques, G. M., M. G. Wells, L. Padman, and T. M. Özgökmen, 2017: Flow splitting in numerical simulations of oceanic dense-water outflows. *Ocean Modell.*, **113**, 66 – 84, doi:https://doi.org/10.1016/j.ocemod.2017.03.011.
- Marsaleix, P., F. Auclair, J. W. Floor, M. J. Herrmann, C. Estournel, I. Pairaud, and C. Ulses, 2008: Energy conservation issues in sigma-coordinate free-surface ocean models. *Ocean Modell.*, **20** (1), 61 – 89, doi:http://dx.doi.org/10.1016/j.ocemod.2007.07.005.
- Petersen, M., and Coauthors, 2018: An evaluation of the ocean and sea ice climate of E3SM using MPAS and interannual CORE-II forcing. URL <https://doi.org/10.5281/zenodo.1194911>, doi:10.5281/zenodo.1194911.
- Petersen, M. R., D. W. Jacobsen, T. D. Ringler, M. W. Hecht, and M. E. Maltrud, 2015: Evaluation of the arbitrary lagrangianeulerian vertical coordinate method in the mpas-ocean model. *Ocean Modell.*, **86**, 93 – 113, doi:https://doi.org/10.1016/j.ocemod.2014.12.004.
- Popinet, S., and G. Rickard, 2007: A tree-based solver for adaptive ocean modelling. *Ocean Modell.*, (16), 224–249.
- Ringler, T., J. Thuburn, J. Klemp, and W. Skamarock, 2010: A unified approach to energy conservation and potential vorticity dynamics for arbitrarily-structured c-grids. *J. Comp. Phys.*, **229** (9), 3065 – 3090, doi:https://doi.org/10.1016/j.jcp.2009.12.007.
- Sein, D. V., and Coauthors, 2017: Ocean modeling on a mesh with resolution following the local rossby radius. *J. Adv. Model. Earth Syst.*, **9** (7), 2601–2614, doi:10.1002/2017MS001099.
- Shao, A., A. Adcroft, R. Hallberg, and S. Griffies, 2018: Improvements to an Extrema-diminishing, Density-preserving Lateral Diffusion Algorithm. *AGU Ocean Sciences meeting, Portland, OR, USA, February 11-16*.
- Sidorenko, D., and Coauthors, 2015: Towards multi-resolution global climate modeling with echam6-fesom. part i: model formulation and mean climate. *Clim. Dynam.*, **44** (3), 757–780, doi:10.1007/s00382-014-2290-6.
- Soufflet, Y., P. Marchesiello, F. Lemarié, J. Jouanno, X. Capet, L. Debreu, and R. Benshila, 2016: On effective resolution in ocean models. *Ocean Modell.*, **98**, 36 – 50, doi:https://doi.org/10.1016/j.ocemod.2015.12.004.
- Ullrich, P. A., C. Jablonowski, J. Kent, P. H. Lauritzen, R. Nair, and M. A. Taylor, 2012: Dynamical core model intercomparison project (dcmip) test case document, URL http://www-personal.umich.edu/~cjablono/DCMIP-2012-TestCaseDocument_v1.7.pdf.
- Vitousek, S., and O. B. Fringer, 2013: Stability and consistency of non-hydrostatic free-surface models using the semi-implicit θ -method. *Int. J. Numer. Meth. Fluids*, **72** (5), 550–582, doi:10.1002/flid.3755.
- Voltzinger, N. E., and A. A. Androsov, 2016: Nonhydrostatic tidal dynamics in the area of a seamount. *Oceanology*, **56** (4), 491–500, doi:10.1134/S0001437016030243.
- Warner, J. C., W. R. Geyer, and H. G. Arango, 2010: Using a composite grid approach in a complex coastal domain to estimate estuarine residence time. *Comput. Geosci.*, **36** (7), 921–935, doi:10.1016/j.cageo.2009.11.008.
- White, L., A. Adcroft, and R. Hallberg, 2009: High-order regridding-remapping schemes for continuous isopycnal and generalized coordinates in ocean models. *J. Comp. Phys.*, **228** (23), 8665 – 8692, doi:https://doi.org/10.1016/j.jcp.2009.08.016.
- Wolfram, P. J., and O. B. Fringer, 2013: Mitigating horizontal divergence checker-board oscillations on unstructured triangular c-grids for nonlinear hydrostatic and nonhydrostatic flows. *Ocean Modell.*, **69**, 64 – 78, doi:https://doi.org/10.1016/j.ocemod.2013.05.007.
- Zhang, Y. J., F. Ye, E. V. Stanev, and S. Grashorn, 2016: Seamless cross-scale modeling with schism. *Ocean Modell.*, **102**, 64 – 81, doi:https://doi.org/10.1016/j.ocemod.2016.05.002.