

Self-similar behaviour of a non-local diffusion equation with time delay

Arnaud Ducrot, Alexandre Genadot

► **To cite this version:**

Arnaud Ducrot, Alexandre Genadot. Self-similar behaviour of a non-local diffusion equation with time delay. 2018. <hal-01964356>

HAL Id: hal-01964356

<https://hal.inria.fr/hal-01964356>

Submitted on 21 Dec 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Self-similar behaviour of a non-local diffusion equation with time delay

Arnaud Ducrot^a and Alexandre Genadot^b

^aemail: arnaud.ducrot@univ-lehavre.fr

Normandie Univ, UNIHAVRE, LMAH, FR-CNRS-3335, ISCN, 76600 Le Havre, France

^bemail: alexandre.genadot@math.u-bordeaux.fr

Univ. Bordeaux, IMB, UMR 5251, F-33076 Bordeaux, France

CNRS, IMB, UMR 5251, F-33400 Talence, France

INRIA Bordeaux-Sud Ouest, Team CQFD, F-33400 Talence, France

December 17, 2018

Abstract

We study the asymptotic behaviour of solutions of a class of linear non-local measure-valued differential equations with time delay. Our main result states that the solutions asymptotically exhibit a parabolic like behaviour in the large times, that is precisely expressed in term of heat kernel. Our proof relies on the study of a – self-similar – rescaled family of solutions. We first identify the asymptotic behaviour of the solutions by deriving a convergence result in the sense of the Young measures. Then we strengthen this convergence by deriving suitable fractional Sobolev compactness estimates. As a by-product, our main result allows to obtain asymptotic results for a class of piecewise constant stochastic processes with memory.

Key words: Non-local diffusion; Time delay; Self-similar behaviour; Fractional Sobolev estimates; Central Limit Theorem.

1 Introduction

Mathematical models with non-local diffusion arise in various applicative fields including physics and biology. Here, by non-local diffusion we mean convolution equation of the form

$$\frac{\partial u(t, x)}{\partial t} = \int_{\mathbb{R}^N} J(x - y) [u(t, y) - u(t, x)] dy, \quad t > 0 \text{ and } x \in \mathbb{R}^N,$$

wherein J denotes a probability kernel on \mathbb{R}^N , for some given and fixed integer $N \geq 1$. As described by Fife in [12], if $u = u(t, x)$ denotes the density of a

population at time t and spatial location $x \in \mathbb{R}^N$, the first term in the right hand side of this equation describes the rate at which individuals jump to x from all other locations y weighted with the probability kernel $J(x-y)$, while the second term $u(t, x) = u(t, x) \int_{\mathbb{R}^N} J(dy)$, corresponds to the rate at which individuals are leaving the location x to move to some other places.

As mentioned above, such a motion equation arises in various applications. We refer to [3, 7, 8, 9, 12, 13, 16, 21] and the references therein for analysis of models coming from physics, mathematical biology and population dynamics. We would like to emphasize that in such models, the jumps of particles or individuals are supposed to be instantaneous.

We can extend the above non-local diffusion equation by taking into account the travel time to jump from y to x . Typically, the travel time depends on the distance to travel ($x-y$) so that the above non-local motion law can be extended as follows. For $t > 0$ and $x \in \mathbb{R}^N$, a typical non-local diffusion equation with time delay reads as

$$\frac{\partial u(t, x)}{\partial t} = \int_{\mathbb{R}^N} J(x-y) [u(t - \tau(x-y), y) - u(t, x)] dy, \quad (1)$$

wherein $\tau(x-y)$ denotes the time needed to jump from y to x . This work is concerned with the asymptotic analysis of such a non-local diffusion equation with time delay. To that aim, we will consider more general version of this problem such as the equation

$$\frac{\partial u(t, x)}{\partial t} = \int_{[-1, 0] \times \mathbb{R}^N} [u(t + \theta, x-y) - u(t, x)] Q(d\theta, dy), \quad t > 0, \quad (2)$$

supplemented with a suitable initial data $u(\theta, x)$ for $\theta \in [-1, 0]$ and $x \in \mathbb{R}^N$. In the above equation, $Q = Q(d\theta, dy)$ denotes a probability distribution on the infinite strip $S := [-1, 0] \times \mathbb{R}^N$. This distribution captures the information about both the jump process and the time needed to perform jumps. For instance, if we choose $Q(d\theta, dy) = \delta_{-\tau(y)}(d\theta) \otimes J(y)dy$, we recover the typical equation presented in (1) above. Below we will describe a first stochastic representation of such a problem.

Observe that when $Q(d\theta, dy) = \delta_0(d\theta) \otimes J(dy)$, Equation (2) becomes

$$\frac{\partial u(t, x)}{\partial t} = \int_{\mathbb{R}^N} [u(t, x-y) - u(t, x)] J(dy).$$

This equation corresponds to the usual random walk equation with instantaneous jumps and associated with the jump measure J . Under suitable assumptions on this jump measure J , namely the existence of second moments and the non degeneracy of the covariance matrix, the positive solutions of this equation satisfy a central limit theorem. We refer to Ignat-Rossi [15] for a detailed study of this equation using Fourier analysis. We also refer to Chasseigne et al in [6] for a refined study in the case where the Fourier transform of the kernel J has lower regularity close to zero and the connexion with the self-similar solutions

of the heat equation with fractional Laplace operator. More general stochastic representations of the solution to equation (2) are considered in Section 2. The corresponding stochastic processes are piecewise constant processes with memory. The main result of the present paper allows to study the asymptotic behaviour of such stochastic processes.

Indeed, the aim of this work is to study the large time behaviour for the delayed equation (2). For the sake of generality and also for future applicative uses, we shall consider a slightly more general version of such an equation. In order to present the equation we will consider throughout this work, we denote by $\mathcal{P}(\mathbb{R}^N)$ the set of Borel probability measure on \mathbb{R}^N . In the sequel, we will simply write \mathcal{P} instead of $\mathcal{P}(\mathbb{R}^N)$ when there is no possible confusion. This space is endowed with the usual metrizable narrow topology (see Appendix A). Let $\alpha : [0, \infty) \times [-1, 0] \rightarrow (0, \infty)$ be a given bounded and continuous function. We consider the following problem:

$$\begin{cases} \frac{\partial}{\partial t} u(t, dx) = \int_S \alpha(t, \theta) [u(t + \theta, dx - y) - u(t, dx)] Q(d\theta, dy), & t > 0, \\ u(\theta, dx) = u^0(\theta, dx), & \forall \theta \in [-1, 0] \text{ and } u^0 \in \mathcal{C}([-1, 0]; \mathcal{P}(\mathbb{R}^N)). \end{cases} \quad (3)$$

Similarly to (2), in the above problem Q belongs to $\mathcal{P}(S)$, the set of Borel probability measures on the strip $S = [-1, 0] \times \mathbb{R}^N$.

Remark that the above equation is posed for probability measures allowing us to handle lattice equations with time delay. Indeed, consider for instance the case where

$$Q(d\theta, dy) = \sum_{j \in \mathbb{Z}} \alpha_j \delta_{-\tau_j}(d\theta) \otimes \delta_j(dy) \text{ with } -\tau_j \in [-1, 0], \alpha_j \geq 0 \text{ and } \sum_{j \in \mathbb{Z}} \alpha_j = 1,$$

then, when equipped with initial data of the form

$$u(\theta, dx) = \sum_{i \in \mathbb{Z}} u_i^0(\theta) \delta_i(dx) \in \mathcal{C}([-1, 0]; \mathcal{P}(\mathbb{R}^N)),$$

the solutions of the above equation takes the form $u(t, dx) = \sum_{i \in \mathbb{Z}} u_i(t) \delta_i(dx)$ where the functions $(u_i(t))_{i \in \mathbb{Z}}$ satisfy the delayed lattice system of equations

$$\begin{cases} u_i'(t) = \sum_{j \in \mathbb{Z}} \alpha_j u_{i-j}(t - \tau_j) - u_i(t), & t > 0, \\ u_i(\theta) = u_i^0(\theta), & \theta \in [-1, 0] \end{cases}, \quad \forall i \in \mathbb{Z}.$$

We now come back to Problem (3). Before stating our main result, we describe the main set of assumptions that will be used to study the asymptotic behaviour of (3).

Assumption 1 *We assume that:*

- (i) *The function $\alpha = \alpha(t, \theta)$ defined from $[0, \infty) \times [-1, 0]$ into $(0, \infty)$ is bounded and continuous and there exist $M > 0$ and $\beta > 0$ such that*

$$|\alpha(t, \theta) - \alpha_\infty(\theta)| \leq M e^{-\beta t}, \quad \forall t \geq 0, \theta \in [-1, 0],$$

wherein $\alpha_\infty = \alpha_\infty(\theta) > 0$ is a continuous function on $[-1, 0]$.

(ii) The function $(\theta, x) \mapsto |x|$ belongs to $L^2(S; Q(d\theta, dx))$. Here we use the symbol $|\cdot|$ to denote the Euclidean norm in \mathbb{R}^N .

Throughout this work, we fix $u^0 \in \mathcal{C}([-1, 0]; \mathcal{P}(\mathbb{R}^N))$ and we consider the solution $u \equiv u(t, dx)$ of (3) associated with the initial data u^0 . Here, by a solution we mean an element $\mu = \mu(t, dx) \in \mathcal{C}([-1, \infty); \mathcal{P}(\mathbb{R}^N))$ that satisfies:

- (i) for all $\theta \in [-1, 0]$, one has $\mu(\theta, dx) = u^0(\theta, dx)$;
- (ii) for any $f \in \mathcal{C}_b(\mathbb{R}^N)$, the space of bounded and continuous functions on \mathbb{R}^N , one has:
 - (iia) the map $t \mapsto \int_{\mathbb{R}^N} f(x)\mu(t, dx)$ is continuously differentiable for $t \in (0, \infty)$;
 - (iib) for all $t > 0$ one has

$$\begin{aligned} \frac{d}{dt} \int_{\mathbb{R}^N} f(x)\mu(t, dx) &= \int_S \alpha(t, \theta) \left[\int_{\mathbb{R}^N} f(x+z)\mu(t+\theta, dx) \right] Q(d\theta, dz) \\ &\quad - \int_{\mathbb{R}^N} f(x)\mu(t, dx) \int_S \alpha(t, \theta) Q(d\theta, dz). \end{aligned}$$

Note that the existence and uniqueness of such a solution simply follows from a usual contraction fixed point argument in the metric space $\mathcal{C}([-1, T]; \mathcal{P}(\mathbb{R}^N))$ for $T > 0$.

Our main result reads as follows.

Theorem 2 (Self-similar behaviour) *Let Assumption 1 be satisfied. Then there exists a vector $\mathbf{K} \in \mathbb{R}^N$ and a probability measure $\pi \in \mathcal{P}(\mathbb{R}^N)$ with zero mean value such that the function $t \mapsto \mu(t, dx) \in \mathcal{C}([-1, \infty); \mathcal{P}(\mathbb{R}^N))$ satisfies the following asymptotic behaviour: for all test functions $f \in \mathcal{C}_b(\mathbb{R}^N)$ one has*

$$\lim_{t \rightarrow \infty} \int_{\mathbb{R}^N} f\left(\frac{x - \mathbf{K}t}{\sqrt{t}}\right) \mu(t, dx) = \int_{\mathbb{R}^N} f(x)\pi(dx).$$

The vector \mathbf{K} is given by

$$\mathbf{K} = \frac{1}{1 + \Gamma} \int_S \alpha_\infty(\theta) z Q(d\theta, dz) \text{ with } \Gamma := \int_S (-\theta) \alpha_\infty(\theta) Q(d\theta, dz), \quad (4)$$

while $\pi(dx) = \Pi(1, dx)$ where $\Pi(t, dx) \in \mathcal{C}(\mathbb{R}^+, \mathcal{P})$ denotes the unique (tempered distribution) solution of the heat equation

$$\partial_t \Pi = \frac{1}{2} \operatorname{div} \left(\frac{D_0}{1 + \Gamma} \nabla \Pi \right) + \delta_{t=0} \otimes \delta_0(dx). \quad (5)$$

Herein D_0 denotes the non-negative symmetric matrix defined by

$$D_0 = \int_S \alpha_\infty(\theta) [z + \theta \overline{\mathbf{K}}] [z + \theta \overline{\mathbf{K}}]^T Q(d\theta, dz). \quad (6)$$

If we denote by $k = \dim \ker D_0 \geq 0$ and $\lambda_1 > 0, \dots, \lambda_{N-k} > 0$ the nonzero eigenvalue of $\frac{1}{1+\Gamma}D_0$ then there exists an orthogonal matrix P such that

$$\frac{1}{1+\Gamma}D_0 = P^T \text{diag}(\lambda_1, \dots, \lambda_{N-k}, 0, \dots, 0) P.$$

Using this orthogonal basis one obtains an explicit formula for $\Pi(t, dx)$ and thus the following explicit formula for $\pi = \Pi_1$:

$$\pi(dy) = P_{\#}^T \left[\left[\bigotimes_{i=1}^{N-k} f_{\lambda_i}(y_i) dy_i \right] \otimes \left[\bigotimes_{i=N-k+1}^N \delta_0(dy_i) \right] \right].$$

Here, $\#$ denotes the usual push forward operator for Borel measures while for each $\lambda > 0$, the function $f_\lambda : \mathbb{R} \rightarrow \mathbb{R}$ denotes centred normal distribution with variance λ , that is

$$f_\lambda(y) = \frac{1}{\sqrt{2\pi\lambda}} \exp\left(-\frac{y^2}{2\lambda}\right).$$

In other words, the measure Π_1 is the multidimensional Gaussian law with variance-covariance matrix given by $\frac{1}{1+\Gamma}D_0$. We refer to [26] for a complete treatment of the relation between Gaussian processes and Fokker-Planck equations of the form (5).

The proof of Theorem 2 will be divided in two parts. Our proof relies on the study of a self-similar rescaled family of solutions. In Section 3, we first identify the asymptotic behaviour of the solutions by deriving a convergence result in the sense of the Young measures. Then, in Section 4, we strengthen this convergence by deriving suitable fractional Sobolev compactness estimates.

Before going to the proof of Theorem 2, we apply it to some example of an hyperbolic equation with time delay. More specifically consider the equation

$$\left[\frac{\partial}{\partial t} + q \frac{\partial}{\partial x} \right] u(t, x) = a \int_{[-1, 0]} u(t + \theta, x) \eta(d\theta) - bu(t, x), \quad t > 0, \quad x \in \mathbb{R}, \quad (7)$$

where $q \in \mathbb{R} \setminus \{0\}$, $a > 0$ and $b \in \mathbb{R}$ are fixed parameters while $\eta \in \mathcal{P}([-1, 0])$ is a given probability measure on the interval $[-1, 0]$. This equation is supplemented with the initial data

$$u(\theta, x) = u^0(\theta, x) \text{ with } u^0 \in \mathcal{C}([-1, 0]; L_+^1(\mathbb{R})) \text{ and } u^0(0, \cdot) \not\equiv 0.$$

A special case of the above equation (with $\eta(d\theta) = \delta_{-1}(d\theta)$ and parameter conditions) has been studied by Laurent et al in [17]. In this paper the authors derived conditions ensuring a parabolic behaviour for the solution of such a problem. Here we will show how our general result, namely Theorem 2, may apply to (7), with general time delay dependence, to obtain similar results as in [17].

Set $v(t, x) = u(t, x + qt)e^{at}$ that satisfies the equation

$$\frac{\partial v(t, x)}{\partial t} = a \int_{[-1, 0]} e^{-b\theta} v(t + \theta, x - q\theta) \eta(d\theta).$$

Set $y = y(t) = \int_{\mathbb{R}} v(t, x) dx$ and observe that it satisfies the linear delay differential equation

$$\begin{aligned} y'(t) &= a \int_{[-1,0]} e^{-b\theta} y(t+\theta) \eta(d\theta), \quad t > 0, \\ y(\theta) &= y^0(\theta) := \int_{\mathbb{R}} u^0(\theta, x) dx, \quad \forall \theta \in [-1, 0]. \end{aligned} \tag{8}$$

Since $u^0(0, \cdot) \not\equiv 0$ then $y^0 \in \mathcal{C}([-1, 0]; \mathbb{R}^+)$ with $y^0(0) \neq 0$ and $y(t) > 0$ for all $t > 0$. Hence, for $t > 0$, the function $w(t, x) = \frac{v(t, x)}{y(t)}$ satisfies

$$\begin{aligned} \frac{\partial w(t, x)}{\partial t} &= a \int_{[-1,0]} \frac{y(t+\theta)}{y(t)} e^{-b\theta} [w(t+\theta, x - q\theta) - w(t, x)] \\ &= \int_S \alpha(t, \theta) [w(t+\theta, x - y) - w(t, x)] Q(d\theta, dy), \end{aligned}$$

with

$$\alpha(t, \theta) = a e^{-b\theta} \frac{y(t+\theta)}{y(t)} \text{ and } Q(d\theta, dy) = \eta(d\theta) \otimes \delta_{q\theta}(dy) \in \mathcal{P}([-1, 0] \times \mathbb{R}).$$

As a consequence Problem (7) re-writes as (3). To conclude this section, it remains to check that Assumption 1 is satisfied for this specific example. To that aim, one may first notice that Assumption 1 (ii) is satisfied. And, Assumption 1 (i) also holds true with the limit function $a_\infty = a_\infty(\theta)$ defined by $\alpha_\infty(\theta) = a e^{-b\theta} e^{\gamma\theta}$, where $\gamma > 0$ is the unique real solution of the equation

$$\gamma = a \int_{[-1,0]} e^{(\gamma-b)\theta} \eta(d\theta).$$

This latter property will be checked in Appendix B. This example will be further developed in the next section.

2 A probabilistic representation for some non-local diffusion equations with delay

In this section, we construct a \mathbb{R}^N -valued càdlàg process whose law satisfies the equation (3). Let $(\Omega, \mathcal{A}, \mathbb{P})$ be a probability space on which is built a nonhomogeneous Poisson point process $(N(t))_{t \in \mathbb{R}_+}$ with intensity function given for $t \in \mathbb{R}_+$ by

$$\lambda(t) = \int_S \alpha(t, \theta) Q(d\theta, dz)$$

and corresponding intensity measure given by $\Lambda((0, t]) = \int_0^t \lambda(s) ds$. As usual, the sequence of times associated to the Poisson point process is defined by $T_0 = 0$ and, for $n \in \mathbb{N}$, by $T_n = \inf\{t \geq 0; N(t) \geq n\}$. Notice that Assumption

1 implies that the intensity function stays positive and converges towards $\lambda_\infty = \int_S \alpha_\infty(\theta)Q(d\theta, dz) > 0$ such that we do have that the T_n 's form a increasing sequence going to infinity, \mathbb{P} -a.s.

We also define, on the same probability space, a sequence $(\Theta_n, Z_n)_{n \in \mathbb{N}}$ of S -valued random variables such that conditional on $(T_n)_{n \in \mathbb{N}}$, the sequence $(\Theta_n, Z_n)_{n \in \mathbb{N}}$ is a sequence of independent random variables with law given, for $n \in \mathbb{N}$, by

$$q_{T_n}(d\theta, dz) = \frac{1}{\lambda(T_n)} \alpha(T_n, \theta)Q(d\theta, dz).$$

Let also $(U(\theta))_{\theta \in [-1, 0]}$ be a \mathbb{R}^N -valued càdlàg process such that for any $\theta \in [-1, 0]$, $U(\theta)$ has for law $u^0(\theta, dx)$. We are now ready to define a \mathbb{R}^N -valued process X with suitable law:

1. For $\theta \in [-1, 0]$, we set $X(\theta) = U(\theta)$ such that $X(T_0) = U(0)$;
2. For $t \in [T_0, T_1)$, $X(t) = X(T_0)$ and then, at time T_1 ,

$$X(T_1) = X(T_1^- + \Theta_1) + Z_1.$$

3. And so on: for $n \in \mathbb{N}$, assume that the process X is built up to the time T_n , then, for $t \in [T_n, T_{n+1})$, $X(t) = X(T_n)$ and $X(T_{n+1}) = X(T_{n+1}^- + \Theta_{n+1}) + Z_{n+1}$.

By construction, the process X is piecewise constant and càdlàg on \mathbb{R}_+ . In the following proposition, we verify that the law of X satisfies the evolution equation (3). In this setting, one obtains the following proposition.

Proposition 3 *Let $f \in C_b(\mathbb{R}^N)$ be given. The function $t \mapsto \mathbb{E}(f(X(t)))$ is continuously differentiable on $(0, \infty)$ and we have, for any $t > 0$,*

$$\frac{d}{dt} \mathbb{E}(f(X(t))) = \lambda(t) \int_S [\mathbb{E}(f(X(t+\theta) + z)) - \mathbb{E}(f(X(t)))] q_t(d\theta, dz). \quad (9)$$

Proof. We use the decomposition of the process induced by the sequence of times $(T_n)_{n \in \mathbb{N}}$. For any $t \geq 0$, we get

$$\mathbb{E}(f(X(t))) = \sum_{i=0}^{\infty} \mathbb{E}(1_{[T_i, T_{i+1})}(t) f(X(T_i))).$$

For any integer i , by conditioning on the history up to the time T_i , we have

$$\mathbb{E}(1_{[T_i, T_{i+1})}(t) f(X(T_i))) = \mathbb{E} \left(1_{T_i \leq t} e^{-\Lambda((T_i, t])} f(X(T_i)) \right).$$

Since $T_0 = 0$ is deterministic, we distinguishes this time and then use the law of T_n for $n \in \mathbb{N}$ to write:

$$\begin{aligned} \mathbb{E}(f(X(t))) &= e^{-\int_0^t \lambda(s) ds} \mathbb{E}(f(U(0))) + \mathbb{E} \left(1_{T_i \leq t} e^{-\Lambda((T_i, t])} f(X(T_i)) \right) \\ &= e^{-\Lambda((0, t])} \mathbb{E}(f(U(0))) \\ &\quad + \sum_{i=1}^{\infty} \int_0^t e^{-\Lambda((s, t])} \mathbb{E}(f(X(T_i)) | T_i = s) \lambda(s) \frac{\Lambda((0, s])^{i-1}}{(i-1)!} e^{-\Lambda((0, s])} ds. \end{aligned}$$

Using the fact that $X(T_i) = X(T_i^- + \Theta_i) + Z_i$, we obtain

$$\begin{aligned} \mathbb{E}(f(X(t))) &= e^{-\Lambda((0,t])} \mathbb{E}(f(U(0))) \\ &+ \sum_{i=1}^{\infty} \int_0^t e^{-\Lambda((s,t])} \int_S \mathbb{E}(f(X(s+\theta) + z)) q_s(d\theta, dz) \\ &\quad \lambda(s) \frac{\Lambda((0,s])^{i-1}}{(i-1)!} e^{-\Lambda((0,s])} ds. \end{aligned}$$

Finally, by summation (using the fact that f is bounded),

$$\begin{aligned} \mathbb{E}(f(X(t))) &= e^{-\Lambda((0,t])} \mathbb{E}(f(U(0))) \\ &+ \int_0^t e^{-\Lambda((s,t])} \lambda(s) \int_S \mathbb{E}(f(X(s+\theta) + z)) q_s(d\theta, dz) ds. \end{aligned}$$

The result follows by derivation. ■

Denoting by $\mu(t, dx)$ the law of $X(t)$, equations (9) and (3) are equivalent. Thus, the process X gives a probabilistic representation of the evolution equation studied in the present paper. Notice that in general, X is not a Markov process: the variables Θ_n 's introduce memory and model the delay in the evolution equation (3). When there is no delay, that is when $Q(d\theta, dz) = \delta_0(d\theta)Q(dz)$, the studied process is simply an inhomogeneous continuous time Markov chain, as already mentioned in the introduction. Let us also notice that this is always possible to recast a non-Markovian process into a Markovian one: for processes very similar to the process X under consideration, this has been done in [22]. However, the fact that the evolution of the law of X satisfies equation (3) allows for a more direct approach.

Theorem 2 is the central limit theorem associated to the process X . Its proof will be the object of Section 3. In probabilistic term, Theorem 2 reads as follows.

Theorem 4 *The process $(\sqrt{t}(X(t)/t - \mathbf{K}))_{t \in (0, \infty)}$ converges in law towards a centred Gaussian random variable with variance-covariance matrix given by*

$$\mathbf{Q} = \frac{\lambda_\infty}{1 - \lambda_\infty \mathbb{E}(\Theta_\infty)} \mathbb{E}((Z_\infty + \Theta_\infty \mathbf{K})(Z_\infty + \Theta_\infty \mathbf{K})^T),$$

where the couple $(\Theta_\infty, Z_\infty)$ has for law $\alpha_\infty(\theta)Q(d\theta, dz)/\lambda_\infty$.

Of course, this central limit theorem implies that the process $(X(t)/t)_{t \in (0, \infty)}$ converges in probability towards \mathbf{K} . We can strengthen this convergence using the explicit construction of the process and obtain the following strong law of large numbers.

Theorem 5 *Assume the result of Theorem 2. Then, the process $(X(t)/t)_{t \in (0, \infty)}$ converges almost-surely towards $\mathbf{K} = \frac{\lambda_\infty \mathbb{E}(Z_\infty)}{1 - \lambda_\infty \mathbb{E}(\Theta_\infty)}$ where the couple $(\Theta_\infty, Z_\infty)$ has for law $\alpha_\infty(\theta)Q(d\theta, dz)/\lambda_\infty$.*

Proof. First, notice that for any $t > 0$,

$$|X(t)| \leq \max_{\theta \in [-1,0]} |U(\theta)| + \sum_{i=1}^{N(t)} |Z_i|.$$

Therefore,

$$\frac{|X(t)|}{t} \leq \frac{1}{t} \max_{\theta \in [-1,0]} |U(\theta)| + \frac{N(t)}{t} \frac{1}{N(t)} \sum_{i=1}^{N(t)} |Z_i|.$$

We have, almost-surely,

$$\lim_{t \rightarrow \infty} \frac{N(t)}{t} = \lim_{t \rightarrow \infty} \frac{\Lambda((0,t])}{t} = \lambda_\infty.$$

Thanks to Assumption 1 and the conditional independence of the Z_i 's, we can use the conditional law of large number, see for instance [23] and obtain that we almost-surely have,

$$\lim_{t \rightarrow \infty} \frac{1}{N(t)} \sum_{i=1}^{N(t)} |Z_i| = \frac{1}{\lambda_\infty} \int_S |z| \alpha_\infty(\theta) Q(d\theta, dz).$$

The process $(X(t)/t)_{t \in (0, \infty)}$ is thus almost-surely asymptotically bounded. Therefore, its convergence in probability towards the deterministic value \mathbf{K} implies its almost-sure convergence towards this same value. ■

Remark that since the process $(X(t)/t)_{t \in (0, \infty)}$ is bounded for t large enough, we can use a slight refinement of equation (9) in order to include an explicit dependence in time and obtain, for $y(t) = \mathbb{E}(X(t)/t)$ with $t > 0$,

$$\begin{aligned} \frac{d}{dt} y(t) &= \lambda(t) \int_S [y(t+\theta) - y(t)] \left[1 + \frac{\theta}{t} \right] q_t(d\theta, dz) \\ &\quad + \frac{1}{t} \left[-y(t) \left(1 - \lambda(t) \int_S \theta q_t(d\theta, dz) \right) + \lambda(t) \int_S z q_t(d\theta, dz) \right]. \end{aligned}$$

Notice that since \mathbf{K} is the limit of $(X(t)/t, t > 0)$ then, by dominated convergence, \mathbf{K} is also the limit of $(y(t), t > 0)$, a fact that is, at first glance, far to be obvious if you only look at the above differential equation with delay.

Let us give a first illustration of the two above results by considering the case where $X(\theta) = 0$ for all $\theta \in [-1, 0]$ and

$$\alpha(t, \theta) = 1, \quad Q(d\theta, dz) = \delta_0(d\theta) \delta_1(dz).$$

Then X is the classical Poisson process satisfying

$$\lim_{t \rightarrow \infty} \frac{X(t)}{t} = 1 \text{ a.s.}, \quad \lim_{t \rightarrow \infty} \sqrt{t} \left(\frac{X(t)}{t} - 1 \right) = \mathcal{N}(0, 1) \text{ in law,}$$

where $\mathcal{N}(0, 1)$ denotes the centred gaussian law with variance 1. Now, with a one unit time delay, namely with

$$\alpha(t, \theta) = 1, \quad Q(d\theta, dz) = \delta_{-1}(d\theta) \delta_1(dz),$$

our results give

$$\lim_{t \rightarrow \infty} \frac{X(t)}{t} = \frac{1}{2} \text{ a.s.}, \quad \lim_{t \rightarrow \infty} \sqrt{t} \left(\frac{X(t)}{t} - \frac{1}{2} \right) = \mathcal{N} \left(0, \frac{1}{8} \right) \text{ in law.}$$

In this example, we see that the delay slow down the process and reduces its scattering.

As a second illustration, we consider the stochastic process associated to the hyperbolic equation with delay (7) considered at the end of the previous section. In this setting, for $t > 0$ and $\theta \in [-1, 0]$, the parameters are

$$\alpha(t, \theta) = ae^{-b\theta} \frac{y(t+\theta)}{y(t)} \text{ and } Q(d\theta, dy) = \eta(d\theta) \otimes \delta_{q\theta}(dy) \in \mathcal{P}([-1, 0] \times \mathbb{R}), \quad (10)$$

such that

$$\lambda(t) = \int_{[-1, 0] \times \mathbb{R}} \alpha(t, \theta) Q(d\theta, dz) = \int_{[-1, 0]} ae^{-b\theta} \frac{y(t+\theta)}{y(t)} \eta(d\theta).$$

For our simulation purpose, as in Laurent et al in [17], we set $\eta(d\theta) = \delta_{-1}(d\theta)$. In this case,

$$\lambda(t) = ae^b \frac{y(t-1)}{y(t)},$$

where

$$\begin{aligned} y'(t) &= ae^b y(t-1), \quad t > 0, \\ y(\theta) &= y^0(\theta) := \int_{\mathbb{R}} u^0(\theta, x) dx, \quad \forall \theta \in [-1, 0], \end{aligned} \quad (11)$$

such that the intensity measure is given by $\Lambda((0, t]) = \log(y(t)/y(0))$. In Figure 1, the law of large numbers as well as the associated central limit theorem are illustrated for this process. We also display, in Figure 2, another illustration of these two limit theorems with $Q(d\theta, dy) = \delta_{-1}(d\theta) \otimes \mathbf{1}_{[-1/2, 1/2]}(y) dy$, all other things being equal.

3 Self-similar rescaling and limit identification

In this section, we deal with rescaled solution and we shall prove that such a family of rescaled solution converges toward the solution of the possibly degenerate parabolic equation (5) for a suitable topology.

Before going further let us introduce some notations that will be used in the sequel. We define $\Gamma > 0$ and the vector $\mathbf{K} \in \mathbb{R}^N$ by

$$\Gamma := \int_S (-\theta) \alpha_\infty(\theta) Q(d\theta, dz) \text{ and } \mathbf{K} := \frac{1}{1+\Gamma} \int_S \alpha_\infty(\theta) z Q(d\theta, dz). \quad (12)$$




Figure 1: Left: 40 trajectories of X (one color per trajectory) with the parameters defined in (10) with $a = 1.01$, $b = 1$, $q = 1$ and initial condition $u^0(\theta, dx) = \mathbf{1}_{[0,1]}(x)dx$ (uniform law on $[0, 1]$). The dashed line has slope $\mathbf{K} \simeq -0.25$. Right: distribution of $Z(100) = \sqrt{100} \frac{X(100)/100 - K}{\sqrt{\mathbf{Q}}}$ ($\sqrt{\mathbf{Q}} \simeq 0.18$), obtained from 1000 trajectories, compared to the density of the normal law.




Figure 2: Left: 40 trajectories of X (one color per trajectory) with the parameters defined in (10) with $a = 1.01$, $b = 1$, $q = 1$ but $Q(d\theta, dy) = \delta_{-1}(d\theta) \otimes \mathbf{1}_{[-1/2, 1/2]}(y)dy$ and initial condition $u^0(\theta, dx) = \mathbf{1}_{[-1/2, 1/2]}(x)dx$ (uniform law on $[-1/2, 1/2]$). The dashed line has slope $\mathbf{K} = 0$. Right: distribution of $Z(100) = \sqrt{100} \frac{X(100)/100 - K}{\sqrt{\mathbf{Q}}}$ ($\sqrt{\mathbf{Q}} \simeq 0.20$), obtained from 1000 trajectories, compared to the density of the normal law.

Remark 6 Note that the vector \mathbf{K} satisfies the following identity, that will be used latter,

$$\int_S \alpha_\infty(\theta) [z + \theta \mathbf{K}] Q(d\theta, dz) = \mathbf{K}.$$

We also define the function $H = H(t) : [-1, \infty) \rightarrow \mathbb{R}^N$ by

$$H(t) = -\frac{1}{1+\Gamma} \int_0^t \int_S \alpha(t, \theta) x Q(d\theta, dx) ds, \quad \forall t \geq -1. \quad (13)$$

As mentioned above we aim at investigating the large behaviour of $t \mapsto \mu(t, dx)$. Thus we need to shift this function in order to follow the mass and to not lose its transportation. As it will appear clearly in the proofs presented below, a suitable shift is to follow the solution along the path $t \mapsto H(t)$ introduced above in (13). Hence we translate the measure solution $t \mapsto \mu(t, dx)$ by introducing the measure

$$\nu(t, \cdot) = \delta_{H(t)} * \mu(t, \cdot) \in \mathcal{P}, \quad \forall t \geq -1. \quad (14)$$

In order to be more precise, the above convolution product, $*$, means that for all $\varphi \in \mathcal{C}_b(\mathbb{R}^N)$:

$$\int_{\mathbb{R}^N} \varphi(x) \nu(t, dx) = \int_{\mathbb{R}^N} \varphi(x + H(t)) \mu(t, dx), \quad \forall t \geq -1.$$

Notice that with the notations of Section 2, $\nu(t, dx)$ is the law of the process $X + H$ at time t .

The following lemma holds true.

Lemma 7 *The map $t \mapsto \nu(t, dx)$ is continuous from $[-1, \infty)$ into \mathcal{P} . It furthermore satisfies, for each test function $\phi \in W^{1,1}(0, \infty; \mathcal{C}_b(\mathbb{R}^N)) \cap L^1(0, \infty; \mathcal{C}_b^1(\mathbb{R}^N))$, the equation*

$$\begin{aligned} & \int_{\mathbb{R}^+ \times \mathbb{R}^N} (\partial_t + H'(t) \nabla) \phi(t, x) \nu(t, dx) dt = - \int_{\mathbb{R}^N} \phi(0, x) \nu(0, dx) \\ & + \int_0^\infty \int_{\mathbb{R}^N} \left[\int_S \alpha(t, \theta) Q(d\theta, dz) \right] \phi(t, x) \nu(t, dx) dt \\ & - \int_0^\infty \int_S \alpha(t, \theta) \int_{\mathbb{R}^N} \phi(t, x + z + G(t, \theta)) \nu(t + \theta, dx) Q(d\theta, dz) dt, \end{aligned}$$

wherein we have set $G(t, \theta) = H(t) - H(t + \theta)$.

Proof. Let us first note that since the map $t \mapsto H(t)$ is continuous, it follows from Portemanteau theorem (see Theorem A.1 in Appendix A) that the map $t \mapsto \nu(t, dx)$ is also continuous.

Let $\phi = \phi(t, x) \in W^{1,1}(0, \infty; \mathcal{C}_b(\mathbb{R}^N)) \cap L^1(0, \infty; \mathcal{C}_b^1(\mathbb{R}^N))$ be a given test function. Then, one has:

$$\begin{aligned} & \int_{\mathbb{R}^+ \times \mathbb{R}^N} (\partial_t + H'(t)\nabla) \phi(t, x) \nu(t, dx) dt \\ &= \int_0^\infty \int_{\mathbb{R}^N} (\partial_t + H'(t)\nabla) \phi(t, x + H(t)) \mu(t, dx) dt \\ &= \int_0^\infty \int_{\mathbb{R}^N} \partial_t \psi(t, x) \mu(t, dx) dt, \end{aligned}$$

wherein we have set $\psi(t, x) := \phi(t, x + H(t))$. It is worth noticing that $\psi \in W^{1,1}(0, \infty; \mathcal{C}_b(\mathbb{R}^N)) \cap L^1(0, \infty; \mathcal{C}_b^1(\mathbb{R}^N))$. Then, integrating by parts yields

$$\begin{aligned} & \int_{\mathbb{R}^+ \times \mathbb{R}^N} (\partial_t + H'(t)\nabla) \phi(t, x) \nu(t, dx) dt \\ &= - \int_{\mathbb{R}^N} \phi(0, x) \nu(0, dx) + \int_0^\infty \int_{\mathbb{R}^N} \left[\int_S \alpha(t, \theta) Q(\theta, dz) \right] \phi(t, x) \nu(t, dx) dt \\ &\quad - \int_0^\infty \int_S \alpha(t, \theta) \int_{\mathbb{R}^N} \phi(t, x + z + H(t)) \mu(t + \theta, dx) Q(d\theta, dz) dt. \end{aligned}$$

On the other hand one has

$$\begin{aligned} & \int_0^\infty \int_S \alpha(t, \theta) \int_{\mathbb{R}^N} \phi(t, x + z + H(t)) \mu(t + \theta, dx) Q(d\theta, dz) dt \\ &= \int_0^\infty \int_S \alpha(t, \theta) \int_{\mathbb{R}^N} \phi(t, x + z + G(t, \theta)) \nu(t + \theta, dx) Q(d\theta, dz) dt. \end{aligned}$$

The result follows by coupling the above computations. \blacksquare

In order to derive the asymptotic self-similar behaviour of $t \mapsto \nu(t, dx)$ for $t \gg 1$ we are interested in the asymptotic as $\lambda \rightarrow \infty$ of the rescaled family of probability measures $p^\lambda(t, dx)$ defined, for $t \geq -\lambda^{-2}$ and for each test function $\varphi \in \mathcal{C}_b(\mathbb{R}^N)$ by

$$\int_{\mathbb{R}^N} \varphi(x) p^\lambda(t, dx) = \int_{\mathbb{R}^N} \varphi\left(\frac{x}{\lambda}\right) \nu(\lambda^2 t, dx), \quad \forall t \geq -\lambda^{-2}. \quad (15)$$

To study its behaviour for $\lambda \gg 1$, we first write down the equation satisfied by p^λ by noticing that

$$\int_{\mathbb{R}^N} \varphi(x) \nu(t, dx) = \int_{\mathbb{R}^N} \varphi(\lambda x) p^\lambda(\lambda^{-2} t, dx), \quad \forall t \geq -1, \quad \forall \varphi \in \mathcal{C}_b(\mathbb{R}^N).$$

Notice that with the notations of Section 2, $p^\lambda(t, dx)$ is the law of the rescaled process $\frac{X(\lambda^2 t) + H(\lambda^2 t)}{\lambda}$.

The equation for p^λ is expressed in the next lemma:

Lemma 8 *Let $\lambda > 0$ be given. Then the function $t \mapsto p^\lambda(t, dx)$ satisfies, for each test function $\phi \in W^{1,1}(0, \infty; \mathcal{C}_b(\mathbb{R}^N)) \cap L^1(0, \infty; \mathcal{C}_b^1(\mathbb{R}^N))$,*

$$\begin{aligned} & \int_{\mathbb{R}^+ \times \mathbb{R}^N} (\partial_t + \lambda H'(\lambda^2 t) \nabla) \phi p^\lambda(t, dx) dt = - \int_{\mathbb{R}^N} \phi(0, x) p^\lambda(0, dx) \\ & + \lambda^2 \int_0^\infty \int_{S \times \mathbb{R}^N} \alpha(\lambda^2 t, \theta) \phi(t, x) p^\lambda(t, dx) Q(d\theta, dz) dt \\ & - \lambda^2 \int_0^\infty \int_{S \times \mathbb{R}^N} \alpha(\lambda^2 t, \theta) \phi(t, x+z) p^\lambda\left(t + \frac{\theta}{\lambda^2}, dx\right) Q^\lambda(\lambda^2 t, \theta; d\theta, dz) dt \end{aligned} \quad (16)$$

wherein, for each $\lambda > 0$, the measure $Q^\lambda(t, \theta; d\theta, dz)$ is defined, for each $t \geq -\lambda^{-2}$, for each $\varphi = \varphi(\theta, z) \in \mathcal{C}_b(S)$, by

$$\int_S \varphi(\theta, z) Q^\lambda(t, \theta; d\theta, dz) = \int_S \varphi\left(\theta, \frac{z}{\lambda} + \frac{1}{\lambda} G(t, \theta)\right) Q(d\theta, dz).$$

The proof of the derivation of this equation directly follows from Lemma 7 using the definition of the map $t \mapsto p^\lambda(t, dx)$.

Here, one may note that we have used a parabolic scaling but the equation is not invariant under this scaling. We are now interested in the asymptotic behaviour of p^λ as $\lambda \rightarrow \infty$ and the main result of this section reads as follows:

Theorem 9 (Young measure convergence) *Let Assumption 1 be satisfied. Recalling the definition of Γ in (4) and of D_0 in (6), the following convergence holds true for each test function $f \in L^1(0, \infty; \mathcal{C}_b(\mathbb{R}^N))$:*

$$\lim_{\lambda \rightarrow \infty} \int_{\mathbb{R}^+ \times \mathbb{R}^N} f(t, x) p^\lambda(t, dx) dt = \int_{\mathbb{R}^+ \times \mathbb{R}^N} f(t, x) \Pi(t, dx) dt,$$

where $\Pi = \Pi(t, dx) \in \mathcal{C}([0, \infty); \mathcal{P})$ is the solution of

$$\partial_t \Pi - \frac{1}{2} \operatorname{div} \left(\frac{1}{1 + \Gamma} D_0 \nabla \Pi \right) = \delta_{t=0} \otimes \delta_0.$$

This result ensures the convergence of p^λ to Π in a rather weak sense, especially in time. This convergence is not sufficient to prove Theorem 2. It will be strengthened in the next section to obtain time pointwise convergence. The proof of the above result will make use of Young measure theory. We refer to reader to [5, 30] for more details and also to Appendix A where basic properties of Young measures are recalled.

Note also that if we choose $f(t, x) = f(x) f_T(t)$ where f is continuous and bounded and f_T is the density of some positive random variable T , the above result implies that the measure $\int_0^\infty p^\lambda(t, dx) f_T(t) dt$ converges in law towards the measure $\int_0^\infty \Pi(t, dx) f_T(t) dt$. If we denote by Z_λ the rescaled process $\left(\frac{X(\lambda^2 t) + H(\lambda^2 t)}{\lambda} \right)_{t \in \mathbb{R}^+}$, one interpretation is to say that $Z_\lambda(T)$, where T is a random variable with density f_T independent of Z_λ , converges in law when λ goes to

infinity towards a random variable with law $\int_0^\infty \Pi(t, dx) f_T(t) dt$. As mentioned above, the objective of Section 4 will be to replace the random variable T by the deterministic time 1. The above theorem also tells us that, in the sense of Young measures, the stochastic process Z_λ converges towards a N -dimensional centred Wiener process with variance-covariance matrix given by $\frac{1}{1+\Gamma} D_0$.

In order to prove Theorem 9, we fix a sequence $\{\lambda_n\}_{n \geq 0} \subset (0, \infty)$ such that

$$\lim_{n \rightarrow \infty} \lambda_n = \infty, \quad (17)$$

and we consider the sequence of Young measures

$$\{t \mapsto p^{\lambda_n}(t, dx)\}_{n \geq 0} \subset L_{\omega^*}^\infty(0, \infty; \mathcal{P}).$$

Then, denoting by \mathcal{M}^+ the set of positive Borel measures on \mathbb{R}^N , due to Lemma A.3 there exists a subsequence, still denoted by λ_n , and $P \equiv P(t, dx) \in L_{\omega^*}^\infty(0, \infty; \mathcal{M}^+)$ such that

$$P(t, dx) \in \mathcal{M}^+ \text{ and } \int_{\mathbb{R}^N} P(t, dx) \leq 1 \text{ a.e. } t \geq 0,$$

and

$$\lim_{n \rightarrow \infty} p^{\lambda_n} = P \text{ weakly* in } (L^1(0, \infty; \mathcal{C}_b(\mathbb{R}^N)))'. \quad (18)$$

Above, $L_{\omega^*}^\infty(0, \infty; \mathcal{M}^+)$ denotes the set of weakly measurable maps from \mathbb{R}^+ into \mathcal{M}^+ and that are essentially bounded (see also Appendix A for more details).

In our next proposition, we will identify the limit measure $P = P(t, dx)$.

Proposition 10 *Let Assumption 1 be satisfied. Then the function $P \equiv P(t, dx)$ is in $L_{\omega^*}^\infty(0, \infty; \mathcal{M}^+)$ defined above is the unique tempered distribution solution of (5), that is $P(t, dx) \equiv \Pi(t, dx)$.*

Since $L^1(0, \infty; \mathcal{C}_0(\mathbb{R}^N))$ is separable, the balls of its dual space are metrizable. This implies the following corollary:

Corollary 11 *Under the assumptions of Proposition 10, one obtains*

$$\lim_{\lambda \rightarrow \infty} p^\lambda = \Pi(t, dx) \text{ weakly* in } (L^1(0, \infty; \mathcal{C}_0(\mathbb{R}^N)))'.$$

Now note that one has

$$\int_{\mathbb{R}^N} \Pi(t, x) dx = 1 \text{ a.e. } t \geq 0.$$

This mass conservation property ensures that the family $\{p_\lambda\}_{\lambda > 0}$ is a tight family of Young measures. Hence Lemma A.4 applies and leads us to the following corollary

Corollary 12 *Under the assumptions of Proposition 10, for each test function $f \in L^1(0, \infty; \mathcal{C}_b(\mathbb{R}^N))$ one has*

$$\lim_{\lambda \rightarrow \infty} \int_{\mathbb{R}^+ \times \mathbb{R}^N} f(t, x) p^\lambda(t, dx) dt = \int_{\mathbb{R}^+ \times \mathbb{R}^N} f(t, x) \Pi(t, dx) dt.$$

To conclude these remarks, one has obtained that proving Proposition 10 is sufficient to complete the proof of Theorem 9. Thus in the remaining of this section we shall focus on proving Proposition 10.

In order to prove Proposition 10, we shall first derive preliminary lemma. In the sequel the notation \mathcal{S} will be used to denote the Schwartz space, that is the set of rapidly decreasing functions, while \mathcal{S}' will be used to denote its dual space, the space of tempered distributions. Next, using these notations, our first lemma reads as follows.

Lemma 13 *Let $F \in \mathcal{C}(\mathbb{R}^+) \cap L^1(\mathbb{R}^+)$ be given. Then the following convergence holds true:*

$$\lim_{\lambda \rightarrow \infty} \lambda^2 F(\lambda^2 t) (p^\lambda(t, dx) \otimes dt) = \left(\int_0^\infty F(s) ds \right) \delta_{t=0} \otimes \delta_0 \text{ in } \mathcal{S}'(\mathbb{R}^+ \times \mathbb{R}^N),$$

that is for each test function $\phi \in \mathcal{S}(\mathbb{R}^+ \times \mathbb{R}^N)$ one has

$$\lim_{\lambda \rightarrow \infty} \lambda^2 \int_{\mathbb{R}^+ \times \mathbb{R}^N} F(\lambda^2 t) \phi(t, x) p^\lambda(t, dx) dt = \left(\int_0^\infty F(s) ds \right) \phi(0, 0).$$

Proof. Let $\phi \equiv \phi(t, x) \in \mathcal{S}(\mathbb{R}^+ \times \mathbb{R}^N)$ be given. Then from the definition of p^λ one obtains:

$$\begin{aligned} \lambda^2 \int_{\mathbb{R}^+ \times \mathbb{R}^N} F(\lambda^2 t) \phi(t, x) p^\lambda(t, dx) dt &= \lambda^2 \int_{\mathbb{R}^+ \times \mathbb{R}^N} F(\lambda^2 t) \phi \left(t, \frac{x}{\lambda} \right) \nu(\lambda^2 t, dx) dt \\ &= \int_{\mathbb{R}^+ \times \mathbb{R}^N} F(t) \phi \left(\frac{t}{\lambda^2}, \frac{x + H(t)}{\lambda} \right) \mu(t, dx) dt \\ &= \int_{\mathbb{R}^+ \times \mathbb{R}^N} F(t) \phi \left(\frac{t}{\lambda^2}, \frac{x + H(t)}{\lambda} \right) (\mu(t, dx) \otimes dt). \end{aligned}$$

Finally since function $(t, x) \mapsto F(t)$ belongs to $L^1(\mathbb{R}^+ \times \mathbb{R}^N; \mu(t, dx) \otimes dt)$, Lebesgue convergence theorem applies and completes the proof of Lemma 13.

■

Now equipped with this lemma we shall first prepare the equation before passing to the limit $\lambda \rightarrow \infty$ and more precisely through the sequence λ_n as $n \rightarrow \infty$.

Lemma 14 (Preparation of the equation) *Let $\phi \in W^{1,1}(0, \infty; \mathcal{C}_b(\mathbb{R}^N)) \cap L^1(0, \infty; \mathcal{C}_b^2(\mathbb{R}^N))$ be given. For all $\lambda > 0$ the function $t \mapsto p^\lambda(t, dx)$ satisfies*

$$\int_{\mathbb{R}^+ \times \mathbb{R}^N} \partial_t \phi dp^\lambda(t, dx) dt = - \int_{\mathbb{R}^N} \phi(0, x) p^\lambda(0, dx) - \mathcal{T}_1^\lambda[\phi] - \mathcal{T}_2^\lambda[\phi] + \mathcal{T}_3^\lambda[\phi], \quad (19)$$

wherein we have set

$$\begin{aligned} \mathcal{T}_1^\lambda[\phi] &= \lambda \int_0^\infty \int_{\mathbb{R}^N} \nabla \phi(t, x) H'(\lambda^2 t) p^\lambda(t, dx) dt \\ &+ \lambda \int_0^\infty \int_S \alpha(\lambda^2 t, \theta) [z + G(\lambda^2 t, \theta)] \left[\int_{\mathbb{R}^N} \nabla \phi(t, x) p^\lambda \left(t + \frac{\theta}{\lambda^2}, dx \right) \right] Q(d\theta, dz) dt, \end{aligned}$$

$$\mathcal{T}_2^\lambda[\phi] = \int_0^\infty \int_S \int_{\mathbb{R}^N} \Phi^\lambda[\phi](t, \theta, x, z) p^\lambda \left(t + \frac{\theta}{\lambda^2}, dx \right) Q(d\theta, dz) dt;$$

with Φ^λ defined by

$$\begin{aligned} \Phi^\lambda[\phi](t, \theta, x, z) \\ = \alpha(\lambda^2 t, \theta) \int_0^1 (1-s) D^2 \phi \left(t, x + s \frac{z + G(\lambda^2 t, \theta)}{\lambda} \right) \cdot [z + G(\lambda^2 t, \theta)]^2 ds; \end{aligned}$$

and

$$\begin{aligned} \mathcal{T}_3^\lambda[\phi] \\ = \lambda^2 \int_0^\infty \int_S \alpha(\lambda^2 t, \theta) \int_{\mathbb{R}^N} \phi(t, x) \left[p^\lambda(t, dx) - p^\lambda \left(t + \frac{\theta}{\lambda^2}, dx \right) \right] Q(d\theta, dz) dt. \end{aligned}$$

The above decomposition directly follows from the formula obtained in (16).

In order to prove Proposition 10 we shall now study the convergence, as $n \rightarrow \infty$, of the different terms arising in the decomposition described in the above lemma with $\lambda = \lambda_n$. Before doing so, let us recall that, due to Assumption 1, one has

$$\alpha(t, \theta) = \alpha_\infty(\theta) + O(e^{-\beta t}) \quad \text{uniformly for } \theta \in [-1, 0] \text{ as } t \gg 1.$$

As a consequence, recalling the definition of $H = H(t)$ in (13), one obtains

$$\begin{aligned} H(t) &= -\mathbf{K}t + O(1) \quad \text{and} \quad H'(t) = -\mathbf{K} + O(e^{-\beta t}) \quad \text{as } t \rightarrow \infty, \\ G(t, \theta) &= \theta \mathbf{K} + O(e^{-\beta t}) \quad \text{uniformly for } \theta \in [-1, 0] \text{ as } t \rightarrow \infty. \end{aligned} \quad (20)$$

These asymptotic expansions will be extensively used in the sequel.

We are now able to investigate the asymptotic behaviour as $\lambda \rightarrow \infty$ of each term arising in the decomposition provided by Lemma 14. To that aim let us fix a test function $\phi \in \mathcal{S}(\mathbb{R}^+ \times \mathbb{R}^N)$. In the sequel of this proof we shall omit to explicitly write down the dependence with respect to ϕ in the decomposition of Lemma 14, that is for $i = 1, \dots, 3$ we shall write \mathcal{T}_i^λ instead of $\mathcal{T}_i^\lambda[\phi]$.

Our convergence proof will be achieved in the next four steps.

Step 0: Recalling (18) one first obtains that

$$\lim_{n \rightarrow \infty} \int_{\mathbb{R}^+ \times \mathbb{R}^N} \partial_t \phi p^{\lambda_n}(t, dx) dt = \int_{\mathbb{R}^+ \times \mathbb{R}^N} \partial_t \phi P(t, dx) dt.$$

Next note that for each $\lambda > 0$ one has

$$\int_{\mathbb{R}^N} \phi(0, x) p^\lambda(0, dx) = \int_{\mathbb{R}^N} \phi \left(0, \frac{x}{\lambda} \right) \mu(0, dx).$$

Hence as $\lambda \rightarrow \infty$ Lebesgue convergence theorem ensures that

$$\lim_{\lambda \rightarrow \infty} \int_{\mathbb{R}^N} \phi(0, x) p^\lambda(0, dx) = \phi(0, 0).$$

Step 1: In this step we investigate the behaviour of \mathcal{T}_1^λ as $\lambda \rightarrow \infty$. To that aim note that one has

$$\begin{aligned} & \lambda \int_S \int_0^\infty \alpha(\lambda^2 t, \theta) [z + G(\lambda^2 t, \theta)] \cdot \left\{ \int_{\mathbb{R}^N} \nabla \phi(t, x) p^\lambda \left(t + \frac{\theta}{\lambda^2}, dx \right) \right\} dt Q(d\theta, dz) \\ &= \lambda \int_S \int_0^\infty \alpha(\lambda^2 t - \theta, \theta) [z + G(\lambda^2 t - \theta, \theta)] \cdot \left\{ \int_{\mathbb{R}^N} \nabla \phi(t, x) p^\lambda(t, dx) \right\} dt Q(d\theta, dz) \\ &+ O\left(\frac{1}{\lambda}\right). \end{aligned}$$

Here the remaining term $O\left(\frac{1}{\lambda}\right)$ depends upon $\|\partial_t \nabla \phi\|_{L^1(0, \infty; W^{1, \infty}(\mathbb{R}^N))}$. As a consequence one obtains that

$$\mathcal{T}_1^\lambda = \lambda \int_{\mathbb{R}^+ \times \mathbb{R}^N} \mathcal{J}_1(\lambda^2 t) \nabla \phi(t, x) p^\lambda(t, dx) dt + O\left(\frac{1}{\lambda}\right),$$

wherein the function $\mathcal{J}_1 : \mathbb{R}^+ \rightarrow \mathbb{R}^N$ is defined by

$$\mathcal{J}_1(t) = H'(t) + \int_S \alpha(t - \theta, \theta) [z + G(t - \theta, \theta)] Q(d\theta, dz).$$

Next note that (20) yields

$$\mathcal{J}_1(t) = O(e^{-\beta t}) \text{ as } t \rightarrow \infty.$$

Hence $\mathcal{J}_1 \in L^1(\mathbb{R}^+)$ and Lemma 13 applies and ensures that $\mathcal{T}_1^\lambda = O\left(\frac{1}{\lambda}\right)$, so that

$$\lim_{\lambda \rightarrow \infty} \mathcal{T}_1^\lambda = 0.$$

Step 2: We are now looking at the second term, namely \mathcal{T}_2^λ . During this step we write Φ^λ instead of $\Phi^\lambda[\phi]$. Next let us first notice that one has

$$\mathcal{T}_2^\lambda = \int_S \left(- \int_{\frac{\theta}{\lambda^2}}^0 + \int_0^\infty \right) \int_{\mathbb{R}^N} \Phi^\lambda \left(t - \frac{\theta}{\lambda^2}, \theta, x, z \right) p^\lambda(t, dx) dt Q(d\theta, dz).$$

Therefore this yields

$$\mathcal{T}_2^\lambda = \int_S \int_0^\infty \int_{\mathbb{R}^N} \Phi^\lambda \left(t - \frac{\theta}{\lambda^2}, \theta, x, z \right) p^\lambda(t, dx) dt Q(d\theta, dz) + O\left(\frac{1}{\lambda^2}\right).$$

Now we claim that:

Claim 15 *The following convergence holds true:*

$$\lim_{\lambda \rightarrow \infty} \int_S \Phi^\lambda \left(t - \frac{\theta}{\lambda^2}, \theta, x, z \right) Q(d\theta, dz) = \frac{1}{2} \operatorname{div} (D_0 \nabla \phi),$$

in $L^1(0, \infty; \mathcal{C}_0(\mathbb{R}^N))$. Here the matrix D_0 is defined by

$$D_0 = \int_S \alpha_\infty(\theta) [z + \theta \mathbf{K}] [z + \theta \mathbf{K}]^T Q(d\theta, dz). \quad (21)$$

The proof of this claim is postponed and we first complete the convergence of $\mathcal{T}_2^{\lambda_n}$. Here recall that $\{\lambda_n\}_{n \geq 0}$ is the sequence chosen at the beginning of this proof (see (17) and (18)). Now because of the Young convergence property $p^{\lambda_n}(t, dx) \rightarrow P(t, dx)$ we obtain that

$$\begin{aligned} & \lim_{n \rightarrow \infty} \int_0^\infty \int_{\mathbb{R}^N} \int_S \Phi^{\lambda_n} \left(t - \frac{\theta}{\lambda_n^2}, \theta, x, z \right) Q(d\theta, dz) p^{\lambda_n}(t, dx) dt \\ &= \frac{1}{2} \int_{\mathbb{R}^+ \times \mathbb{R}^N} \operatorname{div} (D_0 \nabla \phi) P(t, dx) dt. \end{aligned}$$

This re-writes as

$$\lim_{n \rightarrow \infty} \mathcal{T}_2^{\lambda_n} = \frac{1}{2} \int_{\mathbb{R}^+ \times \mathbb{R}^N} \operatorname{div} (D_0 \nabla \phi) P(t, dx),$$

that completes the proof of this step.

It remains to prove Claim 15. To do so, let us observe that there exists some constant $C > 0$ such that for all $(t, x) \in \mathbb{R}^+ \times \mathbb{R}^N$, $(\theta, z) \in S$, $s \in [0, 1]$ and $\lambda \geq 1$ one has, setting $Z = z + G(\lambda^2 t - \theta, \theta)$,

$$\begin{aligned} & \int_S \left| \left[D^2 \phi \left(t - \frac{\theta}{\lambda^2}, x + s \frac{Z}{\lambda} \right) - D^2 \phi(t, x) \right] \cdot Z^2 \right| Q(d\theta, dz) \\ & \leq \frac{C}{\lambda} \int_{[-1, 0] \times \{|z| \leq \lambda^{1/4}\}} [|z|^3 + 1] Q(d\theta, dz) + C \int_{[-1, 0] \times \{|z| \geq \lambda^{1/4}\}} [|z|^2 + 1] Q(d\theta, dz) \\ & \leq \frac{C(1 + \lambda^{3/4})}{\lambda} + C \int_{[-1, 0] \times \{|z| \geq \lambda^{1/4}\}} [|z|^2 + 1] Q(d\theta, dz) \end{aligned}$$

This upper bound converges to zero as $\lambda \rightarrow \infty$ so that

$$\lim_{\lambda \rightarrow \infty} \int_S \left| \left[D^2 \phi \left(t - \frac{\theta}{\lambda^2}, x + s \frac{Z}{\lambda} \right) - D^2 \phi(t, x) \right] \cdot Z^2 \right| Q(d\theta, dz) = 0,$$

uniformly for $(t, x) \in \mathbb{R}^+ \times \mathbb{R}^N$ and also in $L^1(0, \infty; \mathcal{C}_b(\mathbb{R}^N))$ since $\phi \in \mathcal{S}$ and using Lebesgue convergence theorem. To complete the proof of the claim, it is sufficient to show that, for the topology of $L^1(0, \infty; \mathcal{C}_b(\mathbb{R}^N))$, one has

$$\lim_{\lambda \rightarrow \infty} \int_S D^2 \phi(t, x) \cdot Z^2 Q(\theta, dz) = \int_S D^2 \phi(t, x) \cdot [z + \theta \mathbf{K}]^2 Q(d\theta, dz).$$

This latter convergence directly follows from the asymptotic expansion of $G(t, \theta)$ in (20). This complete the proof of Claim 15 by noticing that

$$\int_S D^2 \phi(t, x) \cdot [z + \theta \mathbf{K}]^2 Q(d\theta, dz) = \operatorname{div} (D_0 \nabla \phi),$$

where the matrix D_0 is defined in (21). This completes the proof of the claim.

Step 3: In this last step we investigate the limit, as $\lambda \rightarrow \infty$, of \mathcal{T}_3^λ . In order to deal with this term, note that

$$\begin{aligned}\mathcal{T}_3^\lambda &= \lambda^2 \int_0^\infty \int_S \alpha(\lambda^2 t, \theta) \int_{\mathbb{R}^N} \phi(t, x) p^\lambda(t, dx) Q(d\theta, dz) dt \\ &\quad - \lambda^2 \int_S \int_{\theta/\lambda^2}^0 \alpha(\lambda^2 t - \theta) \int_{\mathbb{R}^N} \phi\left(t - \frac{\theta}{\lambda^2}, x\right) p^\lambda(t, dx) dt Q(d\theta, dz) \\ &\quad - \lambda^2 \int_0^\infty \int_S \alpha(\lambda^2 t - \theta, \theta) \int_{\mathbb{R}^N} \phi\left(t - \frac{\theta}{\lambda^2}, x\right) p^\lambda(t, dx) Q(d\theta, dz) dt.\end{aligned}$$

For the last term, let us observe that one has

$$\begin{aligned}& \int_0^\infty \int_S \alpha(\lambda^2 t - \theta, \theta) \int_{\mathbb{R}^N} \phi\left(t - \frac{\theta}{\lambda^2}, x\right) p^\lambda(t, dx) Q(d\theta, dz) dt \\ &= \int_0^\infty \int_S \alpha(\lambda^2 t - \theta, \theta) \int_{\mathbb{R}^N} \phi(t, x) p^\lambda(t, dx) Q(d\theta, dz) dt \\ &\quad - \int_0^\infty \int_S \frac{\theta}{\lambda^2} \alpha(\lambda^2 t - \theta, \theta) \int_{\mathbb{R}^N} \partial_t \phi(t, x) p^\lambda(t, dx) Q(d\theta, dz) dt + O\left(\frac{1}{\lambda^4}\right).\end{aligned}$$

This allows us to re-write \mathcal{T}_3^λ as follows:

$$\begin{aligned}\mathcal{T}_3^\lambda &= \lambda^2 \int_0^\infty \int_{\mathbb{R}^N} F(\lambda^2 t) p^\lambda(t, dx) dt \\ &\quad - \lambda^2 \int_S \int_{\theta/\lambda^2}^0 \alpha(\lambda^2 t - \theta) \int_{\mathbb{R}^N} \phi\left(t - \frac{\theta}{\lambda^2}, x\right) p^\lambda(t, dx) dt Q(d\theta, dz) \\ &\quad + \int_0^\infty \int_S \theta \alpha(\lambda^2 t - \theta, \theta) \int_{\mathbb{R}^N} \partial_t \phi(t, x) p^\lambda(t, dx) Q(d\theta, dz) dt + O\left(\frac{1}{\lambda^4}\right).\end{aligned}$$

In the above decomposition we have set

$$F(t) = \int_S [\alpha(t, \theta) - \alpha(t - \theta, \theta)] Q(d\theta, dz).$$

Now, recalling (20), one has $F(t) = O(e^{-\beta t})$ as $t \rightarrow \infty$ so that $F \in L^1(\mathbb{R}^+)$ and Lemma 13 applies and ensures that

$$\lim_{\lambda \rightarrow \infty} \lambda^2 \int_0^\infty \int_{\mathbb{R}^N} F(\lambda^2 t) p^\lambda(t, dx) dt = \phi(0, 0) \int_0^\infty F(s) ds.$$

Next, using the same argument as for the proof of Lemma 13, one has

$$\begin{aligned}& \lim_{\lambda \rightarrow \infty} \lambda^2 \int_S \int_{\theta/\lambda^2}^0 \alpha(\lambda^2 t - \theta) \int_{\mathbb{R}^N} \phi\left(t - \frac{\theta}{\lambda^2}, x\right) p^\lambda(t, dx) dt Q(d\theta, dz) \\ &= \phi(0, 0) \int_S \int_\theta^0 \alpha(t - \theta) dt Q(d\theta, dz).\end{aligned}$$

And, for the last term we get, along the sequence $\{\lambda_n\}_{n \geq 0}$ (see (18)),

$$\begin{aligned} & \lim_{n \rightarrow \infty} \int_0^\infty \int_S \theta \alpha(\lambda_n^2 t - \theta, \theta) \int_{\mathbb{R}^N} \partial_t \phi(t, x) p^{\lambda_n}(t, dx) Q(d\theta, dz) dt \\ &= \int_0^\infty \int_S \theta \alpha_\infty(\theta) Q(d\theta, dz) \int_0^\infty \int_{\mathbb{R}^N} \partial_t \phi(t, x) P(t, dx) dt \\ &= -\Gamma \int_0^\infty \int_{\mathbb{R}^N} \partial_t \phi(t, x) P(t, dx) dt. \end{aligned}$$

To summarize we have obtained the following property:

$$\begin{aligned} \lim_{n \rightarrow \infty} \mathcal{T}_3^{\lambda_n} &= \phi(0, 0) \int_0^\infty F(s) ds - \phi(0, 0) \int_S \int_\theta^0 \alpha(t - \theta) dt Q(d\theta, dz) \\ &\quad - \Gamma \int_0^\infty \int_{\mathbb{R}^N} \partial_t \phi(t, x) P(t, dx) dt. \end{aligned}$$

And, to complete this step, note that

$$\begin{aligned} & \int_0^\infty F(s) ds - \int_S \int_\theta^0 \alpha(t - \theta) dt Q(d\theta, dz) \\ &= \lim_{M \rightarrow \infty} \left[\int_S \int_0^M [\alpha(t, \theta) - \alpha(t - \theta, \theta)] Q(d\theta, dz) \int_\theta^0 \alpha(t - \theta) \right] dt \\ &= - \lim_{M \rightarrow \infty} \int_S \int_M^{M-\theta} \alpha(t, \theta) dt Q(d\theta, dz) = \int_S \theta \alpha_\infty(\theta) Q(d\theta, dz) = -\Gamma. \end{aligned}$$

Hence we get

$$\lim_{n \rightarrow \infty} \mathcal{T}_3^{\lambda_n} = -\Gamma \phi(0, 0) - \Gamma \int_0^\infty \int_{\mathbb{R}^N} \partial_t \phi(t, x) P(t, dx) dt.$$

We now conclude the proof of Proposition 10 and thus the one of Theorem 9.

Conclusion of the proof of Proposition 10:

From the four previous steps we have obtained the following convergence: for all $\phi \in \mathcal{S}(\mathbb{R}^+ \times \mathbb{R}^N)$, on the one hand, one has

$$\lim_{n \rightarrow \infty} \int_{\mathbb{R}^+ \times \mathbb{R}^N} \partial_t \phi(t, x) p^{\lambda_n}(t, dx) dt = \int_{\mathbb{R}^+ \times \mathbb{R}^N} \partial_t \phi(t, x) P(t, dx) dt,$$

and, on the other hand, one has

$$\begin{aligned} & \lim_{n \rightarrow \infty} \int_{\mathbb{R}^+ \times \mathbb{R}^N} \partial_t \phi(t, x) p^{\lambda_n}(t, dx) dt = -(1 + \Gamma) \phi(0, 0) \\ & - \frac{1}{2} \int_{\mathbb{R}^+ \times \mathbb{R}^N} \operatorname{div} (D_0 \nabla \phi) P(t, dx) - \Gamma \int_{\mathbb{R}^+ \times \mathbb{R}^N} \partial_t \phi(t, x) P(t, dx) dt. \end{aligned}$$

Hence $P \in \mathcal{S}'(\mathbb{R}^+ \times \mathbb{R}^N)$ is a solution of the equation

$$\partial_t \Pi = \delta_{t=0} \otimes \delta_0 + \frac{1}{2} \operatorname{div} \left(\frac{D_0}{1 + \Gamma} \nabla \Pi \right) \text{ in } \mathcal{S}'(\mathbb{R}^+ \times \mathbb{R}^N).$$

The latter equation has a unique tempered distribution, Π , that satisfies $\Pi \in \mathcal{C}(\mathbb{R}^+, \mathcal{P})$. Finally, since the limit function Π is unique, one obtains $P(t, dx) = \Pi(t, dx)$ and this complete the proof of Proposition 10 and thus the one of Theorem 9 since the sequence (λ_n) is arbitrary.

4 Proof of Theorem 2

In this section, we complete the proof of Theorem 2. To that aim we will prove that the family $\{p^\lambda(t, dx)\}_{\lambda > 0}$ is relatively compact with respect to a stronger topology (in time) than those of the Young measures. We shall more specifically show that this family is relatively compact for the topology of $\mathcal{C}_{\text{loc}}((0, \infty); H_{\text{loc}}^{-\sigma}(\mathbb{R}^N))$ for some parameter $\sigma > 0$ large enough. And, this compactness property will be sufficient to complete the proof of the Theorem 2 by using the identification of the weak limit as described in Theorem 9.

The main result of this section reads as follows.

Theorem 16 *Let $R > 0$, $0 < \varepsilon < T$ be given. Let $\sigma > \frac{N}{2} + 2$ be given. Then there exists $\tilde{\lambda} > 0$ such that the family $\{p^\lambda(t, dx)\}_{\lambda > \tilde{\lambda}}$ is relatively compact in $\mathcal{C}([\varepsilon, T]; H^{-\sigma-1}(B_R))$. Herein $B_R \subset \mathbb{R}^N$ denotes the ball of radius R centred at the origin while for all $s > \frac{N}{2} + 1$, $H^{-s}(B_R) = (H_0^s(B_R))'$ denotes the dual space of the Hilbert space $H_0^s(B_R)$.*

The proof of this key result is postponed and we first complete the proof of Theorem 2.

Proof of Theorem 2. Recalling the definition of $p^\lambda(t, dx)$ given in (15), to prove Theorem 2, it is sufficient to prove that

$$\lim_{t \rightarrow \infty} p^{\sqrt{t}}(1, dx) = \Pi_1(dx) \text{ for the narrow topology of } \mathcal{P}. \quad (22)$$

To prove the above statement, fix $\sigma > \frac{N}{2} + 2$ and consider a sequence $\{\lambda_n\}_{n \geq 0} \subset (0, \infty)$ such that $\lambda_n \rightarrow \infty$ as $n \rightarrow \infty$. Then because of Theorem 9, Theorem 16 and using a diagonal extraction process, there exists a sub-sequence, still denoted by λ_n , such that

$$\lim_{n \rightarrow \infty} p^{\lambda_n}(t, dx) = \Pi_t \text{ locally uniformly for } t \in (0, \infty) \text{ with value in } H^{-\sigma-1}(B_R).$$

Note also that, possibly along a sub-sequence, the sequence of probability measure $\{p^{\lambda_n}(1, dx)\}$ converges to some positive measure with respect to the vague topology of measures. Hence, one obtained that

$$p^{\lambda_n}(1, dx) \rightarrow \Pi_1(dx) \text{ as } n \rightarrow \infty,$$

wherein the above limit holds with respect to the vague topology of measures. Finally, since $\Pi_1(\mathbb{R}^N) = 1$, Π_1 is also a probability measure and the above limit holds for the narrow topology of probability measures. Now, since the sequence $\{\lambda_n\}$ is arbitrary and \mathcal{P} , endowed with the narrow topology, is a metrizable space, one obtains that

$$\lim_{\lambda \rightarrow \infty} p^\lambda(1, dx) = \Pi_1 \text{ for the narrow topology of } \mathcal{P}.$$

Thus (22) holds true by choosing $\lambda = \sqrt{t}$. This completes the proof of Theorem 2.

■

In the remaining parts of this section, we prove Theorem 16. The proof of this result relies on the formulation of the equation for p^λ obtained in Lemma 14. Here we re-write it with a slightly different form, as follows: for $\phi \in W^{1,1}(0, \infty; \mathcal{C}_b(\mathbb{R}^N)) \cap L^1(0, \infty; \mathcal{C}_b^2(\mathbb{R}^N))$ and $\lambda > 0$, the function $t \mapsto p^\lambda(t, dx)$ satisfies

$$\begin{aligned} \int_{\mathbb{R}^+ \times \mathbb{R}^N} \left[T_{\frac{1}{\lambda^2}} \partial_t \phi \right] (t, x) p^\lambda(t, dx) dt &= - \int_{\mathbb{R}^N} \phi(0, x) p^\lambda(0, dx) - \mathcal{T}_1^\lambda[\phi] \\ &\quad - \mathcal{T}_2^\lambda[\phi] + \mathcal{R}^\lambda[\phi], \end{aligned} \quad (23)$$

wherein \mathcal{T}_1^λ and \mathcal{T}_2^λ are defined in Lemma 14, $T_{\frac{1}{\lambda^2}}$ denotes the nonlocal operator defined by

$$\begin{aligned} g(t, x) &= [T_{\frac{1}{\lambda^2}} f](t, x) \\ \Leftrightarrow g(t, x) &= f(t, x) + \lambda^2 \int_S \alpha_\infty(\theta) \int_0^{-\frac{\theta}{\lambda^2}} f(t+l, x) dl Q(d\theta, dz), \end{aligned}$$

while

$$\begin{aligned} \mathcal{R}^\lambda[\phi] &= \lambda^2 \int_0^\infty \int_S [\alpha(\lambda^2 t, \theta) - \alpha_\infty(\theta)] \\ &\quad \int_{\mathbb{R}^N} \phi(t, x) \left[p^\lambda(t, dx) - p^\lambda\left(t + \frac{\theta}{\lambda^2}, dx\right) \right] Q(d\theta, dz) dt \\ &\quad + \lambda^2 \int_S \alpha_\infty(\theta) \int_{\frac{\theta}{\lambda^2}}^0 \int_{\mathbb{R}^N} \phi\left(t - \frac{\theta}{\lambda^2}, x\right) p^\lambda(t, dx) dt Q(d\theta, dz). \end{aligned}$$

Now, in order to prove Theorem 16 we will investigate some regularization properties for the nonlocal operator $T_{\frac{1}{\lambda^2}}$ defined above when $\lambda \gg 1$ is large. And, we will be used to complete the proof of the theorem. In the sequel we will first investigate some properties of the linear operator $T_{\frac{1}{\lambda^2}}$ before going to the proof of Theorem 16.

4.1 Regularity properties

Let $(X, \|\cdot\|_X)$ be a reflexive Banach space. For each $h > 0$, consider the linear operator T_h defined from $\mathcal{S}(\mathbb{R}^+; X')$ into itself by

$$T_h[\phi](t) = \phi(t) + \frac{1}{h} \int_{\mathcal{S}} \alpha_{\infty}(\theta) \int_0^{-h\theta} \phi(t+l) dl Q(d\theta, dz), \quad t \geq 0. \quad (24)$$

Here X' denotes the dual space of X . Then the main result of this sub-section reads as follows:

Theorem 17 (Sobolev semi-norm estimate) *Let $\{u^h\}_{h>0}$ be a family of X -valued tempered distribution on \mathbb{R}^+ , namely $\{u_h\}_{h>0} \subset \mathcal{S}'(\mathbb{R}^+; X)$. We assume that there exist some constant $M > 0$ and $\alpha \in (0, 1)$ such that for all $h > 0$ and all $\phi \in \mathcal{S}(\mathbb{R}^+; X')$ one has:*

$$\begin{aligned} & |\langle T_h \phi', u^h \rangle| \\ & \leq M \left[\|\phi\|_{L^{\infty}(\mathbb{R}^+; X')} + \|\phi\|_{L^1(\mathbb{R}^+; X')} + \sup_{s>0} \frac{1}{s^{\alpha}} \|\phi(\cdot + s) - \phi\|_{L^1(\mathbb{R}^+; X')} \right]. \end{aligned} \quad (25)$$

Herein the symbol $\langle \cdot, \cdot \rangle$ is used to denote the duality pairing between $\mathcal{S}(\mathbb{R}^+; X')$ and $\mathcal{S}'(\mathbb{R}^+; X)$. Then, for each $p > 1$ and $s \in [\alpha, 1)$ with $ps > 1$, one has

$$u^h \in W_{\text{loc}}^{1-s, p'}(\mathbb{R}^+; X), \quad \forall h > 0,$$

where p' denotes the conjugate exponent of p . Moreover for each given $T > 0$, there exists some constant $C = C(p, s, T)$ such that

$$[u^h]_{W^{1-s, p'}(0, T; X)} \leq C, \quad \forall h > 0.$$

In the above estimate, the bracket denotes the X -valued Sobolev semi-norm described below.

The proof of this result relies on several preliminary studies. As announced in Theorem 17, the estimates we shall obtain make use of Banach valued fractional Sobolev spaces. For that purpose let us first introduce further notations, definitions and some well known results and estimates related to Banach valued fractional Sobolev spaces. We refer to [1, 25] and the references therein for more details on such a topic.

Let $(Y, \|\cdot\|_Y)$ be a given Banach space. For each open interval $I \subset \mathbb{R}$, each $p \in [1, \infty]$ and $\theta \in (0, 1)$, we define the Sobolev space $W^{\theta, p}(I; Y)$ as follows:

$$W^{\theta, p}(I; Y) = \{\psi \in L^p(I; Y) : [\psi]_{W^{\theta, p}(I; Y)} < \infty\},$$

where the Sobolev semi-norm $[\cdot]_{W^{\theta, p}(I; Y)}$ is defined by

$$[\psi]_{W^{\theta, p}(I; Y)} = \left(\iint_{I \times I} \frac{\|\psi(t) - \psi(s)\|_Y^p}{|t - s|^{\theta p + 1}} dt ds \right)^{\frac{1}{p}}. \quad (26)$$

The Sobolev norm is then defined as

$$\|\psi\|_{W^{\theta,p}(I;Y)} = \|\psi\|_{L^p(I;Y)} + [\psi]_{W^{\theta,p}(I;Y)}, \quad \forall \psi \in W^{\theta,p}(I;Y).$$

Now we define Sobolev spaces with negative index. When $I \subset \mathbb{R}$ is a given bounded interval, then for $p \in [1, \infty]$ and $\theta \in (0, 1)$ we define the Sobolev space $W^{-1+\theta,p}(I;Y)$ as the image of the distributional derivative ∂_t , that is $\partial_t W^{\theta,p}(I;Y) \subset \mathcal{D}'(I;Y)$, the set of Y -valued distributions. In other word it is defined as the following:

$$u \in W^{-1+\theta,p}(I;Y) \Leftrightarrow \exists v \in W^{\theta,p}(I;Y), \quad u = \partial_t v \text{ in } \mathcal{D}'(I;Y).$$

Such a function v is called a representation of u . The set of the representations of u is given by $v + Y$ where v is a representation of u . The space $W^{1+\theta,p}(I;Y)$ becomes a Banach space when endowed with the norm $\|\cdot\|_{W^{1+\theta,p}(I;Y)}$ defined as

$$\|u\|_{W^{1+\theta,p}(I;Y)} = \inf_{y \in Y} \|v + y\|_{L^p(I;Y)} + [v]_{W^{\theta,p}(I;Y)},$$

where v is a given representation of u . Note that such a definition does not depend upon the choice of the representation v .

Now let us recall a duality representation that will be used to prove Theorem 17. The proof of this result can be found in [1].

Lemma 18 *Let $I \subset \mathbb{R}$ be a given interval. Let $(Y, \|\cdot\|_Y)$ be a given separable Banach space. Let $p \in (1, \infty)$ and $\theta \in (0, 1)$ be given. Then one has*

$$W^{-\theta,p'}(I;Y) = \left(W_0^{\theta,p}(I;Y') \right)'.$$

Here p' denotes the conjugate exponent associated to p , Y' is the dual space of Y while $W_0^{\theta,p}(I;Y')$ is the closure of $\mathcal{D}(I;Y')$ in $W^{\theta,p}(I;Y')$. The above duality representation holds with respect to the duality pairing $\langle \cdot; \cdot \rangle$ on $\mathcal{D}(I;Y) \times \mathcal{D}(I;Y')$ defined by

$$\langle v; v' \rangle = \int_I \langle v(t); v'(t) \rangle_{Y,Y'} dt, \quad \forall (v, v') \in \mathcal{D}(I;Y) \times \mathcal{D}(I;Y').$$

Before recalling some important estimates that will be used in the sequel, let us introduce further notations. Let $I \subset \mathbb{R}$ be a given interval and let $(Y, \|\cdot\|_Y)$ be a Banach space. For each $p \in [1, \infty]$ and $s \in (0, 1)$ let us define the Besov semi-norm for a function $\phi \in L_{\text{loc}}^p(I;Y)$ by

$$[\phi]_{B_{\infty}^{s,p}(I;Y)} = \sup_{h>0} h^{-s} \|\phi(\cdot + h) - \phi\|_{L^p(I_h;Y)} \text{ with } I_h = \{x \in I : x + h \in I\}.$$

We now turn to derive some important estimates. Using straightforward computations we can derive the following estimates:

Lemma 19 *Let $\theta \in (0, 1)$ be given. Let $T > 0$ be given. Then for all $\psi \in \mathcal{D}(0, T; Y)$ one has*

$$[\bar{\psi}]_{W^{\theta,1}(\mathbb{R}; Y)} \leq [\psi]_{W^{\theta,1}(0, T; Y)} + \frac{4T^{1-\theta}}{\theta(1-\theta)} \|\psi\|_{L^\infty(0, T; Y)}.$$

Here $\bar{\psi}$ denotes the extension by zero of ψ .

Next let us recall the estimate derived by Simon in Corollary 24 of [25]:

Lemma 20 *Let $I \subset \mathbb{R}$ be a given interval and let $(Y, \|\cdot\|_Y)$ be a Banach space. Let $\theta \in (0, 1)$ and $p \in [1, \infty)$ be given. Then for each $\psi \in W^{\theta,p}(I, Y)$ one has*

$$[\psi]_{B_\infty^{\theta,p}(I; Y)} \leq \frac{2}{\theta} [\psi]_{W^{\theta,p}(I; Y)}.$$

Now in view of proving Theorem 17, we are concerned in deriving some properties of the linear operator T_h defined above in (24). Roughly speaking, we shall show that such an operator is invertible on $L^1(\mathbb{R}; X')$ and that the inverse operator is a kernel operator that enjoys nice estimates. To that aim, we first investigate some properties of the function $\Delta : \mathbb{C} \rightarrow \mathbb{C}$ defined by

$$\Delta(s) = s - \int_S \alpha_\infty(\theta) [e^{s\theta} - 1] Q(d\theta, dz).$$

The following lemma holds true:

Lemma 21 *The analytic function Δ enjoys the following property: there exists $\varepsilon_0 > 0$ such that*

$$\begin{cases} \Re(s) > -\varepsilon_0, \\ \Delta(s) = 0, \end{cases} \Rightarrow s = 0.$$

Proof. Note that $\Delta(0) = 0$. Next let $s = \alpha + i\omega$ with $\alpha \geq 0$ and ω be such that $\Delta(s) = 0$. Let us show that $\alpha = \omega = 0$.

To do so, let us first observe that the function $x \mapsto \Delta(x)$ is increasing on \mathbb{R} . Hence $s = 0$ is the only real root of Δ . Next the equation $\Delta(\alpha + i\omega) = 0$ re-writes as

$$\begin{cases} \alpha + \int_S \alpha_\infty(\theta) Q(d\theta, dz) = \int_S \alpha_\infty(\theta) e^{\alpha\theta} \cos(\omega\theta) Q(d\theta, dz), \\ \omega - \int_S \alpha_\infty(\theta) e^{\alpha\theta} \sin(\omega\theta) Q(d\theta, dz) = 0. \end{cases}$$

Hence, from the first equation one gets $\Delta(\alpha) \leq 0$, that implies that $\alpha \leq 0$ and thus $\alpha = 0$ since $\alpha \geq 0$. Next the first equation becomes

$$\int_S \alpha_\infty(\theta) [1 - \cos(\omega\theta)] Q(d\theta, dz) = 0,$$

and, since $\alpha_\infty > 0$, this implies that $\cos(\omega\theta) = 1$ Q -a.e. for $(\theta, z) \in S$ and thus $\sin(\omega\theta) = 0$ Q -a.e. for $(\theta, z) \in S$. Plugging this information together with

$\alpha = 0$ into the second equation in the above system yields $\omega = 0$. The above argument shows that

$$\Re(s) \geq 0 \text{ and } \Delta(s) = 0 \Rightarrow s = 0.$$

Recalling that Δ is an analytic function, to complete the proof of the lemma, it is sufficient to note that if $s \in \mathbb{C}$ satisfies $\Delta(s) = 0$ then

$$|\Im(s)| \leq \int_S \alpha_\infty(\theta) e^{\theta \Re(s)} Q(d\theta, dz).$$

Indeed the above estimate ensures that there is no sequence of roots for Δ approaching the imaginary axis. This completes the proof of the lemma. ■

Observe furthermore that $\Delta'(0) = 1 - \int_S \theta \alpha_\infty(\theta) Q(d\theta, dz) > 0$ so that $s = 0$ is a simple root and the function $s \mapsto \frac{s}{\Delta(s)}$ is holomorphic on the half plane $\{s \in \mathbb{C} : \Re(s) > -\varepsilon_0\}$.

Lemma 22 *For each function $\psi \in L^1(\mathbb{R}, X')$, for each parameter $h > 0$, there exists a unique function $\phi = \phi_h \in L^1(\mathbb{R}; X')$ such that*

$$\psi = T_h[\phi].$$

Moreover there exists a real valued function $K \in L^1(\mathbb{R}; \mathbb{R})$ such that there exist $\varepsilon_0 > 0$ small enough and some constant $M_0 > 0$ such that:

$$|K(x)| \leq M_0 e^{-\varepsilon_0 |x|}, \forall x \in \mathbb{R}, \quad K(x) = 0 \text{ a.e. } x \leq 0, \quad (27)$$

and such that for all $\psi \in L^1(\mathbb{R}, X')$, for all $h > 0$, one has

$$\psi = T_h[\phi] \Leftrightarrow \phi(t) = \psi(t) + \frac{1}{h} \int_{\mathbb{R}} K\left(\frac{t-s}{h}\right) \psi(s) ds, \quad t \in \mathbb{R}. \quad (28)$$

In the sequel we shall denote for each $h > 0$:

$$K_h(x) = \frac{1}{h} K(h^{-1}x).$$

Note that the above bound, namely (27), for the convolution kernel K_h ensures that

$$\|K_h\|_{L^1(\mathbb{R})} \leq M_0 \varepsilon_0^{-1}, \quad \forall h > 0.$$

Proof. Let $h > 0$ be given. Let $\psi \in L^1(\mathbb{R}; X')$ be given. We aim at solving the equation $T_h[\phi] = \psi$. We shall denote by \mathcal{F} the Fourier transform. Then applying the Fourier transform yields

$$\mathcal{F}T_h[\phi](\xi) = \left[1 + \frac{1}{h} \int_S \alpha_\infty(\theta) \int_0^{-h\theta} e^{-i\xi l} dl Q(d\theta, dz) \right] \mathcal{F}\phi(\xi) = \frac{\Delta(ih\xi)}{ih\xi} \mathcal{F}\phi(\xi).$$

Now we claim that:

Claim 23 *There exists a function $K : \mathbb{R} \rightarrow \mathbb{R}$ and $\varepsilon > 0$ such that*

$$K(x) = O(e^{-\varepsilon x}), \quad x \rightarrow \infty, \quad K(x) = 0 \text{ for } x \leq 0,$$

and

$$\frac{i\xi}{\Delta(i\xi)} = 1 + \mathcal{F}K(\xi), \quad \forall \xi \in \mathbb{R}.$$

Before proving this claim we first complete the proof of Lemma 22. Indeed, due to the above claim, one has

$$\frac{ih\xi}{\Delta(ih\xi)} = 1 + \mathcal{F}K_h(\xi), \quad \forall \xi \in \mathbb{R}, \quad h > 0.$$

Hence the function $\phi \in L^1(\mathbb{R}; X')$ is uniquely defined by (28) and Lemma 22 follows. ■

It remains to prove Claim 23.

Proof of Claim 23. Recall that the function $\Delta : \mathbb{C} \rightarrow \mathbb{C}$ is defined by

$$\Delta(s) = s - \int_S \alpha_\infty(\theta) [e^{s\theta} - 1] Q(d\theta, dz), \quad \forall s \in \mathbb{C},$$

and that there exists $\varepsilon > 0$ such that $s \mapsto \frac{s}{\Delta(s)}$ is analytic on the half plane $\mathbb{H} := \{s \in \mathbb{C} : \Re(s) > -\varepsilon\}$.

Next setting

$$\gamma := \int_S \alpha_\infty(\theta) Q(d\theta, dz) \text{ and } I(s) = \int_S \alpha_\infty(\theta) e^{s\theta} Q(d\theta, dz),$$

one has

$$\begin{aligned} G(s) &:= \frac{s}{\Delta(s)} = \frac{s + I(s)}{s + \gamma} + \frac{s}{s + \gamma - I(s)} - \frac{s + I(s)}{s + \gamma} \\ &= \frac{s + I(s)}{s + \gamma} - \frac{I(s)(\gamma - I(s))}{\Delta(s)(s + \gamma)}, \quad \forall s \in \mathbb{H}. \end{aligned}$$

Hence one obtains the following decomposition, for any $s \in \mathbb{H}$,

$$G(s) = 1 - \frac{\gamma}{s + \gamma} + \frac{I(s)}{s + \gamma} + R(s) \text{ with } R(s) := \frac{I(s)(I(s) - \gamma)}{\Delta(s)(s + \gamma)}.$$

Up to reducing $\varepsilon > 0$ if necessary, one may suppose that $0 < \varepsilon < \gamma$ so that each term arising in the above decomposition is analytic of the half plane \mathbb{H} .

Next consider the function $E = E(x)$ defined by

$$E(x) = \begin{cases} 0 & \text{if } x \leq 0, \\ e^{-x} & \text{if } x > 0. \end{cases} \quad (29)$$

And first observe that, for all $\xi \in \mathbb{R}$, one has

$$\frac{\gamma}{i\xi + \gamma} = \mathcal{F}[\gamma E(\gamma \cdot)](\xi).$$

Observe that one also has, for any $\xi \in \mathbb{R}$,

$$\frac{I(i\xi)}{i\xi + \gamma} = \mathcal{F} \left[\int_S E(\gamma \cdot + \theta) Q(d\theta, dz) \right] (\xi).$$

On the other hand, observe that the function $I = I(s)$ is bounded on the complex half plane \mathbb{H} . Indeed, one has

$$|I(s)| \leq M := \int_S \alpha_\theta(\theta) e^{-\varepsilon\theta} Q(d\theta, dz), \quad \forall s \in \mathbb{H}.$$

As a consequence, there exists some constant $C > 0$ such that

$$|R(s)| \leq \frac{C}{1 + |s|^2}, \quad \forall s \in \mathbb{H}.$$

As a consequence of this fast decay at infinity and since R is analytic on \mathbb{H} , one obtains, for all $x \in \mathbb{R}$ and all $\kappa \geq -\varepsilon$, that

$$K_R(x) := \mathcal{F}^{-1} R(i \cdot)(x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{ix\xi} R(i\xi) d\xi = \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{\kappa x + ix\xi} R(\kappa + i\xi) d\xi.$$

Hence choosing $\kappa = -\varepsilon$, we conclude that the function K_R has an exponential decay at infinity. In addition, because of the uniform bound in (4.1), Theorem 4.4 in [27] ensures that there exists a tempered distribution T supported in $[0, \infty)$ such that for all $\kappa > 0$ and $\xi \in \mathbb{R}$ one has

$$R(i(\xi + i(\kappa - \varepsilon))) = \mathcal{F}(e^{-\kappa \cdot} T)(\xi).$$

Choosing $\kappa = \varepsilon > 0$ yields $K_R = e^{-\varepsilon \cdot} T$ and one concludes that $\text{supp } K_R \subset [0, \infty)$.

As a consequence of the above steps, one obtained that for each $\xi \in \mathbb{R}$,

$$G(i\xi) = 1 + \mathcal{F} \left[-\gamma E(\gamma \cdot) + \int_S E(\gamma \cdot + \theta) Q(d\theta, dz) + K_R \right] (\xi).$$

Finally, recalling the definition of E in (29), $\text{supp } K_R \subset [0, \infty)$ and the exponential decay of K_R , this completes the proof of Claim 23.

■

Remark 24 Note that if $\psi \in \mathcal{D}(\mathbb{R}; X')$ then, for each $h > 0$, the solution $\phi = T_h^{-1} \psi$ satisfies $\phi \in \mathcal{C}^\infty(\mathbb{R}; X')$. Moreover one has $\text{supp } \phi \subset \text{supp } \psi + [0, \infty)$ and, for each $k \geq 0$, $\phi^{(k)}(x) = O(e^{-\varepsilon_0 x/h})$ as $x \rightarrow \infty$. Hence ϕ belongs to the Schwartz class, namely $T_h^{-1} \mathcal{D}(\mathbb{R}; X') \subset \mathcal{S}(\mathbb{R}; X')$.

Before going to the proof of Theorem 17, we need to derive further regularity estimates described below.

Lemma 25 Let $T > 0$ be given. Then, for each $h > 0$ and each $\psi \in \mathcal{D}(0, T; X')$, the function $\phi_h := T_h^{-1} \psi$ satisfies the following estimates:

(i) $\phi_h = 0$ on $(-\infty, 0]$, $\phi_h \in L^\infty(0, \infty; X')$ and

$$\|\phi_h\|_{L^\infty(0, \infty; X')} \leq \left(1 + \frac{M_0}{\varepsilon_0}\right) \|\psi\|_{L^\infty(0, T; X')};$$

(ii) for each $p \in [1, \infty)$ $\phi_h \in L^p(0, \infty; X')$ and

$$\begin{aligned} & \|\phi_h\|_{L^p(0, \infty; X')} \\ & \leq \left(1 + \frac{M_0}{\varepsilon_0}\right) \|\psi\|_{L^p(0, T; X')} \leq T^{\frac{1}{p}} \left(1 + \frac{M_0}{\varepsilon_0}\right) \|\psi\|_{L^\infty(0, T; X')}; \end{aligned}$$

(iii) for each $\alpha \in (0, 1)$ and $p \in [1, \infty)$ one has

$$[\phi_h]_{B_\infty^{\alpha, p}(0, \infty; X')} \leq \left(1 + \frac{M_0}{\varepsilon_0}\right) [\psi]_{B_\infty^{\alpha, p}(\mathbb{R}, X')}.$$

Proof. Recall that ϕ_h is defined by

$$\phi_h = T_h^{-1}\psi = \psi + K_h * \psi,$$

wherein $*$ denotes the convolution product. Next one has

$$\|\phi_h\|_{L^\infty(0, \infty; X')} \leq (1 + \|K_h\|_{L^1(\mathbb{R})}) \|\psi\|_{L^\infty(0, T; X')} \leq \left(1 + \frac{M_0}{\varepsilon_0}\right) \|\psi\|_{L^\infty(0, T; X')}.$$

This completes the proof of (i).

Next, for $p \in [1, \infty)$, one also has

$$\begin{aligned} & \|\phi_h\|_{L^p(0, \infty; X')} \\ & \leq (1 + \|K_h\|_{L^1(\mathbb{R})}) \|\psi\|_{L^p(0, T; X')} \leq T^{\frac{1}{p}} \left(1 + \frac{M_0}{\varepsilon_0}\right) \|\psi\|_{L^\infty(0, T; X')}, \end{aligned}$$

that proves (ii).

Finally observe that for each $s > 0$ one has

$$\begin{aligned} & \|\phi_h(\cdot + s) - \phi_h\|_{L^p(0, \infty; X')} \\ & \leq \|\psi(\cdot + s) - \psi\|_{L^p(\mathbb{R}; X')} + \|K_h * (\psi(\cdot + s) - \psi)\|_{L^p(\mathbb{R}; X')} \\ & \leq \left(1 + \frac{M_0}{\varepsilon_0}\right) \|\psi(\cdot + s) - \psi\|_{L^p(\mathbb{R}; X')}, \end{aligned}$$

and this completes the proof of (iii). ■

We are now in position to prove Theorem 17.

Proof of Theorem 17. Let $T > 0$ and $\psi \in \mathcal{D}(0, T; X')$ be given. Then for each $h > 0$ set $\phi_h := T_h^{-1}\psi \in \mathcal{S}(\mathbb{R}; X')$ (see Remark 24).

On the one hand, we infer from Lemma 19, 20 and 25 that, for each $h > 0$, one has

$$[\phi_h]_{B_\infty^{\alpha, 1}(0, \infty; X')} \leq \frac{2}{\alpha} \left(1 + \frac{M_0}{\varepsilon_0}\right) \left[[\psi]_{W^{\alpha, 1}(0, T; X')} + \frac{4T^{1-\alpha}}{\alpha(1-\alpha)} \|\psi\|_{L^\infty(0, T; X')} \right].$$

Next, recalling that $\phi_h \in \mathcal{S}(\mathbb{R}, X')$ and $T_h \partial_t = \partial_t T_h$, applying $\phi = \phi_h$ into (25) ensues that there exists some constant $\widehat{M} > 0$ (depending upon T but independent from ψ and h) such that for all $h > 0$ and any $\psi \in \mathcal{D}(0, T; X')$ one has

$$|\langle \partial_t \psi, u^h \rangle_{\mathcal{S}, \mathcal{S}'}| \leq \widehat{M} [\|\psi\|_{L^\infty(0, T; X')} + \|\psi\|_{W^{\alpha, 1}(0, T; X')}].$$

Now let $s \in [\alpha, 1)$ and $p \in (1, \infty)$ be given such that $ps > 1$. Then let us recall that (see for instance Simon in [25]) that the following continuous embedding holds true:

$$W^{s, p}(0, T; X') \hookrightarrow L^\infty(0, T; X') \cap W^{\alpha, 1}(0, T; X').$$

Hence there exists some positive constant, still denoted by \widehat{M} , depending on T such that for all $h > 0$ and $\psi \in \mathcal{D}(0, T; X')$ one has

$$|\langle \partial_t \psi, u^h \rangle_{\mathcal{S}, \mathcal{S}'}| \leq \widehat{M} \|\psi\|_{W^{s, p}(0, T; X')}.$$

As a consequence $\partial_t u^h$, the distribution derivative of u^h , satisfies

$$\partial_t u^h \in (W_0^{s, p}(0, T; X'))' = W^{-s, p'}(0, T; X)$$

and

$$\|\partial_t u^h\|_{W^{-s, p'}(0, T; X)} \leq \widehat{M}, \quad \forall h > 0.$$

Hence, for each $T > 0$ there exists some constant $M = M(T)$ such that for all $h > 0$ one has

$$u^h \in W^{1-s, p'}(0, T; X) \text{ and } [u^h]_{W^{1-s, p'}(0, T; X)} \leq M(T).$$

This completes the proof of Theorem 17. ■

4.2 Proof of Theorem 16

In this section we complete the proof of Theorem 16. To that aim we apply the abstract result derived above in Theorem 17.

Our proof will be split into two steps. In the first step we derive a fractional Sobolev regularity estimate of $\{p^\lambda\}_{\lambda > 0}$ using Theorem 17. In the second step we bootstrap this estimates to conclude the proof of Theorem 16.

In a first step we derive the following lemma.

Lemma 26 *For each $\sigma > 2 + \frac{N}{2}$, each $s \in [\frac{1}{2}, 1)$ and each $p > \frac{1}{s}$, the family of maps $\{t \mapsto p^\lambda(t, dx)\}_{\lambda \geq 1}$ is bounded in $W_{\text{loc}}^{1-s, p'}(0, \infty; H^{-\sigma}(\mathbb{R}^N))$. This means that for each $T > 0$ there exists some constant $M = M(\sigma, T, s, p) > 0$ such that*

$$\|p^\lambda\|_{W^{1-s, p'}; H^{-\sigma}(\mathbb{R}^N)} \leq M, \quad \forall \lambda \geq 1.$$

Proof. Fix $\sigma > 2 + \frac{N}{2}$ so that the following continuous embedding holds true:

$$H^\sigma(\mathbb{R}^N) \hookrightarrow W^{2,\infty}(\mathbb{R}^N). \quad (30)$$

Now observe that the above continuous embedding ensures that for each $\lambda > 0$ and each $\phi \in H^\sigma(\mathbb{R}^N)$, the maps $t \mapsto \int_{\mathbb{R}^N} \phi(x) p^\lambda(t, dx)$ is measurable. Hence, since $H^{-\sigma}(\mathbb{R}^N) = (H^\sigma(\mathbb{R}^N))'$ is separable then the map $t \mapsto p^\lambda(t, dx)$ is strongly measurable from $[0, \infty)$ into $H^{-\sigma}(\mathbb{R}^N)$. Moreover, there exists some constant $C = C(\sigma) > 0$ such that for any $t \geq 0$, $\lambda > 0$ and $\phi \in H^\sigma(\mathbb{R}^N)$:

$$\left| \int_{\mathbb{R}^N} \phi(x) p^\lambda(t, dx) \right| \leq \|\phi\|_{L^\infty(\mathbb{R}^N)} \leq C(\sigma) \|\phi\|_{H^\sigma(\mathbb{R}^N)}.$$

Hence, one first conclude that the family of measures $\{p^\lambda\}_{\lambda > 0}$ is bounded in $L^\infty(0, \infty; H^{-\sigma}(\mathbb{R}^N))$, namely

$$\|p^\lambda\|_{L^\infty(0, \infty; H^{-\sigma}(\mathbb{R}^N))} \leq C(\sigma), \quad \forall \lambda > 0. \quad (31)$$

To go further, recall that p^λ satisfies (23). Hence we will apply Theorem 17 to this formulation. To that aim we estimate the different terms arising in the right hand side of (23).

Estimate for \mathcal{T}_1^λ :

Recall that \mathcal{T}_1^λ is defined in Lemma 14. Note that, due to (20), \mathcal{T}_1^λ re-writes as follows, for all $\phi \in \mathcal{S}(\mathbb{R}^+; H^\sigma(\mathbb{R}^N))$,

$$\begin{aligned} \mathcal{T}_1^\lambda[\phi] &= I_1^\lambda[\phi] - \lambda \int_0^\infty \int_{\mathbb{R}^N} \nabla \phi(t, x) \mathbf{K} p^\lambda(t, dx) dt \\ &\quad + \lambda \int_0^\infty \int_S \alpha_\infty(\theta) [z + \theta \mathbf{K}] \left[\int_{\mathbb{R}^N} \nabla \phi(t, x) p^\lambda \left(t + \frac{\theta}{\lambda^2}, dx \right) \right] Q(d\theta, dz) dt. \end{aligned}$$

Herein $I_1^\lambda[\phi]$ is a remainder that satisfies that there exists some constant $M > 0$ such that for all $\lambda > 0$ and $\phi \in \mathcal{S}(\mathbb{R}^+; H^\sigma(\mathbb{R}^N))$ one has

$$|I_1^\lambda[\phi]| \leq \frac{M}{\lambda} \|\phi\|_{L^1(0, \infty; W^{1, \infty}(\mathbb{R}^N))} \leq \frac{M}{\lambda} \|\phi\|_{L^1(0, \infty; H^\sigma(\mathbb{R}^N))}.$$

Next we re-write \mathcal{T}_1^λ as follows:

$$\begin{aligned} \mathcal{T}_1^\lambda[\phi] &= I^\lambda[\phi] - \lambda \int_0^\infty \int_{\mathbb{R}^N} \nabla \phi(t, x) \mathbf{K} p^\lambda(t, dx) dt \\ &\quad + \lambda \int_0^\infty \int_S \alpha_\infty(\theta) [z + \theta \mathbf{K}] \left[\int_{\mathbb{R}^N} \nabla \phi(s - \frac{\theta}{\lambda^2}, x) p^\lambda(s, dx) \right] Q(d\theta, dz) ds \\ &\quad + I_2^\lambda[\phi] \end{aligned}$$

with

$$I_2^\lambda[\phi] = \lambda \int_S \int_0^{-\frac{\theta}{\lambda^2}} \alpha_\infty(\theta) [z + \theta \mathbf{K}] \left[\int_{\mathbb{R}^N} \nabla \phi(t, x) p^\lambda \left(t + \frac{\theta}{\lambda^2}, dx \right) \right] dt Q(d\theta, dz).$$

And note that, as for I_1^λ , there exists $M > 0$ such that for all $\lambda > 0$ and $\phi \in \mathcal{S}(\mathbb{R}^+; H^\sigma(\mathbb{R}^N))$ one has

$$|I_2^\lambda[\phi]| \leq \frac{M}{\lambda} \|\phi\|_{L^1(0,\infty;W^{1,\infty}(\mathbb{R}^N))} \leq \frac{M}{\lambda} \|\phi\|_{L^1(0,\infty;H^\sigma(\mathbb{R}^N))}.$$

Next recalling the identity for \mathbf{K} in Remark 6, \mathcal{T}_1^λ re-writes, for any $\phi \in \mathcal{S}(\mathbb{R}^+; H^\sigma(\mathbb{R}^N))$ as

$$\begin{aligned} \mathcal{T}_1^\lambda[\phi] &= I^\lambda[\phi] + I_2^\lambda[\phi] + I_3^\lambda[\phi] \text{ with} \\ I_3^\lambda[\phi] &= \lambda \int_0^\infty \int_S \alpha_\infty(\theta) [z + \theta \mathbf{K}] \\ &\quad \left[\int_{\mathbb{R}^N} \left(\nabla \phi(s - \frac{\theta}{\lambda^2}, x) - \nabla \phi(s, x) \right) p^\lambda(s, dx) \right] Q(d\theta, dz) ds. \end{aligned}$$

Hence, one obtains

$$\begin{aligned} |I_3^\lambda[\phi]| &\leq \int_S \sqrt{-\theta} \alpha_\infty(\theta) [|z| + |\theta| |\mathbf{K}|] Q(d\theta, dz) \\ &\quad \sup_{h>0} h^{-\frac{1}{2}} \|\phi(\cdot + h, \cdot) - \phi\|_{L^1(0,\infty;W^{1,\infty}(\mathbb{R}^N))}. \end{aligned}$$

As consequence, we obtains that there exists some constant, still denoted by $M > 0$, such that for any $\lambda > 0$ and $\phi \in \mathcal{S}(\mathbb{R}^+; H^\sigma(\mathbb{R}^N))$

$$|\mathcal{T}_1^\lambda[\phi]| \leq M \left[\lambda^{-1} \|\phi\|_{L^1(0,\infty;H^\sigma(\mathbb{R}^N))} + \|\phi\|_{B_\infty^{\frac{1}{2},1}(0,\infty;H^\sigma(\mathbb{R}^N))} \right].$$

Estimate for \mathcal{T}_2^λ :

Recall that \mathcal{T}_2^λ is defined in Lemma 14. Hence it directly follows from this definition that there exists some constant $C > 0$ such that, for all $\phi \in \mathcal{S}(\mathbb{R}^+; H^\sigma(\mathbb{R}^N))$ and $\lambda > 0$, one has

$$|\mathcal{T}_2^\lambda[\phi]| \leq C \|\phi\|_{L^1(0,\infty;W^{2,\infty}(\mathbb{R}^N))} \leq C \|\phi\|_{L^1(0,\infty;H^\sigma(\mathbb{R}^N))}.$$

Estimate for \mathcal{R}^λ :

Recalling Assumption 1 (i), it readily follows that there exists some constant $C > 0$ such that for all $\lambda > 0$, $\phi \in \mathcal{S}(\mathbb{R}^+; H^\sigma(\mathbb{R}^N))$ one has

$$|\mathcal{R}^\lambda[\phi]| \leq C \|\phi\|_{L^\infty(\mathbb{R}^+ \times \mathbb{R}^N)} \leq C \|\phi\|_{L^\infty(0,\infty;H^\sigma(\mathbb{R}^N))}.$$

Conclusion:

As a consequence of the above estimates, there exists some constant $C > 0$ such that for all $\lambda \geq 1$ and $\phi \in \mathcal{S}(\mathbb{R}^+; H^\sigma(\mathbb{R}^N))$ one has:

$$\begin{aligned} &\left| \left\langle T_{\frac{1}{\lambda^2}} \partial_t \phi, p^\lambda \right\rangle \right| \\ &\leq C \left[\|\phi\|_{L^1(0,\infty;H^\sigma(\mathbb{R}^N))} + \|\phi\|_{B_\infty^{\frac{1}{2},1}(0,\infty;H^\sigma(\mathbb{R}^N))} + \|\phi\|_{L^\infty(0,\infty;H^\sigma(\mathbb{R}^N))} \right]. \end{aligned}$$

Now since $H^\sigma(\mathbb{R}^N) = (H^{-\sigma}(\mathbb{R}^N))'$ is a separable reflexive Banach space, Theorem 17 applies and ensures that for each $T > 0$ there exists $M_T > 0$ such that

$$[p^\lambda]_{W^{1-s,p'}(0,T;H^{-\sigma}(\mathbb{R}^N))} \leq M_T, \forall \lambda \geq 1.$$

Finally the uniform bound (31) and the above semi-norm estimate completes the proof of Lemma 26. ■

Before going to the proof of Theorem 16 we will first prove the following regularity lemma

Lemma 27 *Fix $\sigma > 2 + \frac{N}{2}$. Let $0 < \varepsilon < T$ and $p > 2$ be given. Then there exists $M > 0$ such that for any $\psi \in \mathcal{D}(\varepsilon, T; H^\sigma(\mathbb{R}^N))$ and all λ large enough the following estimate holds true*

$$\left| \int_0^\infty \int_{\mathbb{R}^N} \partial_t \psi(t, x) p^\lambda(t, dx) dt \right| \leq M \|\psi\|_{L^p(\varepsilon, T; H^\sigma(\mathbb{R}^N))}.$$

Proof. Let $\psi \in \mathcal{D}(\varepsilon, T; H^\sigma(\mathbb{R}^N))$ be given. Let $\phi_\lambda \in \mathcal{S}(\mathbb{R}^+; H^\sigma(\mathbb{R}^N))$ be the function defined by

$$T_{\frac{1}{\lambda^2}} \phi_\lambda = \psi, \forall \lambda > 0.$$

Here $T_{\frac{1}{\lambda^2}}$ is the operator defined in (24) with $h = 1/\lambda^2$. Next note that, due to Lemma 22, for all $t \in \mathbb{R}^+$ one has

$$\phi_\lambda(t, \cdot) = \psi(t, \cdot) + \int_0^\infty K_{1/\lambda^2}(s) \psi(t-s, \cdot) ds. \quad (32)$$

Hence $\phi_\lambda(t, \cdot) = 0$ for all $t \in [0, \varepsilon]$.

Next using the same notations as above one obtains due to (32) that the following decomposition holds true:

$$\int_{\mathbb{R}^+} \int_{\mathbb{R}^N} \partial_t \psi p^\lambda(t, dx) dt = -\mathcal{T}_1^\lambda[\phi_\lambda] - \mathcal{T}_2^\lambda[\phi_\lambda] + \mathcal{R}_1^\lambda[\phi_\lambda]. \quad (33)$$

Next we shall estimate each of these terms. We start by the second one by observing that due to the above computations and Lemma 25 (ii), there exists some constant $M > 0$ such that

$$|\mathcal{T}_2^\lambda[\phi_\lambda]| \leq M_2 \|\psi\|_{L^1(\varepsilon, T; H^\sigma(\mathbb{R}^N))}, \forall \lambda > 0.$$

We now estimate the third term arising in (33). To that aim we first choose $\lambda > \frac{1}{\varepsilon}$ so that, recalling that $\phi_\lambda(t, \cdot) = 0$ for $t \in [0, \lambda^{-1}] \subset [0, \varepsilon]$, $\mathcal{R}_1^\lambda[\phi_\lambda]$ re-writes

$$\begin{aligned} \mathcal{R}_1^\lambda[\phi_\lambda] = & \lambda^2 \int_\varepsilon^\infty \int_S [\alpha(\lambda^2 t, \theta) - \alpha_\infty(\theta)] \\ & \int_{\mathbb{R}^N} \phi_\lambda(t, x) \left[p^\lambda(t, dx) - p^\lambda\left(t + \frac{\theta}{\lambda^2}, dx\right) \right] Q(d\theta, dz) dt; \end{aligned}$$

Recalling (1) (i), there exists $M > 0$ such that for all $\lambda > 1/\varepsilon$ one has

$$|\mathcal{R}_1^\lambda[\phi_\lambda]| \leq \lambda^2 M e^{-\beta\varepsilon\lambda^2} \|\phi_\lambda\|_{L^1(\varepsilon, T; H^\sigma(\mathbb{R}^N))} \leq \lambda^2 M e^{-\beta\varepsilon\lambda^2} \|\psi\|_{L^1(\varepsilon, T; H^\sigma(\mathbb{R}^N))}.$$

It remains to estimate the first term in (33), namely $\mathcal{T}_1^\lambda[\phi_\lambda]$. To that aim we will make use of the estimate provided in Lemma 26 above. First note that coupling the estimate in this lemma together with those in Lemma 19, for all $T' > 0$ and $p > 2$ there exists some constant $M = M(T', p) > 0$ such that for any $\lambda \geq 1$ one has

$$[p^\lambda]_{B_{\infty}^{\frac{1}{2}, p'}(0, T'; H^{-\sigma}(\mathbb{R}^N))} \leq M.$$

Using this bound and similar computations as the ones given in the above section, for each $p > 2$ there exists some constant $M > 0$ such that for all $\lambda > 1/\varepsilon$ and any $\psi \in \mathcal{D}(\varepsilon, T; H^\sigma(\mathbb{R}^N))$ one has

$$|\mathcal{T}_1^\lambda[\phi_\lambda]| \leq M \|\psi\|_{L^p(\varepsilon, T; H^\sigma(\mathbb{R}^N))}.$$

Finally, coupling all the above inequalities together with Hölder inequality completes the proof of the lemma. ■

Using the two above lemmas, namely Lemma 26 and Lemma 27 we are in position to conclude the proof of Theorem 16.

Proof of Theorem 16. Note that for any given $R > 0$ and $\sigma > \frac{N}{2} + 2$ then embedding $W_0^{\sigma+1, 2}(BR) \hookrightarrow H^\sigma(R^N)$ is compact. Hence the dual continuous embedding $H^{-\sigma}(R^N) \hookrightarrow H^{-\sigma-1}(BR)$ is also compact. In addition we deduce from the above lemmas that for each $0 < \varepsilon < T$ and each $p > 2$, there exists $\tilde{\lambda} > 0$ large enough such that the family $\{p^\lambda\}_{\lambda > \tilde{\lambda}}$ is bounded in $L^\infty(\varepsilon, T; H^{-\sigma}(R^N))$ while the family $\partial_t p^\lambda_{\lambda > \tilde{\lambda}}$ is bounded in $L^p(\varepsilon, T; H^{-\sigma-1}(BR))$. Thus Aubin-Lions-Simon lemma (see [2, 19, 24]) applies and ensures that Theorem 16 holds true. ■

References

- [1] H. Amann, Compact embeddings for vector-valued Sobolev and Besov spaces, *Glas. Mat.* 35 (2000), 161–177.
- [2] J.-P. Aubin, Un théorème de compacité, *C. R. Acad. Sci. Paris* 256 (1963), 5042–5044.
- [3] P.W. Bates, On some nonlocal evolution equations arising in materials science, In: *Nonlinear dynamics and evolution equations* (Ed. by H. Brunner, X. Zhao and X. Zou), pp. 13-52, *Fields Inst. Commun.*, 48, AMS, Providence, 2006.
- [4] P. Billingsley, *Convergence of Probability Measures*, Wiley, 2nd ed., 1999.

- [5] C. Castaing, P. Raynaud de Fitte and M. Valadier, Young measures on topological spaces: With applications in control theory and probability theory. Mathematics and its Applications, 571. Kluwer Academic Publishers, Dordrecht, 2004. xii+320 pp.
- [6] E. Chasseigne, M. Chaves and J.D. Rossi, Asymptotic behavior for nonlocal diffusion equations, *J. Math. Pures Appl.* 86 (2006), 271–291.
- [7] C. Cosner, J. Dávila and S. Martínez, Evolutionary stability of ideal free nonlocal dispersal, *J. Bio. Dyn.*, 6 (2012), 395–405.
- [8] J. Coville, Remarks on the strong maximum principle for nonlocal operator, *Elec. J. Diff. Eqs.*, 68 (2008), 1–10.
- [9] J. Coville, L. Dupaigne, On a non-local reaction diffusion equation arising in population dynamics, *Proc. Math. Roy. Soc. of Edinburgh Sect. A* 137 (2007), 1–29.
- [10] P. Cembranos and J. Mendoza, Banach Spaces of Vector-Valued Functions, vol. 1676, Springer-Verlag, Berlin, 1997.
- [11] O. Diekmann, S.A. van Gils, S.M. Verduyn Lunel, H.-O. Walther, Delay Equations, Function-, Complex-, and Nonlinear Analysis, Springer-Verlag, New York, 1995.
- [12] P. Fife, Some nonclassical trends in parabolic and parabolic-like evolutions. Trends in nonlinear analysis, 153–191, Springer, Berlin, 2003.
- [13] P. Fife and X. Wang, A convolution model for interfacial motion: the generation and propagation of internal layers in higher space dimensions, *Adv. Differential Equations*, 3 (1998), 85–110.
- [14] J.K. Hale, S.M. Verduyn Lunel, Introduction to Functional Differential Equations, Springer-Verlag, New York, 1993.
- [15] L.I. Ignat, J.D. Rossi, A nonlocal convection-diffusion equation, *J. Funct. Anal.*, 251 (2007), 399–437.
- [16] Y. Jin, X.-Q. Zhao, Spatial dynamics of a periodic population model with dispersal, *Nonlinearity*, 22 (2009), 1167–1189.
- [17] T. Laurent, B. Rider, M. Reed, Parabolic behaviour of a hyperbolic delay equation, *SIAM Journal Math. Anal.* 38 (2006), 1–15.
- [18] W.T. Li, J.B. Wang, X.-Q. Zhao, Spatial dynamics of a nonlocal dispersal population model in a shifting environment, *J. Nonlinear Sci.* <https://doi.org/10.1007/s00332-018-9445-2>
- [19] J.L. Lions, Quelques méthodes de résolution des problèmes aux limites non linéaires, 1969, Paris: Dunod-Gauth. Vill.

- [20] Z. Liu, P. Magal and S. Ruan, Projectors on the generalized eigenspaces for functional differential equations using integrated semigroups, *J. Diff. Eqs*, 244 (2008), 1784–1809.
- [21] F. Lutscher, E. Pachepsky, and M. A. Lewis, The effect of dispersal patterns on stream populations, *SIAM J. Appl. Math.*, 65 (2005), 1305–1327.
- [22] M.M. Meerschaert and P. Straka. Semi-Markov approach to continuous time random walk limit processes, *The Annals of Probability*, 42(4) (2014), 1699–1723.
- [23] D. Majerek, W. Nowak and W. Zieba. Conditional strong law of large number. *Int. J. Pure Appl. Math*, 20(2) (2005), 143–156.
- [24] J. Simon, Compact sets in the space $L^p(O, T; B)$, *Annali di Matematica Pura ed Applicata* 146 (1986), 65–96.
- [25] J. Simon, Sobolev, Besov and Nikolskii fractional spaces: Imbeddings and comparisons for vector valued spaces on an interval, *Ann. Math. Pura Appl.* 157 (1990), 117–148.
- [26] D. W. Stroock and S. S. Varadhan, S. S. (2007). *Multidimensional diffusion processes*. Springer.
- [27] M. Suwa and K. Yoshino, A characterization of tempered distributions with support in a cone by the heat kernel method and its applications, *J. Math. Sci. Univ. Tokyo*, 11 (2004), 75–90.
- [28] H.R. Thieme, Semiflows generated by Lipschitz perturbations of non-densely defined operators, *Differential Integral Equations* 3 (1990), 1035–1066.
- [29] G.F. Webb, *Theory of Nonlinear Age-Dependent Population Dynamics*, Marcel Dekker, New York, 1985.
- [30] M. Valadier, Young measures. In *Methods of Nonconvex Analysis*, A. Cellina Ed., *Lecture Notes in Math.* 1446, Springer Verlag, Berlin 1990, 152–188.

A Various notions of topology on the space of measures

In this appendix we recall different notions of topology that have been used in this work. We refer for instance to the textbook of Billingsley [4] for more details and results on this topic.

Here we denote by \mathcal{M} the set of bounded Borel – signed – measures on \mathbb{R}^N . We also denote by \mathcal{M}^+ and \mathcal{M}^- the set of positive and negative Borel measures respectively.

Total variation norm on \mathcal{M} :

The space \mathcal{M} can be firstly endowed with the usual total variation norm, denoted by $\|\cdot\|_{TV}$ and defined through the Hahn-Jordan decomposition of a signed measure $\mu \in \mathcal{M}$ as

$$\mu = \mu^+ - \mu^- \text{ with } \mu^\pm \in \mathcal{M}^+,$$

and $\|\mu\|_{TV} = \mu^+(\mathbb{R}^N) + \mu^-(\mathbb{R}^N)$. Hence \mathcal{M} endowed with the total variation norm becomes a Banach space.

This strong topology has not been used in this work because translation are usually not continuous for this norm topology. Indeed if $\mathbf{q} \in \mathbb{R}^N \setminus \{0\}$ is given, the map $\theta \in [-1, 0] \mapsto \delta_{\mathbf{q}\theta} \in \mathcal{P}$ is not continuous with respect to this norm topology.

Narrow topology on \mathcal{M} :

The space \mathcal{M} can be endowed with the so-called narrow topology which is defined as the weakest topology on \mathcal{M} such that for each test function $f \in \mathcal{C}_b(\mathbb{R}^N)$ the functional

$$\mu \in \mathcal{M} \mapsto \int_{\mathbb{R}^N} f d\mu \in \mathbb{R},$$

is continuous. The topological space \mathcal{M} endowed with the narrow topology will be denoted by $\mathcal{M}_{\text{narr}}$.

Note that for this topology the map $\theta \mapsto \delta_{\mathbf{q}\theta}$ is continuous.

Let us also recall the so-called Portemanteau theorem:

Theorem A.1 (Portemanteau Theorem) *Let $\{\mu_n\}_{n \geq 0} \subset \mathcal{M}$ be a given sequence and $\mu \in \mathcal{M}$ be given. Then the following properties are equivalent:*

- (i) $\mu_n \rightarrow \mu$ in $\mathcal{M}_{\text{narr}}$;
- (ii) $\int_{\mathbb{R}^N} f d\mu_n \rightarrow \int_{\mathbb{R}^N} f d\mu$ for all Lipschitz continuous function on \mathbb{R}^N ;
- (iii) $\limsup_{n \rightarrow \infty} \int_{\mathbb{R}^N} f d\mu_n \leq \int_{\mathbb{R}^N} f d\mu$ for every upper semi-continuous function f on \mathbb{R}^N bounded from above;
- (iv) $\liminf_{n \rightarrow \infty} \int_{\mathbb{R}^N} f d\mu_n \geq \int_{\mathbb{R}^N} f d\mu$ for every lower semi-continuous function f on \mathbb{R}^N bounded from below;
- (v) $\lim_{n \rightarrow \infty} \mu_n(A) = \mu(A)$ for each $A \in \mathcal{B}(\mathbb{R}^N)$, the Borel sets of \mathbb{R}^N such that $\mu(\partial A) = 0$.

We can now focus on the subset $\mathcal{P} \subset \mathcal{M}$ of probability measures endowed with the induced narrow topology. The topological space $\mathcal{P}_{\text{narr}}$ is metrizable with the distance associated to the dual norm of $W^{1,\infty}(\mathbb{R}^N)$. Such a distance is called bounded Lipschitz distance, denoted by d_{BL} and it reads for each $\mu, \nu \in \mathcal{P}$ as

$$d_{BL}(\mu, \nu) = \|\mu - \nu\|_{W^{-1,\infty}(\mathbb{R}^N)} = \sup_{\|f\|_{W^{1,\infty}(\mathbb{R}^N)} \leq 1} \int_{\mathbb{R}^N} f d(\mu - \nu).$$

Vague topology or Weak* topology of \mathcal{M} :

The set \mathcal{M} can also be endowed with weaker topology. Denote by $\mathcal{C}_0(\mathbb{R}^N)$ the space of continuous functions on \mathbb{R}^N tending to 0 at infinity. Endowed with the usual sup norm, it becomes a separable Banach space. Then the Riesz representation theorem ensures the following dual representation

$$\mathcal{M} = (\mathcal{C}_0(\mathbb{R}^N))'.$$

Hence \mathcal{M} can be endowed with the weak topology, that is also called vague topology. Note that the close balls are compact due to Banach-Alaoglu theorem. It is also important to notice that since $\mathcal{C}_0(\mathbb{R}^N)$ is separable, the closed ball are metrizable and thus sequentially compact. In the sequel the topological space \mathcal{M} endowed with the vague convergence will be denoted by $\mathcal{M}_{\text{vague}}$.

Let us recall the following connexion between narrow and weak topology:

Lemma A.2 *Let $\{\mu_n\}_{n \geq 0} \subset \mathcal{M}^+$ be a given sequence and $\mu \in \mathcal{M}^+$ be given. Assume that*

$$\lim_{n \rightarrow \infty} \mu_n = \mu \text{ in } \mathcal{M}_{\text{vague}} \text{ and } \lim_{n \rightarrow \infty} \int_{\mathbb{R}^N} d\mu_n = \int_{\mathbb{R}^N} d\mu,$$

then $\mu_n \rightarrow \mu$ narrowly, namely in $\mathcal{M}_{\text{narr}}$.

Young measures: The notion of Young measure has also been used in this work. Here we recall some basic facts and properties and we refer the reader to [5, 30] and the references cited therein for more details on Young measures theory.

First we say that a function $m \equiv m_t : \mathbb{R}^+ \rightarrow \mathcal{M}$ is weakly measurable if for each test function $\varphi \in \mathcal{C}_0(\mathbb{R}^N)$ the map $t \mapsto \int_{\mathbb{R}^N} \varphi d m_t$ is measurable from \mathbb{R}^+ to \mathbb{R} . This allows to defined the vector space $L_{\omega^*}^\infty(\mathbb{R}^+; \mathcal{M})$ as the set of weakly measurable map from \mathbb{R}^+ into \mathcal{M} and that is essentially bounded. The elements of $L_{\omega^*}^\infty(\mathbb{R}^+; \mathcal{P})$ are called Young measures.

Now let us recall (see for instance [10]) the following duality representation

$$L_{\omega^*}^\infty(\mathbb{R}^+; \mathcal{M}) = (L^1(\mathbb{R}^+; \mathcal{C}_0(\mathbb{R}^N)))'.$$

Recall also that the Banach space $L^1(\mathbb{R}^+; \mathcal{C}_0(\mathbb{R}^N))$ is separable. Therefore Banach-Alaoglu theorem implies that bounded set in $L_{\omega^*}^\infty(\mathbb{R}^+; \mathcal{M})$ are relatively (sequentially) compact with respect to the weak topology. This leads to the following fundamental compactness lemma for Young measures.

Lemma A.3 *Let $\{m^k\}_{k \geq 0}$ be a sequence of Young measures, that is elements of $L_{\omega^*}^\infty(\mathbb{R}^+; \mathcal{P})$. Then there exists a subsequence $\{k_n\}_{n \geq 0}$ and a map of measures $m^\infty \equiv m_t^\infty \in L_{\omega^*}^\infty(\mathbb{R}^+; \mathcal{M}^+)$ such that*

$$\int_{\mathbb{R}^N} d m_t^\infty \leq 1 \text{ a.e. } t \in \mathbb{R}^+,$$

and such that for all $f \in L^1(\mathbb{R}^+; \mathcal{C}_0(\mathbb{R}^N))$ one has

$$\lim_{n \rightarrow \infty} \int_{\mathbb{R}^+ \times \mathbb{R}^N} f(t, x) m_t^{k_n}(dx) dt = \int_{\mathbb{R}^+ \times \mathbb{R}^N} f(t, x) m_t^\infty(dx) dt. \quad (34)$$

Similarly to Lemma A.2, one has the following result

Lemma A.4 (Tightness Lemma) *With the same notations as in Lemma A.3, if the limit measure satisfies the tightness condition*

$$\int_{\mathbb{R}^N} dm_t^\infty = 1 \text{ a.e. } t \in \mathbb{R}^+,$$

then the above convergence, namely (34) holds for the so-called narrow topology of Young measures, that reads as for all test function $f \in L^1(\mathbb{R}^+; \mathcal{C}_b(\mathbb{R}^N))$ one has

$$\lim_{n \rightarrow \infty} \int_{\mathbb{R}^+ \times \mathbb{R}^N} f(t, x) m_t^{k_n}(dx) dt = \int_{\mathbb{R}^+ \times \mathbb{R}^N} f(t, x) m_t^\infty(dx) dt.$$

B A linear delay differential equation

In this appendix we come back to the linear delay differential (8) and we will prove that

$$\alpha(t, \theta) \rightarrow a_\infty(\theta) \text{ as } t \rightarrow \infty,$$

uniformly for $\theta \in [-1, 0]$ and with an exponential rate, that re-writes as $\frac{y(t+\theta)}{y(t)} \rightarrow e^{\gamma\theta}$ as $t \rightarrow \infty$ uniformly with respect to $\theta \in [-1, 0]$ and exponentially fast.

To that aim, we consider the equation

$$\begin{aligned} y'(t) &= \int_{[-1,0]} y(t+\theta) K(d\theta), \quad t > 0, \\ y(\theta) &= y^0(\theta), \quad \forall \theta \in [-1, 0]. \end{aligned} \tag{35}$$

wherein $K \in \mathcal{M}^+([-1, 0])$ denotes a bounded positive Borel measure on $[-1, 0]$ with $K([-1, 0]) > 0$ and $y^0 \in \mathcal{C}([-1, 0]; \mathbb{R}^+)$ with $y^0(0) > 0$ so that $y(t) > 0$ for all $t > 1$.

Next to investigate the large time behaviour of the above equation let us introduce the history function $u \equiv u(t, \theta)$ defined by

$$u(t, \theta) = y(t + \theta), \quad t \geq 0, \quad \theta \in [-1, 0].$$

This function formally satisfies the following Cauchy problem:

$$\begin{cases} \partial_t u - \partial_\theta u = 0, \quad t > 0, \quad \theta \in [-1, 0], \\ \partial_\theta u(t, 0) = \int_{[-1,0]} u(t, \theta) K(d\theta), \\ u(0, \cdot) = y^0 \in \mathcal{C}([-1, 0]). \end{cases} \tag{36}$$

Such a linear functional differential problem has been extensively studied in the literature. We refer for instance to [20, 28], the monographs [11, 14] and the references cited therein.

Here to be more precise we consider the Banach space $Y = \mathbb{R} \times \mathcal{C}([-1, 0])$ and

$Y_0 = \{0\} \times \mathcal{C}([-1, 0])$ as well as the linear operator $A : D(A) \subset Y \rightarrow Y$ defined by

$$D(A) = \{0\} \times \mathcal{C}^1([-1, 0]), \quad A \begin{pmatrix} 0 \\ \varphi \end{pmatrix} = \begin{pmatrix} -\varphi'(0) + \int_{[-1, 0]} \varphi(\theta) K(d\theta) \\ \varphi' \end{pmatrix}.$$

Then setting $U(t) = \begin{pmatrix} 0 \\ u(t, \cdot) \end{pmatrix}$, Problem (36) re-writes as the following abstract Cauchy problem

$$\frac{dU(t)}{dt} = AU(t), \quad t > 0 \text{ and } U(0) = \begin{pmatrix} 0 \\ y^0 \end{pmatrix} \in Y_0 = \overline{D(A)}.$$

Following [20] (see also the references therein) this problem generates a strongly continuous linear semigroup $\{T(t)\}_{t \geq 0}$ on Y_0 with infinitesimal generator A_0 , the part of A in Y_0 , defined as

$$D(A_0) = \{x \in D(A) : Ax \in Y_0\} \text{ and } A_0x = Ax \quad \forall x \in D(A_0).$$

In addition, the essential growth rate $\omega_{0,ess}(A_0)$ satisfies $\omega_{0,ess}(A_0) = -\infty$ so that, due to usual results for spectral theory (see for instance the monograph [29]), the spectrum of A_0 only consists in point spectrum and one has

$$\sigma(A_0) = \{z \in \mathbb{C} : \Delta(z) = 0\},$$

wherein the function $\Delta : \mathbb{C} \rightarrow \mathbb{C}$ is defined by

$$\Delta(z) = z - \int_{[-1, 0]} e^{\theta z} K(d\theta). \quad (37)$$

In addition, the growth rate $\omega_0(A_0)$ of the linear semigroup $\{T(t)\}_{t \geq 0}$ is obtained by

$$\omega_0(A_0) = \max\{\Re z : z \in \sigma(A_0)\}.$$

Let us now consider the unique $\gamma > 0$ solution of the equation $\Delta(z) = 0$. Next the following lemma holds true:

Lemma B.1 *Let $z \in \mathbb{C}$ be given such that $\Delta(z) = 0$. Then the following properties hold true:*

- (i) *One has $\Re(z) \leq \gamma$ and $\Re(z) = \gamma \implies z = \gamma$.*
- (ii) *If $\Re(z) \geq 0$ then $|\Im(z)| \leq K([-1, 0])$.*

The proof of (i) follows the same lines as the one of Lemma 21 while the proof of (ii) is straightforward.

As a direct corollary of the above lemma, one obtains that

$$\omega_0(A_0) = \gamma,$$

and there exists $\varepsilon > 0$ such that for all $z \in \mathbb{C}$:

$$\Delta(z) = 0 \text{ and } \Re(z) \geq \gamma - \varepsilon \implies z = \gamma. \quad (38)$$

Moreover let us notice that

$$\Delta'(\gamma) = 1 + \int_{[-1,0]} (-\theta)e^{\theta\gamma} K(d\theta) \in [1, \infty).$$

This means that the dominant eigenvalue γ is simple.

And therefore, the usual spectral theory ensures that there exists $\alpha_1 > 0$ and $\beta > 0$ such that

$$y(t) = e^{\gamma t} (\alpha_1 + O(e^{-\beta t})) \text{ as } t \rightarrow \infty.$$

Hence we obtain

$$\frac{y(t+\theta)}{y(t)} \rightarrow e^{\gamma\theta} \text{ as } t \rightarrow \infty,$$

uniformly with respect to $\theta \in [-1, 0]$ and exponentially fast.