
An upscaled DGTD method for time-domain electromagnetics

A. Gobé¹, S. Lanteri¹, C. Scheid² and F. Valentin³

¹ Côte d'Azur University, Inria, CNRS, LJAD, France

² Côte d'Azur University, LJAD, CNRS, Inria, France

³ LNCC - National Laboratory for Scientific Computing, Petrópolis, Brazil

In this work, we address time-dependent electromagnetic wave propagation problems with strong multiscale features with application to nanophotonics, where problems usually involve complex multiscale geometries, heterogeneous materials, and intense, localized electromagnetic fields. Nanophotonics simulations require very fine meshes to incorporate the influence of geometries as well as high order polynomial interpolations to minimize dispersion. Our goal is to design a family of innovative high performance numerical methods perfectly adapted for the simulation of such multiscale problems. For that purpose we extend the Multiscale Hybrid-Mixed (MHM) finite element method, originally proposed for the Laplace problem in [1], to the solution of 2d and 3d transient Maxwell equations with heterogeneous media. The MHM method arises from the decomposition of the exact electromagnetic fields in terms of the solutions of locally independent Maxwell problems. Those problems are tied together with an one field formulation on top of a coarse mesh skeleton. The multiscale basis functions, which are responsible for upscaling, are also driven by local Maxwell problems. A high order Discontinuous Galerkin method (see [2]) in space combined with a second-order explicit leap-frog scheme in time discretizes the local problems. This makes the MHM method effective and parallelizable, and yields a staggered algorithm within a *divide-and-conquer* framework. In this talk we will present the general formulation of this MHM-DGTD method and present some preliminary numerical results in 2d.

REFERENCES

1. C. Harder, D. Paredes, and F. Valentin. A family of Multiscale Hybrid-Mixed finite element methods for the Darcy equation with rough coefficients. *J. Comput. Phys.*, 245:107–130, 2013.
2. S. Descombes, C. Durochat, S. Lanteri, L. Moya, C. Scheid, and J. Viquerat. Recent advances on a DGTD method for time-domain electromagnetics. *Photonics and Nanostructures - Fundamentals and Applications*, 11(4):291–302, 2013.

1. Title of the paper

High order curvilinear DGTD methods for local and nonlocal plasmonics

2. Name, affiliation, and email of each author

- Alexis Gobé and Stéphane Lanteri
Côte d'Azur University, Inria, CNRS, LJAD, France
- Claire Scheid
Côte d'Azur University, LJAD, CNRS, Inria, France
- Frédéric Valentin
LNCC - National Laboratory for Scientific Computing, Petrópolis, Brazil

3. Mailing address

Inria Sophia Antipolis-Méditerranée research center
2004 Route des Lucioles, BP 93
06902 Sophia Antipolis Cedex, France

4. Telephone number

+33-4-92-38-78-34

5. Corresponding author and presenting author

Stéphane Lanteri

6. Topic or session organizer

Multiscale and multiphysics computation and applications
Organized by Prof. Qing Huo Liu (Duke University)