Estimation of Leaf Nitrogen Content of Winter Wheat Based on Akaike's Information Criterion

Haojie Pei, Haikuan Feng, Fuqin Yang, Zhenhai Li, Guijun Yang, Qinglin Niu

► To cite this version:

Haojie Pei, Haikuan Feng, Fuqin Yang, Zhenhai Li, Guijun Yang, et al.. Estimation of Leaf Nitrogen Content of Winter Wheat Based on Akaike's Information Criterion. 11th International Conference on Computer and Computing Technologies in Agriculture (CCTA), Aug 2017, Jilin, China. pp.231-240, 10.1007/978-3-030-06179-1_24 . hal-02111567

HAL Id: hal-02111567

<https://hal.inria.fr/hal-02111567>

Submitted on 26 Apr 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Distributed under a Creative Commons Attribution| 4.0 International License

Estimation of Leaf Nitrogen Content of Winter Wheat based on Akaike's Information Criterion

Haojie Pei^{1,2,3,4}, Haikuan Feng^{1,2,3(✉)}, Fuqin Yang^{1,5}, Zhenhai Li^{1,2,3}, Guijun Yang^{1,2,3}, Qinglin Niu^{1,2,3,4}

¹Key Laboratory of Quantitative Remote Sensing in Agriculture of Ministry of Agriculture P. R. China, Beijing Research Center for Information Technology in Agriculture, Beijing 100097, China

{xmljphj, fenghaikuan123, yangfuqin0202, guijun.yang}@163.com,

lizh323@126.com

²National Engineering Research Center for Information Technology in Agriculture, Beijing, 100097, China

³Beijing Engineering Research Center for Agriculture Internet of Things, Beijing 100097, China

⁴School of Surveying and Land Information Engineering, Henan Polytechnic University, Jiaozuo 454000,

⁵College of Civil Engineering, Henan Institute of Engineering, Zhengzhou 451191, China

China

Abstract. Nitrogen is one of the important indices for evaluation of crop growth and output quality. At present, there are a lot of researches on the estimation of crop nitrogen content, but most of the studies do not consider whether the model established by vegetation index and crop nitrogen content is the best. The purpose of this study was to estimate the nitrogen content of wheat leaves and to establish a method and the optimal model for monitoring nitrogen content in wheat leaves. Spectral reflectance of leaves and concurrent leaf nitrogen content parameters of samples were acquired in during 2013 and 2014 wheat growth season, in Beijing Academy of Agriculture and Forestry Sciences. 17 vegetation indices related to nitrogen content were chosen, and the relationship between related vegetation indices and leaf nitrogen content were built for screening vegetation indices with variable importance projection (VIP). Choosing first 10 different vegetation indices after ranking with VIP value as the independent variable for estimating nitrogen content of leaf in wheat. And the number of vegetation indices was gradually increased from top 4 to 10. The leaf nitrogen content estimation model with different vegetation indices can be built using the integrated model of variable importance projection (VIP) - partial least squares (PLS). At the same time, Akaike's Information Criterion (AIC) value was calculated in different estimation model, and the AIC value of 7 different estimation model was compared. Then the optimal model with 5 vegetation indices was selected, which AIC value is the lowest. The optimal model was validated by leave one out cross-validation method. The result, (1) the comprehensive interpretation ability of the first 10 spectral indices on the nitrogen content of winter wheat leaves was PSSRc, GMI-2, SR705, RI-half, ZM, GMI-1, PSSRb, RI-3dB, VOGc and Clred edge. (2) The optimal model with 5 vegetation indices was selected from 7 models. (3) The decision coefficient (R^2) and root-mean-square error (RMSE) of the optimal model respectively were 0.73 and 0.33. The R^2 and RMSE of wheat by validating were 0.73 and 0.33, respectively. The study showed: (1) The VIP-PLS model has higher ability to estimate the nitrogen content of leaf in wheat, which laying an important foundation for improving the precision of forecasting winter wheat leaf nitrogen content with remote sensing. (2) The AIC method can be used to select the optimal model, and the selected model has the higher predictive ability. And the optimal estimation model of wheat LNC can be obtained based on AIC.

Keywords: leaf nitrogen content, Akaike's Information Criterion, Variable Importance for Projection, partial least squares, vegetation index

1 Introduction

Nitrogen is a "life element" in crop growth and development, rapid and accurate determination of nitrogen is conducive to the real-time management of nitrogen, providing a guarantee for high yield and quality of crops[1-2]. The use of spectral reflectance and spectral index for crop nitrogen estimation has been an important content of agricultural remote sensing research, and domestic and foreign scholars have done a lot of research [3-5]. In the jointing stage of wheat, artificial neural network, support vector regression and random forest were used to establish the estimation model of nitrogen accumulation in wheat leaves [6]. The absorption characteristic spectrum (590 ~ 756nm, 1096 ~ 1295nm, 1295 ~ 1642nm) were used to determine the characteristics of the depth and area of the leaves, and it can be a good total nitrogen content inversion [7]. A number of spectral indices were established using multi-spectral data of high score number satellite to estimate the nitrogen content of wheat leaves [8]. The new red edge bimodal parameters were constructed by using the ground spectrum, and the nitrogen content of wheat leaves was estimated by regression analysis [9]. The red edge index was used to estimate the nitrogen nutrition index [10]. The grey relational analysis was used to analyze the correlation of commonly used vegetation indices, and five indices of good vegetation indices were selected based on experience to participate in partial least squares regression (PLSR) [11,12]. The Manhattan, Euclidean measures and a variety of neural networks were combined into an optimal model to predict the leaf nitrogen content. The results showed that the accuracy of the algorithm could predict leaf nitrogen content to 92.1% [13].

However, most researches do not consider whether the model established between crop nitrogen content and vegetation index is optimal. In this study, the variable importance projection (VIP) is used to sort the selected vegetation indices, and the number of different vegetation indices was selected as the input variables according to the importance of the vegetation index. The number of input variables was gradually increased. Based on the AIC, the optimum nitrogen content of winter wheat leaves estimation model was selected, and the optimal nitrogen content estimation model was selected to estimate the nitrogen content of winter wheat leaf. The purpose of this study was to estimate the nitrogen content of wheat leaves, to establish a method and the optimal model for monitoring nitrogen content in wheat leaves.

2 Material and method

2.1 Overview of the study area

The experiment was conducted at the wheat experiment base in Beijing Academy of Agriculture and Forestry in 2013-2014. The experiment base was located in Haidian District, Beijing, 39 ° 56'N, 116 ° 16'E, the specific location shown in Figure 1. The district has four distinct seasons, cold and dry in winter, high temperature and rainy in summer, the average annual temperature of 12.5 °C. Wheat was sown on September 29, 2013. And the soil was tested. The content of organic matter in the soil was 1.54~2.16 g/kg; the available potassium content was 77.54~216.27 mg/kg, the content of nitrate nitrogen was 6.99~21.40mg/kg, the available phosphorus content was 29.92~136.44mg/kg, the Total nitrogen content is 0.09~0.12g/kg. The tested varieties were Jing 9428 and Nongda 211 respectively. Each plot size 1.2 × 1.5m, planting 33.75g, showing 6 lines, planting density of 6 million / hm². 40.5g

of urea per cell. 2 varieties, 5 water treatment, 4 repeat tests, water stress test from no water to water full saturation test, 0 m³/ha (W1), 225 m³/ha (W2), 450 m³/ha (W3), 675 m³/ha (W4), 900 m³/ha (W5), respectively. The test was conducted on April 20~21, 2014.

Fig.1 Beijing academy of agriculture and forestry sciences

2.2 Leaf spectrum determination

Leaf spectrum measurements were carried out using an American FieldSpec[®] Pro FR spectrometer (ASD Inc. Boulder Colorado, USA) coupled with a hand-held leaf blade spectroscopy detector (ASD leaf clip). The wavelength range is 350~2500 nm, and the interval is 1 nm. Blade holder detector built-in quartz halogenated lamp, guaranteeing that the light source is stable. The blade detector can cover the whole leaf chamber of the leaf blade and has the same detection area. It can eliminate the background reflection, the spectral fluctuation caused by the curved surface of the blade, and the internal variation caused by the leaf blade. Before the measurement of the leaf spectrum, the dust on the surface of the blade was wiped off with a clean, damp cloth, and corrected with a standard whiteboard to reduce the error.

2.3 Leaf nitrogen content determination

There are 40 wheat plots. Each plant is divided into 5 layers. 5 samples were taken per cell, and 5 layers of each plant were combined as samples of each layer, 199 samples in total. The samples were placed in a paper bag, placed in an oven at 105 °C for 30 minutes, and then each organ sample was dried at 80 °C for more than 48 hours until constant mass. Finally, the dried crop leaves were crushed. The nitrogen content (N, %) of the crop leaves was measured using a Kjeldahl nitrogen apparatus (Buchi B-339, Switzerland).

2.4 Method

2.4.1 Selection of Vegetation Index

According to the previous research results and the significant relationship between vegetation index and leaf nitrogen content, 19 vegetation indices related to leaf nitrogen content were selected.

Table 1 Spectral indices for leaf nitrogen content

Spectral indices	Definitions	references
Pigment –specific simple ratio-b	$PSSRb = R_{800} / R_{635}$	[14]
Pigment –specific simple ratio-c	$PSSRc = R_{800} / R_{470}$	[14]
Gitelson and Merzlyak index-1	$GMI - 1 = R_{750} / R_{550}$	[15]
Gitelson and Merzlyak index-2	$GMI - 2 = R_{750} / R_{700}$	[15]
Simple ratio ₇₀₅	$SR_{705} = R_{750} / R_{705}$	[15]
Vogelman index a	$VOG_a = R_{740} / R_{720}$	[16]
Zarco and Miller	$ZM = R_{750} / R_{710}$	[17]
Ratio index-half	$RI - half = R_{747} / R_{708}$	[18]
Ratio index-1dB	$RI - 1dB = R_{735} / R_{720}$	[18]
Ratio index-2dB	$RI - 2dB = R_{738} / R_{720}$	[18]
Ratio index-3dB	$RI - 3dB = R_{741} / R_{717}$	[18]
Simple ratio _(533,565)	$SR(533,565) = R_{533} / R_{565}$	[19]
Vogelman index c	$VOGc = \frac{(R_{734} - R_{747})}{(R_{715} + R_{720})}$	[17]
Normalized difference red edge	$NDRE = \frac{(R_{790} - R_{720})}{(R_{790} + R_{720})}$	[20]
Photochemical reflectance index	$PRI = \frac{(R_{570} - R_{531})}{(R_{570} + R_{531})}$	[21]
Red edge model	$CI_{rededge} = \frac{R_{750}}{R_{720}} - 1$	[22]
MCARI/MTVI2	MCARI/MTVI2	[23]
REP interpolation	$REP = \frac{700 + 40 \times [(R_{670} + R_{780}) / 2 - R_{700}]}{R_{740} - R_{700}}$	[24]
REPgauss	$R(\lambda) = R_s - (R_s - R_0) \exp \left[\frac{-(\lambda_0 - \lambda)^2}{2\sigma^2} \right]$	[25]

2.4.2 Variable Importance Projection

Variable projection importance (VIP) is a variable screening method based on partial least square (PLS), which describes the interpretation of the dependent variable to the dependent variable and ranks the independent variables according to their explanatory power[26,27]. The formula is

$$V_{VIP_j} = \sqrt{\frac{k}{\sum_{h=1}^m r^2(y, c_h)} \sum_{h=1}^m r^2(y, c_h) W_{hj}^2} \quad (1)$$

k--the number of vegetation indices, m--The number of components extracted from the original vegetation index variables, c_h --Principal Component Analysis of Independent Variables of Correlated Vegetation Index, $r(y, c_h)$ --correlative coefficient of dependent variables and principal components of winter wheat, indicating that the ability of **principal** components to explain y(LAI of wheat), W_{hj} --Weight of Vegetation Index on Principal Components. The higher the VIP value, the stronger the vegetation index can explain the **LAI** and vice versa.

2.4.3 Akaike's Information Criterion

The Akaike's Information Criterion [28] is a measure of the goodness of fit of a statistical model that measures the complexity of the model and the goodness of fit of the model. AIC can be expressed as

$$AIC = n \ln S_p^2 + 2(k + 1) \quad (2)$$

n--the number of samples, S_p^2 --Residual Mean Square, k-- Number of variables in the model.

2.4.4 Partial least squares

The partial least squares method is a multivariate statistical data analysis method, which mainly studies the multivariate regression model for single or **multi-variables** on the independent variables. The formula is

$$Y = a_0 + a_1 X_1 + a_2 X_2 + \Lambda + a_n X_n \quad (3)$$

a_0 --The intercept of the regression coefficient, a_i -- regression coefficient, X_i -- Independent variables 1 ~ n.

2.5 Accuracy verification

Leave one out **cross-validation** (LOOCV) each time to select a sample for verification, all other samples as a training sample. The model was validated, **and** the sample errors were calculated until all the samples were involved in the validation. LOOCV is the extreme case of K-cross-validation, and all values are modeled and validated. And it can avoid the problem of selection between modeling sample and verification sample in experimental design. This method can effectively evaluate the reliability and stability of the model.

The determination coefficient (R^2), root mean square error (RMSE) and relative error (RE) were selected as the indices to evaluate the modeling accuracy and validation accuracy. **R^2** is indicating that the predictive value and the measured value of the degree of fit. R^2 is closer to 1, indicating that the

higher the accuracy of the fitting curve. RMSE reflects the deviation between the predictive value and the measured value, the smaller the value, the higher the model accuracy. RE mainly reflects the credibility of the measurement. The smaller the relative error is, the more reliable the measured value. The calculation formula is shown below.

$$R^2 = \frac{(\sum xy - \frac{\sum x \sum y}{q})^2}{(\sum x^2 - \frac{(\sum x)^2}{q})(\sum y^2 - \frac{(\sum y)^2}{q})} \quad (4)$$

$$RMSE = \sqrt{\frac{\sum_{i=1}^q (y_i - \hat{y}_i)^2}{q}} \quad (5)$$

$$RE = \frac{\left| \sum_{i=1}^q \hat{y}_i - \sum_{i=1}^q y_i \right|}{\sum_{i=1}^q y_i} \times 100\% \quad (6)$$

x--vegetation index, y--crop LNC measured value, $\hat{y}$ -- the predicted value of the crop, q—the number of samples.

3. Results and analysis

3.1 Analysis of VIP between vegetation index and LNC of winter wheat

19 hyper-spectral indices (as shown in table 1) were calculated based on hyper-spectral data of leaves in wheat. The relationship between the obtained hyper-spectral indices and the measured nitrogen content of wheat leaves were analyzed by VIP. The results about the ability of vegetation indices to explain nitrogen content in wheat leaves are shown in Table 2. At the flag stage of wheat, between the spectral index and the leaf nitrogen content of the wheat, the maximum VIP corresponding spectral index is PSSRc, and the VIP value is 1.1944. The smallest VIP corresponding spectral index is MCARI/MTVI, and the VIP value is 0.8815. The top 10 spectral indices ordered by the VIP values are, PSSRc, GMI-2, SR705, RI-half, ZM, GMI-1, PSSRb, RI-3dB, VOGc and Cired edge, respectively.

Table 2 VIP and rank of evaluation index for LNC of winter wheat (n=199)

Vegetation index	VIP	Ranking	Vegetation Index	VIP	Ranking
PSSRc	1.1944	1	VOGa	1.1259	11
GMI-2	1.1897	2	RI-2dB	1.1253	12
SR705	1.1843	3	RI-1dB	1.1249	13
RI-half	1.1730	4	NDRE	1.1019	14
ZM	1.1659	5	REP	0.9680	15
GMI-1	1.1631	6	REPGAUSS	0.9643	16
PSSRb	1.1491	7	PRI	0.9565	17
RI-3dB	1.1364	8	SR _(533,565)	0.9465	18
VOGc	1.1360	9	MCARI/MTVI	0.8815	19

3.2 Selection of optimal LNC estimation model for wheat

According to the VIP sequence shown in table 2. The PLS models were constructed by using the different number of vegetation indices as the independent variables, calculating the AIC value of the model respectively. As shown in table 3, when the number of independent variables was 5 vegetation indices, the AIC value was the smallest, and the AIC value was -427.878. The AIC principle of building model was using as few parameters as possible. According to the AIC principle, in the flag stage of wheat, 5 vegetation indices were selected as independent variables to build PLS model, and its equation was an optimal model. The model was $V_{LNC}=4.9467+0.5186V_{PSSRc}-13.9993 V_{GMI-2}+43.7809 V_{SR705}+2.9433 V_{RI-half}-36.7703 V_{ZM}$. The 5 vegetation indices were PSSRc, GMI-2, SR705, RI-half, and ZM. The R^2 、RMSE and RE of the model were 0.79, 0.33 and 0, respectively.

Table 3 Comparison of PLS regression equations built by wheat LNC different independent variables according to VIP (flag leaf stage)

The number of variables	Regression equation	R^2	RMSE	AIC
4	$V_{LNC}=3.8885+0.4881V_{PSSRc}-9.4837 V_{GMI-2}+41.3387 V_{SR705}-34.7129 V_{RI-half}$	0.78	0.33	-424.580
5	$V_{LNC}=4.9467+0.5186V_{PSSRc}-13.9993 V_{GMI-2}+43.7809 V_{SR705}+2.9433 V_{RI-half}-36.7703 V_{ZM}$	0.79	0.33	-427.878
6	$V_{LNC}=5.0651+0.5004V_{PSSRc}-14.1179 V_{GMI-2}+45.0444 V_{SR705}+0.0615 V_{RI-half}-35.3281 V_{ZM}+0.1789 V_{GMI-1}$	0.79	0.33	-425.129
7	$V_{LNC}=5.9001+0.4098V_{PSSRc}-18.0207 V_{GMI-2}+55.1183 V_{SR705}-4.3869 V_{RI-half}-37.2390 V_{ZM}-0.1354 V_{GMI-1}+0.5298V_{PSSRb}$	0.79	0.33	-423.094
8	$V_{LNC}=6.1881+0.4101V_{PSSRc}-17.7359 V_{GMI-2}+53.4222 V_{SR705}-2.7516 V_{RI-half}-37.2390 V_{ZM}-0.1285 V_{GMI-1}+0.5186V_{PSSRb}-1.0977 V_{RI-3dB}$	0.79	0.33	-420.051
9	$V_{LNC}=9.7976+0.4726V_{PSSRc}-19.9056 V_{GMI-2}+69.2828 V_{SR705}+58.1725 V_{RI-half}-152.2723 V_{ZM}-0.1294 V_{GMI-1}+0.1085V_{PSSRb}+36.2288 V_{RI-3dB}-121.1323V_{VOGc}$	0.79	0.32	-422.513
10	$V_{LNC}=5.7079+0.5456V_{PSSRc}-19.7727V_{GMI-2}+76.3950V_{SR705}+38.7431V_{RI-half}-142.8775V_{ZM}+2.6587V_{GMI-1}-0.0605V_{PSSRb}+43.8932V_{RI-3dB}-88.1755V_{VOGc}-3.2832V_R$	0.79	0.32	-420.121

SI(543,728)

3.3 Verification of optimal LNC model for winter wheat

In order to verify the accuracy of VIP-PLS-AIC to predict the nitrogen status of wheat leaves, the data of wheat flag date of 2013~2014 were used respectively. For the nitrogen content of wheat leaves, the VIP-PLS-AIC model of 5 vegetation indices was validated. The results showed that R^2 and RMSE

were 0.79 and 0.33 respectively. The predicted values and the measured values of the scatter plot have high consistency, indicating that the use of VIP-PLS-AIC for winter wheat leaf nitrogen status estimation with high accuracy and good reliability as shown in figure 2.

Fig.2 Comparison of predicted and measured Wheat LNC based on VIP -PLS-AIC

4 Conclusion

The relationship between vegetation index and LNC was evaluated by VIP. The vegetation index with the highest VIP value was PSSRc, and the vegetation index with the lowest VIP value was MCARI / MTVI.

The number of vegetation indices **was** increased as the input factors, PLS regression model was used to establish the regression model of seven wheat LNC. The optimal LNC model was selected by AIC criterion. The results show that the PLS model with 5 vegetation indices are **the** optimal model for estimating LNC.

In wheat LNC model, the R^2 of VIP-PLS-AIC model set and validation set are 0.79, and RMSE is 0.33. The results of modeling and validation show that the importance of ordination between vegetation index and wheat LNC was better by VIP. And the optimal estimation model of wheat LNC can be obtained based on AIC.

Acknowledgments

This work was supported in part by the National Key Research and Development Programs (2016YFD0300603-5) ,National Natural Science Foundation of China (No.41601346) .

Reference

1. Wang J H, Zhao C J, Huang W J, et al. Quantitative Remote Sensing of Agriculture[M]. Beijing: Science Press, 2008.
2. Pinter Jr P J, Hatfield J L, Schepers J S, et al. Remote sensing for crop management[J]. Photogrammetric Engineering & Remote Sensing, 2003, 69(6): 647-664.
3. Hansen P M, Schjoerring J K. Reflectance measurement of canopy biomass and nitrogen status in wheat crops using normalized difference vegetation indices and partial least squares regression[J]. Remote sensing of environment, 2003, 86(4): 542-553.
4. Feng W, Yao X, Zhu Y, et al. Monitoring leaf nitrogen status with hyperspectral reflectance in wheat[J]. European Journal of Agronomy, 2008, 28(3): 394-404.
5. Clevers J, Kooistra L. Using hyperspectral remote sensing data for retrieving total canopy chlorophyll and

- nitrogen content[C]//Hyperspectral Image and Signal Processing: Evolution in Remote Sensing (WHISPERS), 2011 3rd Workshop on. IEEE, 2011: 1-4.
6. Cui Rixian, Liu Yadong, Fu Jindong. Estimation of Winter Leaf Nitrogen Accumulation using Machine Learning Algorithm and Visible Spectral[J]. Spectroscopy and Spectral Analysis, 2016, 36(6):1837-1842.
 7. Zhang Xia, Liu Liangyun, Zhao Chunjiang, et al. Estimating Wheat Nitrogen Concentration with High Spectral Resolution Image[J]. Journal of Remote Sensing, 2003, 7(3):176-181.
 8. Li Fenling, Chang Qingrui, Shen Jian, et al. Remote sensing estimation of winter wheat leaf nitrogen content based on GF-1 satellite data[J]. Transactions of the Chinese Society of Agricultural Engineering , 2016, 32(9):157-164.
 9. Fei Wei, Zhu Yan, Yao Xia, et al. Monitoring nitrogen accumulation in wheat leaf with red edge characteristics parameters[J]. Transactions of the Chinese Society of Agricultural Engineering, 2009, 25(11):194-201.
 10. Wang Renhong, Song Xiaoyu, Li Zhenhai, et al. Estimation of winter wheat nitrogen nutrition index using hyperspectral remote sensing[J]. Transactions of the Chinese Society of Agricultural Engineering, 2014, 30(19):191-198.
 11. Li Z H, Xu X G, Jin X L, et al. Remote Sensing Prediction of Winter Wheat Protein Content Based on Nitrogen Translocation and GRA-PLS Method.[J]. Scientia Agricultura Sinica, 2014, 47(19): 3780-3790.
 12. Jin X L, Xu X G, Wang J H, et al. Hyperspectral Estimation of Leaf Water Content for Winter Wheat Based on Relational Analysis (GRA)[J]. Spectroscopy and Spectral Analysis, 2012, 32(11): 3103-3106.
 13. Gautam R, Panigrahi S, Franzen D. Neural network optimisation of remotely sensed maize leaf nitrogen with a genetic algorithm and linear programming using five performance parameters [J]. Biosystems Engineering, 2006, 95(3): 359-370.
 14. Blackburn G A. Spectral indices for estimating photosynthetic pigment concentrations: a test using senescent tree leaves[J]. International Journal of Remote Sensing, 1998, 19(4): 657-675.(2,3,4,18,19,20)
 15. Gitelson A A, Merzlyak M N. Remote sensing of chlorophyll concentration in higher plant leaves[J]. Advances in Space Research, 1998, 22(5): 689-692.(6,7,8)
 16. Vogelmann J E, Rock B N, Moss D M. Red edge spectral measurements from sugar maple leaves[J]. Title REMOTE SENSING, 1993, 14(8): 1563-1575.(9)
 17. Zarco-Tejada P J, Miller J R, Noland T L, et al. Scaling-up and model inversion methods with narrowband optical indices for chlorophyll content estimation in closed forest canopies with hyperspectral data[J]. Geoscience and Remote Sensing, IEEE Transactions on, 2001, 39(7): 1491-1507.(10,24)
 18. Gupta R K, Vijayan D, Prasad T S. Comparative analysis of red-edge hyperspectral indices[J]. Advances in Space Research, 2003, 32(11): 2217-2222.(11-14)
 19. Tian Y C, Yao X, Yang J, et al. Assessing newly developed and published vegetation indices for estimating rice leaf nitrogen concentration with ground-and space-based hyperspectral reflectance[J]. Field Crops Research, 2011, 120(2): 299-310.(15)
 20. Rodriguez D, Fitzgerald G J, Belford R, et al. Detection of nitrogen deficiency in wheat from spectral reflectance indices and basic crop eco-physiological concepts[J]. Crop and Pasture Science, 2006, 57(7): 781-789.(27)
 21. Penuelas J, Baret F, Filella I. Semi-empirical indices to assess carotenoids/chlorophyll a ratio from leaf spectral reflectance[J]. Photosynthetica, 1995, 31: 221-230.(28,30)
 22. Dash J, Curran P J. Evaluation of the MERIS terrestrial chlorophyll index (MTCI). Advances in Space Research, 2007, 39(1): 100-104.(38)
 23. Eitel J U H, Long D S, Gessler P E, et al. Using in-situ measurements to evaluate the new RapidEye™ satellite series for prediction of wheat nitrogen status[J]. International Journal of Remote Sensing, 2007, 28(18):4183-4190.
 24. Guyot G, Baret F, Major D J. High spectral resolution: Determination of spectral shifts between the red and the near infrared[J]. International Archives of Photogrammetry and Remote Sensing, 1988, 11(750-760).
 25. Zhu Y, Li Y, Feng W, et al. Monitoring leaf nitrogen in wheat using canopy reflectance spectra[J]. Canadian Journal of Plant Science, 2006, 86(4): 1037-1046.
 26. Zeng T, Ju C Y, Cai T J, et al. Selection of parameters for Estimating Canopy Closure Density Using Variable Importance of Projection Criterion[J]. Journal of Beijing Forestry University, 2010, 32(6): 37-41.
 27. Yang F Q, Feng H K, Li Z H, et al. Estimation of Leaf Area Index of Winter Wheat Based on Akaike's Information Criterion[J]. Transactions of The Chinese Society of Agricultural Machinery, 2015, 46(11): 112-120.
 28. Akaike H.(1973) Information theory and an extension of the maximum likelihood principle[J], 2nd International Symposium of Information Theory, B. N. Petrov and F.Csaki, eds., Akademiai Kiado, Budapest, 267-281.