

Computability on quasi-Polish spaces

Mathieu Hoyrup, Cristóbal Rojas, Victor Selivanov, Donald Stull

► **To cite this version:**

Mathieu Hoyrup, Cristóbal Rojas, Victor Selivanov, Donald Stull. Computability on quasi-Polish spaces. DCFS 2019 - 21st International Conference on Descriptive Complexity of Formal Systems, Jul 2019, Kosice, Slovakia. pp.171-183. hal-02118947v2

HAL Id: hal-02118947

<https://hal.inria.fr/hal-02118947v2>

Submitted on 7 May 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Computability on quasi-Polish spaces^{*}

Mathieu Hoyrup¹, Cristóbal Rojas^{**2}, Victor Selivanov^{***3}, Donald M. Stull^{†1}

¹ Université de Lorraine, CNRS, Inria, LORIA, F 54000 Nancy, France
`mathieu.hoyrup@inria.fr`, `donald.stull@inria.fr`

² Universidad Andres Bello, Santiago, Chile
`crojas@mat-unab.cl`

³ A.P. Ershov Institute of Informatics Systems, Novosibirsk
and Kazan Federal University, Russia
`vseliv@iis.nsk.su`


Abstract. We investigate the effectivizations of several equivalent definitions of quasi-Polish spaces and study which characterizations hold effectively. Being a computable effectively open image of the Baire space is a robust notion that admits several characterizations. We show that some natural effectivizations of quasi-metric spaces are strictly stronger.

1 Introduction

Classical descriptive set theory (DST) [11] deals with hierarchies of sets, functions and equivalence relations in Polish spaces. Theoretical Computer Science, in particular Computable Analysis [21], motivated an extension of the classical DST to non-Hausdorff spaces; a noticeable progress was achieved for the ω -continuous domains and quasi-Polish spaces [18, 3]. The theory of quasi-Polish spaces is already a well-established part of classical DST [3, 5]

Theoretical Computer Science and Computable Analysis especially need an effective DST for some effective versions of the mentioned classes of topological spaces. A lot of useful work in this direction was done in Computability Theory but only for the discrete space \mathbb{N} , the Baire space \mathcal{N} , and some of their relatives [16, 6]. For a systematic work to develop the effective DST for effective Polish spaces see e.g. [15, 14, 7]. There was also some work on the effective DST for effective domains and approximation spaces [18, 19, 2, 20].

In this paper we continue the search of a “correct” version of a computable quasi-Polish space initiated in [20, 13]. By a correct version we mean one having properties similar to effective versions of those in the classical case: the

^{*} This project has received funding from the European Unions Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 731143 

^{**} This author was supported by Marie Curie RISE project CID

^{***} This author was supported by Inria program Invited Researcher, and the Regional Mathematical Center of Kazan Federal University (project 1.13556.2019/13.1 of the Ministry of Education and Science of Russian Federation)

[†] This author was supported by Inria post-doc program

computable quasi-Polish spaces have to subsume the well established classes of computable Polish spaces and computable ω -continuous domains and to admit a good enough effective DST.

We identify effective versions of quasi-Polish spaces satisfying these specifications. One of them is the class of computable effectively open images of the Baire space identified and studied in [20]. We provide some characterizations of this class which are effective versions of the corresponding known characterizations of quasi-Polish spaces in [3]. However we show that some natural effectivizations of complete quasi-metric spaces are strictly stronger.

The results of this paper were obtained in September 2018 during a research stay of the second and third authors in Inria, Nancy. On the final stage of preparing this paper the preprint [4] appeared where some of our results were obtained independently (using a slightly different approach and terminology), notably Definition 4.1 and Theorem 4.1.

In order to make our discussion of effective spaces closer to the corresponding classical theory, we use an approach based on the canonical embeddings of cb_0 -spaces into the Scott domain $\mathcal{P}(\omega)$ and on the computability in this domain. This approach (which emphasizes the notion of effective continuity rather than the equivalent notion of computability w.r.t. admissible representations more popular in Computable Analysis) was promoted in [12, 19].

We start in the next section with recalling definitions of some notions of effective spaces and of effective DST in such spaces; we also try to simplify and unify rather chaotic terminology in this field. In Sections 2 and 3 we establish the main technical tools used in the sequel. In Section 4 we propose a definition of effective quasi-Polish spaces and prove characterizations of this notion. In Section 5 we propose two effective notions of quasi-metric space and prove that they differ from the notion of effective quasi-Polish space.

2 Preliminaries

Here we recall some known notions and facts, with a couple of new observations.

Notions similar to those considered below were introduced (sometimes independently) and studied in [9], [12], [19] under different names. We use a slightly different terminology, trying to simplify it and make it closer to that of classical topology. Note that the terminology in effective topology is still far from being fixed. All topological spaces considered in this paper are assumed to be countably based. Such a space satisfying the T_0 -separation axiom is sometimes called a cb_0 -space, for short. We recall that $(W_e)_{e \in \mathbb{N}}$ is some effective enumeration of the computably enumerable (c.e.) subsets of \mathbb{N} .

Definition 2.1. *An effective topological space is a countably-based T_0 topological space coming with a numbering $(B_i^X)_{i \in \mathbb{N}}$ of a basis, such that there is a computable function $f : \mathbb{N}^2 \rightarrow \mathbb{N}$ such that $B_i^X \cap B_j^X = \bigcup_{k \in W_{f(i,j)}} B_k^X$.*

Many popular spaces (e.g., the discrete space \mathbb{N} of naturals, the space of reals \mathbb{R} , the domain $\mathcal{P}(\omega)$, the Baire space $\mathcal{N} = \mathbb{N}^{\mathbb{N}}$, the Cantor space $\mathcal{C} = 2^{\mathbb{N}}$ and the

Baire domain $\mathbb{N}^{\leq \mathbb{N}} = \mathbb{N}^* \cup \mathbb{N}^{\mathbb{N}}$ of finite and infinite sequences of naturals) are effective topological spaces in an obvious way. The effective topological space \mathbb{N} is trivial topologically but very interesting for Computability Theory. We use some almost standard notation related to the Baire space. In particular, $[\sigma]$ denotes the basic open set induced by $\sigma \in \mathbb{N}^*$ consisting of all $p \in \mathcal{N}$ having σ as an initial segment; we sometimes call such sets cylinders. Let ε denote the empty string in \mathbb{N}^* .

In [17, 19] the effective Borel and effective Hausdorff hierarchies in arbitrary effective topological spaces X were introduced. Also the effective Luzin hierarchy is defined naturally [20]. Below we use the simplified notation for levels of these hierarchies like $\Pi_n^0(X)$, $\Sigma_n^1(X)$ or $\Sigma_n^{-1}(X)$ (which naturally generalizes the notation in computability theory) and some of their obvious properties. We will also use the expression *effective open set* for sets in the class $\Sigma_1^0(X)$, which are the sets $\bigcup_{i \in W} B_i^X$ for some c.e. set $W \subseteq \mathbb{N}$.

Definition 2.2. *If X, Y are effective topological spaces then a function $f : X \rightarrow Y$ is **computable** if the sets $f^{-1}(B_i^Y)$ are uniformly effective open sets.*

As observed in [20], for any effective topological space X , the equality relation $=_X$ on X is in $\Pi_2^0(X \times X)$. The argument in [20] shows that also the specialization partial order \leq_X has the same descriptive complexity. In particular, every singleton is in the boldface class $\mathbf{\Pi}_2^0(X)$.

With any effective topological space X we associate the *canonical embedding* $e : X \rightarrow \mathcal{P}(\omega)$ defined by $e(x) = \{n \mid x \in B_n^X\}$ (in [19] the canonical embedding was denoted as O_ξ ; we changed the notation here to make it closer to that of the paper [5] which is cited below). The canonical embedding is a computable homeomorphism between X and the subspace $e[X]$ of $\mathcal{P}(\omega)$. It can be used to study computability on cb_0 -spaces [12, 19] using the fact that the computable functions on $\mathcal{P}(\omega)$ coincide with the enumeration operators [16].

The more popular and general approach to computability on topological spaces is based on representations [21]. The relation between the two approaches is based on the so called enumeration representation $\rho : \mathcal{N} \rightarrow \mathcal{P}(\omega)$ defined by $\rho(x) = \{n \mid \exists i(x(i) = n + 1)\}$. The function ρ is a computable effectively open surjection. The canonical embedding e induces the **standard representation** $\rho_X = e^{-1} \circ \rho_A$ of X where $A = e(X)$ and ρ_A is the restriction of ρ to $\rho^{-1}(A)$. The function ρ_X is a computable effectively open surjection. We will implicitly identify any effective topological space X with its image under the canonical embedding, so that X is a subspace of $\mathcal{P}(\omega)$, and ρ_X is the restriction of ρ to $\rho^{-1}(X)$.

Note that for effective topological spaces X and Y , $f : X \rightarrow Y$ is computable iff there exists a computable function $F : \text{dom}(\rho_X) \rightarrow \text{dom}(\rho_Y)$ such that $\rho_Y \circ F = f \circ \rho_X$.

3 Results on $\mathbf{\Pi}_2^0$ -sets

This section contains the technical tools that will be used to prove the characterizations of effective quasi-Polish spaces.

Definition 3.1. Let X be an effective topological space. We say that $A \subseteq X$ is **computably overt** if the set $\{i \in \mathbb{N} : B_i^X \cap A \neq \emptyset\}$ is c.e.

Observe that the overt information does not uniquely determine the set, but only its closure. In the literature, overt and computably overt sets are often assumed to be closed. It is important to note that in this paper, no such assumption is made.

We recall that if X is an effective topological space then a set is in $\Pi_2^0(X)$ if it is an intersection of Boolean combinations of uniformly effective open subsets of X . We prove an effective version of Theorem 68 in [3].

Lemma 3.1. Let X be an effective topological space. For $A \subseteq X$,

- $A \in \Pi_2^0(X)$ iff $\rho_X^{-1}(A) \in \Pi_2^0(\text{dom}(\rho_X))$,
- A is computably overt iff $\rho_X^{-1}(A)$ is computably overt.

Proof. If $A \in \Pi_2^0(X)$ then one easily obtains $\rho_X^{-1}(A) \in \Pi_2^0(\text{dom}(\rho_X))$. We now prove that if $\rho_X^{-1}(A) \in \Sigma_2^0(\text{dom}(\rho_X))$ then $A \in \Sigma_2^0(X)$, which implies the same result for the class Π_2^0 . Let $\rho_X^{-1}(A) = \text{dom}(\rho_X) \cap \bigcup_n U_n \setminus V_n$ where $U_n, V_n \in \Sigma_1^0(\mathcal{N})$ uniformly and $V_n \subseteq U_n$. Then the set

$$A' := \bigcup_{\sigma, n} \rho_X([\sigma] \cap U_n) \setminus \rho_X([\sigma] \cap V_n)$$

belongs to $\Sigma_2^0(X)$, because the image of a $\Sigma_1^0(\mathcal{N})$ -set is a $\Sigma_1^0(X)$ -set uniformly. We show that $A = A'$. The inclusion $A' \subseteq A$ is straightforward. For the other inclusion, let $x \in A$. One has $\rho_X^{-1}(x) \in \Pi_2^0(\mathcal{N})$ so $\rho_X^{-1}(x)$ is quasi-Polish so it is a Baire space ([3]). One has $\rho_X^{-1}(x) \subseteq \bigcup_n U_n \setminus V_n$. By Baire category, there exists n such that $\rho_X^{-1}(x) \cap U_n \setminus V_n$ is somewhere dense in $\rho_X^{-1}(x)$, i.e. there exists $\sigma \in \mathbb{N}^*$ such that $\emptyset \neq \rho_X^{-1}(x) \cap [\sigma] \subseteq U_n \setminus V_n$. As a result, $x \in \rho_X([\sigma] \cap U_n) \setminus \rho_X([\sigma] \cap V_n) \subseteq A'$.

If A is computably overt then $[\sigma] \cap \rho_X^{-1}(A) \neq \emptyset$ iff $\rho_X([\sigma]) \cap A \neq \emptyset$ which is c.e. as $\rho_X([\sigma]) \in \Sigma_1^0(X)$, uniformly in σ .

If $\rho_X^{-1}(A)$ is computably overt then $[\sigma] \cap A \neq \emptyset$ iff $\rho_X^{-1}([\sigma]) \cap \rho_X^{-1}(A) \neq \emptyset$ is a c.e. relation as $\rho_X^{-1}([\sigma]) \in \Sigma_1^0(X)$, uniformly in i .

An important property of computably overt Π_2^0 -sets is that they contain computable points. It is a crucial ingredient in the next results.

Proposition 3.1 ([10]). In a computable Polish space, a Π_2^0 -set is computably overt if and only if it contains a dense computable sequence.

Moreover, the next result shows that in a computably overt Π_2^0 -set, not only can one find an effective indexing over \mathbb{N} of a dense set of elements, but one can even find an effective indexing over \mathcal{N} of *all* its elements.

Lemma 3.2. Let $A \subseteq \mathcal{N}$ be non-empty. The following are equivalent:

- (i) A is a computably overt Π_2^0 -set,

(ii) *There exists a computable effectively open surjective map $f : \mathcal{N} \rightarrow A$.*

When we write that $f : \mathcal{N} \rightarrow A$ is open, we mean that for each $\sigma \in \mathbb{N}^*$, there exists an open set $U_\sigma \subseteq \mathcal{N}$ such that $f([\sigma]) = A \cap U_\sigma$. f is effectively open when U_σ is effectively open, uniformly in σ .

Proof. Assume (i). Let $A = \bigcap_n A_n$ where A_n are uniformly effective open sets. We can assume w.l.o.g. that $A_{n+1} \subseteq A_n$.

One can build a computable sequence $(u_\sigma)_{\sigma \in \mathbb{N}^*}$ such that $u_\epsilon = \epsilon$ and:

- If τ properly extends σ then u_τ properly extends u_σ ,
- If $|\sigma| = n$ then $[u_\sigma] \subseteq A_n$,
- $[u_\sigma]$ intersects A ,
- $[u_\sigma] \cap A$ is contained in $\bigcup_{i \in \mathbb{N}} [u_{\sigma \cdot i}]$.

We build this sequence inductively in σ . Given u_σ intersecting A with $|\sigma| = n$, one can compute a covering of $[u_\sigma] \cap A_{n+1}$ with cylinders properly extending u_σ and extract the cylinders intersecting A . Let $(u_{\sigma \cdot i})_{i \in \mathbb{N}}$ be some computable enumeration of them.

We now define f . For each $p \in \mathcal{N}$, the sequence $u_{p \upharpoonright_n}$ converges to some $q \in \mathcal{N}$. We define $f(p) = q$. One easily checks that the function $f : \mathcal{N} \rightarrow A$ is computable, onto and effectively open as $f([\sigma]) = [u_\sigma] \cap A$.

Now assume (ii). The function f has a computable right-inverse, i.e. $g : A \rightarrow \mathcal{N}$ such that $f \circ g$ is the identity on A . Indeed, given $p \in A$, one can enumerate all the cylinders intersecting $f^{-1}(p)$ as $[\sigma] \cap f^{-1}(p) \neq \emptyset$ iff $p \in f([\sigma])$ which can be recognized as $f([\sigma])$ is effectively open. Hence one can progressively build an element of the closed set $f^{-1}(p)$.

The function g is a partial computable function from $\mathcal{N} \rightarrow \mathcal{N}$. Its domain is Π_2^0 and contains A . One has $p \in A \iff p$ belongs to the domain of g and $p = f \circ g(p)$. As a result, A is Π_2^0 . The image under f of a dense computable sequence in \mathcal{N} is a dense computable sequence in A , so the set of cylinders intersecting A is c.e.

This result can be extended to subsets of $\mathcal{P}(\omega)$.

Lemma 3.3. *Let $A \subseteq \mathcal{P}(\omega)$ be non-empty. The following are equivalent:*

- (i) *A is a computably overt Π_2^0 -set,*
- (ii) *There exists a computable effectively open surjective map $f : \mathcal{N} \rightarrow A$.*

Proof. If A is a computably overt Π_2^0 -set then so is $\rho^{-1}(A)$, so there exists a computable effectively open onto function $f : \mathcal{N} \rightarrow \rho^{-1}(A)$. The function $\rho \circ f$ satisfies the required conditions.

Conversely, assume that $f : \mathcal{N} \rightarrow A$ is a computable effectively open surjective function.

Claim. There exists a computable function $g : \rho^{-1}(A) \rightarrow \mathcal{N}$ such that $f \circ g = \rho$.

Proof (of the claim). Given $p \in \rho^{-1}(A)$, let $A_p = \{q \in \mathcal{N} : f(q) = \rho(p)\}$. The set A_p is a computably overt Π_2^0 -set relative to p . Indeed, it is Π_2^0 relative to p because equality is Π_2^0 in $\mathcal{P}(\omega)$. It is computably overt relative to p because a cylinder $[\sigma]$ intersects A_p iff $\rho(p) \in f([\sigma])$ which is a c.e. relation in p as f is effectively open. As a result, by relativizing Proposition 3.1, one can compute an element in A_p . Everything is uniform in p , so there is a computable function g mapping each $p \in \rho^{-1}(A)$ to an element of A_p , hence $f \circ g(p) = \rho(p)$.

Now, one has $q \in \rho^{-1}(A)$ iff $g(q)$ is defined and $f \circ g(q) = \rho(q)$. Both relations are Π_2^0 , so $\rho^{-1}(A) \in \Pi_2^0(\mathcal{N})$ hence $A \in \Pi_2^0(\mathcal{P}(\omega))$ by Lemma 3.1. Moreover, A is computably overt because for each basic open set B of $\mathcal{P}(\omega)$, $B \cap A \neq \emptyset$ iff $f^{-1}(B) \neq \emptyset$, which is a c.e. relation.

4 Effective quasi-Polish spaces

According to Theorem 23 of [3], the quasi-Polish spaces (defined originally as the countably based completely quasi-metrizable spaces) coincide with the continuous open images of the Baire space. Effectivizing this definition, we obtain the following candidate for a notion of effective quasi-Polish space.

Definition 4.1. *An effective topological space X is an **effective quasi-Polish space** if X is the image of \mathcal{N} under a computable effectively open map, or X is empty.*

Of course this notion is preserved by computable homeomorphisms (bijections that are computable in both directions).

Theorem 4.1. *Let X be an effective topological space with its standard representation ρ_X . The following statements are equivalent:*

1. X is effective quasi-Polish,
2. The image of X under its canonical embedding in $\mathcal{P}(\omega)$ is a computably overt Π_2^0 -subset of $\mathcal{P}(\omega)$,
3. $\text{dom}(\rho_X)$ is a computably overt Π_2^0 -subset of \mathcal{N} .

Proof. The equivalence 1. \iff 2. is the content of Lemma 3.3. The equivalence 2. \iff 3. is the content of Lemma 3.1 for the space $\mathcal{P}(\omega)$.

We also formulate the effective version of Theorem 21 of [3].

Theorem 4.2. *Let X be an effective quasi-Polish space. A subspace $Y \subseteq X$ is an effective quasi-Polish space iff Y is a computably overt Π_2^0 -subset of X .*

Proof. Via the canonical embedding, we have $Y \subseteq X \subseteq \mathcal{P}(\omega)$. We start by assuming that Y is a computably overt Π_2^0 -subset of X . As $Y \in \Pi_2^0(X)$ and $X \in \Pi_2^0(\mathcal{P}(\omega))$, one has $Y \in \Pi_2^0(\mathcal{P}(\omega))$. To show that Y is computably overt in $\mathcal{P}(\omega)$, simply observe that for a basic open set B of $\mathcal{P}(\omega)$, B intersects Y iff $B^X := B \cap X$ intersects Y . It is a c.e. relation as Y is computably overt in X .

If Y is effective quasi-Polish then it is a computably overt Π_2^0 -subset of $\mathcal{P}(\omega)$ so it is a computably overt Π_2^0 -subset of X , which is a subspace of $\mathcal{P}(\omega)$.

Sometimes it is easier to work with a computably admissible representation other than the standard representation.

Theorem 4.3. *Let X be an effective topological space. If X admits a computably admissible representation whose domain is a computably overt Π_2^0 -subset of \mathcal{N} , then X is an effective Polish space.*

Proof. Let δ be a computably admissible representation of X such that $\text{dom}(\delta)$ is a computably overt Π_2^0 -subset of \mathcal{N} . By definition of computably admissible, δ is computably equivalent to ρ_X , i.e. there exist partial computable functions $F, G : \subseteq \mathcal{N} \rightarrow \mathcal{N}$ satisfying $\rho_X = \delta \circ F$ and $\delta = \rho_X \circ G$. We show that $\text{dom}(\rho_X)$ is a computably overt Π_2^0 -set.

We recall that ρ_X is the restriction of the representation ρ of $\mathcal{P}(\omega)$ to $\rho^{-1}(X)$. We show that $\text{dom}(\rho_X) = \rho^{-1}(X)$ is a computably overt Π_2^0 -set. Let $p \in \mathcal{N}$. One has $p \in \text{dom}(\rho_X) = \rho^{-1}(X)$ iff $F(p)$ is defined, $F(p) \in \text{dom}(\delta)$ and $\rho(G \circ F(p)) = \rho(p)$. All these conditions are Π_2^0 , so $\rho^{-1}(X)$ belongs to $\Pi_2^0(\mathcal{N})$.

One has $[\sigma] \cap \text{dom}(\rho_X) \neq \emptyset$ iff $\delta^{-1}(\rho([\sigma])) \neq \emptyset$ which is c.e. in σ as $\text{dom}(\delta)$ is computably overt.

5 Effective quasi-metric spaces

We now propose two effective versions of quasi-metric spaces and compare them with the notion of effective quasi-Polish space. A quasi-metric on a set X is a function $d : X \times X \rightarrow \mathbb{R}_{\geq 0}$ satisfying:

- $d(x, z) \leq d(x, y) + d(y, z)$,
- $x = y$ iff $d(x, y) = d(y, x) = 0$.

The quasi-metric d induces a metric $\hat{d}(x, y) = \max(d(x, y), d(y, x))$.

Definition 5.1. *A computable quasi-metric space is a triple (X, d, S) where d is a quasi-metric on X and $S = \{s_i\}_{i \in \mathbb{N}}$ is a \hat{d} -dense sequence such that $d(s_i, s_j)$ are uniformly computable.*

We recall that a real number x is **right-c.e.** if $x = \inf_i q_i$ for some computable sequence of rationals $(q_i)_{i \in \mathbb{N}}$.

Definition 5.2. *A right-c.e. quasi-metric space is a triple (X, d, S) where d is a quasi-metric on X and $S = \{s_i\}_{i \in \mathbb{N}}$ is a \hat{d} -dense sequence such that $d(s_i, s_j)$ are uniformly right-c.e.*

Every right-c.e. quasi-metric space is an effective topological space with the basis of balls $B(s, r) = \{x \in X : d(s, x) < r\}$ with $s \in S$ and r positive rational. To see this, we consider formal inclusion between balls: $B(s, q) \sqsubseteq B(t, r)$ iff $d(t, s) + q < r$. Formal inclusion is c.e. and $B_i \cap B_j = \bigcup_{k: B_k \sqsubseteq B_i \text{ and } B_k \sqsubseteq B_j} B_k$, so the axiom of effective topological spaces is satisfied. As a result, any such space has its standard representation δ_S . We define another representation.

Definition 5.3. The **Cauchy representation** δ_C is defined in the following way: a point $x \in X$ is represented by any sequence $s_n \in S$ such that $d(s_n, s_{n+1}) < 2^{-n}$ and s_n converges to x in the metric \hat{d} .

Theorem 5.1. On a right-c.e. quasi-metric space, the Cauchy representation is computably equivalent to the standard representation.

Proof. For the proof we will also consider a slightly different representation δ'_C where x is represented by any sequence s_n such that $d(s_n, x) < 2^{-n}$ and s_n converges to x in \hat{d} .

We prove the following computable reductions: $\delta_C \leq \delta'_C \leq \delta_S \leq \delta_C$.

Proof of $\delta_C \leq \delta'_C$. Assume we are given a δ_C -name of x , which is essentially a sequence $(s_n)_{n \in \mathbb{N}}$ such that $d(s_n, s_{n+1}) < 2^{-n}$ and $\lim_{n \rightarrow \infty} \hat{d}(s_n, x) = 0$. One easily checks that the sequence $(s_{n+1})_{n \in \mathbb{N}}$ is a δ'_C -name for x .

Proof of $\delta'_C \leq \delta_S$. Assume we are given a δ'_C -name of x , which is essentially a sequence s_n such that $d(s_n, x) < 2^{-n}$ and $\lim_{n \rightarrow \infty} \hat{d}(s_n, x) = 0$. We show that we can enumerate the basic balls containing x . Indeed, we show that $x \in B(s, r)$ if and only if there exists n such that $d(s, s_n) < r - 2^{-n}$. First assume that the latter inequality holds. By the triangle inequality,

$$d(s, x) \leq d(s, s_n) + d(s_n, x) < r - 2^{-n} + 2^{-n} = r.$$

Conversely, if $d(s, x) < r$ then as $\hat{d}(s_n, x)$ converges to 0, for sufficiently large n one has $d(x, s_n) + 2^{-n} < r - d(s, x)$ so $d(s, s_n) \leq d(s, x) + d(x, s_n) < r - 2^{-n}$.

Proof of $\delta_S \leq \delta_C$. Assume we are given an enumeration of the basic balls containing x , call it U_1, U_2, U_3, \dots . We build a sequence $(s_n)_{n \in \mathbb{N}}$ as follows.

We take s_0 such that $d(s_0, x) < 1$, which we can find by looking for a ball of radius 1 containing x . Once s_0, \dots, s_n have been defined, we look for s_{n+1} satisfying:

- $s_{n+1} \in U_1 \cap \dots \cap U_{n+1}$,
- $d(s_n, s_{n+1}) < 2^{-n}$,
- $d(s_{n+1}, x) < 2^{-n-1}$.

Such a point must exist, as if x' is sufficiently \hat{d} -close to x , the first and third conditions are satisfied, and $d(s_n, x') \leq d(s_n, x) + d(x, x') < 2^{-n} + d(x, x')$ by induction hypothesis, so $d(s_n, x') < 2^{-n}$ if $d(x, x')$ is sufficiently small. Such a point can be effectively found, d is right-c.e. on S .

The sequence $(s_n)_{n \in \mathbb{N}}$ satisfies the conditions of being a δ_C -name of x . Indeed, $d(x, s_n)$ converge to 0, as for each rational ϵ there exists $s \in S$ such that $\hat{d}(s, x) < \epsilon$, so the ball $B(s, \epsilon)$ appears as some U_i , so for $n \geq i$, $d(x, s_n) \leq d(x, s) + d(s, s_n) < 2\epsilon$.

We recall that a quasi-metric d is (Smyth-)complete if every Cauchy sequence converges in the metric \hat{d} , and that a space is quasi-Polish iff it is completely quasi-metrizable [3]. One direction of this equivalence admits an effective version.

Theorem 5.2. *Every right-c.e. quasi-metric space that is complete is an effective quasi-Polish space.*

Proof. The domain of the Cauchy representation is a computably overt Π_2^0 -set. Indeed, the relation $\forall n, d(s_n, s_{n+1}) < 2^{-n}$ is Π_2^0 , and any finite sequence satisfying this condition can be extended (to an ultimately constant sequence, e.g.). As the Cauchy representation is computably equivalent to the standard representation, we can apply Theorem 4.3.

Proposition 5.1. *In a right-c.e. quasi-metric space, the following conditions are equivalent for a point x :*

- x is computable,
- The numbers $d(s, x)$ are right-c.e., uniformly in $s \in \mathcal{S}$.

Proof. One has $x \in B(s, r) \iff d(s, x) < r$. The first relation is c.e. iff x is computable. The second relation is c.e. iff $d(s, x)$ is right-c.e.

We recall that a real number x is left-c.e. if $-x$ is right-c.e.

Proposition 5.2. *In a computable quasi-metric space, the following conditions are uniformly equivalent for a point x :*

- x is computable,
- The numbers $d(s, x)$ are right-c.e., uniformly in $s \in \mathcal{S}$,
- The numbers $d(s, x)$ and $d(x, s)$ are right-c.e. and left-c.e. respectively, uniformly in $s \in \mathcal{S}$.

Proof. We only have to prove that for a computable point x , the numbers $d(x, s)$ are uniformly left-c.e. Let $(s_n)_{n \in \mathbb{N}}$ be a computable δ_C -name of x . We show that $d(x, s) = \sup_n d(s_n, s) - 2^{-n}$ which is left-c.e., uniformly in s .

Indeed, $d(x, s) \geq d(s_n, s) - d(s_n, x)$, and as $d(x, s_n) \leq \hat{d}(x, s_n)$ converges to 0, $d(x, s) \leq d(x, s_n) + d(s_n, s)$ is arbitrarily close to $d(s_n, s)$.

6 Separation

Classically, a space is quasi-Polish if and only if it is completely quasi-metrizable [3]. However the proof is not constructive. We know that each right-c.e. quasi-metric space that is complete is an effective quasi-Polish space, but that the converse fails. For this, we fully characterize the effective notions of quasi-Polish space in a restricted case.

Let $[0, 1]_{<}$ come with the quasi-metric $d(x, y) = \max(0, x - y)$, with the rational points as \hat{d} -dense sequence. It is a computable quasi-metric space that is complete. For $\alpha \in (0, 1)$, the subspace $[\alpha, 1]_{<}$ is an effective topological subspace of $[0, 1]_{<}$. We investigate when it is an effective quasi-Polish space, a computably quasi-metrizable space, and a right-c.e. quasi-metrizable space.

Proposition 6.1. *The space $[\alpha, 1]_{<}$ is an effective quasi-Polish space iff α is left-c.e. relative to the halting set.*

Proof. Observe that the set $\{q : (q, 1] \cap A \neq \emptyset\}$ is always c.e. Therefore, $[\alpha, 1]_<$ is an effective quasi-Polish space if and only if $[\alpha, 1] \in \Pi_2^0([0, 1]_<)$ by Theorem 4.2 (the c.e. conditions is always satisfied as observed above). This is equivalent to the existence of uniformly right-c.e. numbers r_i such that $[\alpha, 1] = \bigcap_i (r_i, 1]$, i.e. $\alpha = \sup_i r_i$. This is equivalent to α being left-c.e. relative to the halting set.

Proposition 6.2. *The space $[\alpha, 1]_<$ admits a computably equivalent computable quasi-metric structure if and only if α is right-c.e.*

Proof. Assume first that α is right-c.e. There is a computable enumeration $S = \{q_i\}_{i \in \mathbb{N}}$ of the rational numbers in $(\alpha, 1]$. The quasi-metric $d(x, y) = \max(0, x - y)$ is computable on S .

Conversely, assume a computable quasi-metric d with an associate set $S = \{s_i\}_{i \in \mathbb{N}}$. We now prove that the points s_i are uniformly computable real numbers, which implies that $\alpha = \inf_s s_i$ is right-c.e. The function mapping a real number $x \in [\alpha, 1]$ to $d(x, s_i)$ is left-c.e. (x is given using the standard Cauchy representation). Indeed, from x one can compute a name of x in $[0, 1]_<$, from which one can compute a name of x in $[\alpha, 1]_<$ and we can apply the uniform relative version of Proposition 5.2.

The left-c.e. function $x \mapsto d(x, s_i)$ is non-decreasing. Indeed, for $x \leq x'$, one has $d(x, s_i) \leq d(x, x') + d(x', s_i) = d(x', s_i)$. Therefore, it can be extended to a left-c.e. non-decreasing function f over $[0, 1]$. Indeed, if $f_0 : [0, 1] \rightarrow \mathbb{R}$ is a left-c.e. function such that $f_0(x) = d(x, s_i)$ for $x \in [\alpha, 1]$, then $f(x) := \inf\{f_0(x') : x' \in [x, 1]\}$ is left-c.e. non-decreasing and agrees with $d(x, s_i)$ on $[\alpha, 1]$.

As a result, for $q \in \mathbb{Q}$, $q > s_i$ if and only if $f(q) > 0$ which is a c.e. condition, so s_i is right-c.e. Of course, s_i is left-c.e. as it is a computable point of $[\alpha, 1]_<$.

Proposition 6.3. *The space $[\alpha, 1]_<$ admits a computably equivalent right-c.e. quasi-metric structure if and only if α is left-c.e. or right-c.e.*

Proof. If α is right-c.e. then there is a computable quasi-metric structure by Proposition 6.2. If α is left-c.e. then we can take $S = \{s_i\}_{i \in \mathbb{N}}$ with $s_i = \max(q_i, \alpha)$, where $(q_i)_{i \in \mathbb{N}}$ is a computable enumeration of the rational numbers in $[0, 1]$. We can take the restriction of the quasi-metric $d(x, y) = \max(0, x - y)$. It is right-c.e. on S . To approximate $d(s_i, s_j)$ from the right, do the following: if $q_i \leq q_j$ then output 0 (correct as $s_i \leq s_j$ in that case). If $q_i > q_j$ then start approximating $d(q_i, s_j)$ from the right (possible as s_j is left-c.e.) and switching to 0 if we eventually see that $q_i < \alpha$.

Conversely, assume a right-c.e. metric structure (d', S) . Given $q \in \mathbb{Q} \cap [\alpha, 1]$, $d'(s_i, q)$ is uniformly right-c.e. Indeed, each such q is a computable point of the right-c.e. quasi-metric space $([\alpha, 1]_<, d', S)$, so by Proposition 5.1, $d'(s_i, q)$ is right-c.e.

Claim. Given $s \in S$, $\epsilon > 0$, one can compute $\delta > 0$ such that $B'(s, \delta) \subseteq B(s, \epsilon)$.

Proof (of the claim). The identity from the quasi-metric space $[\alpha, 1]_<$ to the quasi-metric space is computable, so $B(s, \epsilon)$ is effectively open in $[\alpha, 1]_<$, hence can be expressed as a union of d' -balls. One can find one of them, $B'(t, r)$,

containing s . One has $d'(t, s) < r$ and $d'(t, s)$ is right-c.e., so one can compute $\delta > 0$ such that $d'(t, s) + \delta < r$. One has $B'(s, \delta) \subseteq B'(t, r) \subseteq B(s, \epsilon)$.

Let $\delta_{s, \epsilon}$ be obtained from the previous Claim. Consider the set $E = \{q \in \mathbb{Q} \cap [0, 1] : \exists s \in S, \epsilon > 0, d'(s, q) < \delta_{s, \epsilon} \text{ and } d(s, q) > \epsilon\}$. It is a c.e. set. It is disjoint from $[\alpha, 1]$: if $q \in [\alpha, 1]$ and $q \in B'(s, \delta_{s, \epsilon})$ then $q \in B(s, \epsilon)$. As a result, $\sup E$ is left-c.e. and $\sup E \leq \alpha$. If α is not left-c.e. then $\sup E < \alpha$. As a result, we can fix some rational number q_0 between $\sup E$ and α , and work with rationals above q_0 only, so that they do not belong to E .

Let $F = \{(q, \epsilon) : q \in \mathbb{Q} \cap [q_0, 1], \epsilon > 0, \exists s \in S, \text{ such that } d'(s, q) < \delta_{s, \epsilon}\}$. F is c.e. so $I := \inf\{q + \epsilon : (q, \epsilon) \in F\}$ is right-c.e. If $q > \alpha$ then there must exist $s \in S$ such that $s \leq q$, i.e. $d'(s, q) = 0$, so $(q, \epsilon) \in F$ for every $\epsilon > 0$. As a result, $I \leq \alpha$. If α is not right-c.e. then $I < \alpha$.

Take $(q, \epsilon) \in F$ such that $q + \epsilon < \alpha$. Let $s \in S$ witness that $(q, \epsilon) \in F$. One has $d'(s, q) < \delta_{s, \epsilon}$ and $d(s, q) \geq d(\alpha, q) > \epsilon$ so $q \in E$, giving a contradiction.

Therefore, α is left-c.e. or right-c.e.

Corollary 6.1. *There exists an effective quasi-Polish space which cannot be presented as a right-c.e. quasi-metric space.*

Proof. Take α that is left-c.e. relative to the halting set but neither left-c.e. nor right-c.e., and apply Propositions 6.1 and 6.3.

7 Discussion and open questions

By a *computable Polish space* we mean an effective topological space X induced by a computable complete metric space (X, d, S) [21, 15, 8]. Most of the popular Polish spaces are computable.

By a *computable ω -continuous domain* [1] we mean a pair (X, b) where X is an ω -continuous domain and $b : \mathbb{N} \rightarrow X$ is a numbering of a domain base in X modulo which the approximation relation \ll is c.e. Any computable ω -continuous domain (X, b) has the induced effective base β where $\beta_n = \{x \mid b_n \ll x\}$. Most of the popular ω -continuous domains are computable.

By Theorem 1 in [20], both the computable Polish spaces and computable ω -continuous domains are computable effective images of the Baire space, hence they are effective quasi-Polish, hence the notion of effective quasi-Polish space introduced in this paper is a reasonable candidate for capturing the computable quasi-Polish spaces. By Theorem 4 in [13] (which extends Theorem 4 in [20]), any effective quasi-Polish space satisfies the effective Suslin-Kleene theorem. By Theorem 5 in [20], any effective quasi-Polish space satisfies the effective Hausdorff theorem.

It seems that our search, as well as the independent search in [4] resulted in natural and convincing candidates for capturing the computable quasi-Polish spaces. Nevertheless, many interesting closely related questions remain open. Since the class of quasi-Polish spaces admits at least ten seemingly different characterizations [3], the status of effective analogues of these characterizations deserves additional investigation. In particular, this concerns the characterization

of quasi-Polish spaces as the subspaces of non-compact elements in (ω -algebraic or ω -continuous) domains.

References

1. Abramsky, S., Jung, A.: Domain theory. In: Abramsky, S., Gabbay, D., Maibaum, T.S.E. (eds.) *Handbook of Logic in Computer Science Volume 3*, pp. 1–168. Oxford University Press (1994)
2. Becher, V., Grigorieff, S.: Borel and hausdorff hierarchies in topological spaces of choquet games and their effectivization. *Mathematical Structures in Computer Science* **25**(7), 1490–1519 (2015)
3. de Brecht, M.: Quasi-polish spaces. *Ann. Pure Appl. Logic* **164**(3), 356–381 (2013)
4. de Brecht, M., Pauly, A., Schröder, M.: Overt choice (2019), arXiv: 1902.05926
5. Chen, R.: Notes on quasi-polish spaces (2018), arXiv: 1809.07440
6. Gao, S.: *Invariant Descriptive Set Theory*. CRC Press, New York (2009)
7. Gregoriades, V.: Classes of polish spaces under effective Borel isomorphism. *Memiors of the American Mathematical Society* **240**(1135) (2016)
8. Gregoriades, V., Kispéter, T., Pauly, A.: A comparison of concepts from computable analysis and effective descriptive set theory. *Mathematical Structures in Computer Science* **27**(8), 1414–1436 (2017)
9. Grubba, T., Schröder, M., Weihrauch, K.: Computable metrization. *Math. Log. Q.* **53**(4-5), 381–395 (2007)
10. Hoyrup, M.: Genericity of weakly computable objects. *Theory Comput. Syst.* **60**(3), 396–420 (2017)
11. Kechris, A.S.: *Classical Descriptive Set Theory*. Springer (Jan 1995)
12. Korovina, M.V., Kudinov, O.V.: Towards computability over effectively enumerable topological spaces. *Electr. Notes Theor. Comput. Sci.* **221**, 115–125 (2008)
13. Korovina, M.V., Kudinov, O.V.: On higher effective descriptive set theory. In: Kari, J., Manea, F., Petre, I. (eds.) *Unveiling Dynamics and Complexity - CiE 2017*, Turku, Finland, June 12-16, 2017, Proceedings. *Lecture Notes in Computer Science*, vol. 10307, pp. 282–291. Springer (2017)
14. Louveau, A.: Recursivity and compactness. In: *Higher Set Theory*. vol. 669, pp. 303–337 (1978)
15. Moschovakis, Y.: *Descriptive Set Theory*. *Mathematical Surveys and Monographs*, American Mathematical Society (2009), (Second edition)
16. Rogers, H.J.: *Theory of Recursive Functions and Effective Computability*. MIT Press, Cambridge, MA, USA (1987)
17. Selivanov, V.: On index sets in the Kleene-Mostowski hierarchy. *Trans. Inst. Math.* pp. 135–158 (1982), russian
18. Selivanov, V.L.: Towards a descriptive set theory for domain-like structures. *Theor. Comput. Sci.* **365**(3), 258–282 (2006)
19. Selivanov, V.L.: On the difference hierarchy in countably based t_0 -spaces. *Electr. Notes Theor. Comput. Sci.* **221**, 257–269 (2008)
20. Selivanov, V.L.: Towards the effective descriptive set theory. In: Beckmann, A., Mitrana, V., Soskova, M.I. (eds.) *Evolving Computability - CiE 2015*, Bucharest, Romania, June 29 - July 3, 2015. Proceedings. *Lecture Notes in Computer Science*, vol. 9136, pp. 324–333. Springer (2015)
21. Weihrauch, K.: *Computable Analysis*. Springer, Berlin (2000)