

Semi-conical eigenvalue intersections and the ensemble controllability problem for quantum systems

Nicolas Augier, Ugo Boscain, Mario Sigalotti

► **To cite this version:**

Nicolas Augier, Ugo Boscain, Mario Sigalotti. Semi-conical eigenvalue intersections and the ensemble controllability problem for quantum systems. *Mathematical Control and Related Fields*, AIMS, 2020, 10, pp.877-911. hal-02277653

HAL Id: hal-02277653

<https://hal.inria.fr/hal-02277653>

Submitted on 4 Sep 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Semi-conical eigenvalue intersections and the ensemble controllability problem for quantum systems

Nicolas Augier*, Ugo Boscain^{†§}, Mario Sigalotti^{‡§}

September 5, 2019

Abstract

We study one-parametric perturbations of finite dimensional real Hamiltonians depending on two controls, and we show that generically in the space of Hamiltonians, conical intersections of eigenvalues can degenerate into *semi-conical* intersections of eigenvalues. Then, through the use of normal forms, we study the problem of ensemble controllability between the eigenstates of a generic Hamiltonian.

1 Introduction

Controlling parametrized families of quantum systems with a common control signal is a critical task for many applications in quantum control (see [16] and references therein), notably in Nuclear Magnetic Resonance [17].

For a general closed quantum system under the action of a control u and depending on a parameter z , the corresponding controlled equation is of the form

$$i \frac{d\psi}{dt}(t) = H(u(t), z)\psi(t), \quad \psi(t) \in \mathcal{H}, \quad (1)$$

with $H(u, z)$ self-adjoint on the complex Hilbert space \mathcal{H} for every value of u and z .

The parameter z can be used either to describe a family of physical systems on which acts a common field driven by u or a physical systems for which the value of one parameter is not known precisely.

The controllability properties of systems of this form has been studied both for discrete and continuous sets of parameters. The case of a finite set of systems is characterized in [14]. In [10] the asymptotic ensemble stabization is studied for countable sets of parameters. In [22], [5] a proof of a strong notion of ensemble controllability has been obtained for a two-level system. Numerical ensemble control in the case of a continuum of parameters has been throughly studied for two-level systems [32, 9, 29].

Adiabatic control is a powerful technique which can be used to handle perturbations and uncertainties. One of its main advantages is that it provides explicit and regular control laws. It has hence been successfully applied to obtain control strategies such as the chirp pulses (see, for instance [8, 28]) for spin 1/2 systems with dispersed Larmor frequency. Another nowadays classical application of adiabatic control to ensemble controllability are the so-called counterintuitive pulses for the STIRAP

*CMAP, École Polytechnique, Institut Polytechnique de Paris

†CNRS

‡Inria Paris

§Sorbonne Université, Inria, Université de Paris, CNRS, Laboratoire Jacques-Louis Lions, Paris, France

process [15, 27]. A generalization of this approach has been proposed in [21]. These techniques use in an explicit or implicit way the existence of conical intersections between the eigenvalues of the Hamiltonian corresponding to each of the systems of the ensemble. A conical intersection (also called diabolic point) is a cone-like singularity of the spectrum of $H(u, z)$, seen as a function of the control u (see Figure 1). They are generic in the sense that they are the least degenerate singularities of the spectrum of a Hamiltonian and have been studied since the beginning of quantum mechanics [33]. They play an important role in the context of semiclassical analysis [12, 13]. Adiabatic paths through conical intersections can be used to induce superpositions of eigenstates, as shown in [7] and to obtain tests for exact controllability when \mathcal{H} is finite-dimensional and for approximate controllability when \mathcal{H} is infinite-dimensional [6]. Results for ensemble control beyond the quantum control setting can be found in [2, 20, 23, 26].

In our paper [4] we proposed a framework for the adiabatic ensemble control of a continuum of n -level systems with real Hamiltonian, driven by two controls and having conical intersections between the eigenvalues. The main idea was that, if a system corresponding to a fixed parameter has conical intersections between two eigenvalues, then a small perturbation of the parameter yields a curve of conical intersections, each point of the curve corresponding to exactly one value of the parameter. One can then follow adiabatically such curves in the space of controls and obtain a population transfer between the two levels for the whole ensemble of systems.

The argument sketched above works under the assumption that for all values of the parameter eigenvalue intersections remain conical and follow a smooth curve. These assumptions are satisfied for generic small parametric perturbations. For generic large perturbations it may happen that conicity of eigenvalue intersections is lost at isolated points of the curve. The goal of this paper is to extend the analysis to this case. In particular, we

- characterize typical non-conical intersections and give normal forms for them;
- study the evolution of the system corresponding to adiabatic paths in the space of controls passing through such intersections;
- conclude on the ensemble controllability of generic one-parameter systems presenting typical intersections (conical and non-conical).

The results of this paper concern n -level systems with a real Hamiltonian and they can be generalized to systems evolving in an infinite-dimensional Hilbert space. As explained in Section 6, thanks to the adiabatic decoupling, their study can be reduced to the case of zero-trace two-level systems. Consider then an equation of the form

$$i \frac{d\psi}{dt}(t) = H(u(t), v(t))\psi(t), \quad \psi(t) \in \mathbb{C}^2, \quad (u(t), v(t)) \in \mathbb{R}^2, \quad (2)$$

with

$$H(u, v) = \begin{pmatrix} f_1(u, v) & f_2(u, v) \\ f_2(u, v) & -f_1(u, v) \end{pmatrix}, \quad f = (f_1, f_2) \in C^\infty(\mathbb{R}^2, \mathbb{R}^2).$$

As in [4], we restrict our attention to real Hamiltonians, which are relevant in many physical systems, for instance for Galerkin approximations of the Schrödinger equation $i\partial_t\psi(x, t) = (-\Delta + V(x) + u(t)W(x))\psi(x, t)$, where x belongs to a bounded set of \mathbb{R}^n and V, W are regular enough real functions. The spectrum of $H(u, v)$ is $\{\pm\sqrt{f_1(u, v)^2 + f_2(u, v)^2}\}$ and, in particular, it is degenerate if and only if $f(u, v) = (0, 0)$, that is, if (u, v) is an *eigenvalue intersection*. Denote by $\lambda^+(u, v) = \sqrt{f_1(u, v)^2 + f_2(u, v)^2}$ the largest eigenvalue of $H(u, v)$ and notice that the gap (denoted $\text{Gap}(u, v)$) between the two eigenvalues of $H(u, v)$ is equal to $2\lambda^+(u, v)$. An eigenvalue intersection (u, v) is said to be *conical* if

$$\chi(f) := \det(\nabla f_1, \nabla f_2)$$

is nonzero at (u, v) , where ∇ denotes the gradient with respect to the variables u and v , and *semi-conical* if $\nabla f_1(u, v)$ and $\nabla f_2(u, v)$ are collinear, are not both zero, and the directional derivative $\partial_\eta \chi(f)$ along $\eta = (-\partial_2 f_j(u, v), \partial_1 f_j(u, v))$ is nonzero if $j \in \{1, 2\}$ is such that $\eta \neq 0$. The direction spanned by η is called the *non-conical direction* at (u, v) . If (u, v) is a conical intersection, then

$$\frac{1}{C} \|(u', v') - (u, v)\| \leq \text{Gap}(u', v') \leq C \|(u', v') - (u, v)\| \quad (3)$$

for some $C > 0$ and for all (u', v') in a neighborhood of (u, v) . If, instead, (u, v) is a semi-conical intersection, then an inequality of the type (3) holds along any line passing through (u, v) in a direction transversal to the non-conical direction. Along the non-conical direction η we have

$$\frac{1}{C} t^2 \leq \text{Gap}((u, v) + t\eta) \leq C t^2$$

for some $C > 0$ and for all t in a neighborhood of 0.




Figure 1: Conical intersection as a function of the controls $(u, v) \in \mathbb{R}^2$.




Figure 2: Semi-conical intersection of eigenvalues as a function of the controls $(u, v) \in \mathbb{R}^2$.

Intersections with the previous properties appear for instance in STIRAP processes with two successive states having the same energy, that is, when, for $(u, v) \in \mathbb{R}^2$, $H(u, v) = \begin{pmatrix} E & u & 0 \\ u & E & v \\ 0 & v & E' \end{pmatrix}$, where $E < E'$. On Figure 3, we have plotted the spectrum of $H(u, v)$ as a function of (u, v) for such a Hamiltonian. We notice graphically that there is a semi-conical intersection between the first and second levels, and two conical intersections between the second and third levels.

Let us now consider a one-parameter family of two-level systems as above, that is, the *Ensemble Schrödinger Equation*

$$i \frac{d\psi}{dt}(t) = H(u(t), v(t), z)\psi(t), \quad \psi(t) \in \mathbb{C}^2, \quad (u(t), v(t)) \in \mathbb{R}^2, \quad (4)$$

with

$$H(u, v, z) = \begin{pmatrix} f_1(u, v, z) & f_2(u, v, z) \\ f_2(u, v, z) & -f_1(u, v, z) \end{pmatrix}, \quad f = (f_1, f_2) \in C^\infty(\mathbb{R}^3, \mathbb{R}^2).$$

The spectrum of $H(u, v, z)$ is $\{\lambda^\pm(u, v, z) = \pm \sqrt{f_1(u, v, z)^2 + f_2(u, v, z)^2}\}$ and, in particular, it is degenerate if and only if $f(u, v, z) = (0, 0)$. In order to extend the definition of conical and semi-conical intersections for a one-parameter Hamiltonian, we need to add to the previous definitions




Figure 3: Semi-conical intersection for the STIRAP as a function of the controls $(u, v) \in \mathbb{R}^2$.

some regularity assumptions with respect to the perturbation parameter z . Let (u, v, z) be a point such that $f(u, v, z) = (0, 0)$. It is said to be conical for the family (*F-conical*) if (u, v) is conical for $f(\cdot, \cdot, z)$ and $\partial_3 f(u, v, z) \neq (0, 0)$. It is said to be semi-conical for the family (*F-semi-conical*) if it is semi-conical for $f(\cdot, \cdot, z)$ and f is a submersion at (u, v, z) . The requirement that f is a submersion guarantees that the set

$$Z(f) = \{(u, v, z) \mid f(u, v, z) = (0, 0)\}$$

is a smooth curve in the neighborhood of a semi-conical point. In the following we denote by $Z_{\text{nc}}(f)$ the set of non-conical intersections in $Z(f)$. The following lemma is a consequence of the results in Section 3.

Lemma 1.1. *Let $(\bar{u}, \bar{v}, \bar{z})$ be a *F-semi-conical* intersection. Then $(\bar{u}, \bar{v}, \bar{z})$ is isolated in $Z_{\text{nc}}(f)$, $Z(f)$ is a smooth curve locally near $(\bar{u}, \bar{v}, \bar{z})$, whose tangent is not vertical. Moreover, the tangent at (\bar{u}, \bar{v}) to the projection on the plane (u, v) of such a curve coincides with the non-conical direction corresponding to $(\bar{u}, \bar{v}, \bar{z})$.*

We focus in what follows on generic properties for systems of the type (4). This means that we look for properties which hold for all f in a subset of $C^\infty(\mathbb{R}^3, \mathbb{R}^2)$ with “small” complement. For a precise definition, we refer to Section 2.1.

Theorem 1.2. *Generically with respect to $f \in C^\infty(\mathbb{R}^3, \mathbb{R}^2)$, for any connected component γ of $Z(f)$,*

- (i) γ is a one-dimensional submanifold of \mathbb{R}^3 ;
- (ii) The projection $\pi(\gamma)$ of γ on the plane (u, v) is a C^∞ embedded curve of \mathbb{R}^2 ;
- (iii) $(Z(f) \setminus Z_{\text{nc}}(f)) \cap \gamma$ is made of *F-conical* intersections and $Z_{\text{nc}}(f) \cap \gamma$ is made of *F-semi-conical* intersections only.

The following theorem resumes the main properties of the control strategy that we study in the paper.

Theorem 1.3. *Assume that $Z(f)$ has a single connected component γ . Assume moreover that*

1. γ satisfies properties (i)–(iii) of Theorem 1.2;
2. $\pi(\gamma)$ has no self-intersections.

Take two conical intersections $(u_0, v_0, z_0), (u_1, v_1, z_1)$ in γ with $z_0 < z_1$. Consider a regular C^4 path $(u, v) : [0, 1] \rightarrow \mathbb{R}^2$ such that $(u, v)(t_0) = (u_0, v_0)$, $(u, v)(t_1) = (u_1, v_1)$ for some $0 < t_0 < t_1 < 1$. Assume, moreover, that $(u, v)(0) = (u, v)(1) =: (\bar{u}, \bar{v})$, that $(u, v)(t) \in \pi(\gamma)$ if and only if $t \in [t_0, t_1]$, that $z \in [z_0, z_1]$ for every z and t such that $(u(t), v(t), z) \in \gamma$. For every $z \in [z_0, z_1]$, let ϕ_-^z and ϕ_+^z be two normalized eigenvectors of $H(\bar{u}, \bar{v}, z)$ corresponding to $\lambda^-(\bar{u}, \bar{v}, z)$ and $\lambda^+(\bar{u}, \bar{v}, z)$, respectively. Then there exists $C > 0$ such that for every $z \in [z_0, z_1]$ and every $\epsilon > 0$, the solution ψ of $i\dot{\psi}(t) = H_f(u(\epsilon t), v(\epsilon t), z)\psi(t)$, $\psi(0) = \phi_-^z$, satisfies

$$\left\| \psi \left(\frac{1}{\epsilon} \right) - e^{i\xi} \phi_+^z \right\| < C\epsilon^{\frac{1}{3}},$$

for some $\xi \in \mathbb{R}$, possibly depending on ϵ and z .




Figure 4: A curve (u, v) as in the statement of Theorem 1.3

The paper is organized as follows: In Section 2 we provide normal forms for the Hamiltonians yielding the different types of eigenvalue intersections introduced above. In Section 3, we prove Theorem 1.2 and we study the singularities of the projection $\pi(f)$ of $Z(f)$ on the control plane. In Section 4, we study the dynamical properties of an isolated semi-conical intersection of eigenvalues and we prove Theorem 1.3. Then, in Section 6, we extend Theorem 1.3 to systems with more than two levels.

2 Basic facts and normal forms

2.1 Generic families of 2-level Hamiltonians

Consider a smooth function $f = (f_1, f_2) : \mathbb{R}^2 \times \mathbb{R}^l \rightarrow \mathbb{R}^2$ with $l = 0$ or $l = 1$. Denote by (e_1, \dots, e_{2+l}) the canonical basis of \mathbb{R}^{2+l} . Given a vector $\eta \in \mathbb{R}^{2+l}$ and a smooth function $g : \mathbb{R}^{2+l} \rightarrow \mathbb{R}^q$, $q \in \mathbb{N}$, we write $\partial_\eta g$ for the directional derivative of g in the direction η and ∂_i for ∂_{e_i} , $i = 1, \dots, 2+l$. For $x \in \mathbb{R}^{2+l}$ and $h \in T_x \mathbb{R}^{2+l} \approx \mathbb{R}^{2+l}$, denote the differential of f at x applied to h by $Df_x(h)$.

In the following, we study generic situations in the cases $l = 0$ and $l = 1$. The coordinates (x_1, x_2) play the role of controls, and are denoted by (u, v) , while—in the case $l = 1$ —the coordinate x_3 is a parameter and is denoted by z . The space $C^\infty(\mathbb{R}^{2+l}, \mathbb{R}^2)$ is endowed in what follows with the C^∞ -Whitney topology. We say that a property satisfied by $f \in C^\infty(\mathbb{R}^{2+l}, \mathbb{R}^2)$ is *generic* if it is satisfied in an open and dense subset of the space $C^\infty(\mathbb{R}^{2+l}, \mathbb{R}^2)$.

2.1.1 The single system case $l = 0$

Consider a two-dimensional real Hamiltonian of the form

$$H_f(u, v) = \begin{pmatrix} f_1(u, v) & f_2(u, v) \\ f_2(u, v) & -f_1(u, v) \end{pmatrix},$$

where $f_1, f_2 : \mathbb{R}^2 \rightarrow \mathbb{R}$ are smooth functions depending on 2 control variables (u, v) and $f = (f_1, f_2)$. Denote by $\chi(f)(u, v)$ the Jacobian of f . Notice that the eigenvalues of H_f are $\lambda^+ = \sqrt{f_1^2 + f_2^2}$ and $\lambda^- = -\sqrt{f_1^2 + f_2^2}$. Define $\text{Gap} = \lambda^+ - \lambda^- = 2\sqrt{f_1^2 + f_2^2}$.

Definition 2.1. Consider $f \in C^\infty(\mathbb{R}^2, \mathbb{R}^2)$ and let $(\bar{u}, \bar{v}) \in \mathbb{R}^2$ be such that $f(\bar{u}, \bar{v}) = (0, 0)$. We say that the eigenvalue intersection (\bar{u}, \bar{v}) is

- conical in the direction $\nu \in \mathbb{R}^2$ if $\partial_\nu f(\bar{u}, \bar{v}) \neq (0, 0)$;
- conical if $\chi(f)(\bar{u}, \bar{v}) \neq 0$;
- semi-conical if $\nabla f_1(\bar{u}, \bar{v})$ and $\nabla f_2(\bar{u}, \bar{v})$ are collinear, are not both zero, and the directional derivative of $\chi(f)(\bar{u}, \bar{v})$ along $\eta = (-\partial_2 f_j(\bar{u}, \bar{v}), \partial_1 f_j(\bar{u}, \bar{v}))$ is nonzero if $j \in \{1, 2\}$ is such that $\eta \neq 0$. The direction spanned by η is called the non-conical direction at (\bar{u}, \bar{v}) .

Remark 2.2. The definition of conical intersection for $f \in C^\infty(\mathbb{R}^2, \mathbb{R}^2)$ given above is equivalent to the one used in [7], namely, a point (\bar{u}, \bar{v}) such that $f(\bar{u}, \bar{v}) = (0, 0)$ and there exists $c > 0$ such that, for every $\eta \in \mathbb{R}^2$ of norm 1 and $\delta > 0$ small enough, we have $\frac{\text{Gap}((\bar{u}, \bar{v}) + \delta\eta)}{\delta} \geq c$.

The following proposition states that semi-conical points are isolated zeros of f . Although the proof could be deduced rather straightforwardly from the definition, we postpone it for simplicity to Section 2.3, where we base it on normal forms.

Proposition 2.3 (Semi-conical intersections are isolated). Consider a semi-conical point (\bar{u}, \bar{v}) for f . Then there exists a neighborhood V of (\bar{u}, \bar{v}) in \mathbb{R}^2 such that for every $(u, v) \in V \setminus \{(\bar{u}, \bar{v})\}$, $f(u, v) \neq (0, 0)$.

We introduce here the transversality argument used in the following of the section to prove genericity of several properties. As an illustration, we recall how such an argument can be used to prove that conical intersections are generic. Denote by $J^1(\mathbb{R}^2, \mathbb{R}^2)$ the space of 1-jets of functions from \mathbb{R}^2 to \mathbb{R}^2 . For every $(u, v) \in \mathbb{R}^2$ and $f \in C^1(\mathbb{R}^2, \mathbb{R}^2)$, we write $j^1(f)(u, v) \in J^1(\mathbb{R}^2, \mathbb{R}^2)$ to denote the 1-jet of f at (u, v) . Define

$$S_r = \{j^1(f)(0) \mid f \in C^\infty(\mathbb{R}^2, \mathbb{R}^2) \text{ } f(0) = 0, \text{ rank}(Df(0)) = r\}, \quad r = 0, 1.$$

It is easy to check that S_0, S_1 are two submanifolds of $J^1(\mathbb{R}^2, \mathbb{R}^2)$ of codimension 6 and 3, respectively. One can easily show that the algebraic subset $S_0 \cup S_1$ of $J^1(\mathbb{R}^2, \mathbb{R}^2)$ admits a Whitney stratification (see [19] Part I, Chapter 1) whose strata have codimension strictly larger than the dimension of \mathbb{R}^2 . By Thom's transversality theorem (see, e.g., [18]) used in combination with [19, §1.3.2],

$$\begin{aligned} U &= \{f \in C^\infty(\mathbb{R}^2, \mathbb{R}^2) \mid j^1(f)(\mathbb{R}^2) \cap (S_1 \cup S_2) = \emptyset\} \\ &= \{f \in C^\infty(\mathbb{R}^2, \mathbb{R}^2) \mid j^1(f)(\mathbb{R}^2) \cap S_1 = \emptyset\} \cap \{f \in C^\infty(\mathbb{R}^2, \mathbb{R}^2) \mid j^1(f)(\mathbb{R}^2) \cap S_2 = \emptyset\} \end{aligned}$$

is an open and dense subset of $C^\infty(\mathbb{R}^2, \mathbb{R}^2)$.

2.1.2 The ensemble case $l = 1$

Consider a two-dimensional real Hamiltonian of the form

$$H_f(u, v, z) = \begin{pmatrix} f_1(u, v, z) & f_2(u, v, z) \\ f_2(u, v, z) & -f_1(u, v, z) \end{pmatrix},$$

where $f_1, f_2 : \mathbb{R}^3 \rightarrow \mathbb{R}$ are smooth functions depending on two control variables (u, v) and one parameter z . Define the smooth function $f = (f_1, f_2) : \mathbb{R}^3 \rightarrow \mathbb{R}^2$. An *eigenvalue intersection* is a point (u, v, z) such that $f(u, v, z) = (0, 0)$.

Definition 2.4. For $i, j \in \{1, 2, 3\}$, let $\chi_{ij}(f)$ be the Jacobian of the restriction of f to the plane parallel to $\text{span}(e_i, e_j)$, i.e.,

$$\chi_{ij}(f)(u, v, z) = \begin{vmatrix} \partial_i f_1(u, v, z) & \partial_j f_1(u, v, z) \\ \partial_i f_2(u, v, z) & \partial_j f_2(u, v, z) \end{vmatrix}.$$

By a slight abuse of notation, we set $\chi(f) = \chi_{12}(f)$.

In order to extend the definition of conical and semi-conical intersections to parametrized Hamiltonians, we need to add to the previous definitions some regularity assumptions with respect to the parameter z .

Definition 2.5. Let (u, v, z) be a point such that $f(u, v, z) = (0, 0)$. It is said to be *conical* for the family (F-conical) if (u, v) is conical for $f(\cdot, \cdot, z)$ and $\partial_3 f(u, v, z) \neq (0, 0)$. It is said to be *semi-conical* for the family (F-semi-conical) if it is semi-conical for $f(\cdot, \cdot, z)$ and f is a submersion at (u, v, z) .

Proposition 2.6. Generically with respect to $f \in C^\infty(\mathbb{R}^3, \mathbb{R}^2)$, f is a submersion at every point of $Z(f)$ and the set $Z(f) = \{(u, v, z) \in \mathbb{R}^3 \mid f(u, v, z) = 0\}$ is a submanifold of \mathbb{R}^3 of dimension 1.

Proof. Define

$$\Sigma_r = \{j^1(f)(0) \in J^1(\mathbb{R}^3, \mathbb{R}^2) \mid f \in C^\infty(\mathbb{R}^3, \mathbb{R}^2), f(0) = 0, \text{rank}(Df(0)) = r\}, \quad r = 0, 1.$$

Notice that Σ_0 and Σ_1 are smooth submanifolds of $J^1(\mathbb{R}^3, \mathbb{R}^2)$ of codimensions 8 and 4, respectively. One can easily show that the algebraic subset $\Sigma_0 \cup \Sigma_1$ of $J^1(\mathbb{R}^3, \mathbb{R}^2)$ admits a Whitney stratification whose strata have codimension strictly larger than 3. Transversality theory then allows to conclude that $U = \{f \in C^\infty(\mathbb{R}^3, \mathbb{R}^2) \mid j^1(f)(\mathbb{R}^3) \cap (\Sigma_0 \cup \Sigma_1) = \emptyset\}$ is an open and dense subset of $C^\infty(\mathbb{R}^3, \mathbb{R}^2)$. Hence, f is generically a submersion at every point $(u, v, z) \in Z(f)$. The proposition is proved. \square

In the next two propositions we provide a geometric description of the curve $Z(f)$ and we show its links with the conicity properties of f .

Proposition 2.7. *A point $(\bar{u}, \bar{v}, \bar{z}) \in Z(f)$ is conical for $f(\cdot, \cdot, \bar{z})$ if and only if f is a submersion at $(\bar{u}, \bar{v}, \bar{z})$ such that $Z(f)$ is locally near $(\bar{u}, \bar{v}, \bar{z})$ a one-dimensional submanifold transversal to the plane of \mathbb{R}^3 of equation $z = \bar{z}$.*

Proof. Let $(\bar{u}, \bar{v}, \bar{z}) \in Z(f)$ be conical for $f(\cdot, \cdot, \bar{z})$. By definition, we have $\chi(f)(\bar{u}, \bar{v}, \bar{z}) \neq 0$, hence f is a submersion at $(\bar{u}, \bar{v}, \bar{z})$. It follows that $Z(f)$ is locally near $(\bar{u}, \bar{v}, \bar{z})$ a one-dimensional submanifold of \mathbb{R}^3 . Fix $\bar{t} \in \mathbb{R}$ and a local smooth regular parametrization $c(t) = (u(t), v(t), z(t))_{t \in \mathbb{R}}$ of $c \subset Z(f)$ such that $c(\bar{t}) = (\bar{u}, \bar{v}, \bar{z})$. Assume for the sake of contradiction that $\dot{z}(\bar{t}) = 0$. Differentiating the condition $f(c(t)) = 0$, we have

$$\begin{cases} \dot{u}(\bar{t})\partial_1 f_1(\bar{u}, \bar{v}, \bar{z}) + \dot{v}(\bar{t})\partial_2 f_1(\bar{u}, \bar{v}, \bar{z}) = 0 \\ \dot{u}(\bar{t})\partial_1 f_2(\bar{u}, \bar{v}, \bar{z}) + \dot{v}(\bar{t})\partial_2 f_2(\bar{u}, \bar{v}, \bar{z}) = 0 \end{cases} \quad (5)$$

Hence $\chi(f)(\bar{u}, \bar{v}, \bar{z}) = 0$, that is impossible.

Conversely, consider a submersion f at $(\bar{u}, \bar{v}, \bar{z})$ such that $Z(f)$ is locally near $(\bar{u}, \bar{v}, \bar{z})$ a one-dimensional submanifold transversal to the plane of \mathbb{R}^3 of equation $z = \bar{z}$. For the sake of contradiction, assume that $(\bar{u}, \bar{v}, \bar{z})$ is non conical for $f(\cdot, \cdot, \bar{z})$. By definition, there exists a direction $\eta \in \mathbb{R}^2 \setminus \{(0, 0)\}$ such that $\partial_{(\eta, 0)} f(\bar{u}, \bar{v}, \bar{z}) = 0$. Fix $\bar{t} \in \mathbb{R}$ and a local smooth regular parametrization $c(t) = (u(t), v(t), z(t))_{t \in \mathbb{R}}$ of $c \subset Z(f)$ such that $c(\bar{t}) = (\bar{u}, \bar{v}, \bar{z})$. Differentiating the condition $f(c(t)) = 0$, we have $(\dot{u}(\bar{t}), \dot{v}(\bar{t}), \dot{z}(\bar{t})) \in \ker Df_{(\bar{u}, \bar{v}, \bar{z})}$. Since f is a submersion at $(\bar{u}, \bar{v}, \bar{z})$, we deduce that $(\dot{u}(\bar{t}), \dot{v}(\bar{t}), \dot{z}(\bar{t}))$ is collinear to $(\eta, 0)$. Hence we get $\dot{z}(\bar{t}) = 0$, which is impossible. \square

Proposition 2.8. *Assume that f is a submersion locally near $(\bar{u}, \bar{v}, \bar{z})$ and that $(\bar{u}, \bar{v}, \bar{z})$ is non-conical for $f(\cdot, \cdot, \bar{z})$ in the direction $\eta \in \mathbb{R}^2 \setminus \{(0, 0)\}$. Fix $\bar{t} \in \mathbb{R}$ and a local smooth regular parametrization $c(t) = (u(t), v(t), z(t))_{t \in \mathbb{R}}$ of $c \subset Z(f)$ such that $c(\bar{t}) = (\bar{u}, \bar{v}, \bar{z})$ and $\dot{z}(\bar{t}) = 0$. Then we have the equivalence*

$$\ddot{z}(\bar{t}) = 0 \iff \partial_{(\eta, 0)} \chi(f)(\bar{u}, \bar{v}, \bar{z}) = 0.$$

In particular, if $(\bar{u}, \bar{v}, \bar{z})$ is F -semi-conical for f then $\ddot{z}(\bar{t}) \neq 0$.

Proof. Without loss of generality, we can assume $(\bar{u}, \bar{v}, \bar{z}) = (0, 0, 0)$. Under the assumption that f is a submersion, we have $(\partial_1 f_1(0), \partial_2 f_1(0)) \neq (0, 0)$ or $(\partial_1 f_2(0), \partial_2 f_2(0)) \neq (0, 0)$. Without loss of generality, assume $\partial_1 f_1(0) = r \cos(\theta)$ and $\partial_2 f_1(0) = r \sin(\theta)$ where $r > 0$ and $\theta \in [0, 2\pi]$. Define the

matrix $R_\theta = \begin{pmatrix} -\sin(\theta) & -\cos(\theta) & 0 \\ \cos(\theta) & -\sin(\theta) & 0 \\ 0 & 0 & 1 \end{pmatrix}$. For every $(u, v, z) \in \mathbb{R}^3$, define $\tilde{f}(u, v, z) = (f \circ R_\theta)(u, v, z)$.

By simple computations, we have $\partial_1 \tilde{f}(0) = 0$ and $\partial_1 \chi(\tilde{f}) = \partial_{(\eta, 0)} \chi(f)$. Hence, it is sufficient to prove the proposition for $\eta = (1, 0)$.

Notice that

$$\partial_1 \chi(f)(0) = \begin{vmatrix} \partial_{11} f_1(0) & \partial_{21} f_1(0) \\ \partial_{11} f_2(0) & \partial_{21} f_2(0) \end{vmatrix} + \begin{vmatrix} \partial_1 f_1(0) & \partial_{21} f_1(0) \\ \partial_1 f_2(0) & \partial_{21} f_2(0) \end{vmatrix} = \begin{vmatrix} \partial_{11} f_1(0) & \partial_{21} f_1(0) \\ \partial_{11} f_2(0) & \partial_{21} f_2(0) \end{vmatrix}, \quad (6)$$

where we used that $\partial_1 f_1(0) = \partial_1 f_2(0) = 0$.

Since f is a submersion at 0, the equality $\left. \frac{d}{dt} f(c(t)) \right|_{t=\bar{t}} = 0$ implies that $\dot{c}(\bar{t}) = a e_1$ for some $a \neq 0$. The equality $\left. \frac{d^2}{dt^2} f(c(t)) \right|_{t=\bar{t}} = 0$ can be rewritten as $Df_{c(\bar{t})}(\ddot{c}(\bar{t})) + D^2 f_{c(\bar{t})}(\dot{c}(\bar{t}), \dot{c}(\bar{t})) = 0$, that is, for $t = \bar{t}$,

$$\begin{cases} \ddot{y}(\bar{t})\partial_2 f_1(0) + \ddot{z}(\bar{t})\partial_3 f_1(0) + a^2 \partial_{11} f_1(0) = 0, \\ \ddot{y}(\bar{t})\partial_2 f_2(0) + \ddot{z}(\bar{t})\partial_3 f_2(0) + a^2 \partial_{11} f_2(0) = 0, \end{cases}$$

which can be rewritten as

$$\begin{pmatrix} \partial_{11}f_1(0) & \partial_2f_1(0) \\ \partial_{11}f_2(0) & \partial_2f_2(0) \end{pmatrix} \begin{pmatrix} a^2 \\ \ddot{y}(\bar{t}) \end{pmatrix} = -\ddot{z}(\bar{t})\partial_3f(0). \quad (7)$$

Since f is a submersion at 0 and $\partial_1f(0) = 0$, we have that $\partial_3f(0)$ is nonzero. The conclusion then follows from (6) and (7). \square

Remark 2.9. *If $(\bar{u}, \bar{v}, \bar{z})$ is F -semi-conical and $(u(t), v(t), z(t))_{t \in \mathbb{R}}$ such that $(u(\bar{t}), v(\bar{t}), z(\bar{t})) = (\bar{u}, \bar{v}, \bar{z})$ is a smooth and regular local parametrization of $Z(f)$, then $\dot{z}(\bar{t}) = 0$ and $\ddot{z}(\bar{t}) \neq 0$. In particular, F -semi-conical intersections are isolated in \mathbb{R}^3 .*

The following two propositions guarantee that for a generic f , all intersections are either F -conical or F -semi-conical.

Proposition 2.10. *For a generic $f \in C^\infty(\mathbb{R}^3, \mathbb{R}^2)$, for every $(u, v, z) \in \mathbb{R}^3$ such that (u, v, z) is a conical intersection for $f(\cdot, \cdot, z)$, we have that (u, v, z) is a F -conical intersection for f .*

Proof. The set $Q = \{j^1(f)(0) \in J^1(\mathbb{R}^3, \mathbb{R}^2) \mid f(0) = 0, \partial_3f(0) = 0\}$ is a Whitney stratified subset of $J^1(\mathbb{R}^3, \mathbb{R}^2)$ of codimension 4. By transversality theory the set $\{f \in C^\infty(\mathbb{R}^3, \mathbb{R}^2) \mid j^1(f)(\mathbb{R}^3) \cap Q = \emptyset\}$ is an open and dense subset of $C^\infty(\mathbb{R}^3, \mathbb{R}^2)$. \square

Proposition 2.11. *For a generic $f \in C^\infty(\mathbb{R}^3, \mathbb{R}^2)$, for every $(u, v, z) \in \mathbb{R}^3$ such that (u, v, z) is a non-conical intersection for $f(\cdot, \cdot, z)$, we have that (u, v, z) is a F -semi-conical intersection for f .*

Proof. Set

$$S_j = \{j^2(f)(0) \in J^2(\mathbb{R}^3, \mathbb{R}^2) \mid f(0) = (0, 0), \partial_1f_j(0) = \partial_2f_j(0) = 0\}, \quad j = 1, 2.$$

Then S_1 and S_2 are smooth subspaces of $J^2(\mathbb{R}^3, \mathbb{R}^2)$ of codimension 4. Define

$$\eta = (-\partial_2f_1(0), \partial_1f_1(0), 0),$$

$$S_3 = \{j^2(f)(0) \in J^2(\mathbb{R}^3, \mathbb{R}^2) \mid f(0) = 0, (\partial_1f_1(0), \partial_2f_1(0)) \neq 0, \chi(f)(0) = 0, \partial_\eta\chi(f)(0) = 0\}$$

and

$$\tilde{S}_3 = \{j^2(f)(0) \in J^2(\mathbb{R}^3, \mathbb{R}^2) \mid f(0) = 0, \chi(f)(0) = 0, \partial_\eta\chi(f)(0) = 0\}.$$

We are going to prove that S_3 is a smooth submanifold of $J^2(\mathbb{R}^3, \mathbb{R}^2)$ of codimension 4, that is, that the equalities $f(0) = 0$, $\chi(f)(0) = 0$ and $\partial_\eta\chi(f)(0) = 0$ define independent equations in $J^2(\mathbb{R}^3, \mathbb{R}^2)$ under the condition $(\partial_1f_1(0), \partial_2f_1(0)) \neq 0$. The equality $f(0) = 0$ is clearly independent from the two others. Using the property that $\partial_1f_1(0) \neq 0$ or $\partial_2f_1(0) \neq 0$, one easily establishes that $\chi(f)(0) = 0$ and $\partial_\eta\chi(f)(0) = 0$ define independent equations in $J^2(\mathbb{R}^3, \mathbb{R}^2)$.

One then can easily prove that the algebraic subset $S = S_1 \cup S_2 \cup \tilde{S}_3 = S_1 \cup S_2 \cup S_3$ of $J^2(\mathbb{R}^3, \mathbb{R}^2)$ admits a Whitney stratification whose strata have a codimension strictly larger than 3. By transversality theory we get that $O = \{f \in C^\infty(\mathbb{R}^3, \mathbb{R}^2) \mid j^2(f)(\mathbb{R}^3) \cap S = \emptyset\}$ is an open and dense subset of $C^\infty(\mathbb{R}^3, \mathbb{R}^2)$. \square

2.2 Admissible transformations

The aim of this section is to define admissible transformations in order to get normal forms for the Hamiltonians $H_f = \begin{pmatrix} f_1 & f_2 \\ f_2 & -f_1 \end{pmatrix}$ defined for $f \in C^\infty(\mathbb{R}^2, \mathbb{R}^2)$ and $f \in C^\infty(\mathbb{R}^3, \mathbb{R}^2)$. Consider the *Schrödinger Equation*, defined for $f \in C^\infty(\mathbb{R}^2, \mathbb{R}^2)$ by

$$i \frac{d\psi(t)}{dt} = H_f(u(t), v(t))\psi(t), \quad \psi(t) \in \mathbb{C}^2, \quad (8)$$

and the *Ensemble Schrödinger Equation*, defined for $f \in C^\infty(\mathbb{R}^3, \mathbb{R}^2)$ by

$$i \frac{d\psi(t)}{dt} = H_f(u(t), v(t), z)\psi(t), \quad \psi(t) \in \mathbb{C}^2. \quad (9)$$

The control functions u, v are in $L^\infty(\mathbb{R}, \mathbb{R})$ and the perturbation z belongs to $[z_0, z_1]$ where $z_0, z_1 \in \mathbb{R}$.

The three transformations correspond to equivalent representations of the dynamical systems (8) and (9) achieved, respectively, by time-reparameterization, state-space diffeomorphism, and independent diffeomorphic transformations of both the space of controls and the space of perturbations.

Definition 2.12. We say that two elements f and \tilde{f} of $C^\infty(\mathbb{R}^2, \mathbb{R}^2)$ (respectively, $C^\infty(\mathbb{R}^3, \mathbb{R}^2)$) are time-equivalent at 0 if there exists a nowhere-vanishing function $\xi \in C^\infty(\mathbb{R}^2, \mathbb{R})$ such that $\tilde{f}(u, v) = \xi(u, v)f(u, v)$ (respectively, $\tilde{f}(u, v, z) = \xi(u, v)f(u, v, z)$) in a neighborhood of 0.

Remark 2.13. A time-equivalence as introduced in Definition 2.12 with $\xi > 0$ corresponds to a time-change in Equation (8). As for the case $\xi < 0$, consider $f \in C^\infty(\mathbb{R}^2, \mathbb{R}^2)$, $\psi_0, \psi_1 \in \mathbb{C}^2$ and a control path $(u(\cdot), v(\cdot))$ such that the solution $\psi : [0, T] \rightarrow \mathbb{C}^2$ of Equation (8) with $\psi(0) = \psi_0$ satisfies $\psi(T) = \psi_1$. Then the solution $\tilde{\psi}$ of Equation (8) associated with $(u(T - \cdot), v(T - \cdot))$ such that $\tilde{\psi}(0) = \tilde{\psi}_1$ satisfies $\tilde{\psi}(T) = \tilde{\psi}_0$ (where we denote by \bar{x} the complex-conjugate of $x \in \mathbb{C}^2$). Hence the equations (8) and

$$i \frac{d\psi(t)}{dt} = -H_f(u(t), v(t))\psi(t), \quad \psi(t) \in \mathbb{C}^2,$$

have the same controllability properties. Hence time-equivalence is justified for a function $\xi \in C^\infty(\mathbb{R}^2, (-\infty, 0))$. The same argument is also valid for $f \in C^\infty(\mathbb{R}^3, \mathbb{R}^2)$.

Definition 2.14. We say that two elements f and \tilde{f} of $C^\infty(\mathbb{R}^3, \mathbb{R}^2)$ or $C^\infty(\mathbb{R}^2, \mathbb{R}^2)$ are left-equivalent if there exists $P \in O_2(\mathbb{R})$ independent of u, v, z such that $H_f = PH_{\tilde{f}}P^{-1}$.

Remark 2.15. Let f be in $C^\infty(\mathbb{R}^3, \mathbb{R}^2)$ or $C^\infty(\mathbb{R}^2, \mathbb{R}^2)$. Considering

$$P_{\theta, \zeta} = \begin{pmatrix} \cos(\theta) & -\zeta \sin(\theta) \\ \sin(\theta) & \zeta \cos(\theta) \end{pmatrix} \in O_2(\mathbb{R}),$$

where $\theta \in \mathbb{S}^1$ and $\zeta = \pm 1$, the associated left-equivalence transforms $f = (f_1, f_2)$ into

$$\tilde{f} = (\cos(2\theta)f_1 - \zeta \sin(2\theta)f_2, \zeta \cos(2\theta)f_2 + \sin(2\theta)f_1).$$

Remark 2.16. Let f be in $C^\infty(\mathbb{R}^3, \mathbb{R}^2)$ or $C^\infty(\mathbb{R}^2, \mathbb{R}^2)$. If $t \mapsto \psi(t)$ is a solution of Equation (8) or (9) associated with f and with initial condition $\psi(0) = \psi_0$, then $t \mapsto Y(t) = P\psi(t)$ is a solution of Equation (9) associated with \tilde{f} satisfying $Y(0) = P\psi_0$. Hence, transitions for Y between the eigenstates of $H_{\tilde{f}} = PH_fP^{-1}$ correspond to transitions for ψ between the eigenstates of H_f .

Definition 2.17. We say that two elements f and \tilde{f} of $C^\infty(\mathbb{R}^2, \mathbb{R}^2)$ are right-equivalent at 0 if there exists a diffeomorphism $\phi \in C^\infty(\mathbb{R}^2, \mathbb{R}^2)$ such that $\phi(0) = 0$ and $\tilde{f} = f \circ \phi$ in a neighborhood of 0.

Definition 2.18. We say that two elements f and \tilde{f} of $C^\infty(\mathbb{R}^3, \mathbb{R}^2)$ are right-equivalent at 0 if there exists a diffeomorphism $\phi \in C^\infty(\mathbb{R}^3, \mathbb{R}^3)$ of the form $\phi : (u, v, z) \mapsto (\phi_1(u, v), \phi_2(u, v), \phi_3(z))$, where $\phi_1, \phi_2 \in C^\infty(\mathbb{R}^2, \mathbb{R})$ and $\phi_3 \in C^\infty(\mathbb{R}, \mathbb{R})$, satisfying $\phi(0) = 0$ and $\tilde{f} = f \circ \phi$ in a neighborhood of 0.

Combining the previous three definitions we introduce the following notion of equivalence.

Definition 2.19. We say that two elements f and \tilde{f} of $C^\infty(\mathbb{R}^2, \mathbb{R}^2)$ are equivalent at 0 if there exists $(\phi, \theta, \xi) \in C^\infty(\mathbb{R}^2, \mathbb{R}^2) \times \mathbb{S}^1 \times C^\infty(\mathbb{R}^2, \mathbb{R} \setminus \{0\})$ with ϕ as in Definition 2.18, and $\zeta = \pm 1$ such that for every (u, v, z) in a neighborhood of 0,

$$\begin{cases} \tilde{f}_1(u, v) = \xi(u, v)(\cos(2\theta)f_1 \circ \phi(u, v) - \zeta \sin(2\theta)f_2 \circ \phi(u, v)), \\ \tilde{f}_2(u, v) = \xi(u, v)(\sin(2\theta)f_1 \circ \phi(u, v) + \zeta \cos(2\theta)f_2 \circ \phi(u, v)). \end{cases}$$

Definition 2.20. We say that two elements f and \tilde{f} of $C^\infty(\mathbb{R}^3, \mathbb{R}^2)$ are equivalent at 0 if there exists $(\phi, \theta, \xi) \in C^\infty(\mathbb{R}^3, \mathbb{R}^3) \times \mathbb{S}^1 \times C^\infty(\mathbb{R}^2, \mathbb{R} \setminus \{0\})$ with ϕ as in Definition 2.18, and $\zeta = \pm 1$ such that for every (u, v, z) in a neighborhood of 0,

$$\begin{cases} \tilde{f}_1(u, v, z) = \xi(u, v)(\cos(2\theta)f_1 \circ \phi(u, v, z) - \zeta \sin(2\theta)f_2 \circ \phi(u, v, z)), \\ \tilde{f}_2(u, v, z) = \xi(u, v)(\sin(2\theta)f_1 \circ \phi(u, v, z) + \zeta \cos(2\theta)f_2 \circ \phi(u, v, z)). \end{cases}$$

An essential feature of admissible transformations is the following proposition, which is obtained by a direct application of the definitions.

Proposition 2.21. • Let $f, \tilde{f} \in C^\infty(\mathbb{R}^2, \mathbb{R}^2)$ be equivalent. Then 0 is conical for f if and only if it is conical for \tilde{f} and 0 is semi-conical for f if and only if it is semi-conical for \tilde{f} .

- Let $f, \tilde{f} \in C^\infty(\mathbb{R}^3, \mathbb{R}^2)$ Then 0 is F -conical for f if and only if it is F -conical for \tilde{f} and 0 is F -semi-conical for f if and only if it is F -semi-conical for \tilde{f} .

2.3 Normal forms for the single system case

2.3.1 Conical intersection

Define $f \in C^\infty(\mathbb{R}^2, \mathbb{R}^2)$ such that $\chi(f)(0) \neq 0$. In this case, f is a diffeomorphism in a neighborhood of 0. Hence f is right-equivalent to $\text{Id} : \mathbb{R}^2 \rightarrow \mathbb{R}^2$. The normal form provides the well-known Hamiltonian $H(u, v) = \begin{pmatrix} u & v \\ v & -u \end{pmatrix}$, for $u, v \in \mathbb{R}^2$.

2.3.2 Semi-conical intersection

The main result of this section is the following theorem.

Theorem 2.22. Assume that 0 is semi-conical for $f \in C^\infty(\mathbb{R}^2, \mathbb{R})$. Then f is equivalent to $(u, v) \mapsto \begin{pmatrix} h(u)u \\ u + v^2 \end{pmatrix}$ where $h : \mathbb{R} \rightarrow \mathbb{R}$ is a smooth function satisfying $h(0) = 1$.

Remark 2.23. As claimed in Proposition 2.3, it follows from the normal form of Theorem 2.22 that semi-conical intersections are isolated (as eigenvalue intersections) in \mathbb{R}^2 .

The **algorithm** that we will refer as **(A)** to get the normal form is the following:

- **STEP 1:** By a left-equivalence we transform f_1 and f_2 into two functions \tilde{f}_1 and \tilde{f}_2 such that $\nabla \tilde{f}_1(0) = \nabla \tilde{f}_2(0) \neq 0$.
- **STEP 2:** By a right-equivalence, we bring the non-conical direction to $\text{span}(e_2)$.
- **STEP 3:** By a further right-equivalence then a time-equivalence we transform f into the announced form.

2.3.3 Proof of Theorem 2.22: STEP 1

Proposition 2.24. *Consider $f \in C^\infty(\mathbb{R}^2, \mathbb{R}^2)$ having a semi-conical intersection at 0. Then there exists \tilde{f} left-equivalent to f such that $\nabla \tilde{f}_1(0) = \nabla \tilde{f}_2(0) \neq 0$.*

Proof. Without loss of generality, we can assume $\nabla f_1(0) \neq 0$. Define $\alpha \in \mathbb{R}$ such that $\nabla f_2(0) = \alpha \nabla f_1(0)$.

Denote by \tilde{f} the function obtained by applying to f the left-equivalence associated with $\theta \in \mathbb{S}^1$ and $\zeta = 1$ as in Remark 2.15. Hence,

$$\nabla \tilde{f}_1 = \cos(2\theta) \nabla f_1 - \sin(2\theta) \nabla f_2, \quad \nabla \tilde{f}_2 = \cos(2\theta) \nabla f_2 + \sin(2\theta) \nabla f_1.$$

We have $\nabla \tilde{f}_1(0) = \nabla \tilde{f}_2(0)$ if and only if $\langle \begin{pmatrix} \cos(2\theta) \\ \sin(2\theta) \end{pmatrix}, \begin{pmatrix} 1 - \alpha \\ -(1 + \alpha) \end{pmatrix} \rangle = 0$. It is clearly possible to choose $\theta \in \mathbb{S}^1$ satisfying the previous condition, the proposition is proved. \square

2.3.4 Proof of Theorem 2.22: STEP 2

Proposition 2.25. *Assume that 0 is semi-conical for $f \in C^\infty(\mathbb{R}^2, \mathbb{R}^2)$. There exists a right-equivalence $\phi : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ of f such that $\tilde{f} = f \circ \phi$ satisfies*

$$\partial_2 \tilde{f}_1(0, 0) = \partial_2 \tilde{f}_2(0, 0) = 0$$

and

$$\partial_1 \tilde{f}_1(0, 0) \neq 0, \quad \partial_1 \tilde{f}_2(0, 0) \neq 0.$$

Proof. Consider $r_1, r_2 \neq 0$ and $\beta_1 \in [0, 2\pi]$ such that

$$\partial_2 f_1(0, 0) = r_1 \cos(\beta_1), \quad \partial_1 f_1(0, 0) = r_1 \sin(\beta_1), \quad \partial_2 f_2(0, 0) = r_2 \cos(\beta_1), \quad \partial_1 f_2(0, 0) = r_2 \sin(\beta_1).$$

Introducing the right-equivalence $\phi(u, v) = \begin{pmatrix} -\sin(\beta_1) & \cos(\beta_1) \\ -\cos(\beta_1) & -\sin(\beta_1) \end{pmatrix} \begin{pmatrix} u \\ v \end{pmatrix}$ and $\tilde{f} = (f \circ \phi)$,

$$\text{we have } D\tilde{f}(0, 0) = \begin{pmatrix} -r_1 & 0 \\ -r_2 & 0 \end{pmatrix}. \quad \square$$

Propositions 2.24 and 2.25 lead us to consider the next condition:

$$f(0) = 0, \quad \partial_2 f(0) = 0, \quad \partial_1 f_1(0) = \partial_1 f_2(0) \neq 0, \quad \partial_2 \chi(f)(0) \neq 0. \quad (\mathbf{SC})$$

2.3.5 Proof of Theorem 2.22: STEP 3

Proposition 2.26. *Let $f \in C^\infty(\mathbb{R}^2, \mathbb{R}^2)$ satisfy Condition (\mathbf{SC}) . Then there exists $h \in C^\infty(\mathbb{R}, \mathbb{R})$ satisfying $h(0) = 1$ such that f is right-equivalent to $(u, v) \mapsto \begin{pmatrix} h(u)u \\ u + v^2 \end{pmatrix}$ or $(u, v) \mapsto \begin{pmatrix} h(u)u \\ u - v^2 \end{pmatrix}$.*

Proof. Because of the condition $\partial_1 f_1(0) \neq 0$, the map $\Phi : (u, v) \mapsto (f_1(u, v), v)$ is a diffeomorphism in a neighborhood of 0 and $g = f \circ \Phi^{-1}$ is right-equivalent to f . Locally near 0 we have

$$g_1(u, v) = u, \quad g_2(u, v) = f_2(G(u, v), v),$$

where G is a smooth function satisfying $\partial_1 G(u, v) = \frac{1}{\partial_1 f_1(G(u, v), v)}$ and $\partial_2 G(u, v) = -\frac{\partial_2 f_1(G(u, v), v)}{\partial_1 f_1(G(u, v), v)}$. Hence,

$$\begin{aligned} \partial_1 g_2(u, v) &= \partial_1 f_2(G(u, v), v) \partial_1 G(u, v), \\ \partial_2 g_2(u, v) &= \partial_1 f_2(G(u, v), v) \partial_2 G(u, v) + \partial_2 f_2(G(u, v), v). \end{aligned}$$

The condition $\partial_1 f_1(0) \neq 0 \neq \partial_1 f_2(0)$ implies that $\partial_1 g_2(0) \neq 0$. Moreover,

$$\begin{aligned} \partial_{22} g_2(u, v) &= \partial_{22} G(u, v) \partial_1 f_2(G(u, v), v) + \partial_{22} f_2(G(u, v), v) \\ &\quad + \partial_2 G(u, v)^2 \partial_{11} f_2(G(u, v), v) + 2 \partial_2 G(u, v) \partial_{12} f_2(G(u, v), v). \end{aligned}$$

Evaluating at 0 and noticing that $\partial_1 G(0) = \frac{1}{\partial_1 f_1(0)}$ and $\partial_2 G(0) = 0$, we get $\partial_2 g_2(0) = 0$ and

$$\partial_{22} g_2(0) = \partial_{22} G(0) \partial_1 f_2(0) + \partial_{22} f_2(0) = \frac{\partial_2 \chi(f)(0)}{\partial_1 f_1(0)} \neq 0,$$

since f satisfies Condition **(SC)**. By the Implicit Function Theorem, $\partial_2 g_2(u, v)$ vanishes on a smooth curve $v = \eta(u)$ in a neighborhood of 0. By Lemma B.2, there exist two smooth functions $m \in C^\infty(\mathbb{R}^2, \mathbb{R})$ and $f_0 \in C^\infty(\mathbb{R}, \mathbb{R})$ such that $g_2(u, v) = (v - \eta(u))^2 m(u, v) + f_0(u)$ in a neighborhood of 0. The conditions $\partial_1 g_2(0) \neq 0$ and $\partial_{22} g_2(0) \neq 0$ yield $f_0'(0) \neq 0$ and $m(0) \neq 0$, respectively. Applying first the right-equivalence $(u, v) \mapsto \begin{pmatrix} u \\ (v - \eta(u)) \sqrt{|m(u, v)|} \end{pmatrix}$ and then $(u, v) \mapsto \begin{pmatrix} f_0(u) \\ v \end{pmatrix}$, we deduce that f is right-equivalent to $(u, v) \mapsto \begin{pmatrix} h(u)u \\ u + \text{sign}(m(0, 0))v^2 \end{pmatrix}$, for some smooth function h obtained by inversion of f_0 . Noticing that the quantity $\frac{\partial_1 f_1(0)}{\partial_1 f_2(0)}$ is invariant by right-equivalence, this provides $h(0) = \frac{\partial_1 f_1(0)}{\partial_1 f_2(0)} = 1$. \square

Proof of Theorem 2.22. According to Propositions 2.25 and 2.26, we are left to prove that

$$f : (u, v) \mapsto \begin{pmatrix} h(u)u \\ u - v^2 \end{pmatrix}$$

is right-time-equivalent to

$$(u, v) \mapsto \begin{pmatrix} \tilde{h}(u)u \\ u + v^2 \end{pmatrix},$$

where \tilde{h} is in $C^\infty(\mathbb{R}, \mathbb{R})$. Indeed, we can apply the right-equivalence $(u, v) \mapsto (-u, v)$, then the time-equivalence associated with $\xi \equiv -1$. The result follows defining $\tilde{h}(u) = h(-u)$. \square

2.4 Normal forms for the ensemble case

Before discussing separately the conical and the semi-conical cases, let us present a useful technical result.

Lemma 2.27. *For every $f \in C^\infty(\mathbb{R}^3, \mathbb{R}^2)$ such that $\partial_3 f_2(0) \neq 0$, let $\beta(f) = \frac{\partial_3 f_1}{\partial_3 f_2}(0)$. Then $\beta(f)$ is invariant by right-equivalence.*

Proof. Let \tilde{f} be right-equivalent to f and let $\phi_1 \in C^\infty(\mathbb{R}^2, \mathbb{R})$, $\phi_2 \in C^\infty(\mathbb{R}^2, \mathbb{R})$, $\phi_3 \in C^\infty(\mathbb{R}, \mathbb{R})$ be such that $\phi(u, v, z) = (\phi_1(u, v), \phi_2(u, v), \phi_3(z))$ is a right-equivalence between f and \tilde{f} . Then,

$$\frac{\partial_3 \tilde{f}_1}{\partial_3 \tilde{f}_2}(0) = \frac{\partial_3 f_1(0) \phi_3'(0)}{\partial_3 f_2(0) \phi_3'(0)} = \beta(f),$$

using the fact $\phi_3'(0) \neq 0$ because ϕ is a diffeomorphism. \square

2.4.1 Conical case

Theorem 2.28. *Let $f \in C^\infty(\mathbb{R}^3, \mathbb{R}^2)$. Then 0 is F-conical for f if and only if there exist $h_1, h_2 \in C^\infty(\mathbb{R}^3, \mathbb{R})$ satisfying $h_1(0) = h_2(0) = 1$, such that f is equivalent to*

$$(u, v, z) \mapsto \begin{pmatrix} h_1(u, v, z)(z - u) \\ h_2(u, v, z)(z - v) \end{pmatrix}.$$

Proof. By the same argument as in the proof of Proposition 2.24, there exists \tilde{f} left-equivalent to f such that $\partial_3 \tilde{f}_1(0) \neq 0$, $\partial_3 \tilde{f}_2(0) \neq 0$ and $\beta(\tilde{f}) = 1$. Using the fact that $\tilde{f}(0) = 0$, we deduce that \tilde{f}_1 and \tilde{f}_2 vanish respectively on two smooth surfaces whose equations are of the form $z = \eta_1(u, v)$ and $z = \eta_2(u, v)$, where η_1, η_2 are smooth functions vanishing at $(0, 0)$. By Lemma B.1, there exist two smooth scalar functions $(u, v, z) \mapsto \phi_1(u, v, z)$, $(u, v, z) \mapsto \phi_2(u, v, z)$ such that

$$\tilde{f}_1(u, v, z) = \phi_1(u, v, z)(z - \eta_1(u, v)), \quad \tilde{f}_2(u, v, z) = \phi_2(u, v, z)(z - \eta_2(u, v)).$$

Differentiating these expressions and evaluating them at 0 we get that $\chi(\tilde{f})(0) = \phi_1(0)\phi_2(0)\chi(\eta)(0)$, where $\eta = (\eta_1, \eta_2)$. Hence, by F-conicity of 0, $\phi_1(0) \neq 0$, $\phi_2(0) \neq 0$, and $\chi(\eta)(0) \neq 0$. In particular, η is a diffeomorphism in a neighborhood of $(0, 0)$. Then \tilde{f} is right-equivalent to

$$\tilde{f} \circ \mu^{-1}(u, v, z) = \begin{pmatrix} h_1(u, v, z)(u - z) \\ h_2(u, v, z)(v - z) \end{pmatrix},$$

where $\mu : (u, v, z) \mapsto (\eta_1(u, v), \eta_2(u, v), z)$ and $h_1, h_2 \in C^\infty(\mathbb{R}^3, \mathbb{R})$ satisfy $h_1(0) \neq 0$ and $h_2(0) \neq 0$. By Lemma 2.27, $\frac{h_1(0)}{h_2(0)} = \beta(\tilde{f}) = 1$. By applying a time equivalence associated with $\xi \equiv \frac{1}{h_1(0)}$, we conclude the proof of the theorem. \square

2.4.2 Semi-conical case

Theorem 2.29. *Let $f \in C^\infty(\mathbb{R}^3, \mathbb{R}^2)$. Then 0 is F-semi-conical for $f \in C^\infty(\mathbb{R}^3, \mathbb{R}^2)$ if and only if there exist $h_1, h_2 \in C^\infty(\mathbb{R}^3, \mathbb{R}^2)$ satisfying $h_1(0) = h_2(0) = 1$ and $m \in C^\infty(\mathbb{R}, \mathbb{R})$ satisfying $m(0) \notin \{-1, 0\}$ such that f is equivalent to*

$$(u, v, z) \mapsto \begin{pmatrix} h_1(u, v, z)(z - m(u)u) \\ h_2(u, v, z)(z + u + v^2) \end{pmatrix}.$$

Before proving the theorem, let us make some general considerations and provide an intermediate result in Proposition 2.30.

First remark that, up to a left-equivalence, we can assume that

$$\partial_1 f_1(0) \neq 0, \quad \partial_1 f_2(0) \neq 0, \quad \partial_3 f_1(0) = \partial_3 f_2(0) \neq 0.$$

In particular, $\beta(f) = 1$. In order to impose the non-conical direction to be in the $\text{span}(e_2)$ -direction, we use the same right-equivalence of the plane (u, v) as in the first step of the algorithm **(A)** in Section 2.3 (see Proposition 2.25). As a result, we end up with \hat{f} equivalent to f and such that

$$\begin{aligned} f(0) = 0, \quad \partial_2 f(0) = 0, \quad \partial_1 f_1(0) \neq 0, \quad \partial_1 f_2(0) \neq 0, \quad \partial_3 f_1(0) = \partial_3 f_2(0) \neq 0, \\ \chi_{13}(f)(0) \neq 0, \quad \partial_2 \chi(f)(0) \neq 0. \end{aligned} \tag{SCP}$$

Notice that the condition $\chi_{13}(f)(0) \neq 0$ can then be rewritten as $\partial_1 f_1(0) \neq \partial_1 f_2(0)$.

Proposition 2.30. *Let $f \in C^\infty(\mathbb{R}^3, \mathbb{R}^2)$ satisfy (SCP) at 0. Then there exist $h_1, h_2 \in C^\infty(\mathbb{R}^3, \mathbb{R})$ non-vanishing at 0 and $m \in C^\infty(\mathbb{R}, \mathbb{R})$ such that $\frac{h_1(0)}{h_2(0)} = 1$, $m(0) = -\frac{\partial_1 f_1(0)}{\partial_1 f_2(0)} \notin \{-1, 0\}$, and f is right-equivalent to*

$$(u, v, z) \mapsto \begin{pmatrix} h_1(u, v, z)(z - m(u)u) \\ h_2(u, v, z)(z + u \pm v^2) \end{pmatrix},$$

where the sign depends on f .

Proof. Using the fact that $f(0) = 0$ and the conditions $\partial_3 f_1(0) \neq 0 \neq \partial_3 f_2(0)$, we can deduce that f_1 and f_2 are smooth functions vanishing, in the neighborhood of the origin, on two smooth surfaces whose equations are, respectively, $z = \eta_1(u, v)$ and $z = \eta_2(u, v)$, where $\eta_1, \eta_2 : \mathbb{R}^2 \rightarrow \mathbb{R}$ are smooth functions vanishing at 0. By Lemma B.1, there exist two smooth functions $\phi_1, \phi_2 : \mathbb{R}^3 \rightarrow \mathbb{R}$ such that

$$f_1(u, v, z) = \phi_1(u, v, z)(z - \eta_1(u, v)), \quad f_2(u, v, z) = \phi_2(u, v, z)(z - \eta_2(u, v)).$$

Differentiating these expressions with respect to z , we deduce that $\phi_1(0) \neq 0 \neq \phi_2(0)$. Differentiating f_1 and f_2 with respect to y , we get then from (SCP) that $\partial_1 \eta_1(0) \neq 0 \neq \partial_1 \eta_2(0)$ and $\partial_2 \eta_1(0) = \partial_2 \eta_2(0) = 0$. Applying the right-equivalence associated with the inverse of $(u, v) \mapsto \begin{pmatrix} \eta_1(u, v) \\ v \end{pmatrix}$, we get that f is right-equivalent to

$$(u, v, z) \mapsto \begin{pmatrix} \phi_1(G(u, v), v, z)(z - u) \\ \phi_2(G(u, v), v, z)(z - \eta_2(G(u, v), v)) \end{pmatrix},$$

for some smooth function G such that $\partial_1 G(0) \neq 0$. Set $\tilde{\eta}(u, v) = \eta_2(G(u, v), v)$. Then $\tilde{\eta}(0) = 0$ and

$$\partial_1 \tilde{\eta}(u, v) = \partial_1 G(u, v) \partial_1 \eta_2(G(u, v), v), \quad \partial_2 \tilde{\eta}(u, v) = \partial_1 \eta_2(G(u, v), v) \partial_2 G(u, v) + \partial_2 \eta_2(G(u, v), v).$$

Evaluating at zero, we get that $\partial_1 \tilde{\eta}(0) \neq 0$ and $\partial_2 \tilde{\eta}(0) = 0$. Differentiating once more and using the hypothesis $\partial_2 \chi(f)(0) \neq 0$, we have $\partial_{22} \tilde{\eta}(0) \neq 0$. By the same arguments as in the proof of Theorem 2.22, f is right-equivalent to

$$(u, v, z) \mapsto \begin{pmatrix} h_1(u, v, z)(z - m(u)u) \\ h_2(u, v, z)(z + u \pm v^2) \end{pmatrix}$$

where $h_1, h_2 \in C^\infty(\mathbb{R}^3, \mathbb{R}^2)$ and $m \in C^\infty(\mathbb{R}, \mathbb{R})$. Noticing that the quantities $\frac{\partial_1 f_1(0)}{\partial_1 f_2(0)}$ and $\beta(f)$ are invariant by right-equivalence, we get $\frac{h_1(0)}{h_2(0)} = \beta(f) = 1$ and $m(0) = -\frac{\partial_1 f_1(0)}{\partial_1 f_2(0)} \notin \{-1, 0\}$ because f satisfies (SCP) at 0. \square

Proof of Theorem 2.29. First notice that if f is of the form

$$f : (u, v, z) \mapsto \begin{pmatrix} h_1(u, v, z)(z - m(u)u) \\ h_2(u, v, z)(z + u - v^2) \end{pmatrix}$$

then there exist $\tilde{h}_1, \tilde{h}_2 \in C^\infty(\mathbb{R}^3, \mathbb{R})$ and $\tilde{m} \in C^\infty(\mathbb{R}, \mathbb{R})$ such that f is right-time equivalent to

$$(u, v, z) \mapsto \begin{pmatrix} \tilde{h}_1(u, v, z)(z - \tilde{m}(u)u) \\ \tilde{h}_2(u, v, z)(z + u + v^2) \end{pmatrix}.$$

Indeed, applying the right-equivalence $(u, v, z) \mapsto (-u, v, -z)$ and the time-equivalence associated with $\xi : (u, v) \mapsto -1$ the claim follows with $\tilde{h}_i(u, v, z) = h_i(-u, v, -z)$, $i \in \{1, 2\}$, and $\tilde{m}(u) = m(-u)$. Theorem 2.29 hence follows from Proposition 2.30. \square

3 Generic global properties of the singular locus

3.1 Proof of Lemma 1.1 and Theorem 1.2

Let $f \in C^\infty(\mathbb{R}^3, \mathbb{R}^2)$, $Z(f) = \{(u, v, z) \mid f(u, v, z) = (0, 0)\}$, and denote by $Z_{\text{nc}}(f)$ the set of non-conical intersections in $Z(f)$. Let $\pi(f)$ be the orthogonal projection of $Z(f)$ onto the plane (u, v) .

Proposition 3.1. *Assume that $(\bar{u}, \bar{v}, \bar{z})$ is a F -semi-conical intersection for $f \in C^\infty(\mathbb{R}^3, \mathbb{R}^2)$. Then $\pi(f)$ is tangent at (\bar{u}, \bar{v}) to the non-conical direction of f at $(\bar{u}, \bar{v}, \bar{z})$.*

Proof. By Proposition 2.7, $Z(f)$ is locally near $(\bar{u}, \bar{v}, \bar{z})$ a smooth curve that we parameterize by $c(t) = (u(t), v(t), z(t))_{t \in \mathbb{R}}$, with $c(0) = (\bar{u}, \bar{v}, \bar{z})$ and $\dot{z}(0) = 0$. The condition $f(c(t)) \equiv 0$ implies that $Df_{c(t)}(\dot{c}(t)) \equiv 0$. In particular, $(\dot{u}(0), \dot{v}(0))$ is collinear to $\eta = \begin{pmatrix} -\partial_2 f_1(\bar{u}, \bar{v}, \bar{z}) \\ \partial_1 f_1(\bar{u}, \bar{v}, \bar{z}) \end{pmatrix}$, which is the non-conical direction of f at $(\bar{u}, \bar{v}, \bar{z})$. \square

Proposition 3.2. *Assume that $f \in C^\infty(\mathbb{R}^3, \mathbb{R}^2)$ is a submersion at every point of $Z(f)$. Then $\pi(f)$ has no cuspidal point.*

Proof. Fix a local smooth regular parametrization $c(t) = (u(t), v(t), z(t))_{t \in \mathbb{R}}$ of $c \subset Z(f)$. It is sufficient to show that there exist no $t \in \mathbb{R}$ such that $\dot{u}(t) = \dot{v}(t) = 0$. By the same arguments as those used in the proof of Proposition 2.7, we get that $\dot{u}(t) = 0$ implies that $\begin{vmatrix} \partial_2 f_1(c(t)) & \partial_3 f_1(c(t)) \\ \partial_2 f_2(c(t)) & \partial_3 f_2(c(t)) \end{vmatrix} = 0$, while $\dot{v}(t) = 0$ implies that $\begin{vmatrix} \partial_1 f_1(c(t)) & \partial_3 f_1(c(t)) \\ \partial_1 f_2(c(t)) & \partial_3 f_2(c(t)) \end{vmatrix} = 0$. If the two determinants simultaneously vanish then f is not a submersion at $c(t)$. \square

Propositions 3.1 and 3.2, together with Remark 2.9, prove Lemma 1.1. As for Theorem 1.2, it follows from Proposition 3.2 together with Propositions 2.6, 2.10, and 2.11.

3.2 Generic self-intersections of $\pi(f)$

By a multi-jet version of the transversality arguments already used in the previous sections (see, for instance, [18, §4, Theorem 4.13]) one can deduce the following result.

Proposition 3.3. *Generically with respect to $f \in C^\infty(\mathbb{R}^3, \mathbb{R}^2)$,*

1. $\pi(f)$ has no triple points;
2. Let (u, v) and (\tilde{u}, \tilde{v}) be two double points of $\pi(f)$ and let $z_1 \neq z_2$ and $\tilde{z}_1 \neq \tilde{z}_2$ be such that $f(u, v, z_1) = f(u, v, z_2) = f(\tilde{u}, \tilde{v}, \tilde{z}_1) = f(\tilde{u}, \tilde{v}, \tilde{z}_2) = 0$. Then $z_i \neq \tilde{z}_j$ for every $i, j \in \{1, 2\}$;
3. Let (u, v) and $z \neq \tilde{z}$ satisfy $f(u, v, z) = f(u, v, \tilde{z}) = 0$. Then (u, v, z) and (u, v, \tilde{z}) are F -conical for f ;
4. Let (u, v, z) and $(\tilde{u}, \tilde{v}, \tilde{z})$ be two non-conical intersections for f . Then $z \neq \tilde{z}$.

4 Adiabatic control through a semi-conical intersection of eigenvalues

Let us consider $f \in C^\infty(\mathbb{R}^2, \mathbb{R}^2)$, its associated Hamiltonian $H_f = \begin{pmatrix} f_1 & f_2 \\ f_2 & -f_1 \end{pmatrix}$, and a control path $(u(t), v(t))_{t \in [0, 1]}$. Denote by $\lambda^-(u, v)$ and $\lambda^+(u, v)$ the smallest and the largest eigenvalue of $H_f(u, v)$, respectively.

In the following, we denote by $\phi_-(u(t), v(t))$ (respectively, $\phi_+(u(t), v(t))$) a real normalized eigenvector of $H_f(u(t), v(t))$ associated with $\lambda_-(u(t), v(t))$ (respectively, $\lambda_+(u(t), v(t))$). If $f(u(t), v(t)) \neq 0$ then $\lambda_-(u(t), v(t))$ and $\lambda_+(u(t), v(t))$ are simple and the choice of $\phi_\pm(u(t), v(t))$ is unique up to multiplication by -1 . If $(u(t), v(t))_{t \in [0,1]}$ does not cross $Z(f)$ then $t \mapsto \phi_\pm(u(t), v(t))$ and $t \mapsto \lambda^\pm(u(t), v(t))$ can be chosen with the same regularity as $(u(t), v(t))_{t \in [0,1]}$. It is a classical fact that this may not be the case when $(u(t), v(t))_{t \in [0,1]}$ crosses $Z(f)$. However, we are going to prove the existence of a C^k basis of eigendirections of H_f along a C^{k+2} path $(u(t), v(t))_{t \in [0,1]}$ passing through a semi-conical intersection in a conical or a non-conical direction.

4.1 Adiabatic dynamics

Let $f \in C^\infty(\mathbb{R}^2, \mathbb{R}^2)$. Consider a smooth regular control path $(u(t), v(t))_{t \in [0,1]}$ such that there exist $P \in C^2([0, 1], \text{SO}_2(\mathbb{R}))$ and $\lambda \in C^k([0, 1], \mathbb{R})$ such that $\{\lambda(t), -\lambda(t)\}$ is the spectrum of $H_f(u(t), v(t))$, and the columns of P form a basis of eigenvectors of $H_f(u(t), v(t))$ for every $t \in [0, 1]$. We can write, for every $t \in [0, 1]$, $P(t) = \begin{pmatrix} \cos(\theta(t)) & -\sin(\theta(t)) \\ \sin(\theta(t)) & \cos(\theta(t)) \end{pmatrix}$ where $\theta \in C^2([0, 1], \mathbb{R})$.

Let us study the dynamics of

$$i \frac{d\psi_\epsilon(t)}{dt} = H_f(u(\epsilon t), v(\epsilon t))\psi_\epsilon(t), \quad \psi_\epsilon(0) = \tilde{\psi}_0, \quad (10)$$

where $t \in [0, \frac{1}{\epsilon}]$ and $\tilde{\psi}_0$ is independent of ϵ .

Defining $Y_\epsilon(\tau) = P(\tau)\psi_\epsilon(\frac{\tau}{\epsilon})$ for every $\tau \in [0, 1]$, we have

$$i \frac{dY_\epsilon(\tau)}{d\tau} = \left(\frac{1}{\epsilon} \begin{pmatrix} \lambda(\tau) & 0 \\ 0 & -\lambda(\tau) \end{pmatrix} + \begin{pmatrix} 0 & i\dot{\theta}(\tau) \\ -i\dot{\theta}(\tau) & 0 \end{pmatrix} \right) Y_\epsilon(\tau). \quad (11)$$

Thanks to the further change of variables $\tilde{Y}_\epsilon(\tau) = \begin{pmatrix} e^{\frac{i}{\epsilon} \int_0^\tau \lambda(s) ds} & 0 \\ 0 & e^{-\frac{i}{\epsilon} \int_0^\tau \lambda(s) ds} \end{pmatrix} Y_\epsilon(\tau)$, the dynamics are transformed into

$$\frac{d\tilde{Y}_\epsilon(\tau)}{d\tau} = \begin{pmatrix} 0 & i\dot{\theta}(\tau) e^{\frac{2i}{\epsilon} \int_0^\tau \lambda(s) ds} \\ -i\dot{\theta}(\tau) e^{-\frac{2i}{\epsilon} \int_0^\tau \lambda(s) ds} & 0 \end{pmatrix} \tilde{Y}_\epsilon(\tau). \quad (12)$$

Based on Corollaries A.3 and A.7, one gets the following result.

Theorem 4.1 (Adiabatic Theorem). *Let $k \in \mathbb{N}$ and assume that $\lambda : [0, 1] \rightarrow \mathbb{R}$ is C^k and $\theta : [0, 1] \rightarrow \mathbb{R}$ is C^2 . Let $\psi_\epsilon : [0, 1] \rightarrow \mathbb{C}^2$ be the solution of Equation (10). Assume that there exists $c > 0$ such that*

$$\left| \int_0^t e^{\frac{2i}{\epsilon} \int_0^s \lambda(x) dx} ds \right| \leq c\epsilon^{1/(k+1)}, \quad \forall t \in [0, 1]. \quad (13)$$

Then $\tilde{Y}_\epsilon(\tau) = \tilde{Y}_\epsilon(0) + O(\epsilon^\eta)$ uniformly w.r.t. $\tau \in [0, 1]$. In particular, if $\tilde{\psi}_0 = \begin{pmatrix} \cos(\theta(0)) \\ \sin(\theta(0)) \end{pmatrix}$, then $\psi_\epsilon(\frac{1}{\epsilon}) = e^{i\eta} \begin{pmatrix} \cos(\theta(1)) \\ \sin(\theta(1)) \end{pmatrix} + O(\epsilon^{\frac{1}{k+1}})$, where η possibly depends on ϵ .

4.2 Regularity of the eigenpairs along smooth control paths

The main goal of this section is to study the regularity of eigenpairs of H_f along smooth curves passing through a semi-conical intersection for f . Using the normal form obtained in Section 2.3, we can restrict our attention to the case where f has the form $f : (u, v) \mapsto \begin{pmatrix} uh(u) \\ u + v^2 \end{pmatrix}$, where $h \in C^\infty(\mathbb{R}, \mathbb{R})$ is such that $h(0) = 1$.

4.2.1 Conical directions

We recall here the following regularity result which is a special case of [11, Proposition 3.1] and [4, Lemma 3.2].

Proposition 4.2 (Eigenpairs in the conical directions). *Consider a Hamiltonian H_f where $f : (u, v) \mapsto \begin{pmatrix} uh(u) \\ u + v^2 \end{pmatrix}$ and $h \in C^\infty(\mathbb{R}, \mathbb{R})$ is such that $h(0) = 1$. Let $\ell \in \mathbb{N}$, $t_0 \in (0, 1)$, and $(u(t), v(t))_{t \in [0, 1]}$ be a $C^{\ell+1}$ path such that $(u(t), v(t)) = 0$ if and only if $t = t_0$ and $\dot{u}(t_0) \neq 0$. Define $\lambda_0, \lambda_1 : [0, 1] \rightarrow \mathbb{R}$ by*

$$\begin{aligned} \lambda_0(t) &= \lambda_-(u(t), v(t)), \quad \lambda_1(t) = \lambda_+(u(t), v(t)), & \text{for } t < t_0, \\ \lambda_0(t) &= \lambda_+(u(t), v(t)), \quad \lambda_1(t) = \lambda_-(u(t), v(t)), & \text{for } t \geq t_0. \end{aligned}$$

Then λ_0 and λ_1 are $C^{\ell+1}$ on $[0, 1]$. Moreover, there exist $\Phi_0, \Phi_1 \in C^\ell([0, 1], \mathbb{R}^2)$ such that $\Phi_j(t)$ is a normalized eigenvector of $H_f(u(t), v(t))$ corresponding to the eigenvalue $\lambda_j(t)$ for $j \in \{0, 1\}$ and $t \in [0, 1]$.

The following proposition states that the limit eigenvectors along a C^2 curve crossing conically a semi-conical intersection do not depend on the choice of the curve.

Proposition 4.3 (Limit eigenvectors along a conical direction). *Consider f , $(u(t), v(t))_{t \in [0, 1]}$, t_0 , and Φ_0, Φ_1 as in Proposition 4.2, where the latter are uniquely defined up to multiplication to -1 . Then $\Phi_0(t_0)$ and $\Phi_1(t_0)$ depend only on the sign of $\dot{u}(t_0)$. More precisely, $\Phi_0(t_0) = \frac{1}{\sqrt{1+\bar{V}^2}} \begin{pmatrix} -1 \\ \bar{V} \end{pmatrix}$ and $\Phi_1(t_0) = \frac{1}{\sqrt{1+\bar{V}^2}} \begin{pmatrix} \bar{V} \\ 1 \end{pmatrix}$ with $\bar{V} = -(1 + \text{sign}(\dot{u}(t_0))\sqrt{2})$.*

Proof. By definition of Φ_0, Φ_1 , we have

$$\Phi_0(t) = \phi_-(u(t), v(t)), \quad \Phi_1(t) = \phi_+(u(t), v(t)), \quad \text{for } t < t_0.$$

Hence, up to multiplication by -1 ,

$$\Phi_0(t) = \frac{1}{\sqrt{1+V(t)^2}} \begin{pmatrix} -1 \\ V(t) \end{pmatrix}, \quad \Phi_1(t) = \frac{1}{\sqrt{1+V(t)^2}} \begin{pmatrix} V(t) \\ 1 \end{pmatrix}, \quad \text{for } t < t_0,$$

where $V(t) = 0$ if $u(t) + v(t)^2 = 0$ and

$$V(t) = \frac{-u(t)h(u(t)) + \sqrt{u(t)^2h(u(t))^2 + (u(t) + v(t)^2)^2}}{u(t) + v(t)^2}$$

otherwise. Then, as $t \rightarrow t_0^-$,

$$\begin{aligned} V(t) &= \frac{-\dot{u}(t_0)h(0)(t-t_0) + o(t-t_0) + \sqrt{\dot{u}(t_0)^2(t-t_0)^2(h(0)^2 + 1) + o((t-t_0)^2)}}{\dot{u}(t_0)(t-t_0) + o(t-t_0)} \\ &= -\frac{\dot{u}(t_0)(t-t_0) + |t-t_0||\dot{u}(t_0)|\sqrt{2}}{\dot{u}(t_0)(t-t_0)} + o(1) \\ &= -(1 + \text{sign}(\dot{u}(t_0))\sqrt{2}) + o(1). \end{aligned}$$

Since Φ_0, Φ_1 are continuous on $[0, 1]$ by Proposition 4.2, the conclusion follows. \square

4.2.2 Non-conical direction

Proposition 4.4 (Continuity of the eigenstates in the non-conical direction). *Let f be as in Proposition 4.2. Let $\ell \in \mathbb{N}$, $t_0 \in (0, 1)$, and $(u(t), v(t))_{t \in [0, 1]}$ be a $C^{\ell+2}$ path such that $(u(t), v(t)) = 0$ if and only if $t = t_0$ and $\dot{u}(t_0) = 0$ (i.e., $(u(t), v(t))$ passes through 0 in the non-conical direction at $t = t_0$ as on Figure 5). Then there exist $\lambda_0, \lambda_1 \in C^{\ell+2}([0, 1], \mathbb{R}^2)$, $\Phi_0, \Phi_1 \in C^\ell([0, 1], \mathbb{R}^2)$ such that $\lambda_0(t) = \lambda_-(u(t), v(t))$, $\lambda_1(t) = \lambda_+(u(t), v(t))$, $\Phi_0(t) = \phi_-(u(t), v(t))$ and $\Phi_1(t) = \phi_+(u(t), v(t))$ for every $t \in [0, t_0) \cup (t_0, 1]$. Moreover, defining $\beta = \frac{\ddot{u}(t_0)}{2} + \dot{v}(t_0)^2$, we have $\Phi_0(t_0) = \frac{1}{\sqrt{1+\bar{V}^2}} \begin{pmatrix} -1 \\ \bar{V} \end{pmatrix}$ and $\Phi_1(t_0) = \frac{1}{\sqrt{1+\bar{V}^2}} \begin{pmatrix} \bar{V} \\ 1 \end{pmatrix}$, where $\bar{V} = -\frac{\ddot{u}(t_0) - \sqrt{\frac{\ddot{u}(t_0)^2}{4} + \beta^2}}{\beta}$ if $\beta \neq 0$. If $\beta = 0$, we have $\Phi_0(t_0) = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$ and $\Phi_1(t_0) = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$.*

Proof. The condition $\dot{u}(t_0) = 0$ provides $u(t) + v(t)^2 = (t - t_0)^2(\frac{\ddot{u}(t_0)}{2} + \dot{v}(t_0)^2) + o((t - t_0)^2) = \beta(t - t_0)^2 + o((t - t_0)^2)$ when $t \rightarrow t_0$.

By direct computations, we show that $\frac{d^2 \lambda_{\pm}(u(t), v(t))}{dt^2} = \pm \sqrt{\beta^2 + \ddot{u}(t_0)^2} + o(1)$ when $t \rightarrow t_0$. Hence $t \mapsto \lambda_{\pm}(u(t), v(t))$ can be extended in a C^2 function at $t = t_0$ by fixing $\frac{d^2 \lambda_{\pm}(u(t), v(t))}{dt^2} = 2 \pm \sqrt{\beta^2 + \ddot{u}(t_0)^2}$. The $C^{\ell+2}$ regularity follows by higher order analog computations.

Define, for every $t \in [0, 1]$ such that $u(t) + v(t)^2 \neq 0$, $V(t) = \frac{-u(t)h(u(t)) + \sqrt{u(t)^2 h(u(t))^2 + (u(t) + v(t)^2)^2}}{u(t) + v(t)^2}$. Reasoning as in the proof of Proposition 4.3, we must prove that V can be extended as a C^ℓ function at t_0 by setting $V(t_0) = \bar{V}$.

First case: $\beta \neq 0$. Assuming that $\beta \neq 0$, we have, as $t \rightarrow t_0$,

$$V(t) = -\frac{\frac{\ddot{u}(t_0)}{2}(t - t_0)^2 - \sqrt{\frac{(t-t_0)^4}{4}\ddot{u}(t_0)^2 + \beta^2(t-t_0)^4 + o((t-t_0)^4)}}{\beta(t-t_0)^2 + o((t-t_0)^2)} = -\frac{\frac{\ddot{u}(t_0)}{2} - \sqrt{\frac{\ddot{u}(t_0)^2}{4} + \beta^2}}{\beta} + o(1).$$

The continuity of V is proved in the case $\beta \neq 0$. The same computations show that V is C^ℓ at t_0 if $(u(t), v(t))_{t \in [0, 1]}$ is $C^{\ell+2}$.

Second case: $\beta = 0$. In the case $\beta = 0$, consider the left equivalence \tilde{f} of f associated with $\theta = \frac{\pi}{4}$ and $\zeta = -1$ as in Remark 2.15, so that, we have $H_{\tilde{f}} = P_{\theta, \zeta} H_f P_{\theta, \zeta}^{-1}$. If $f = (f_1, f_2) \in C^\infty(\mathbb{R}^2, \mathbb{R}^2)$, then we have $\tilde{f} = (f_2, f_1)$. Define, for every $t \in [0, 1]$ such that $u(t)h(u(t)) \neq 0$, $\tilde{V}(t) = \frac{-(u(t) + v(t)^2) + \sqrt{u(t)^2 h(u(t))^2 + (u(t) + v(t)^2)^2}}{u(t)h(u(t))}$. We have easily $\lim_{t \rightarrow t_0} \tilde{V}(t) = -1$. Hence we can define continuous eigenvectors $\tilde{\Phi}_0$ and $\tilde{\Phi}_1$ of $H_{\tilde{f}}$ along $(u(t), v(t))_{t \in [0, 1]}$, respectively equal, up to phases, to $\frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ 1 \end{pmatrix}$ and $\frac{1}{\sqrt{2}} \begin{pmatrix} -1 \\ 1 \end{pmatrix}$ at $t = t_0$. Knowing that the eigenvectors of H_f are equal, up to phases, to $P_{\theta, \zeta} \tilde{\Phi}_j$, for $j \in \{0, 1\}$, we can deduce the continuity of Φ_0 and Φ_1 at $t = t_0$, and that $\Phi_0(t_0) = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$ and $\Phi_1(t_0) = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$. \square

4.3 Dynamical properties at semi-conical intersections of eigenvalues

By using the previous results, we get the following adiabatic approximations along curves going through a semi-conical intersection, either along conical directions (Proposition 4.5) or along the non-conical direction (Proposition 4.6).

Proposition 4.5. *Let f and $(u(t), v(t))_{t \in [0, 1]}$ be as in Proposition 4.2. Consider a solution ψ_ϵ of Equation (10) such that $\tilde{\psi}_0 = \phi_-(u(0), v(0))$. Then $\langle \psi_\epsilon(\frac{1}{\epsilon}), \phi_-(u(1), v(1)) \rangle = O(\sqrt{\epsilon})$.*

Proof. First notice that λ_- and λ_+ are C^2 separately on $[0, t_0]$ and $[t_0, 1]$. Moreover, Corollary A.7 proves that they satisfy (23) with $k = 1$ on $[0, t_0]$ and $[t_0, 1]$. By Proposition 4.3, moreover, there exists a C^2 basis of eigenvectors defined on $[0, 1]$ with $\lim_{t \rightarrow t_0^-} \phi_{\pm}(u(t), v(t)) = \lim_{t \rightarrow t_0^+} \phi_{\mp}(u(t), v(t))$. By applying Theorem 4.1 on the interval $[0, t_0]$, then on the interval $[t_0, 1]$, we get the result. \square

Proposition 4.6. *Let f and $(u(t), v(t))_{t \in [0,1]}$ be as in Proposition 4.4 with $l \geq 2$ (see Figure 5). Consider a solution ψ_{ϵ} of Equation (14) such that $\psi_{\epsilon}(0) = \phi_-(u(0), v(0))$. Then $\langle \psi_{\epsilon}(\frac{1}{\epsilon}), \phi_+(u(1), v(1)) \rangle = O(\epsilon^{1/3})$.*

Proof. Let λ_0 be as in Proposition 4.4 and define $\varphi(\tau) = \int_0^{\tau} \lambda_0(s) ds$ for $\tau \in [0, 1]$. Notice that λ_0 is at least C^2 by Proposition 4.4 and that φ satisfies $\varphi(t_0) = \varphi'(t_0) = \varphi''(t_0) = 0$ and $\varphi^{(3)}(t_0) \neq 0$. Hence, by Lemma A.4, λ_0 satisfies the estimate (23) with $k = 2$. The result follows by applying Theorem 4.1 in combination with Proposition 4.4. \square




Figure 5: A control path passing at a semi-conical intersection in the non-conical direction as a function of the controls $(u, v) \in \mathbb{R}^2$.

5 Control of an ensemble of systems

The main goal of this section is to prove the controllability result stated in Theorem 1.3.

5.1 Ensemble adiabatic dynamics

Let $f \in C^\infty(\mathbb{R}^3, \mathbb{R}^2)$. Consider a smooth regular control path $(u(t), v(t))_{t \in [0,1]}$. In analogy with the previous sections, denote by $\lambda_z^\pm(u(t), v(t))$ the eigenvalues of $H_f(u(t), v(t), z)$. Similarly, let $\phi_z^\pm(u(t), v(t))$ be two real normalized eigenvector of $H_f(u(t), v(t), z)$ at $(u(t), v(t))$ associated with $\lambda_z^\pm(u(t), v(t))$, uniquely defined up to a sign.

Let $\mathcal{Z} = [z_0, z_1]$ be a compact interval of \mathbb{R} . Assume that for every $z \in \mathcal{Z}$ there exist $P_z \in C^1([0, 1], \text{SO}_2(\mathbb{R}))$ and $\lambda_z \in C^1([0, 1], \mathbb{R})$ such that $\{\lambda_z(t), -\lambda_z(t)\}$ is the spectrum of $H_f(u(t), v(t), z)$ and the columns of P_z form a basis of eigenvectors of $H_f(u(t), v(t), z)$ for every $t \in [0, 1]$. We can write, for every $t \in [0, 1]$, $P_z(t) = \begin{pmatrix} \cos(\theta_z(t)) & -\sin(\theta_z(t)) \\ \sin(\theta_z(t)) & \cos(\theta_z(t)) \end{pmatrix}$ where $\theta_z \in C^1([0, 1], \mathbb{R})$.

Let us study the dynamics of

$$i \frac{d\psi_\epsilon^z(t)}{dt} = H_f(u(\epsilon t), v(\epsilon t), z) \psi_\epsilon^z(t), \quad \psi_\epsilon^z(0) = \tilde{\psi}_0^z, \quad (14)$$

where $t \in [0, \frac{1}{\epsilon}]$ and $\tilde{\psi}_0^z$ is independent of ϵ .

Defining $Y_\epsilon^z(\tau) = P_z(\tau) \psi_\epsilon^z(\frac{\tau}{\epsilon})$ for every $\tau \in [0, 1]$, we have

$$i \frac{dY_\epsilon^z(\tau)}{d\tau} = \left(\frac{1}{\epsilon} \begin{pmatrix} \lambda_z(\tau) & 0 \\ 0 & -\lambda_z(\tau) \end{pmatrix} + \begin{pmatrix} 0 & i\dot{\theta}_z(\tau) \\ -i\dot{\theta}_z(\tau) & 0 \end{pmatrix} \right) Y_\epsilon^z(\tau). \quad (15)$$

Thanks to the further change of variable $\tilde{Y}_\epsilon^z(\tau) = \begin{pmatrix} e^{\frac{i}{\epsilon} \int_0^\tau \lambda_z(s) ds} & 0 \\ 0 & e^{-\frac{i}{\epsilon} \int_0^\tau \lambda_z(s) ds} \end{pmatrix} Y_\epsilon^z(\tau)$, the dynamics is transformed into

$$\frac{d\tilde{Y}_\epsilon^z(\tau)}{d\tau} = \begin{pmatrix} 0 & i\dot{\theta}_z(\tau) e^{\frac{2i}{\epsilon} \int_0^\tau \lambda_z(s) ds} \\ -i\dot{\theta}_z(\tau) e^{-\frac{2i}{\epsilon} \int_0^\tau \lambda_z(s) ds} & 0 \end{pmatrix} \tilde{Y}_\epsilon^z(\tau). \quad (16)$$

Based on Corollary A.8, one get the following result.

Theorem 5.1 (Parametric Adiabatic Theorem). *Let $k \in \mathbb{N}$. For every $z \in [z_0, z_1]$, assume that $\lambda_z : [0, 1] \rightarrow \mathbb{R}$ is C^k in $[0, 1]$ and θ_z is C^2 in $[0, 1]$. Let $\psi_\epsilon^z(t)$ be the solution of Equation (14). Assume that there exists $c > 0$ such that for every t in $[0, 1]$, and for every $z \in [z_0, z_1]$,*

$$\left| \int_0^t e^{\frac{2i}{\epsilon} \int_0^s \lambda_z(x) dx} ds \right| \leq c \epsilon^{\frac{1}{k+1}}, \quad \forall \epsilon > 0, \quad (17)$$

and that $(t, z) \mapsto \dot{\theta}_z(t)$ and $(t, z) \mapsto \ddot{\theta}_z(t)$ are uniformly bounded with respect to $(t, z) \in [0, 1] \times [z_0, z_1]$. Then we have $\tilde{Y}_\epsilon^z(\tau) = \tilde{Y}_\epsilon^z(0) + O(\epsilon^{\frac{1}{k+1}})$, uniformly w.r.t. $\tau \in [0, 1]$ and $z \in [z_0, z_1]$. In particular, if $\tilde{\psi}_0^z = \begin{pmatrix} \cos(\theta_z(0)) \\ \sin(\theta_z(0)) \end{pmatrix}$, then we have $\psi_\epsilon^z(\frac{1}{\epsilon}) = e^{in} \begin{pmatrix} \cos(\theta_z(1)) \\ \sin(\theta_z(1)) \end{pmatrix} + O(\epsilon^{\frac{1}{k+1}})$, uniformly w.r.t. $z \in [z_0, z_1]$, where η is possibly depending on ϵ, z .

5.2 Controllability properties between the eigenstates for the normal forms

Let $f : (u, v, z) \mapsto \begin{pmatrix} h_1(u, v, z)(z - m(u)u) \\ h_2(u, v, z)(z + u + v^2) \end{pmatrix}$, where $h_1, h_2 \in C^\infty(\mathbb{R}^3, \mathbb{R})$ and $m \in C^\infty(\mathbb{R}, \mathbb{R})$ satisfy $h_1(0) = h_2(0) \neq 0$ and $m(0) \notin \{-1, 0\}$. Recall that f has a F-semi-conical intersection at 0.

Consider a compact neighborhood \mathcal{S} of 0 in \mathbb{R}^3 on which the product $h_1 h_2$ does not vanish and m is different from 0 and -1 . Assume that \mathcal{S} writes $\mathcal{S} = U \times [z_0, z_1]$, where U is a compact neighborhood of 0 in \mathbb{R}^2 and $z_0 < 0 < z_1$. Define $\mathcal{C} = Z(f) \cap \mathcal{S}$ and, for every $(u, v, z) \in \mathcal{S}$, $h(u, v, z) = \frac{h_1(u, v, z)}{h_2(u, v, z)}$. Notice that $(u, v, z) \in \mathcal{S}$ is in \mathcal{C} if and only if $m(u)u = z = -u - v^2$. Up to restricting U , we can assume that $u \rightarrow (m(u) + 1)u$ is monotone, so that

$$(m(u) + 1)u = -v^2 \quad (18)$$

defines a smooth submanifold of U which contains the projection $\pi(\mathcal{C})$ of \mathcal{C} onto the plane of controls $(u, v) \in \mathbb{R}^2$.

Without loss of generality, assume that $m(0) > -1$ (the case $m(0) < -1$ being analogous). According to (18), this means that $\pi(\mathcal{C})$ lies in the intersection of U with the left half-plane. Notice that the sign of $z = m(u)u$ on \mathcal{C} is the opposite as the sign of $m(0)$.

5.2.1 Uniform adiabatic estimates when $(u(t), v(t))_{t \in [0,1]} \subset \pi(\mathcal{C})$

Assume that $(u, v) : [0, 1] \rightarrow U$ is a regular C^∞ control path satisfying the following conditions, referred to as **(C)**:

- $(u, v) \subset \pi(\mathcal{C})$
- $(u, v)(0) = (u_0, v_0)$ where $(u_0, v_0, z_0) \in \mathcal{C}$ is a F-conical intersection for f ;
- $(u, v)(1) = (0, 0)$.

Under the previous assumptions, for every $z \in [z_0, 0]$, we can consider t_z as the unique element in $[0, 1]$ that satisfies $(u(t_z), v(t_z), z) \in \mathcal{C}$. On the other hand, for $z \in (0, z_1]$ there exist no $t \in [0, 1]$ such that $(u(t), v(t), z) \in \mathcal{C}$. By the regularity of (u, v) , the application $[z_0, 0] \ni z \mapsto t_z$ is C^∞ . Moreover, as a direct consequence of equation (18), which holds on \mathcal{C} , we have $f_1(u(t), v(t), z) = h(u(t), v(t), z)f_2(u(t), v(t), z)$ for every $t \in [0, 1]$ and $z \in [z_0, z_1]$.

Definition 5.2. For any $z \in [z_0, z_1]$, define $V_z : [0, 1] \rightarrow \mathbb{R}$ as follows: if $z \in [z_0, 0]$, let $V_z(t) = -h(u(t), v(t), z) - \sqrt{1 + h(u(t), v(t), z)^2}$ for $t < t_z$ and $V_z(t) = -h(u(t), v(t), z) + \sqrt{1 + h(u(t), v(t), z)^2}$ for $t \geq t_z$; if $z \in [0, z_1]$, let $V_z(t) = -h(u(t), v(t), z) - \sqrt{1 + h(u(t), v(t), z)^2}$ for every $t \in [0, 1]$.

By Propositions 4.2 and 4.4, we get the following result on the regularity of eigenpairs.

Proposition 5.3 (Regularity of eigenpairs for a control path $(u, v) \subset \pi(\mathcal{C})$). Let (u, v) satisfy **(C)**. For $z \in [0, z_1]$, define, for every $t \in [0, 1]$, $\Phi_0^z(t) = \phi_-(u(t), v(t))$, $\Phi_1^z(t) = \phi_+(u(t), v(t))$, $\lambda_0^z(t) = \lambda_-^z(u(t), v(t))$, and $\lambda_1^z(t) = \lambda_+^z(u(t), v(t))$. For $z \in [z_0, 0]$, define $\lambda_0^z, \lambda_1^z : [0, 1] \rightarrow \mathbb{R}$ by

$$\begin{aligned} \lambda_0^z(t) &= \lambda_-^z(u(t), v(t)), & \lambda_1^z(t) &= \lambda_+^z(u(t), v(t)), & \text{for } t < t_z, \\ \lambda_0^z(t) &= \lambda_+^z(u(t), v(t)), & \lambda_1^z(t) &= \lambda_-^z(u(t), v(t)), & \text{for } t \geq t_z, \end{aligned}$$

and $\Phi_0^z, \Phi_1^z : [0, 1] \rightarrow \mathbb{R}^2$ by

$$\begin{aligned} \Phi_0^z(t) &= \phi_-^z(u(t), v(t)), & \Phi_1^z(t) &= \phi_+^z(u(t), v(t)), & \text{for } t < t_z, \\ \Phi_0^z(t) &= \phi_+^z(u(t), v(t)), & \Phi_1^z(t) &= \phi_-^z(u(t), v(t)), & \text{for } t \geq t_z. \end{aligned}$$

Then, for every $z \in [z_0, z_1]$, λ_0^z, λ_1^z and Φ_0^z, Φ_1^z are C^∞ on $[0, 1]$. Moreover, Φ_0^z and Φ_1^z can be written as $\Phi_0^z = \frac{1}{\sqrt{1+V_z^2}} \begin{pmatrix} -1 \\ V_z \end{pmatrix}$ and $\Phi_1^z = \frac{1}{\sqrt{1+V_z^2}} \begin{pmatrix} V_z \\ 1 \end{pmatrix}$, where $V_z \in C^\infty([0, 1], \mathbb{R})$ is defined as in Definition 5.2.

A direct corollary of Proposition 5.3 is the following.

Proposition 5.4. For any $z \in [z_0, z_1]$, let $\theta_z = \arctan(V_z) \in C^\infty([0, 1], \mathbb{R})$, where V_z is defined as in Definition 5.2. Then $(t, z) \mapsto \dot{\theta}_z(t)$ and $(t, z) \mapsto \ddot{\theta}_z(t)$ are bounded w.r.t. $(t, z) \in [0, 1] \times [z_0, z_1]$.

Proof. By definition of V_z , for every $(t, z) \in [0, 1] \times [z_0, z_1]$, $\dot{\theta}_z(t)$ and $\ddot{\theta}_z(t)$ depend only on $\frac{d}{dt}h(u(t), v(t), z)$ and $\frac{d^2}{dt^2}h(u(t), v(t), z)$, which are uniformly bounded w.r.t. $(t, z) \in [0, 1] \times [z_0, z_1]$ because $h \in C^\infty(\mathcal{S}, \mathbb{R})$. \square

Remark 5.5. *In the particular (non-generic) case in which h is constant, the curve (u, v) is non-mixing for all $z \in [z_0, z_1]$, in the sense developed in [7]. Non-mixing curves are characterized by an enhanced adiabatic approximation with respect to general curves passing through an eigenvalue intersection.*

Proposition 5.6. *The functions λ_0^z, λ_1^z , defined as in Proposition 5.3, satisfy (17) with $k = 2$.*

Proof. As a first step of the proof, let us show the following local estimate: There exist $t_1 \in [0, 1)$, a nonempty compact neighborhood $W \subset [z_0, z_1]$ of 0, and $C_1 > 0$ independent of z such that for every $t \in [t_1, 1]$, $j \in \{0, 1\}$, and $z \in W$ we have

$$\left| \int_{t_1}^t e^{\frac{2i}{\epsilon} \int_0^s \lambda_j^z(r) dr} ds \right| \leq C_1 \epsilon^{\frac{1}{3}}, \quad \forall \epsilon > 0. \quad (19)$$

According to Corollary A.8, it is enough to prove that there exist t_1 and W as above such that

$$|\ddot{\lambda}_z(t)| > c, \quad \forall z \in W, t \in [t_1, 1] \setminus \{t_z\}, \quad (20)$$

where $c > 0$ is independent of $z \in W$. Notice that $\lambda_z(t) = |z - m(u(t))u(t)| \sqrt{h_1(u(t), v(t), z)^2 + h_2(u(t), v(t), z)^2}$. By hypothesis (C), $u(1) = \dot{u}(1) = 0$ and $\ddot{u}(1) = 0$. Hence,

$$\left. \frac{d}{dt}(z - m(u(t))u(t)) \right|_{t=1} = 0, \quad \left. \frac{d^2}{dt^2}(z - m(u(t))u(t)) \right|_{t=1} = (z - m(0))\ddot{u}(1),$$

and, in particular, for $z = 0$ we have

$$\lim_{z \rightarrow 0} \left| \frac{d^2}{dt^2} \lambda_z(t) \right|_{t=1} = |m(0)\ddot{u}(1)| > 0.$$

Inequality (20), and hence the required local estimate (19), follow by a continuity argument. Notice that, up to restricting W or increasing t_1 , we can assume that $\{t_z \mid z \in W \cap [z_0, 0]\} = [t_1, 1]$.

Let us now extend (19) to $z \in [z_0, z_1]$ and $t \in [0, 1]$. For $z \in [0, z_1] \setminus W$, there exists $c_1 > 0$ (independent of z) such that $|\lambda_0^z(t)| > c_1 > 0$ for every $t \in [0, 1]$. Hence, by applying Lemma A.9, we have $|\int_0^t e^{\frac{2i}{\epsilon} \int_0^s \lambda_0^z(r) dr} ds| \leq C_1 \epsilon$, where $C_1 > 0$ is independent of $(t, z) \in [0, 1] \times ([0, z_1] \setminus W)$.

For every z in $[z_0, 0)$ we have $\dot{\lambda}_0^z(t_z) \neq 0$. By continuity of the applications $z \mapsto t_z$ and $(t, z) \mapsto \dot{\lambda}_0^z(t)$, there exist $\alpha, c_2 > 0$ such that $|\dot{\lambda}_0^z(t)| > c_2 > 0$ for every $z \in [z_0, 0]$ such that $t_z \leq (t_1 + 1)/2$ and every $t \in [t_z - \alpha, t_z + \alpha]$. By continuity of the application $(t, z) \mapsto \lambda_0^z(t)$, moreover, we get the existence of $c_3 > 0$ such that $|\lambda_0^z(t)| > c_3 > 0$ for every $z \in [z_0, 0] \setminus W$ and every $t \in [0, 1] \setminus [t_z - \alpha, t_z + \alpha]$, also for every $z \in [z_0, 0] \cap W$ and every $t \in [0, t_1] \setminus [t_z - \alpha, t_z + \alpha]$.

For $z \in [z_0, 0] \setminus W$ and $t \in [0, 1]$, we write

$$\begin{aligned} \int_0^t e^{\frac{2i}{\epsilon} \int_0^s \lambda_0^z(r) dr} ds &= \int_{[0, t] \cap [0, t_1 - \alpha]} e^{\frac{2i}{\epsilon} \int_0^s \lambda_0^z(r) dr} ds + \int_{[0, t] \cap [t_1 - \alpha, t_1 + \alpha]} e^{\frac{2i}{\epsilon} \int_0^s \lambda_0^z(r) dr} ds \\ &\quad + \int_{[0, t] \cap [t_1 + \alpha, 1]} e^{\frac{2i}{\epsilon} \int_0^s \lambda_0^z(r) dr} ds, \end{aligned}$$

and we conclude, up a change of time variable, by applying Corollary A.8 (on $[0, t] \cap [t_1 - \alpha, t_1 + \alpha]$, with $k = 2$) and Lemma A.9 (on $[0, t] \cap [0, t_1 - \alpha]$ and $[0, t] \cap [t_1 + \alpha, 1]$).

We conclude similarly for $z \in W \cap [z_0, 0]$, by splitting $[0, 1]$ in the intervals $[0, \min(t_1, t_z - \alpha)]$, $[\min(t_1, t_z - \alpha), t_1]$, and $[t_1, 1]$ and by applying Corollary A.8, Lemma A.9, and (19). \square

Proposition 5.4 and 5.6 allow us to apply Theorem 5.1 and deduce the following ensemble adiabatic approximation result.

Theorem 5.7 (Semi-conical case). *Let (u, v) be a regular C^∞ control path satisfying condition (C). Let ψ_ϵ^z be the solution of Equation (14), where $\tilde{\psi}_0^z = \phi_-^z(u(0), v(0))$ for every $z \in (z_0, z_1]$ and $\tilde{\psi}_0^{z_0} = \lim_{t \rightarrow 0^+} \phi_-^{z_0}(u(t), v(t))$. Set $T_\epsilon(z) = |\langle \psi_\epsilon^z(\frac{1}{\epsilon}), \phi_+^z(u(1), v(1)) \rangle|$. Then*

$$\lim_{\epsilon \rightarrow 0} T_\epsilon(z) = \begin{cases} 0 & \text{if } z \in [0, z_1] \cup \{z_0\}, \\ 1 & \text{if } z \in (z_0, 0), \end{cases}$$

the convergence being uniform w.r.t. $z \in [z_0, z_1]$. More precisely, we have $T_\epsilon(z) = O(\epsilon^{1/3})$ for $z \in [0, z_1] \cup \{z_0\}$.

5.3 The control path (u, v) exits from $\pi(f)$.

By similar arguments as those developed in [4], we get the following proposition.

Proposition 5.8 (Conical exit). *Let $f \in C^\infty(\mathbb{R}^3, \mathbb{R}^2)$. Let (u_1, v_1, z_1) be a F -conical intersection for f . Let N be a neighborhood of (u_1, v_1, z_1) in \mathbb{R}^3 such that $Z(f) \cap N$ is made of F -conical intersections only and $\pi(Z(f) \cap N)$ is a C^∞ submanifold of \mathbb{R}^2 . Let $(u_0, v_0, z_0) \in Z(f) \cap N$ be such that $z_0 < z_1$. Consider a regular C^3 control path $(u(t), v(t))_{t \in [0, 1]}$ and a time $t_1 \in (0, 1)$ such that $(u(0), v(0)) = (u_0, v_0)$, $(u(t_1), v(t_1)) = (u_1, v_1)$, $(u(t), v(t)) \in \pi(Z(f) \cap N)$ for $t \in [0, t_1]$, and $(u(t), v(t)) \notin \pi(f)$ for $t > t_1$. For every $z \in \mathbb{R}$, consider $\theta_z \in C^2([0, 1], \mathbb{R})$ and $\lambda_z \in C^2([0, 1], \mathbb{R})$ as in Theorem 5.1. Then $(t, z) \mapsto \dot{\theta}_z(t)$ and $(t, z) \mapsto \ddot{\theta}_z(t)$ are uniformly bounded with respect to $(t, z) \in [0, 1] \times [z_0, z_1]$, and there exists $c > 0$ such that for every $z \in (z_0, z_1]$ and for every t in $[0, 1]$,*

$$\left| \int_0^t e^{\frac{2i}{\epsilon} \int_0^s \lambda_z(x) dx} ds \right| \leq c\epsilon^{\frac{1}{2}}, \quad \forall \epsilon > 0.$$

Corollary 5.9. *Let $f, (u, v)$ as in Proposition 5.8. Define ψ_ϵ^z as the solution of (14) with $\tilde{\psi}_0^z = \phi_-^z(u(0), v(0))$ for $z \neq 0$ and $\tilde{\psi}_0^{z_0} = \lim_{t \rightarrow 0} \phi_-^{z_0}(u(t), v(t))$. Then for every $z \in (z_0, z_1]$ and for every t in $[0, 1]$,*

$$|\langle \psi_\epsilon^z(1/\epsilon), \phi_+^z(u(1), v(1)) \rangle| = 1 + O(\epsilon^{1/2}).$$

Moreover, for every $\bar{z} \leq z_0$, we have, uniformly w.r.t. $z \in [\bar{z}, z_0]$,

$$|\langle \psi_\epsilon^z(1/\epsilon), \phi_+^z(u(1), v(1)) \rangle| = O(\epsilon^{1/2}).$$

5.4 Proof of Theorem 1.3

Proof of Theorem 1.3. Consider a regular C^4 control path $(\eta(t))_{t \in [0, 1]} = (u(t), v(t))_{t \in [0, 1]}$ such that $\eta(0) = \eta(1)$, $\eta(t_0) = (u_0, v_0)$, $\eta(t_1) = (u_1, v_1)$, $\eta(t) \in \pi(\gamma)$ for $t \in [t_0, t_1]$, and $\eta(t) \notin \pi(f)$ for $t \notin [t_0, t_1]$. Under these hypotheses, we can define, for every $z \in \mathbb{R}$, $\theta_z \in C^2([0, 1], \mathbb{R})$ and $\lambda_z \in C^2([0, 1], \mathbb{R})$ along the path η , as required in Theorem 5.1.

For $t \in [t_0, t_1]$, the hypothesis of non-existence of self-intersections for $\pi(\gamma)$ guarantees that we can apply the same arguments as those used in the proof of Proposition 5.6 for the normal form in order to get

$$\left| \int_{t_0}^t e^{\frac{2i}{\epsilon} \int_0^s \lambda_z(r) dr} ds \right| \leq C\epsilon^{\frac{1}{3}}, \quad \forall \epsilon > 0$$

where $C > 0$ is independent of $(t, z) \in [t_0, t_1] \times [z_0, z_1]$. Moreover by Proposition 5.4, $(t, z) \mapsto \dot{\theta}_z(t)$ and $(t, z) \mapsto \ddot{\theta}_z(t)$ are bounded on $[t_0, t_1] \times [z_0, z_1]$.

Under the assumptions that $z \in [z_0, z_1]$ for every z and t such that $(u(t), v(t), z) \in \gamma$ and that (u_1, v_1, z_1) is a F -conical intersection, we can apply Proposition 5.8 and get that $(t, z) \mapsto \dot{\theta}_z(t)$ and

$(t, z) \mapsto \ddot{\theta}_z(t)$ are uniformly bounded with respect to $(t, z) \in [t_1, 1] \times [z_0, z_1]$, and there exists $c > 0$ such that for every $z \in [z_0, z_1]$ and for every t in $[t_1, 1]$,

$$\left| \int_{t_1}^t e^{\frac{2i}{\epsilon} \int_0^s \lambda_z(x) dx} ds \right| \leq c\epsilon^{\frac{1}{2}}, \quad \forall \epsilon > 0.$$

By similar arguments for $t \in [0, t_0]$, we obtain that on the whole interval $t \in [0, 1]$, $(t, z) \mapsto \dot{\theta}_z(t)$ and $(t, z) \mapsto \theta_z(t)$ are uniformly bounded with respect to $(t, z) \in [0, 1] \times [z_0, z_1]$, and by triangular inequality, there exists $\tilde{C} > 0$ such that for every $z \in [z_0, z_1]$ and for every t in $[0, 1]$,

$$\left| \int_0^t e^{\frac{2i}{\epsilon} \int_0^s \lambda_z(x) dx} ds \right| \leq \tilde{C}\epsilon^{\frac{1}{3}}, \quad \forall \epsilon > 0.$$

We get the expected result by applying Theorem 5.1. □

6 Extension to n -level systems

The goal of this section is to extend Theorem 1.3 of ensemble controllability between the eigenfunctions to the case of n -level systems.

6.1 Generic assumptions on n -level Hamiltonians and adiabatic decoupling

In this section, we show that the study of a n -level real Hamiltonian can be reduced locally to the study of a 2-level Hamiltonian in the adiabatic regime and that such a transformation preserves the codimension of the generic conditions expressed in Section 2.1. Such a reduction allows us to define a semi-conical intersection model for a n -level real Hamiltonian.

For every $H \in S_n(\mathbb{R})$ denote by $(\lambda_j(H))_{j=1}^n$ the spectrum of H , where $j \mapsto \lambda_j(H)$ is the nondecreasing sequence of eigenvalues of H repeated according to their multiplicities. We write $(\phi_1(H), \dots, \phi_n(H))$ to denote an orthonormal basis of associated eigenvectors.

Next lemma is a classical result of continuity of the spectrum (see, for instance [25]).

Lemma 6.1. *Let $H_0 \in S_n(\mathbb{R})$ and $j \in \{1, \dots, n-1\}$ be such that $\lambda_j(H_0), \lambda_{j+1}(H_0)$ are separated from the rest of the spectrum of H_0 . Then, there exists a neighborhood V of H_0 in $S_n(\mathbb{R})$ and a Jordan curve c in \mathbb{C} separating $\{\lambda_q(H) \mid q \in \{j, j+1\}, H \in V\}$ from $\cup_{H \in V} (\text{Spectrum}(H) \setminus \{\lambda_j(H), \lambda_{j+1}(H)\})$.*

From now on, we consider H_0, j, c, V verifying Lemma 6.1. For all $H \in V$, we consider $P_{j,j+1}(H) = \frac{1}{2i\pi} \int_c (H - \xi)^{-1} d\xi$. Notice that, $P_{j,j+1}(H)$ is a real matrix because H is real. By construction of c , $V \ni H \mapsto P_H$ is smooth. Up to reducing V , for every H we can consider an orthogonal mapping $I(H) : \mathbb{R}^2 \rightarrow \text{Im}(P_{j,j+1}(H))$ such that $V \ni H \mapsto I(H)$ is smooth. For every $H \in V$ define $\pi_{j,j+1}(H) = \text{Im}(P_{j,j+1}(H))$, $I^{-1}(H)$ as the inverse of $I(H)$ on $\pi_{j,j+1}(H)$ and

$$F(H) = I^{-1}(H)HI(H) \in S_2(\mathbb{R}).$$

Notice that $I^{-1}(H) = \mathcal{I}(H)$.

Consider $H \in C^\infty(\mathbb{R}^k, S_n(\mathbb{R}))$ such that $H(0) = H_0$, and denote by W a neighborhood of 0 in \mathbb{R}^k such that $H(u) \in V$ for every $u \in W$. Define $h \in C^\infty(\mathbb{R}^k, S_2(\mathbb{R}))$ such that for every $u \in W$, $h(u) = (F \circ H)(u)$. We say that h is a *reduced Hamiltonian* for H . Notice that if $\phi \in \mathbb{C}^2$ is an eigenvector of $h(u)$ associated with the eigenvalue $\lambda \in \mathbb{R}$ then $I(H(u))\phi$ is an eigenvector of $H(u)$ associated with the same eigenvalue λ . We deduce from this, as it has been already used in [4], that the regularity of the eigenpairs of H with respect to $u \in W$ can be deduced from the regularity of those of h .

Proposition 6.2. F is a submersion from V to $S_2(\mathbb{R})$.

Proof. Consider $A \in V$. Define $\psi_1 = I(A)e_1, \psi_2 = I(A)e_2$ where (e_1, e_2) is the canonical basis of \mathbb{C}^2 .

- Define $H = h_{11}\psi_1{}^t\psi_1 + h_{22}\psi_2{}^t\psi_2 + h_{12}\psi_1{}^t\psi_2 + h_{12}\psi_2{}^t\psi_1$ with $h_{11}, h_{22}, h_{12} \in \mathbb{R}$. By direct computations, we get

$${}^tI(A)HI(A) = \begin{pmatrix} h_{11} & h_{12} \\ h_{12} & h_{22} \end{pmatrix}.$$

Hence, the application $S_n(\mathbb{R}) \ni H \mapsto {}^tI(A)HI(A) \in S_2(\mathbb{R})$ is surjective.

- For $H \in S_n(\mathbb{R})$ such that $A + H \in V$,

$$\begin{aligned} F(A + H) &= {}^tI(A + H)(A + H)I(A + H) \\ &= F(A) + {}^tI(A)HI(A) + {}^tDI_A(H)AI(A) + {}^tI(A)ADI_A(H) + o(H). \end{aligned}$$

Hence, $\forall H \in S_n(\mathbb{R})$, $DF_A(H) = {}^tI(A)HI(A) + {}^tDI_A(H)AI(A) + {}^tI(A)ADI_A(H)$. Let us consider $H = h_{11}\psi_1{}^t\psi_1 + h_{22}\psi_2{}^t\psi_2 + h_{12}\psi_1{}^t\psi_2 + h_{12}\psi_2{}^t\psi_1$. Then, we have $DF_A(H) = {}^tI(A)HI(A)$. Hence, F is a submersion. □

Using classical facts on the composition of k -jets (see [24]), we obtain the following result.

Proposition 6.3. Consider $\mathcal{F} : J^2(W, S_n(\mathbb{R})) \rightarrow J^2(W, S_2(\mathbb{R}))$ such that for every $u \in W$, $j^2(h)(u) = \mathcal{F}(j^2(H)(u))$. Then \mathcal{F} is a submersion.

It follows that if S is a codimension q smooth submanifold of $J^2(W, S_2(\mathbb{R}))$, then $\mathcal{F}^{-1}(S)$ is a codimension q smooth submanifold of $J^2(W, S_n(\mathbb{R}))$. This can be used to deduce generic properties for $H \in C^\infty(W, S_n(\mathbb{R}))$ from generic properties for $h \in C^\infty(W, S_2(\mathbb{R}))$.

6.2 Adiabatic decoupling

We present here some results of adiabatic decoupling, adapted from [31].

Theorem 6.4 (Adiabatic decoupling). *Let $H \in C^\infty(\mathbb{R}^k, S_n(\mathbb{R}))$ and $j \in \{1, \dots, n-1\}$ be such that $\{\lambda_q(H(u)) \mid q \in \{j, j+1\}\}$ is separated from $\text{Spectrum}(H(u)) \setminus \{\lambda_q(H(u)) \mid q \in \{j, j+1\}\}$ for u in a neighborhood W of 0 in \mathbb{R}^k . Define, for every $u \in W$, $I(H(u))$ and $h(u)$ as in Section 6.1. Consider a C^2 regular path $u : [0, 1] \rightarrow W$ such that there exist $\ell \in \mathbb{N}$ and C^ℓ functions $\Lambda_j, \Lambda_{j+1} : [0, 1] \rightarrow \mathbb{R}$ such that for every $t \in [0, 1]$, $\{\Lambda_j(t), \Lambda_{j+1}(t)\} = \{\lambda_j(H(u(t))), \lambda_{j+1}(H(u(t)))\}$ and that h admits C^2 eigenvectors along u .*

Assume that there exists $c > 0$ such that

$$\left| \int_0^t e^{\frac{i}{\epsilon} \int_0^s (\Lambda_{j+1}(x) - \Lambda_j(x)) dx} ds \right| \leq c\epsilon^{\frac{1}{\ell+1}}, \quad \forall t \in [0, 1]. \quad (21)$$

Let $\tilde{\psi}_0 \in \mathbb{C}^2$. Then the solutions ψ_ϵ and $\tilde{\psi}_\epsilon$ of, respectively,

$$i \frac{d\psi}{dt} = H(u(\epsilon t))\psi, \quad \psi(0) = I(H(u(0)))\tilde{\psi}_0, \quad \text{and} \quad i \frac{d\tilde{\psi}}{dt} = h(u(\epsilon t))\tilde{\psi}, \quad \tilde{\psi}(0) = \tilde{\psi}_0,$$

are such that $\psi_\epsilon(1/\epsilon)$ is $O(\epsilon^{\frac{1}{\ell+1}})$ -close to $I(H(u(1)))\tilde{\psi}_\epsilon(1/\epsilon)$.

Proof. Define for $q \in \{1, 2\}$, and for every $u \in W$, $\psi_q(u) = I(H(u))e_q$, where (e_1, e_2) is the canonical basis of \mathbb{C}^2 . Define $\psi_\epsilon^{\text{eff}}(t)$ as the solution for $t \in [0, \frac{1}{\epsilon}]$, of

$$i \frac{d\psi_\epsilon^{\text{eff}}(t)}{dt} = \left(h(u(\epsilon t)) - i\epsilon \begin{pmatrix} 0 & \langle \dot{\psi}_j(u(\epsilon t)), \psi_{j+1}(u(\epsilon t)) \rangle \\ \langle \dot{\psi}_{j+1}(u(\epsilon t)), \psi_j(u(\epsilon t)) \rangle & 0 \end{pmatrix} \right) \psi_\epsilon^{\text{eff}}(t), \quad (22)$$

with $\psi_\epsilon^{\text{eff}}(0) = \tilde{\psi}_0$.

By [31, Theorem 1.4], there exists $C > 0$, such that, for every $t \in [0, \frac{1}{\epsilon}]$,

$$\|\psi_\epsilon(t) - I(h(u(\epsilon t)))\psi_\epsilon^{\text{eff}}(t)\| \leq C\epsilon, \quad \forall \epsilon > 0.$$

Under the assumptions of the theorem, we can consider a C^2 basis of eigenvectors of h along u . Hence, by the same arguments as those used in Section 4.1 in order to prove Theorem 4.1, there exists $c > 0$ such that, for every $t \in [0, \frac{1}{\epsilon}]$,

$$\|\tilde{\psi}_\epsilon(t) - \psi_\epsilon^{\text{eff}}(t)\| \leq c\epsilon^{\frac{1}{\ell+1}}, \quad \forall \epsilon > 0.$$

We get the expected result by triangular inequality. \square

In the ensemble case, using estimates that are uniform with respect to the parameter z , we get the following extension of Theorem 6.4.

Theorem 6.5 (Adiabatic decoupling for parametric systems). *Let $H \in C^\infty(\mathbb{R}^{k+1}, S_n(\mathbb{R}))$ and $j \in \{1, \dots, n-1\}$ be such that $\{\lambda_q(H(u, z)) \mid q \in \{j, j+1\}\}$ is separated from $\text{Spectrum}(H(u, z)) \setminus \{\lambda_q(H(u, z)) \mid q \in \{j, j+1\}\}$ for u in a neighborhood W of 0 in \mathbb{R}^k and $z \in [z_0, z_1]$. Define, for every $(u, z) \in W \times [z_0, z_1]$, $I(H(u, z))$ and $h(u, z)$ as in Section 6.1. Consider a C^2 regular path $u : [0, 1] \rightarrow W$ and $\tilde{\psi}_0^z \in \mathbb{C}^2$, for every $z \in [z_0, z_1]$. Let $\ell \in \mathbb{N}$ and assume that for every $z \in [z_0, z_1]$, there exist C^ℓ functions $\Lambda_j^z, \Lambda_{j+1}^z : [0, 1] \rightarrow \mathbb{R}$ such that for every $t \in [0, 1]$, $\{\Lambda_j^z(t), \Lambda_{j+1}^z(t)\} = \{\lambda_j^z(u(t)), \lambda_{j+1}^z(u(t))\}$ and that, for every $z \in [z_0, z_1]$, $h(u(\cdot), z)$ admits C^2 eigenvectors Φ_j^z, Φ_{j+1}^z such that $(t, z) \mapsto \frac{d\Phi_q^z(t)}{dt}$ and $(t, z) \mapsto \frac{d^2\Phi_q^z(t)}{dt^2}$ are bounded uniformly with respect to $(t, z) \in [0, 1] \times [z_0, z_1]$, for every $q \in \{j, j+1\}$. Assume that there exists $c > 0$ such that*

$$\left| \int_0^t e^{\frac{i}{\epsilon} \int_0^s (\Lambda_{j+1}^z(x) - \Lambda_j^z(x)) dx} ds \right| \leq c\epsilon^{\frac{1}{\ell+1}}, \quad \forall t \in [0, 1], \quad \forall z \in [z_0, z_1]. \quad (23)$$

Then the solutions ψ_ϵ^z and $\tilde{\psi}_\epsilon^z$ of, respectively,

$$i \frac{d\psi^z}{dt} = H(u(\epsilon t))\psi^z, \quad \psi^z(0) = I(H(u(0), z))\tilde{\psi}_0^z, \quad \text{and} \quad i \frac{d\tilde{\psi}^z}{dt} = h(u(\epsilon t), z)\tilde{\psi}^z, \quad \tilde{\psi}^z(0) = \tilde{\psi}_0^z,$$

are such that $\psi^z(1/\epsilon)$ is $O(\epsilon^{\frac{1}{\ell+1}})$ -close to $I(H(u(1), z))\tilde{\psi}^z(1/\epsilon)$, uniformly w.r.t. $z \in [z_0, z_1]$.

6.3 Semi-conical intersections for n -level quantum systems

Let $H \in C^\infty(\mathbb{R}^k, S_n(\mathbb{R}))$ and $j \in \{1, \dots, n-1\}$ be such that $\{\lambda_q(H(u)) \mid q \in \{j, j+1\}\}$ is separated from $\text{Spectrum}(H(u)) \setminus \{\lambda_q(H(u)) \mid q \in \{j, j+1\}\}$ for u in a neighborhood W of 0 in \mathbb{R}^k . Define, for every $u \in W$, $I(H(u))$ and $h(u)$ as in Section 6.1. Define for $q \in \{1, 2\}$, and for every $u \in W$, $\psi_q(u) = I(H(u))e_q$, where (e_1, e_2) is the canonical basis of \mathbb{C}^2 . Then we have the identity

$$h(u) = \begin{pmatrix} \langle \psi_1(u), H(u)\psi_1(u) \rangle & \langle \psi_1(u), H(u)\psi_2(u) \rangle \\ \langle \psi_1(u), H(u)\psi_2(u) \rangle & \langle \psi_2(u), H(u)\psi_2(u) \rangle \end{pmatrix}.$$

Definition 6.6. By removing the trace of $H(u)$, define the reduced zero-trace Hamiltonian of H as, for every $u \in W$,

$$h_{\text{red}}(u) = \begin{pmatrix} f_1(u) & f_2(u) \\ f_2(u) & -f_1(u) \end{pmatrix},$$

with $f_1(u) = \frac{1}{2}(\langle \psi_1(u), H(u)\psi_1(u) \rangle - \langle \psi_2(u), H(u)\psi_2(u) \rangle)$ and $f_2(u) = \langle \psi_1(u), H(u)\psi_2(u) \rangle$.

Assume $k = 2$. By a slight abuse of notations, write $u := (u, v) \in U$, where U is a connected open neighborhood of the origin in \mathbb{R}^2 .

Next proposition, which follows by direct computations, states that conicity properties do not depend on the choice of the unitary transformation $I(H(u, v)) : \mathbb{C}^2 \rightarrow \text{Im}(P_{j,j+1}(H(u, v)))$.

Proposition 6.7. Let $f \in C^\infty(U, \mathbb{R}^2)$ and $R \in C^\infty(U, O_2(\mathbb{R}))$. Define $\tilde{f} \in C^\infty(U, \mathbb{R}^2)$ such that $H_{\tilde{f}}(u, v) = R(u, v)H_f(u, v)^t R(u, v)$ for every (u, v) in U . Then

- 0 is conical for f if and only if 0 is conical for \tilde{f} ;
- 0 is semi-conical for f if and only if 0 is semi-conical for \tilde{f} . Moreover, their non-conical directions are the same.

Similarly, one can show that in the ensemble case, F -conical intersections and F -semi-conical intersections are invariant under such a orthogonal mapping, possibly depending on the parameter z . We define semi-conical intersections of eigenvalues for $H(\cdot) \in C^\infty(U, S_n(\mathbb{R}))$ and F -conical (respectively F -semi-conical intersections) for $H(\cdot) \in C^\infty(U \times \mathbb{R}, S_n(\mathbb{R}))$ (see Section 2.24 for precise definitions of these notions for two level systems) as follows.

Definition 6.8. Let $j \in \{1, \dots, n-1\}$.

- We say that $(\bar{u}, \bar{v}) \in U$ is a semi-conical intersection for $H \in C^\infty(U, S_n(\mathbb{R}))$ between the levels j and $j+1$ if and only if there exists a unitary mapping $I(H(u, v)) : \mathbb{C}^2 \rightarrow \text{Im}(P_{j,j+1}(H(u, v)))$, C^∞ with respect to $(u, v) \in U$, such that (\bar{u}, \bar{v}) is a semi-conical intersection for the associated reduced Hamiltonian $h_{\text{red}} \in C^\infty(U, S_2(\mathbb{R}))$.
- We say that $(\bar{u}, \bar{v}, \bar{z}) \in U \times \mathbb{R}$ is a F -conical (respectively F -semi-conical) intersection for $H \in C^\infty(U \times \mathbb{R}, S_n(\mathbb{R}))$ between the levels j and $j+1$ if and only if there exists a unitary mapping $I(H(u, v, z)) : \mathbb{C}^2 \rightarrow \text{Im}(P_{j,j+1}(H(u, v, z)))$, C^∞ with respect to $(u, v, z) \in U \times \mathbb{R}$, such that $(\bar{u}, \bar{v}, \bar{z})$ is a F -conical (respectively F -semi-conical) intersection for the associated reduced Hamiltonian $h_{\text{red}} \in C^\infty(U \times \mathbb{R}, S_2(\mathbb{R}))$.

By Proposition 6.3, we get that F -conical intersections and F -semi-conical intersections as defined in Definition 6.8 are generic for $H \in C^\infty(\mathbb{R}^3, S_n(\mathbb{R}))$ endowed with the Whitney topology.

Remark 6.9. For $j \in \{1, \dots, n-1\}$ set $Z_j = \{(u, v, z) \in U \times \mathbb{R} \mid \lambda_j(u, v, z) = \lambda_{j+1}(u, v, z)\}$. By Definition 6.8, we have the expected result (see Proposition 2.8 for the same property for two-level systems) that if $(\bar{u}, \bar{v}, \bar{z})$ is a F -semi-conical intersection between the levels j and $j+1$, then Z_j is tangent to the plane $z = \bar{z}$ at the point $(\bar{u}, \bar{v}, \bar{z})$ and, considering a local smooth and regular parametrization $(u(t), v(t), z(t))_{t \in [0,1]}$ of Z_j and $\bar{t} \in [0, 1]$ such that $(u(\bar{t}), v(\bar{t}), z(\bar{t})) = (\bar{u}, \bar{v}, \bar{z})$, we have $\dot{z}(\bar{t}) = 0$ and $\ddot{z}(\bar{t}) \neq 0$.

6.4 Controllability result

We consider the controlled Schrödinger equation in \mathbb{C}^n , $n \in \mathbb{N}$,

$$i \frac{d\psi(t)}{dt} = H(u(t), v(t), z)\psi(t). \quad (24)$$

Definition 6.10. Let $z_0, z_1 \in \mathbb{R}$. We say that system (24) is ensemble approximately controllable between eigenstates if for every $\epsilon > 0$, $j, k \in \{1, \dots, n\}$, and $(u_0, v_0), (u_1, v_1) \in U$ such that $\lambda_j(u_0, v_0, z)$ and $\lambda_k(u_0, v_0, z)$ are simple for every $z \in [z_0, z_1]$, there exists a control $(u(\cdot), v(\cdot)) \in L^\infty([0, T], U)$ such that for every $z \in [z_0, z_1]$ the solution of (24) with initial condition $\psi^z(0) = \phi_j^z(u_0, v_0)$ satisfies $\|\psi^z(T) - e^{i\theta} \phi_k^z(u_1, v_1)\| < \epsilon$ for some $\theta \in \mathbb{R}$ (possibly depending on z and ϵ).

For $j \in \{1, \dots, n-1\}$, let us denote by γ_j the set $\{(u, v, z) \in U \times [z_0, z_1] \mid \lambda_j(u, v, z) = \lambda_{j+1}(u, v, z)\}$. Let, moreover, $\gamma_0 = \gamma_n = \emptyset$. Denote by π the projection $\pi : (u, v, z) \mapsto (u, v)$.

Assumption A_j . There exists a connected component $\hat{\gamma}_j$ of γ_j such that

- $\hat{\gamma}_j$ is a one-dimensional submanifold of \mathbb{R}^3 made of F -conical intersections and F -semi-conical intersections only;
- There exist $(u_0, v_0) \in U$ and $(u_1, v_1) \in U$ such that $(u_0, v_0, z_0), (u_1, v_1, z_1) \in \hat{\gamma}_j$ are F -conical intersections for H ;
- $\pi(\hat{\gamma}_j)$ is a C^∞ embedded curve of \mathbb{R}^2 without self-intersections, which is contained in $U \setminus (\pi(\gamma_{j-1}) \cup \pi(\gamma_{j+1}))$.

Using the control strategy proposed in Theorem 1.3 and the result of adiabatic decoupling proposed in Theorem 6.5, we get the following result.

Theorem 6.11. Consider a C^∞ map $U \times [z_0, z_1] \ni (u, v, z) \mapsto H(u, v, z) \in S_n(\mathbb{R})$. Let assumption A_j be satisfied for every $j \in \{1, \dots, n-1\}$. Then system (24) is ensemble approximately controllable between eigenstates.

A Averaging theorems and estimates of oscillatory integrals

The following theorem is a quantitative version in $u(n)$ of a more general averaging result stated in [3, Lemma 8.2]. Its proof is similar to the proof of [3, Lemma 8.2] using an explicit inequality that yields the speed of convergence of order $O(\epsilon)$.

Theorem A.1. Consider A and $(A_\epsilon)_{\epsilon>0}$ in $C^\infty([0, 1], u(n))$ and assume that $A_\epsilon(\tau)$ is uniformly bounded w.r.t. (τ, ϵ) . Denote the flow of the equation $\frac{dx(\tau)}{d\tau} = A(\tau)x(\tau)$ at time τ by $P_\tau \in U(n)$ and the flow of the equation $\frac{dx(\tau)}{d\tau} = A_\epsilon(\tau)x(\tau)$ at time τ by $P_\tau^\epsilon \in U(n)$. If

$$\int_0^\tau A_\epsilon(s) ds = \int_0^\tau A(s) ds + O(\epsilon)$$

then

$$P_\tau^\epsilon = P_\tau + O(\epsilon),$$

both estimates being uniform w.r.t. $\tau \in [0, 1]$.

Remark A.2. Notice that in [3, Lemma 8.2], the hypothesis that $\|A_\epsilon\|_\infty$ is bounded w.r.t. ϵ is not explicitly mentioned, but it is necessary for concluding that $P_\tau^\epsilon \rightarrow P_\tau$ as $\epsilon \rightarrow 0$.

A direct consequence of Theorem A.1 is the following.

Corollary A.3 (Quantum two-level systems averaging). Let $v, \varphi : [0, 1] \rightarrow \mathbb{R}$ be two smooth functions and, for every $\epsilon > 0$, denote by P_τ^ϵ the flow at time $\tau \in [0, 1]$ of

$$i \frac{dX}{d\tau} = \begin{pmatrix} 0 & v(\tau) e^{i\varphi(\tau)} \\ v(\tau) e^{-i\varphi(\tau)} & 0 \end{pmatrix} X(\tau).$$

If $|\int_0^\tau v(s)e^{\frac{i}{\epsilon}\varphi(s)}ds| \leq c\epsilon^q$, where q is a positive real number and $c > 0$ is independent of ϵ, τ , then P_τ^ϵ satisfies $P_\tau^\epsilon = \text{Id} + O(\epsilon^q)$.

We recall a classical result (see [30]) which is useful to estimate oscillatory integrals.

Lemma A.4 (Van Der Corput). *Let $k \in \mathbb{N}$ and $\varphi : [a, b] \rightarrow \mathbb{R}$ be smooth and such that $|\varphi^{(k)}(x)| \geq 1$ for all $x \in [a, b]$. Assume either that $k \geq 2$ or that $k = 1$ and φ' is monotone. Then*

$$\left| \int_a^b e^{i\varphi(x)/\epsilon} dx \right| \leq c_k \epsilon^{1/k},$$

where c_k is independent of φ and ϵ .

In the case $k = 1$, if φ' is not monotone we may lose the uniformity of the estimate with respect to the phase φ . However, we can recover by a direct integration by parts the following estimate.

Lemma A.5 (The case $k = 1$). *Let $\varphi : [a, b] \rightarrow \mathbb{R}$ be smooth and such that $|\varphi'(x)| \geq 1$ for all $x \in [a, b]$. Then*

$$\left| \int_a^b e^{i\varphi(x)/\epsilon} dx \right| \leq 2\epsilon + \epsilon \int_a^b \left| \frac{d}{dx} \frac{1}{\varphi'(x)} \right| dx.$$

By integration by parts we also get the following results.

Corollary A.6. *Let φ and k be as in Lemma A.4. Let, moreover, $v : [a, b] \rightarrow \mathbb{R}$ be smooth. Then*

$$\left| \int_a^b v(x)e^{i\varphi(x)/\epsilon} dx \right| \leq c_k \epsilon^{1/k} \left[|v(b)| + \int_a^b |v'(x)| dx \right]$$

where c_k is the constant obtained in Lemma A.4.

Corollary A.7. *Let φ be as in Lemma A.5 and consider a smooth function $v : [a, b] \rightarrow \mathbb{R}$. Then*

$$\left| \int_a^b v(x)e^{i\varphi(x)/\epsilon} dx \right| \leq c\epsilon$$

where c is independent of ϵ .

Corollary A.8. *Consider an open subset \mathcal{Z} of \mathbb{R} . Assume that $\varphi : [a, b] \times \mathcal{Z} \rightarrow \mathbb{R}$ and $v : [a, b] \times \mathcal{Z} \rightarrow \mathbb{R}$ are real-valued and smooth with respect to the first variable $x \in [a, b]$. Assume that there exists $k > 1$ such that $|\frac{\partial^k \varphi}{\partial x^k}(x, y)| \geq 1$ for all $x \in [a, b]$ and $y \in \mathcal{Z}$. Then*

$$\left| \int_a^b v(x, y)e^{i\varphi(x, y)/\epsilon} dx \right| \leq c_k \epsilon^{1/k} \left[|v(b, y)| + \int_a^b \left| \frac{\partial v}{\partial x}(x, y) \right| dx \right]$$

where c_k is the constant obtained in the Lemma A.4 (independent of φ, y and ϵ). If we assume that v and $\frac{\partial v}{\partial x}$ are uniformly bounded on $[a, b] \times \mathcal{Z}$, then

$$\left| \int_a^b v(x, y)e^{i\varphi(x, y)/\epsilon} dx \right| \leq d_k \epsilon^{1/k}$$

where d_k depends on v and is independent of $\varphi, y \in \mathcal{Z}$ and ϵ .

Next lemma is a direct consequence of Lemma A.5.

Lemma A.9. *Consider a compact subset \mathcal{Z} of \mathbb{R} . Consider two real-valued and smooth functions $\varphi : [a, b] \times \mathcal{Z} \rightarrow \mathbb{R}$ and $v : [a, b] \times \mathcal{Z} \rightarrow \mathbb{R}$. Assume that $|\frac{\partial \varphi}{\partial x}(x, y)| \geq 1$ for all $x \in [a, b]$ and $y \in \mathcal{Z}$. Then*

$$\left| \int_a^b v(x, y)e^{i\varphi(x, y)/\epsilon} dx \right| \leq d\epsilon$$

where d depends on v and φ and is independent of $y \in \mathcal{Z}$ and ϵ .

B Two useful lemmas

We recall some classical results that are derived from [1, §9].

Lemma B.1. *Let $n \in \mathbb{N}$ and let $\mathbb{R}^n \times \mathbb{R} \ni (x, y) \mapsto F(x, y) \in \mathbb{R}$ be a smooth function vanishing on the graph $y = \eta(x)$, where $\eta : \mathbb{R}^n \rightarrow \mathbb{R}$ is a smooth function. Then for every point $x_0 \in \mathbb{R}^n$ there exist a neighborhood W of $(x_0, \eta(x_0))$ and a smooth function $\varphi : W \rightarrow \mathbb{R}$ such that*

$$\forall (x, y) \in W, \quad F(x, y) = (y - \eta(x))\varphi(x, y).$$

Lemma B.2. *Let $n \in \mathbb{N}$ and let $F : \mathbb{R}^n \times \mathbb{R} \rightarrow \mathbb{R}$ be a smooth function such that $\frac{\partial F}{\partial y}$ is vanishing on the smooth hypersurface $y = \eta(x)$. Then for every point $x_0 \in \mathbb{R}^n$ there exist a neighborhood W of $(x_0, \eta(x_0))$ that can be written as $W = W_1 \times W_2$ where W_1 is an open subset of \mathbb{R}^n and W_2 is an open subset of \mathbb{R} , and smooth functions $\varphi : W \rightarrow \mathbb{R}$ and $f_0 : W_1 \rightarrow \mathbb{R}$ such that*

$$\forall (x, y) \in U, F(x, y) = (y - \eta(x))^2\varphi(x, y) + f_0(x).$$

Acknowledgements: This work was supported by the ANR project SRGI ANR-15-CE40-0018 and by the ANR project Quaco ANR-17-CE40-0007-01.

References

- [1] A. Agrachev, D. Barilari, and U. Boscain. *A Comprehensive Introduction to sub-Riemannian Geometry*, volume 181 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 2020.
- [2] A. Agrachev, Y. Baryshnikov, and A. Sarychev. Ensemble controllability by Lie algebraic methods. *ESAIM Control Optim. Calc. Var.*, 22(4):921–938, 2016.
- [3] A. A. Agrachev and Y. L. Sachkov. *Control theory from the geometric viewpoint*, volume 87 of *Encyclopaedia of Mathematical Sciences*. Springer-Verlag, Berlin, 2004. Control Theory and Optimization, II.
- [4] N. Augier, U. Boscain, and M. Sigalotti. Adiabatic ensemble control of a continuum of quantum systems. *SIAM J. Control Optim.*, 56(6):4045–4068, 2018.
- [5] K. Beauchard, J.-M. Coron, and P. Rouchon. Controllability issues for continuous-spectrum systems and ensemble controllability of Bloch equations. *Comm. Math. Phys.*, 296(2):525–557, 2010.
- [6] U. Boscain, J.-P. Gauthier, F. Rossi, and M. Sigalotti. Approximate controllability, exact controllability, and conical eigenvalue intersections for quantum mechanical systems. *Comm. Math. Phys.*, 333(3):1225–1239, 2015.
- [7] U. V. Boscain, F. Chittaro, P. Mason, and M. Sigalotti. Adiabatic control of the Schrödinger equation via conical intersections of the eigenvalues. *IEEE Trans. Automat. Control*, 57(8):1970–1983, 2012.
- [8] S. Chelkowski, A. D. Bandrauk, and P. B. Corkum. Efficient molecular dissociation by a chirped ultrashort infrared laser pulse. *Phys. Rev. Lett.*, 65:2355–2358, Nov 1990.

- [9] C. Chen, D. Dong, R. Long, I. R. Petersen, and H. A. Rabitz. Sampling-based learning control of inhomogeneous quantum ensembles. *Phys. Rev. A*, 89:023402, Feb 2014.
- [10] F. C. Chittaro and J.-P. Gauthier. Asymptotic ensemble stabilizability of the Bloch equation. *Systems Control Lett.*, 113:36–44, 2018.
- [11] F. C. Chittaro and P. Mason. Approximate controllability via adiabatic techniques for the three-inputs controlled Schrödinger equation. *SIAM J. Control Optim.*, 55(6):4202–4226, 2017.
- [12] Y. Colin de Verdière. The level crossing problem in semi-classical analysis. I. The symmetric case. In *Proceedings of the International Conference in Honor of Frédéric Pham (Nice, 2002)*, volume 53, pages 1023–1054, 2003.
- [13] Y. Colin de Verdière. The level crossing problem in semi-classical analysis. II. The Hermitian case. *Ann. Inst. Fourier (Grenoble)*, 54(5):1423–1441, xv, xx–xxi, 2004.
- [14] G. Dirr. Ensemble controllability of bilinear systems. *Oberwolfach Rep.*, 9(1):661–732, 2012.
- [15] U. Gaubatz, P. Rudecki, S. Schiemann, and K. Bergmann. Population transfer between molecular vibrational levels by stimulated raman scattering with partially overlapping laser fields. a new concept and experimental results. *The Journal of Chemical Physics*, 92(9):5363–5376, 1990.
- [16] S. J. Glaser, U. Boscain, T. Calarco, C. P. Koch, W. Köckenberger, R. Kosloff, I. Kuprov, B. Luy, S. Schirmer, T. Schulte-Herbrüggen, D. Sugny, and F. K. Wilhelm. Training Schrödinger’s cat: quantum optimal control. *The European Physical Journal D*, 69(12):279, Dec 2015.
- [17] S. J. Glaser, T. Schulte-Herbrüggen, M. Sieveking, O. Schedletzky, N. C. Nielsen, O. W. Sørensen, and C. Griesinger. Unitary control in quantum ensembles: Maximizing signal intensity in coherent spectroscopy. *Science*, 280(5362):421–424, 1998.
- [18] M. Golubitsky and V. Guillemin. *Stable mappings and their singularities*. Springer-Verlag, New York-Heidelberg, 1973. Graduate Texts in Mathematics, Vol. 14.
- [19] M. Goresky and R. MacPherson. *Stratified Morse Theory*. Springer Berlin Heidelberg, Berlin, Heidelberg, 1988.
- [20] U. Helmke and M. Schönlein. Uniform ensemble controllability for one-parameter families of time-invariant linear systems. *Systems Control Lett.*, 71:69–77, 2014.
- [21] Z. Leghtas, A. Sarlette, and P. Rouchon. Adiabatic passage and ensemble control of quantum systems. *Journal of Physics B: Atomic, Molecular and Optical Physics*, 44(15):154017, 2011.
- [22] J.-S. Li and N. Khaneja. Ensemble control of Bloch equations. *IEEE Trans. Automat. Control*, 54(3):528–536, 2009.
- [23] J.-S. Li and J. Qi. Ensemble control of time-invariant linear systems with linear parameter variation. *IEEE Trans. Automat. Control*, 61(10):2808–2820, 2016.
- [24] P. Libermann and C.-M. Marle. *Symplectic geometry and analytical mechanics*, volume 35 of *Mathematics and its Applications*. D. Reidel Publishing Co., Dordrecht, 1987. Translated from the French by Bertram Eugene Schwarzbach.
- [25] M. Reed and B. Simon. *Methods of Modern Mathematical Physics. IV. Analysis of Operators*. Academic Press, New York, 1978.

- [26] M. Schönlein and U. Helmke. Controllability of ensembles of linear dynamical systems. *Math. Comput. Simulation*, 125:3–14, 2016.
- [27] E. A. Shapiro, V. Milner, and M. Shapiro. Complete transfer of populations from a single state to a preselected superposition of states using piecewise adiabatic passage: Theory. *Phys. Rev. A*, 79:023422, Feb 2009.
- [28] B. W. Shore. *The Theory of Coherent Atomic Excitation, Volume 1, Simple Atoms and Fields*. July 1990.
- [29] T. E. Skinner, T. O. Reiss, B. Luy, N. Khaneja, and S. J. Glaser. Application of optimal control theory to the design of broadband excitation pulses for high-resolution NMR. *Journal of Magnetic Resonance*, 163(1):8 – 15, 2003.
- [30] E. M. Stein. *Harmonic analysis: real-variable methods, orthogonality, and oscillatory integrals*, volume 43 of *Princeton Mathematical Series*. Princeton University Press, Princeton, NJ, 1993. With the assistance of Timothy S. Murphy, Monographs in Harmonic Analysis, III.
- [31] S. Teufel. *Adiabatic perturbation theory in quantum dynamics*, volume 1821 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 2003.
- [32] L. Van Damme, Q. Ansel, S. J. Glaser, and D. Sugny. Robust optimal control of two-level quantum systems. *Phys. Rev. A*, 95:063403, Jun 2017.
- [33] J. von Neumann and E. P. Wigner. *Über das Verhalten von Eigenwerten bei adiabatischen Prozessen*, pages 294–297. Springer Berlin Heidelberg, Berlin, Heidelberg, 1993.