



**HAL**  
open science

## Role of cognitive resources on everyday functioning among oldest-old physically frail

Lucile Dupuy, Bernard N'kaoua, Patrick Dehail, H el ene Sauz eon

► **To cite this version:**

Lucile Dupuy, Bernard N'kaoua, Patrick Dehail, H el ene Sauz eon. Role of cognitive resources on everyday functioning among oldest-old physically frail. *Aging Clinical and Experimental Research*, 2019, 10.1007/s40520-019-01384-3 . hal-02353741

**HAL Id: hal-02353741**

**<https://inria.hal.science/hal-02353741>**

Submitted on 7 Nov 2019

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destin ee au d ep ot et  a la diffusion de documents scientifiques de niveau recherche, publi es ou non,  emanant des  tablissements d'enseignement et de recherche fran ais ou  trangers, des laboratoires publics ou priv es.

1 **Role of cognitive resources on everyday functioning among oldest old physically frail**

2  
3 DUPUY, Lucile<sup>1</sup> \*

4 Email : lucile.dupuy@u-bordeaux.fr

5  
6 N'KAOUA Bernard<sup>2</sup>

7 Email : bernard.nkaoua@u-bordeaux.fr

8  
9 DEHAIL, Patrick<sup>2</sup>

10 Email : patrick.dehail@chu-bordeaux.fr

11  
12 SAUZEON, Hélène<sup>2,3</sup>

13 Email : helene.sauzeon@u-bordeaux.fr

14  
15 1. University of Bordeaux, USR 3413 SANPSY, 33076 Bordeaux Cedex, France

16  
17 2. University of Bordeaux, EA 4136, Handicap, Activity, Cognition, Health, 33000  
18 Bordeaux, France

19  
20 3. Centre Inria Bordeaux Sud-Ouest, Flowers Research Group, 33400 Talence, France

21  
22  
23 **\* Corresponding author:**

24 Lucile.dupuy@u-bordeaux.fr

25 Lucile Dupuy, Ph.D

26 SANPSY – CNRS USR 3413 – Sommeil, Addiction et Neuropsychiatrie

27 University of Bordeaux

28 Site Carreire – Zone Nord, Bat 3B, 3<sup>rd</sup> floor

29 Place Amélie Raba Léon

30 33076 Bordeaux Cedex, France

31 ORCID : 0000-0001-8107-9758

32

1 **Abstract:**

2 **Background:** Everyday functioning becomes a challenge with aging, particularly among frail  
3 oldest-old adults. Several factors have been identified as influencing everyday activities  
4 realization, including physical and cognitive functioning. However, the influence of cognitive  
5 resources as a compensatory factor in the context of physical frailty deserves further  
6 consideration.

7 **Aims:** This study aims to investigate in older adults physically frail the possible compensatory  
8 role of cognitive resources to perform everyday tasks.

9 **Methods:** Two groups of community-dwelling old participants (n=26 per group) matched for  
10 their age and cognitive resources, have been drawn according to their level of physical  
11 functioning. Two measures of everyday functioning have been assessed: one self-reported by  
12 the participant (the IADL scale) and one performance-based measure (the TIADL tasks).

13 **Results:** Participants performed equally the TIADL tasks irrespective of their physical  
14 condition. Contrariwise, participants with low physical functioning reported more everyday  
15 difficulties than their counterparts with a high level of physical functioning. Additionally,  
16 regressions analyses revealed differential influence of cognitive resources on performance and  
17 reported measures of everyday functioning.

18 **Discussion:** Our data suggests that cognitive resources are more strongly involved in the  
19 performance-based IADL measure in situation of physical frailty. Additionally, for participants  
20 with low physical functioning, lower cognitive resources are associated with more perceived  
21 difficulties in everyday life.

22 **Conclusion:** These results highlight the compensatory role of cognitive resources in physically  
23 frail older adults, and suggest that an overestimation of everyday difficulties compared to  
24 performance on IADL tasks is an early indicator of physical decline and cognitive  
25 compensation.

26

27

28 **Keywords:** aging; physical functioning, cognitive resources, daily life activities, compensation

## 1 BACKGROUND

2 Aging is a multifactorial process, which is influenced by a large number of physical,  
3 psychological and social variables. While many persons experience healthy aging without  
4 significant impairments, sensory, motor and cognitive decline can occur with age. In any case,  
5 the capacity to perform activities of daily living can be affected. In this study, we focus on the  
6 compensatory role of cognitive resources on independent everyday functioning among older  
7 adults who have reduced physical functioning.  
8

### 9 **Independent everyday functioning and its assessment**

10 Independent everyday functioning, commonly called functional status, refers to the individual's  
11 abilities to autonomously perform activities of daily life (ADL)[1, 2]. ADLs include basic  
12 (BADLs) and instrumental (IADLs) activities of daily living[3, 4]. The former refer to basic,  
13 physical, self-care tasks, such as ambulating, dressing, grooming, toileting, eating, etc. The  
14 latter activities are more complex self-care tasks, such as meal preparation, medication and  
15 financial management, etc. Hence, IADLs entail more cognitively complex tasks than  
16 BADLs[3]. The IADL-related abilities are often assessed through self-reports or proxy ratings  
17 of an individual's ability to perform activities. IADL questionnaires commonly used in the older  
18 adult population include the Multilevel Assessment Instrument[5], the SF-36[6] and the OARS  
19 Multidimensional Functional Assessment Questionnaire[7]. As these questionnaires can  
20 generate biases and inaccuracies in the informant's perceptions, they are increasingly being  
21 complemented with objective performance measures of physical and cognitive tasks important  
22 for everyday functioning, such as the Timed Instrumental Activities of Daily Living test  
23 (TIADL[8]; for a review of such ADL measures, see[9]). To be noted, while with cognitively  
24 impaired older adults, a discrepancy has been observed between self-reported and objective  
25 measure of everyday functioning[10], in cognitively healthy old people, self-reported ADL  
26 measures still remain strongly related to objective measures of everyday functioning[11, 12].  
27

### 28 **Main underpinnings of age-related changes in independent everyday functioning**

29 Autonomy in daily living has been shown to be influenced by a number of factors. First, ADL  
30 performance is related to socio-demographic variables, such as age, gender, education, and  
31 marital status[1, 13, 14]. For instance, over a longitudinal study with more than 3,000  
32 participants, authors found that women with an advanced age and lower education were more  
33 likely to develop poorer functional status in the next 4 years [13].

34 Cognitive functioning is also well documented as major underpinnings of functional status[11,  
35 15, 16]. Indeed, as noted earlier, BADLs and even more so IADLs require the involvement of  
36 cognitive resources to remember, plan, focus, read or count during everyday activities  
37 realization. Notably, better performance with IADL has been shown to be associated with better  
38 prospective memory (*i.e.*, remember and execute delayed intentions)[13], better inductive  
39 reasoning (*i.e.*, find a common rule from several examples)[16] and better vocabulary[11]. To  
40 note, cognitive functioning in older adults is usually assessed with specialized tools providing  
41 a global measure of cognition (such as the DRS-2 [19]) or with specific measures of cognitive  
42 functions sensitive to aging such as executive functioning (e.g., the FAB [18]).

43 Similarly, there is ample evidence for a positive relationship between poorer physical aptitude  
44 (often called physical frailty[20, 21]) and higher-level ADL loss[22–25]. Indeed, muscle

1 strength, balance, vision, or hearing, are commonly decreasing with age, which may lead to  
2 serious ADL limitations. For instance, an impaired balance renders the gait uncertain (*i.e.*, with  
3 possible negative outcomes like falls or even sickness), which, in turn, may cause a person to  
4 avoid or reduce their mobility[26].

5 Taken together, these findings stress that the ADL decline with aging is multi-determined and  
6 that the understanding of managing aspects of physical and cognitive functioning into ADLs is  
7 likely to be a fruitful way for giving insight into everyday functioning in later life.

8  
9

### 10 **Relationships between Cognitive and Physical functioning to perform ADLs in aging**

11 Beside their independent influence over everyday functioning, there is growing evidence  
12 suggesting that cognitive and physical functioning tend to be closely intertwined in late  
13 adulthood. Notably, in studies using dual tasks[27–29] or even naturalistic tasks like planning  
14 a route while walking[30] it is observed that older adults tend to “prioritize” sensorimotor  
15 processing over cognitive processing. In other words, when dual “cognitive-sensorimotor”  
16 abilities are necessary, for instance when walking while reading a map, older adults tend to  
17 reduce their performance on the cognitive task (*i.e.*, planning) to maintain their sensorimotor  
18 performance (*i.e.*, walking)[30]. Besides, some authors even argue that walking in old age is  
19 *per se* a dual task[29]. This age-related condition directly impacts ADLs which require  
20 simultaneous coordination of physical and cognitive functions.

21 The age-related changes in the coordination of physical and cognitive functions can be  
22 related to the issue of compensation: a strategy responding to functional decline in old age that  
23 is described by the model of selective optimization with compensation (SOC)[31]. Applied to  
24 the situation of gait and balance loss in later life, compensation means that due to a significantly  
25 reduced physical aptitude in daily life, older adults are forced to rely more intensively on  
26 cognitive resources to conduct ADLs[32]. Notably, the study of Heyl and Wahl[2] tested the  
27 involvement of cognitive resources to compensate sensory impairment in the context of ADL  
28 tasks. Indeed, these authors found that cognitive resources and behavior-related everyday  
29 functioning are more closely related in older adults having sensory impairments as compared  
30 to sensory unimpaired counterparts. Authors discussed their results in the light of the SOC  
31 model, and suggest the promotion of cognitive training to support everyday life of elders living  
32 with sensory impairment.

33 In this line of research, the present study attempts to provide further evidence for the  
34 involvement of cognitive resources in the maintenance of ADLs in physically frail oldest-old  
35 adults. The population selected is a sample of cognitively healthy older adults who are divided  
36 into two groups on the basis of their performance on tests assessing their physical functioning  
37 (in terms of mobility, corporal balance, body mass and sensory functions). The expected results  
38 are that older adults having physical disturbances rely more on their cognitive resources to  
39 successfully perform ADL tasks for coping with their physical decline. However, as they are  
40 cognitively spared, they should accurately self-perceive their limitations for carrying out ADL  
41 tasks.

42

43

## 1 **METHODS**

### 2 **Recruitment and final participants**

3 To test our research assumptions, it was essential to include older participants with physical  
4 limitations. For this reason, we collaborated with three public home services for elders  
5 randomly selected among all the Gironde municipalities with a location criterion as follows:  
6 one urban, one semi-urban and one rural location. In France, the level of public financial support  
7 for home service is determined by the functional status. As a result, each file of a home-service  
8 beneficiary is documented by the results of a geriatric assessment performed by medical  
9 consultants, according to the AGGIR scale (gerontological ISO norms for autonomy practiced  
10 in France to establish a person's functional status). Within our sample, the individuals  
11 presenting a dependency syndrome (GIR-score inferior to 4) were excluded from the study.  
12 Among the remaining individuals, we conducted around a hundred of phone calls and 86 older  
13 adults accepted to participate to the study. They underwent a battery of tests. All the interviews  
14 were done at the person's home across two sessions. Thirty-four elders were excluded from the  
15 study because of their MMSE score ( $< 27$ ) to avoid pathological cognitive impairment, such as  
16 dementia cases. Then, we created our two groups of subjects regarding their physical  
17 functioning scores (see below for the calculation of this score). Eventually, the study sample  
18 consisted of 52 community-dwelling old adults aged between 73 to 94 years (mean age  $82.2 \pm$   
19  $4.7$ ); 9 males and 43 females; they were still autonomous and without cognitive impairment.

### 21 **Assessment of Physical functioning**

22 Tasks were selected from widely used clinical and research scales for assessing physical  
23 functioning and subsequent frailty [33, 34] as follows:

24 *Five Chair Stands* (lower body strength): The participant is asked to stand up from a chair five  
25 times without using their arms. The time is recorded and the test is scored from 4 - the  
26 participant takes less than 11.1 sec to complete the task to 0 - the participant is unable to  
27 perform task.

28 *Static Balance Testing* consists of three sorts of standing: side-by-side stand, semi-tandem stand  
29 and tandem stand; each of them scored from 4 - the participant holds the three standing  
30 positions for more than 10 sec; to 0 - the participant did not attempt any standing position.

31 *Timed Get Up and Go Test* (agility and dynamic balance). This test consists of rising from a  
32 chair, walking three meters, turning around, walking back to the chair, and sitting down. Time  
33 in seconds to complete the task is recorded. The task is scored as followed: 1 - the task is  
34 completed in more than 30 sec, 2- the task is completed from 20 to 30 sec, and 3 - the task is  
35 completed in less than 10 sec (in this case, mobility is considered normal).

36 *Gait Speed Test* corresponds to a timed 4-meter walk. It is scored from 4 - time is less than 4.82  
37 sec - to 0 - the participant was unable to do the walk.

38 The score from these four tests ranged from 0 to 13 with higher values indicating greater  
39 mobility function.

40 *Body mass* is also an important component of physical frailty[21]. Thus, two indices have been  
41 scored based on the Mini-Nutritional Assessment[35]. First, the Body Mass Index (BMI),  
42 scored from 0 to 3 with higher values indicating higher BMI values. Second, the brachial and  
43 calf perimeters are scored from 0 to 2 with higher values indicating higher lean mass values.

1 Summed, the two indices provide a score from 0 to 5, with a higher score indicating a better  
2 body mass.

3 Finally, *sensory abilities*, particularly visual acuity and hearing were assessed with a three-point  
4 Likert-type scale, ranging from 0 to 2 (where 0 corresponds to the highest sensory loss). So,  
5 sensory scale provided score ranged from 0 to 4 with higher scores indicating better sensory  
6 functions.

7 So, the score from all test ranged from 0 to 24 with higher values indicating greater physical  
8 functioning (see Table 1). From this score, participants were then divided into two subgroups  
9 based on their physical performance (see below): participants above the physical functioning  
10 score median were considered as ‘high physical functioning’ (n=26, 5 males and 21 females),  
11 whereas other participants were considered as ‘low physical functioning’ (n=26, 4 males and  
12 22 females). These two groups were equivalent in terms of socio-demographic characteristics,  
13 cognitive functioning (including Mini-Mental State Evaluation, MMSE[36]; and Cognitive  
14 resource score described below), and cognitive complaint (assessed by Cognitive Difficulties  
15 Scale, CDS[37]) (see Table 1).  
16  
17

18 *Table 1: Characteristics of the two groups of older participants according to the level of*  
19 *physical functioning (High vs. Low Physical Functioning)*  
20

	Physical Functioning		Group comparison	Effect size ( $\eta^2$ ) Obs. power ( $\beta$ )
	Low Mean (SD)	High Mean (SD)		
Age	83.12 (.88)	81.12 (.95)	$t(50) = 1.54; p = .13$	$\eta^2 = 0.045$
Gender (% female)	80.77	84.61	$t(50) = .36; p = .72$	$\eta^2 = 0.002$
Education years	9.81 (.45)	8.96 (.49)	$t(50) = 1.27; p = .21$	$\eta^2 = 0.031$
Family Status (% married)	23.08	19.23	$t(50) = .333; p = .74$	$\eta^2 = 0.002$
MMSE (max. = 30)	28.26 (0.24)	28.35 (.25)	$t(50) = -.26;$ $p = .78$	$\eta^2 = 0.001$
Physical functioning score (max. = 24)	13.40 (.77)	20.62 (.32)	<b><math>t(50) = -8.67;</math></b> <b><math>p &lt; .0001</math></b>	<b><math>\eta^2 = 0.600</math></b> <b><math>\beta = 1.000</math></b>
Cognitive Resources score (max. = 174)	146.24 (3.31)	150.30 (1.68)	$t(50) = -1.809;$ $p = .28$	$\eta^2 = 0.061$
CDS (max. =144)	42.95 (4.26)	34.33 (3.68)	$t(50) = 1.53;$ $p = .13$	$\eta^2 = 0.045$

21 *Note.* SD=standard deviation; MMSE=Mini-Mental Status Exam; CDS=Cognitive Difficulties Scale.  
22

### 23 **Assessment of cognitive resources**

24 *General cognitive functioning:* The Dementia Rating Scale-2 (DRS-2)[19] has been used. It  
25 assesses five cognitive domains, including attention (e.g., forward and backward digit span,  
26 ability to follow commands), initiation – perseveration (semantic fluency, motor fluency and  
27 perseveration), abstraction (conceptualization from verbal and non-verbal stimuli), visual-  
28 constructional abilities (copy of geometric figures and signature writing) and verbal as well as  
29 non-verbal memory (recall and recognition). This scale gives a score between 0 and 144 (where  
30 144 is the best score).

1 *Executive functioning*: The Frontal Assessment Battery (FAB)[18] has been administrated. It  
2 probes several domains including conceptualization, mental flexibility, motor programming,  
3 resistance to interference, self-regulation, inhibitory control, and environmental autonomy.  
4 FAB gives a score ranging from 0 to 18 (where 18 is the maximum score).  
5 The cognitive resource measure refers to the sum of scores obtained on each scale (with a  
6 maximum score of 162). As indicated in Table 1, the two groups of old participants did not  
7 differ for the cognitive resource measures.

## 8 **Assessment of everyday functioning**

9 *Performance based assessment*. We administrated the TIADL Tasks[8], composed of five  
10 timed tasks that simulate everyday instrumental activities of daily living: communication  
11 (finding a telephone number in the telephone directory), finance (finding and counting out  
12 correct change of money), cooking (finding and reading the ingredients on three food cans),  
13 shopping (finding two specific items in an array of food items) and medicine (finding and  
14 reading the directions on medicine containers). Each task is scored as 1 – completed without  
15 errors and within the time limit, 2 – completed with minor errors, or 3 – not completed within  
16 the time limit, or completed with major errors. Thus, the TIADL gives a score range from 5 to  
17 15, higher scores indicating more difficulties to perform the tasks.

18 *Self-report assessment*. We assess a 24-items scale based on ADL and IADL items [5], where  
19 the answer is based on a 5-point Likert-type format varying from 0 – not at all difficult, to 4 –  
20 very difficult. Then, we selected the 15 IADL items to identify the self-reported IADL score.  
21 Thus, the self-reported IADL gives a score range from 0 to 60, higher scores indicating more  
22 complaint about IADL tasks.

23 *Assessment of Self-perceived health*. We assess the health-related quality of life using the Short  
24 Form-36 (SF-36) questionnaire[6]. This questionnaire consists of 36 items, covering eight  
25 dimensions (physical functioning, physical limitations, body pain, general health, vitality,  
26 social functioning, limitations due to emotional problems, and mental health) and provides two  
27 summary scores, namely physical score and mental score.

## 28 **Statistical analyses**

29 First, group comparisons (Low Physical Functioning vs. High Physical Functioning) have been  
30 performed with the *Student's t test* procedure on each measures of everyday functioning (Table  
31 2). For these tests, t-values, p-values (with a significance level  $<.05$ ) and effect sizes ( $\eta^2$ ) were  
32 reported. As our final sample ended up relatively small, we added post-hoc power sensitivity  
33 analyses (observed power  $\beta$ ) to prevent from Type-I errors (*i.e.*, false positive conclusions)  
34 when significance was observed.

35 Second, global correlations including all participants have been carried out between measures  
36 of everyday functioning and cognitive functioning scores (Table 3). Third, to assess the  
37 influence of the Cognitive Resource factor on IADL measures, two multiple linear regression  
38 analyses were carried out in the two groups independently, with the following statistical design:  
39 physical functioning score and cognitive resource score as predictors, and TIADL and self-  
40 reported IADL scores as criterion variables. In these four regressions, percentage of the  
41 variance explained by the model (Adjusted  $R^2$ ), and unstandardized (B) and standardized ( $\beta$ )  
42

1 regression coefficients for the variables entered into the model were reported (Table 4). Each  
 2 effect size was computed with  $\eta^2$ . All data analysis was performed using SPSS 20.0.

3

4 **RESULTS**

5 **Effects of physical functioning level on everyday functioning (Table 2)**

6 In terms of *performance-based assessment*, the Low and High Physical Functioning groups had  
 7 nearly similar performance on the TIADL test ( $t(50) = .968$  ;  $p > .300$ ;  $\eta^2 = 0.018$ ). Interestingly,  
 8 in terms of *self-report assessment*, self-reported difficulties to perform everyday activities (self-  
 9 report IADL score:  $t(50) = 3.27$ ;  $p = .002$ ) are higher for the Low Physical Functioning group  
 10 than for the High Physical Functioning group. The strong observed power ( $\beta = 89.5\%$ ) and  
 11 effect size ( $\eta^2 = 0.177$ ) values suggest that the difference between the two groups is not a false  
 12 positive.

13 Concerning the *assessment of self-perceived health*, the two groups of participants did not differ  
 14 in terms of self-perceived mental health, *i.e.*, SF-36’s mental sub-score ( $t(50) = -1.32$  ;  $p > .100$ ;  
 15  $\eta^2 = 0.034$ ). By contrast, compared to the High Physical Functioning group, the Low Physical  
 16 Functioning group performed lower on the SF-36’s physical sub-score ( $t(50) = -3.76$ ;  $p < .001$ ),  
 17 indicating a decreased self-perceived physical health, which does not seem to be due to a false  
 18 positive result ( $\eta^2 = 0.261$  ;  $\beta = 95.8\%$ ).

19

20 *Table 2: Performance-based and self-reported measures of Everyday functioning for the two*  
 21 *groups of older participants (High vs. Low Physical Functioning group).*

22

	Physical Functioning		Group Comparison	Effect size ( $\eta^2$ ) Obs. power ( $\beta$ )
	Low Mean (SD)	High Mean (SD)		
TIADL (max. = 15)	6.15 (0.466)	5.65 (0.221)	$t(50) = 0.968$ ; $p = .337$	$\eta^2 = 0.018$
Self-reported IADL (max. = 60)	14.46 (1.72)	7.50 (1.25)	$t(50) = 3.275$ ; $p = .002$	$\eta^2 = 0.177$ $\beta = 0.895$
SF-36 physical (max. = 100)	37.28 (4.04)	57.24 (3.45)	$t(50) = -3.76$ ; $p < .001$	$\eta^2 = 0.261$ $\beta = 0.958$
SF-36 mental (max. = 100)	55.07 (4.56)	63.41 (4.36)	$t(50) = -1.32$ ; $p > .100$	$\eta^2 = 0.034$

23 *Note.* SD=standard deviation; IADL=Instrumental Activities of Daily Living; SF-36=Short Form-36.

24

25 **Global relationships between the everyday functioning measures and the cognitive**  
 26 **resources measures (Table 3)**

27 As illustrated in Table 3, the performance-based and self-reported measures of IADL are  
 28 significantly related with a positive coefficient ( $r = .391$ ;  $p = .004$ ), which means that more  
 29 difficulties in the TIADL tasks are associated with more reported difficulties in everyday  
 30 functioning. Similarly, the physical and mental sub-scores of the SF-36 were strongly related  
 31 with a positive relationship ( $r = .513$ ;  $p < .001$ ).

32

1 *Table 3: Inter-correlations between measures of everyday functioning and cognitive and*  
 2 *physical measures for all participants group.*

	TIADL	Self-report IADL	SF-36 physical	SF-36 mental	Cognitive Resources	Physical Functioning
TIADL	-	<b>.391**</b>	.042	-.058	<b>-.748***</b>	<b>-.343*</b>
Self-reported IADL		-	<b>-.405**</b>	-.220	<b>-.483***</b>	<b>-.533**</b>
SF-36 physical			-	<b>.513***</b>	.062	<b>-.440**</b>
SF-36 mental				-	.253	.074
Cognitive Resources					-	<b>.329*</b>

4 *Notes. \*  $p < .05$ , \*\*\*  $p < .001$ . Significant correlations are bolded*

5  
 6 Regarding the relationships with cognitive resources, both performance-based and self-reported  
 7 measures appeared strongly correlated with cognitive resources but with negative coefficients  
 8 (TIADL:  $r = -.748$ ;  $p < .001$ ; self-reported IADL:  $r = -.483$ ;  $p < .001$ ), which means that greater  
 9 everyday difficulties (observed or reported) are associated with lower cognitive resources.  
 10 Similarly, physical functioning score emerged negatively correlated with TIADL and self-  
 11 reported IADL scores (TIADL:  $r = -.343$ ;  $p = .013$ ; self-perceived IADL:  $r = -.533$ ;  $p < .001$ ),  
 12 suggesting that better physical abilities are associated with lower observed or reported everyday  
 13 difficulties.

#### 15 **Impact of cognitive resources on the relation between physical level and ADL (Table 4)**

16 To assess the relative influence of physical functioning and cognitive resources to manage  
 17 everyday functioning (and highlight a possible compensatory effect of cognitive resources  
 18 when physical functioning is limited), multiple regression analyses have been performed on  
 19 each measure of IADL (TIADL; self-reported IADL) (Table 4).

21 *Table 4: The unstandardized (B), standardized regression coefficients ( $\beta$ ), and percentage of*  
 22 *variance explained (Adjusted  $R^2$ ) for the variables entered into the four models*

Variables	B	SE B	$\beta$	<i>p</i>	Adj $R^2$ (%)
<i>Regression 1 : High PF group – TIADL as criterion variable</i>					
Physical	.002	.115	.002	.989	3.6
<b>Cognitive</b>	<b>-.080</b>	.022	<b>-.605</b>	<b>.001</b>	27.5
<i>Regression 2 : High PF group – self-reported IADL as criterion variable</i>					
<b>Physical</b>	<b>-2.071</b>	.648	<b>-.533</b>	<b>.004</b>	26.3
Cognitive	-.174	.124	-.233	.175	5.2
<i>Regression 3 : Low PF group – TIADL as criterion variable</i>					
Physical	-.103	.084	-.169	.232	16.2
<b>Cognitive</b>	<b>-.100</b>	.019	<b>-.711</b>	<b>&lt; .001</b>	43.2
<i>Regression 4 : Low PF group – self-reported IADL as criterion variable</i>					
Physical	-.281	.422	-.126	.513	6.0
<b>Cognitive</b>	<b>-.254</b>	.098	<b>-.488</b>	<b>.017</b>	18.0

24 *Notes. SE: Standard Error; \*  $p < .05$ , \*\*\*  $p < .001$ . Significant predictors are bolded*

26 The four models appeared significant (*Regression 1:  $F(2, 25) = 6.640$ ;  $p = .005$ ; Regression 2:*  
 27  *$F(2, 25) = 6.744$ ;  $p = .005$ ; Regression 3:  $F(2, 25) = 19.294$ ;  $p < .001$ ; Regression 4:  $F(2,$*

1 25) = 4.948;  $p = .016$ ). In both groups, when the TIADL score is entered as the dependent  
2 variable (Regression 1 and 3), cognitive resources score was a significant predictor, with a  
3 negative relationship with the criterion variable, meaning that more cognitive resources are  
4 associated with better performance at the TIADL (*i.e.*, lower scores). In the Low PF group, the  
5 adjusted  $R^2$  of the cognitive score coefficient is higher compared to the High PF group,  
6 suggesting a stronger importance in variance explanation. In both models, physical functioning  
7 score was not a significant predictor. When the self-reported score is entered as the dependent  
8 variable, physical score emerged as a significant predictor in the High PF group (Regression 2)  
9 with a negative relationship, indicating that lower physical performance is associated with  
10 increased complaint about everyday functioning (*i.e.*, higher scores), and cognitive score  
11 remained non-significant. In the Low PF group however (Regression 4), physical score  
12 appeared non-significant but cognitive score emerged as a significant predictor, with lower  
13 cognitive resources associated with an increased complaint about everyday life.

## 14 **DISCUSSION**

16 Our study is the first to attempt to demonstrate the compensatory role of cognitive resources  
17 among physically frail older adults in everyday functioning. For this purpose, we have recruited  
18 cognitively healthy elders, who differed only depending on their physical functioning level  
19 (Low Physical Functioning *vs.* High Physical Functioning). Performance-based and self-  
20 reported measures of everyday functioning have been collected.

21 A major result is that cognitively healthy elders exhibited similar performance-based  
22 everyday functioning irrespective of their level of physical functioning. In other words, old  
23 participants with low physical functioning successfully handled their physical limitations to  
24 adequately achieve the IADL tasks. Nevertheless, these participants reported more problems  
25 regarding IADL realization and physical health than their counterparts with high physical  
26 functioning. This indicated that even if elders with low physical functioning remain able to  
27 properly perform IADL tasks, they accurately perceive their physical limitations for carrying  
28 out such tasks. These results are fully consistent with our expectations and the literature. Indeed,  
29 this result reinforces the critical role of physical functioning[22, 24, 26] and cognitive health[15,  
30 16, 22] in self-reported everyday functioning and physical health. Additionally, our results  
31 emphasize the possible incongruence between performance and self-reported measure of IADL  
32 in physically frail elders. Indeed, normal cognitive aging studies mostly give evidence for  
33 congruence between performance-based and self-reported scores of everyday difficulties[11,  
34 12], which is perfectly illustrated in our old participants with high physical functioning.  
35 However, when an incongruence between performance and reported IADL (and more precisely  
36 an overestimation of difficulties compared to the actual performance) is observed among  
37 cognitively healthy elders, while some authors proposed the influence of age[38, 39],  
38 cognition[10, 39] or culture[40], this can be an early indicator of physical decline and cognitive  
39 compensation process involved. Hence, ADL performance during performance-based  
40 assessments probably relies more heavily on cognitive resources than on physical functioning,  
41 which is supported by the strong correlation observed between cognitive resources and TIADL  
42 scores (irrespective of physical functioning conditions). Such pattern also underlines the limit  
43 of performance-based assessments, which exclude individual's chosen routines, strategies and  
44 environmental cues that typically facilitate ADL in everyday life[4], and suggest the

1 administration of both performance-based and self-reported measures of everyday difficulties.  
2 To sum up, it is clear that the incongruence reported for cognitive healthy elders with low  
3 physical functioning deserves to be further studied; notably with regard to the potential role of  
4 cognitive functioning as a moderator of physical loss on the performance-based measures of  
5 IADL.

6 Along the line with research by Heyl and Wahl[2] on sensory deficit, we specially  
7 addressed the moderating effect of cognitive resources on physical loss during IADL measures.  
8 Regression results highlighted that the influence of cognitive resource on performance-based  
9 IADL is significantly modified in respect with the level of physical functioning (*i.e.*, adjusted  
10  $R^2$  of the Cognitive resources score with TIADL score as a criterion variable). Importantly, the  
11 importance of cognitive resources to predict the variation of performance on IADL tasks is  
12 significantly increased by a low physical functioning condition compared to the high physical  
13 functioning condition. Consequently, the evidence of a compensatory role of cognitive resource  
14 in performance-based IADL for physically frail elders is provided from this angle. Additionally,  
15 our compensatory assumption is supported by a second major result. The significant predictors  
16 of the self-reported IADL measure differ depending on physical functioning level. The  
17 regressions analyses revealed that the variation of IADL difficulties reported by participants  
18 with low physical functioning is significantly predicted by their level of cognitive resources.  
19 Contrariwise, in participants with high physical functioning, this variation is only predicted by  
20 their level of physical functioning. This means that self-reported IADL varies as follows: for  
21 the old-old participants with high physical functioning, having physical difficulties (even a few)  
22 is the main reason why they would report IADL problems; however, for older adults with low  
23 physical functioning, this is the lack of enough cognitive resources that would predict their self-  
24 reported IADL problems. In other words, in the situation of physical loss, cognitively healthy  
25 older adults who have the least cognitive resources are acutely aware of the cognitive effort  
26 they exert to achieve nominal ADL performance. Thus, the additional cognitive effort for  
27 maintaining IADL performance in physically frail older adults is probably experienced as more  
28 effortful among those having less cognitive resources than those with higher ones. This result  
29 goes further the recently published paper showing that cognitive status indirectly affects  
30 physical aging in the apparition of disability[23], since it proposed an explanation for this  
31 indirect effect, *i.e.*, compensation, as suggested in the SOC Model[2, 31].

32 Some limitations of this study can be noted. The first is related to the modest size of the  
33 sample included in the study (even if we did prevent from false positive conclusions by running  
34 power sensitivity analyses). With a larger sample size, statistical power of our results relative  
35 to compensatory effect could be reinforced. Second, some studies[15, 16] found differential  
36 impact of different cognitive functions on self-reported and performance-based assessment of  
37 everyday functioning, while we used a global measure of cognitive resources. Further works  
38 could investigate the relations between specific cognitive functions (*e.g.*, mental flexibility or  
39 inhibitory control) and measures of everyday functioning for defining more finely which  
40 cognitive resource is involved in the cognitive compensation in situations of physical loss.  
41 Third, we used scores at the TIADL tasks to measure IADL performance (as suggested by the  
42 designer of the test[8]), but finer grain measure of the performance, such as the time used or the

1 type of error (as adapted in[15]) could give further insight of compensation strategies involved  
2 in everyday life.

3 To conclude, our study is the first to highlight a compensatory role of cognitive  
4 resources in response to physical loss, among older-old adults with normal cognitive aging.  
5 Several of our results support this compensation view and give some insights on the  
6 intertwining with aging between physical and cognitive functioning to move forward the field  
7 of aging and everyday functioning. Additionally, underpinning Poli et al.'s recommendations  
8 [25] in their recent study, our findings stress the need for cognitive and psychosocial  
9 interventions for elders with physical loss, in order to counteract their vulnerability induced by  
10 their continuous cognitive efforts in everyday life. In doing so, we would further support the  
11 preservation of older adults' autonomy and promote successful aging.

## 12 **Data availability**

13 The data that support the findings of this study are available from the corresponding author  
14 upon reasonable request.

## 15 **Compliance with ethical standards**

16 **Funding** This work was supported by the National French Institute of Informatics and  
17 Mathematics (Inria), the Public funds *Conseil Régional d'Aquitaine*, CNSA (*Caisse Nationale*  
18 *de la Solidarité pour l'Autonomie*), *Conseil Départemental de la Gironde*, and CARSAT  
19 *Aquitaine (Caisse d'Assurance Retraite et de la Santé au Travail)* and the *Fondation pour la*  
20 *Recherche Médicale*, (grant number : DOC20161136217; PI: Sauzéon).

21 **Conflict of interest** On behalf of all authors, the corresponding author states that there is no  
22 conflict of interest.

23 **Statement of human and animal rights** According to Helsinki declaration, approval was  
24 sought and obtained from the ethics committee of the Bordeaux University.

25 **Informed consent** All participants provided a written consent form prior to participation in the  
26 study.

## 27 **Acknowledgments**

28 The authors are very grateful to the services of UDCCAS Gironde that allowed them access to  
29 the elderly.

## 30 **References**

- 31 1. Diehl M (1998) Everyday Competence in Later Life: Current Status and Future  
32 Directions. *The Gerontologist* 38:422–433. <https://doi.org/10.1093/geront/38.4.422>
- 33 2. Heyl V, Wahl H-W (2012) Managing daily life with age-related sensory loss: Cognitive  
34 resources gain in importance. *Psychol Aging* 27:510–521.  
35 <https://doi.org/10.1037/a0025471>

- 1 3. Lawton MP, Brody EM (1969) Assessment of Older People: Self-Maintaining and  
2 Instrumental Activities of Daily Living. *The Gerontologist* 9:179–186
- 3 4. Gold DA (2012) An examination of instrumental activities of daily living assessment in  
4 older adults and mild cognitive impairment. *J Clin Exp Neuropsychol* 34:11–34.  
5 <https://doi.org/10.1080/13803395.2011.614598>
- 6 5. Lawton MP, Moss M, Fulcomer M, Kleban MH (1982) A Research and Service  
7 Oriented Multilevel Assessment Instrument. *J Gerontol* 37:91–99.  
8 <https://doi.org/10.1093/geronj/37.1.91>
- 9 6. Ware JE, Sherbourne CD (1992) The MOS 36-Item Short-Form Health Survey (SF-36):  
10 I. Conceptual Framework and Item Selection. *Med Care* 30:473–483
- 11 7. Fillenbaum GG (2013) *Multidimensional Functional Assessment of Older Adults : The  
12 Duke Older Americans Resources and Services Procedures*. Psychology Press
- 13 8. Owsley C, Sloane M, McGwin Jr. G, Ball K (2002) Timed Instrumental Activities of  
14 Daily Living Tasks: Relationship to Cognitive Function and Everyday Performance  
15 Assessments in Older Adults. *Gerontology* 48:254–265.  
16 <https://doi.org/10.1159/000058360>
- 17 9. Moore DJ, Palmer BW, Patterson TL, Jeste DV (2007) A review of performance-based  
18 measures of functional living skills. *J Psychiatr Res* 41:97–118.  
19 <https://doi.org/10.1016/j.jpsychires.2005.10.008>
- 20 10. Suchy Y, Kraybill ML, Franchow E (2011) Instrumental activities of daily living among  
21 community-dwelling older adults: Discrepancies between self-report and performance  
22 are mediated by cognitive reserve. *J Clin Exp Neuropsychol* 33:92–100.  
23 <https://doi.org/10.1080/13803395.2010.493148>
- 24 11. Burton CL, Strauss E, Hultsch DF, Hunter MA (2006) Cognitive Functioning and  
25 Everyday Problem Solving in Older Adults. *Clin Neuropsychol* 20:432–452.  
26 <https://doi.org/10.1080/13854040590967063>
- 27 12. Bravell ME, Zarit SH, Johansson B (2011) Self-reported activities of daily living and  
28 performance-based functional ability: a study of congruence among the oldest old. *Eur J  
29 Ageing* 8:199–209. <https://doi.org/10.1007/s10433-011-0192-6>
- 30 13. Pruchno RA, Wilson-Genderson M (2015) A Longitudinal Examination of the Effects of  
31 Early Influences and Midlife Characteristics on Successful Aging. *J Gerontol B Psychol  
32 Sci Soc Sci* 70:850–859. <https://doi.org/10.1093/geronb/gbu046>
- 33 14. Mejía ST, Ryan LH, Gonzalez R, Smith J (2017) Successful Aging as the Intersection of  
34 Individual Resources, Age, Environment, and Experiences of Well-being in Daily  
35 Activities. *J Gerontol B Psychol Sci Soc Sci* 72:279–289.  
36 <https://doi.org/10.1093/geronb/gbw148>
- 37 15. Hering A, Kliegel M, Rendell PG, et al (2018) Prospective Memory Is a Key Predictor  
38 of Functional Independence in Older Adults. *J Int Neuropsychol Soc JINS* 24:640–645.  
39 <https://doi.org/10.1017/S1355617718000152>

- 1 16. Gross AL, Rebok GW, Unverzagt FW, et al (2011) Cognitive Predictors of Everyday  
2 Functioning in Older Adults: Results From the ACTIVE Cognitive Intervention Trial. *J*  
3 *Gerontol Ser B* 66B:557–566. <https://doi.org/10.1093/geronb/gbr033>
- 4 17. Wechsler D (2008) Wechsler adult intelligence scale - Fourth Edition (WAIS-IV). 498
- 5 18. Dubois B, Slachevsky A, Litvan I, Pillon B (2000) The FAB: A frontal assessment  
6 battery at bedside. *Neurology* 55:1621. <https://doi.org/10.1212/WNL.55.11.1621>
- 7 19. Jurica PJ, Leitten CL, Mattis S (1988) DRS-2 dementia rating scale-2: professional  
8 manual. Psychological Assessment Resources
- 9 20. Park H, Jang I-Y, Lee HY, et al (2019) Screening Value of Social Frailty and Its  
10 Association with Physical Frailty and Disability in Community-Dwelling Older Koreans:  
11 Aging Study of PyeongChang Rural Area. *Int J Environ Res Public Health* 16:.  
12 <https://doi.org/10.3390/ijerph16162809>
- 13 21. Fried LP, Tangen CM, Walston J, et al (2001) Frailty in Older Adults: Evidence for a  
14 Phenotype. *J Gerontol Ser A* 56:M146–M157. <https://doi.org/10.1093/gerona/56.3.M146>
- 15 22. Shimada H, Makizako H, Doi T, et al (2015) Incidence of Disability in Frail Older  
16 Persons With or Without Slow Walking Speed. *J Am Med Dir Assoc* 16:690–696.  
17 <https://doi.org/10.1016/j.jamda.2015.03.019>
- 18 23. Makino K, Ihira H, Mizumoto A, et al (2018) Structural analysis of impact of physical,  
19 cognitive and social status on the incidence of disability in community-dwelling people  
20 aged  $\geq 75$  years. *Geriatr Gerontol Int* 18:1614–1619. <https://doi.org/10.1111/ggi.13539>
- 21 24. Seidel D, Brayne C, Jagger C (2011) Limitations in physical functioning among older  
22 people as a predictor of subsequent disability in instrumental activities of daily living.  
23 *Age Ageing* 40:463–469. <https://doi.org/10.1093/ageing/afr054>
- 24 25. Poli S, Cella A, Puntoni M, et al (2017) Frailty is associated with socioeconomic and  
25 lifestyle factors in community-dwelling older subjects. *Aging Clin Exp Res* 29:721–728.  
26 <https://doi.org/10.1007/s40520-016-0623-5>
- 27 26. Owsley C, Allman R, Gossman M, et al (2018) Mobility impairment and its  
28 consequences in the elderly. In: *The Gerontological Prism: Developing Interdisciplinary*  
29 *Bridges: Developing Interdisciplinary Bridges*, CRC Press. Jeffrey Michael Clair and  
30 Richard M. Alzman
- 31 27. Li KZH, Lindenberger U, Freund AM, Baltes PB (2001) Walking While Memorizing:  
32 Age-Related Differences in Compensatory Behavior. *Psychol Sci* 12:230–237.  
33 <https://doi.org/10.1111/1467-9280.00341>
- 34 28. Poole VN, Lo O-Y, Wooten T, et al (2019) Motor-Cognitive Neural Network  
35 Communication Underlies Walking Speed in Community-Dwelling Older Adults. *Front*  
36 *Aging Neurosci* 11:.. <https://doi.org/10.3389/fnagi.2019.00159>
- 37 29. Wollesen B, Wanstrath M, van Schooten KS, Delbaere K (2019) A taxonomy of  
38 cognitive tasks to evaluate cognitive-motor interference on spatiotemporal gait

- 1 parameters in older people: a systematic review and meta-analysis. *Eur Rev Aging Phys*  
2 *Act* 16:12. <https://doi.org/10.1186/s11556-019-0218-1>
- 3 30. Taillade M, Sauzéon H, Arvind Pala P, et al (2013) Age-related wayfinding differences  
4 in real large-scale environments: detrimental motor control effects during spatial  
5 learning are mediated by executive decline? *PloS One* 8:e67193.  
6 <https://doi.org/10.1371/journal.pone.0067193>
- 7 31. Baltes PB, Baltes MM (1993) Psychological perspectives on successful aging: The  
8 model of selective optimization with compensation. In: *Successful Aging: Perspectives*  
9 *from the Behavioral Sciences*. Cambridge University Press, pp 1–35
- 10 32. Zhang W, Radhakrishnan K (2018) Evidence on selection, optimization, and  
11 compensation strategies to optimize aging with multiple chronic conditions: A literature  
12 review. *Geriatr Nurs N Y N* 39:534–542. <https://doi.org/10.1016/j.gerinurse.2018.02.013>
- 13 33. Guralnik JM, Simonsick EM, Ferrucci L, et al (1994) A Short Physical Performance  
14 Battery Assessing Lower Extremity Function: Association With Self-Reported Disability  
15 and Prediction of Mortality and Nursing Home Admission. *J Gerontol* 49:M85–M94.  
16 <https://doi.org/10.1093/geronj/49.2.M85>
- 17 34. Bruyère O, Buckinx F, Beaudart C, et al (2017) How clinical practitioners assess frailty  
18 in their daily practice: an international survey. *Aging Clin Exp Res* 29:905–912.  
19 <https://doi.org/10.1007/s40520-017-0806-8>
- 20 35. Vellas B, Guigoz Y, Garry PJ, et al (1999) The mini nutritional assessment (MNA) and  
21 its use in grading the nutritional state of elderly patients. *Nutrition* 15:116–122.  
22 [https://doi.org/10.1016/S0899-9007\(98\)00171-3](https://doi.org/10.1016/S0899-9007(98)00171-3)
- 23 36. Folstein MF, Folstein SE, McHugh PR (1975) “Mini-mental state.” *J Psychiatr Res*  
24 12:189–198. [https://doi.org/10.1016/0022-3956\(75\)90026-6](https://doi.org/10.1016/0022-3956(75)90026-6)
- 25 37. McNair DM, Kahn RJ, Crook SFT, Bartus R (1983) The cognitive difficulties scale.  
26 *Assess Geriatr Psychopharmacol* 137:143
- 27 38. Wettstein M, Schilling OK, Wahl HW (2016) “Still feeling healthy after all these years”:  
28 The paradox of subjective stability versus objective decline in very old adults’ health and  
29 functioning across five years. *Psychol Aging* 31:815–830.  
30 <https://doi.org/10.1037/pag0000137>
- 31 39. Fors S, Thorslund M, Parker MG (2006) Do actions speak louder than words? Self-  
32 assessed and performance-based measures of physical and visual function among old  
33 people. *Eur J Ageing* 3:15–21. <https://doi.org/10.1007/s10433-006-0021-5>
- 34 40. Spitzer S, Weber D (2019) Who is telling the truth? Biases in self-reported physical and  
35 cognitive health status of older Europeans. *IIASA Work Pap WP-19-002*

36