



Sport Psychology: Technologies Ahead

Camille Jeunet, Denis Hauw, José del R. Millán

► **To cite this version:**

Camille Jeunet, Denis Hauw, José del R. Millán. Sport Psychology: Technologies Ahead. *Frontiers in Sports and Active Living*, Frontiers Media S.A., 2020, 2, 10.3389/fspor.2020.00010 . hal-02476860

HAL Id: hal-02476860

<https://hal.inria.fr/hal-02476860>

Submitted on 13 Feb 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Sport Psychology: Technologies Ahead

Camille Jeunet^{1,2*}, Denis Hauw^{3*} and Jose del R. Millán^{2,4,5*}

¹ CLLE Lab, CNRS, Univ. Toulouse Jean Jaurès, Toulouse, France, ² École Polytechnique Fédérale de Lausanne, Geneva, Switzerland, ³ Institut des Sciences du Sport, Université de Lausanne, Lausanne, Switzerland, ⁴ Department of Electrical and Computer Engineering, The University of Texas at Austin, Austin, TX, United States, ⁵ Department of Neurology, The University of Texas at Austin, Austin, TX, United States

Keywords: training, motor-imagery, brain-computer interface (BCI), brain stimulation, neurophenomenology, virtual reality, sport

1. VIRTUAL REALITY AND MOTOR IMAGERY

Several factors may make it difficult, if not impossible, for athletes to train in real sport conditions. These include on the one hand extrinsic factors such as unfavorable weather conditions or costs related to the facilities and materials used, and on the other hand factors intrinsic to the athlete, such as injuries.

When confronted to such situations, it becomes necessary for the athletes to be provided with alternative training solutions in order to maintain their level of performance. In this section, we will introduce two of these alternative solutions, namely virtual reality (VR) and motor imagery (MI). The first alternative training solution we will mention here is Virtual Reality (VR). VR is a computer-based simulation that is both interactive and immersive. It provides the opportunity to design fully controlled training procedures that are both adaptable to each athlete and ecological. Coupled with the fact that VR enables athletes to train in environments that may not be readily accessible, these properties make VR a very relevant and promising training tool to improve sport performance. Several studies have already revealed that athletes who had trained in a virtual environment obtained performances similar to those who had trained in real sport conditions (Bideau et al., 2004, 2010; Vignais et al., 2015) (for a review, see Neumann et al., 2018) suggesting that VR can provide a sufficiently high level of qualitative feedback to be used as a training tool (e.g., see Miles et al., 2012 for a review of virtual environments designed for ball sport training). Nonetheless, Neumann et al. (2018) stress the fact that the benefits of VR for enhancing physiological, psychological and motor performance can be modulated by different factors related to the athletes themselves, but also to the VR system and environment.

The second alternative training solution is Motor Imagery. MI has been defined as a mental representation of an action without any concomitant body movement (Guillot and Collet, 2008). More specifically, Morris et al. (2005) have stated that in sports, MI “may be considered as the creation or recreation of an experience generated from memorial information, involving quasi-sensorial, quasi-perceptual, and quasi-affective characteristics, that is under the volitional control of the imager, and which may occur in the absence of the real stimulus antecedents normally associated with the actual experience.” MI is extensively practiced by athletes, not only when they cannot access real sport conditions, but also as a complementary training procedure or for preparation in competition. Formal studies have repeatedly shown that, indeed, this practice enabled the improvement motor performance (Guillot and Collet, 2008), and notably of strength (Lebon et al., 2010). In the last decade, extensive research has been led in order to better understand (Moran et al., 2019) and model (Guillot and Collet, 2008) the mechanisms underlying the influence that MI has on motor performance, but also to better understand how the sport that is practiced influences MI abilities (Di Corrado et al., 2019), in order then to design efficient training

OPEN ACCESS

Edited by:

Adelaida María A. M. Castro Sánchez,
University of Almería, Spain

Reviewed by:

Marinella Coco,
University of Catania, Italy

*Correspondence:

Camille Jeunet
camille.jeunet@univ-tlse2.fr
Denis Hauw
Denis.Hauw@unil.ch
Jose del R. Millán
jose.millan@austin.utexas.edu

Specialty section:

This article was submitted to
Movement Science and Sport
Psychology,
a section of the journal
Frontiers in Sports and Active Living

Received: 16 July 2019

Accepted: 24 January 2020

Published: 13 February 2020

Citation:

Jeunet C, Hauw D and Millán JdR
(2020) Sport Psychology:
Technologies Ahead.
Front. Sports Act. Living 2:10.
doi: 10.3389/fspor.2020.00010

procedures. Nonetheless, the main limitation of current MI training procedures is the absence of feedback. In other words, when they practice MI, athletes are provided with no information regarding their performance—which may be measured, for instance, by the extent to which they modulate their sensorimotor rhythms (SMR). As a consequence, they cannot assess the efficiency of their MI training nor know how to adapt in order to improve this capacity. One promising solution to overcome this limitation consists in measuring the athletes' brain activity while they perform MI, and in providing them with real-time feedback indicating how well they modulate the target brain rhythms, e.g., their SMR. A basic form of such paradigm is called neurofeedback. Next section briefly describes the benefits of technologies called Brain-Computer Interfaces (BCIs), that are based on neurofeedback, and how to modulate brain activity by applying small electrical currents.

2. BRAIN-COMPUTER INTERFACES AND BRAIN STIMULATION

In a brain-computer interfaces (BCIs), neural signals recorded from the brain are fed into a decoding algorithm that translates these signals into outputs (e.g., Millán, 2019 and references therein). Compared to neurofeedback—where users modulate a single, usually predefined, physiological component—a BCIs rely on multivariate brain feature processing algorithms for feature selection and decoding, which makes it possible to control in real time a variety of brain-actuated devices. Although BCIs are being extensively explored as new control and communication modalities for people with physical disabilities, including new forms of sports (Perdikis et al., 2018), as well as neurorehabilitation (Biasiucci et al., 2018; Cervera et al., 2018), their use in the framework of sports is still at its infancy. Here we shortly discuss two initial attempts. Jeunet et al. (2018) hypothesized that a BCI based on EEG signatures of athletes' cognitive abilities could help them to train those abilities and contribute to further improve their athletic performance. They present preliminary results on the possibility of using such a kind of BCI aiming to increase soccer goalkeepers' performance through the improvement of their covert visuo-spatial attention (CVSA) abilities—which is key for these athletes and is characterized by modulations of patterns of alpha synchronization over parieto-occipital areas. Our second example combines a BCI with another technology covered in this paper, namely VR. Pereira et al. (2018) investigated EEG correlates of realistic training and competition scenarios while novice shooters were immersed in a 10 m Olympic pistol shooting environment. In the training scenario, shooters remained alone; while in the competition scenario there were other competitors and an audience. Authors found a differential between-subject effect of competition on μ (8–12 Hz) oscillatory activity during aiming; compared to training, the more the subject was able to desynchronize their μ rhythm during competition, the better was their shooting performance. These results provide evidence that μ desynchronization has a positive effect on performance during competition.

Rather than training subjects to learn to modulate their brain rhythms and acquire BCI skills, an alternative is to use brain stimulation techniques, in particular non-invasive modalities such as transcranial direct current stimulation (tDCS). tDCS applies low-amplitude constant currents via scalp electrodes and modify neuronal transmembrane polarization if applied sufficiently long (10–20 min), thereby inducing cortical excitability changes in humans (Nitsche and Paulus, 2000). Modulatory effects of tDCS depend on the current direction: anodal stimulation has an excitatory effect, while cathodal stimulation yields an inhibitory effect. Thus, anodal tDCS over the motor cortex has shown to improve motor learning in healthy subjects (Reis et al., 2009). These initial findings are triggering substantial attention in sports, although early attempts report a mixture of positive (yet without proper control studies) and negative outcomes. As an example, Angius et al. (2016) found that anodal tDCS over the motor cortex led to a reduced perception of effort and increased endurance in amateur cyclist when the cathode was placed on the contralateral shoulder.

3. BACK TO THE BEGINNING: NEUROPHENOMENOLOGY

One future challenge is to link EEG with the understanding of various levels of organization of human activity during sports in ecological settings. This means that (a) phenomenological dimensions of athletes' activity (i.e., being ready to move, feeling comfortable when acting, having good sensations when performing) might be taking into account, (b) portable electroencephalogram systems could record athletes' brain signals during performance, and (c) advanced bioengineering techniques to decode these signals in real time as well as analyzing the experiences with advanced phenomenological techniques should be developed. While neurophenomenological methods have already been used to investigate the neural correlates of several sense-making processes in subjective experience in human activities such as meditation or epileptic seizure (e.g., Petitmengin et al., 2006), rare attempts have been made in sports.

In an exploratory study, we combined brain activity using EEG (third person data) with first-person qualitative analysis of pre-reflective experience emerging during a standing split task in dance. We expected that brain activity signatures underlying the consciousness of various efficient ways of being-and-acting in this task would be identified. Data was collected through (a) video tapes of the athletes' behavior while performing the sporting task, (b) video-recorded and transcribed verbalizations and commentaries elicited post-action during self-confrontation interviews, and (c) 64-electrode EEG recordings made during sport performance. Qualitative analyses were performed on the data from the self-confrontation interviews using the course-of-action method, which consists in identifying elementary units of meaning and their components (involvement, focus and general feelings) (e.g., Hauw and Durand, 2007; Hauw, 2018). This method identified different sequences of organization of the athletes' activities. Results illustrated that each participant typically organized their activity around three or more sequences, but that these sequences were very distinct according to their

expertise level. Advanced EEG analysis allowed us to study neural activity despite the signal contamination due to movement-related artifacts during the task execution. Preliminary results showed the possibility of extracting neural activity that was correlated with movement preparation (Bereitschafts potential).

In conclusion, despite the emergence of some research on the topic, the field of neurotechnologies for sport psychology is still in its infancy and has to face a number of technological difficulties. The conceptual and methodological design might be a relevant direction for future development for sports learning and training.

REFERENCES

- Angius, L., Pageaux, B., Hopker, J., Marcora, S., and Mauger, A. (2016). Transcranial direct current stimulation improves isometric time to exhaustion of the knee extensors. *Neuroscience* 339, 363–375. doi: 10.1016/j.neuroscience.2016.10.028
- Biasiucci, A., Leeb, R., Iturrate, I., Perdikis, S., Al-Khodairy, A., Corbet, T., et al. (2018). Brain-actuated functional electrical stimulation elicits lasting arm motor recovery after stroke. *Nat. Commun.* 9:2421. doi: 10.1038/s41467-018-04673-z
- Bideau, B., Kulpa, R., Vignais, N., Brault, S., Multon, F., and Craig, C. (2010). Using virtual reality to analyze sports performance. *IEEE Comput. Graph. Appl.* 30, 14–21. doi: 10.1109/MCG.2009.134
- Bideau, B., Multon, F., Kulpa, R., Fradet, L., Arnaldi, B., and Delamarche, P. (2004). Using virtual reality to analyze links between handball thrower kinematics and goalkeeper's reactions. *Neurosci. Lett.* 372, 119–122. doi: 10.1016/j.neulet.2004.09.023
- Cervera, M., Soekadar, S., Ushiba, J., Millán, J. d. R., Liu, M., Birbaumer, N., et al. (2018). Brain-computer interfaces for post-stroke motor rehabilitation: a meta-analysis. *Ann. Clin. Transl. Neurol.* 5, 651–663. doi: 10.1002/acn3.544
- Di Corrado, D., Guarnera, M., Vitali, F., Quartiroli, A., and Coco, M. (2019). Imagery ability of elite level athletes from individual vs. team and contact vs. no-contact sports. *PeerJ* 7:e6940. doi: 10.7717/peerj.6940
- Guillot, A., and Collet, C. (2008). Construction of the motor imagery integrative model in sport: a review and theoretical investigation of motor imagery use review and theoretical investigation of motor imagery use. *Int. Rev. Sport Exerc. Psychol.* 1, 31–44. doi: 10.1080/17509840701823139
- Hauw, D. (2018). Enaction et intervention en psychologie du sport chez les sportifs élités et en formation. *Can. J. Behav. Sci.* 50, 54–64. doi: 10.1037/cbs0000094
- Hauw, D., and Durand, M. (2007). Situated analysis of elite trampolinists' problems in competition using retrospective interviews. *J. Sport Sci.* 25, 173–183. doi: 10.1080/02640410600624269
- Jeunet, C., Tonin, L., Albert, L., Chavarriaga, R., Bideau, B., Argelaguet, F., et al. (2020). Uncovering EEG correlates of covert attention in soccer goalkeepers: towards innovative sport training procedures. *Sci. Rep.* 10:1705. doi: 10.1038/s41598-020-58533-2
- Lebon, F., Collet, C., and Guillot, A. (2010). Benefits of motor imagery training on muscle strength. *J. Strength Cond. Res.* 24, 1680–1687. doi: 10.1519/JSC.0b013e3181d8e936
- Miles, H. C., Pop, S. R., Watt, S. J., Lawrence, G. P., and John, N. W. (2012). A review of virtual environments for training in ball sports. *Comput. Graph.* 36, 714–726. doi: 10.1016/j.cag.2012.04.007
- Millán, J. d. R. (2019). The human-computer connection: an overview of brain-computer interfaces. *Métode* 9, 135–141. doi: 10.7203/metode.9.12639
- Moran, A., Campbell, M., and Toner, J. (2019). Exploring the cognitive mechanisms of expertise in sport: Progress and prospects. *Psychol. Sport Exerc.* 42, 8–15. doi: 10.1016/j.psychsport.2018.12.019
- Morris, T., Spittle, M., and Watt, A. P. (2005). *Imagery in Sport*. Champaign, IL: Human Kinetics.
- Neumann, D. L., Moffitt, R. L., Thomas, P. R., Loveday, K., Watling, D. P., Lombard, C. L., et al. (2018). A systematic review of the application of interactive virtual reality to sport. *Virtual Real.* 22, 183–198. doi: 10.1007/s10055-017-0320-5
- Nitsche, M., and Paulus, W. (2000). Excitability changes induced in the human motor cortex by weak transcranial direct current stimulation. *J. Physiol.* 527, 633–639. doi: 10.1111/j.1469-7793.2000.t01-1-00633.x
- Perdikis, S., Tonin, L., Saeedi, S., Schneider, C., and Millán, J. d. R. (2018). The Cybathlon BCI race: successful longitudinal mutual learning with two tetraplegic users. *PLoS Biol.* 16:e2003787. doi: 10.1371/journal.pbio.2003787
- Pereira, M., Argelaguet, F., Millán, J. d. R., and Lécuyer, A. (2018). Novice shooters with lower pre-shooting alpha power have better performance during competition in a virtual reality scenario. *Front. Psychol.* 9:527. doi: 10.3389/fpsyg.2018.00527
- Petitmengin, C., Baulac, M., and Navarro, V. (2006). Seizure anticipation: are neurophenomenological approaches able to detect preictal symptoms? *Epilepsy Behav.* 9, 298–306. doi: 10.1016/j.yebeh.2006.05.013
- Reis, J., Schambra, H. M., Cohen, L. G., Buch, E. R., Fritsch, B., Zarahn, E., et al. (2009). Noninvasive cortical stimulation enhances motor skill acquisition over multiple days through an effect on consolidation. *Proc. Natl. Acad. Sci. U.S.A.* 106, 1590–1595. doi: 10.1073/pnas.0805413106
- Vignais, N., Kulpa, R., Brault, S., Presse, D., and Bideau, B. (2015). Which technology to investigate visual perception in sport: video vs. virtual reality. *Hum. Mov. Sci.* 39, 12–26. doi: 10.1016/j.humov.2014.10.006

AUTHOR CONTRIBUTIONS

CJ, DH, and JM all took part in the writing of this opinion paper.

FUNDING

This work was supported by the Swiss National Foundation (SNF) for scientific research and by the EPFL/Inria International Lab.

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2020 Jeunet, Hauw and Millán. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.