



HAL
open science

Diagnosing the ocean-atmosphere coupling schemes by using a mathematically consistent Schwarz iterative method

Olivier Marti, Sébastien Nguyen, Pascale Braconnot, Sophie Valcke, Florian Lemarié, Eric Blayo

► To cite this version:

Olivier Marti, Sébastien Nguyen, Pascale Braconnot, Sophie Valcke, Florian Lemarié, et al.. Diagnosing the ocean-atmosphere coupling schemes by using a mathematically consistent Schwarz iterative method. [Research Report] Research activities in Earth system modelling. Working Group on Numerical Experimentation. Report No. 51. WCRP Report No.4/2021. WMO, Geneva. 2021. hal-03352312

HAL Id: hal-03352312

<https://hal.inria.fr/hal-03352312>

Submitted on 23 Sep 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Diagnosing the ocean-atmosphere coupling schemes by using a mathematically consistent Schwarz iterative method

Olivier Marti¹, Sébastien Nguyen¹, Pascale Braconnot¹, Sophie Valcke², Florian Lemarié³,
and Eric Blayo³

¹Laboratoire des Sciences du Climat et de l'Environnement, LSCE/IPSL, CEA-CNRS-UVSQ,
Université Paris-Saclay, Gif-sur-Yvette, France

²CECI, Université de Toulouse, CNRS, CERFACS, Toulouse, France

³Univ. Grenoble Alpes, Inria, CNRS, Grenoble INP, LJK, 38000, Grenoble, France

Correspondence: olivier.marti@lsce.ipsl.fr

1 Introduction

Coupling algorithms implemented in coupled general circulation models (CGCMs) are driven by the necessity to conserve energy and mass at the air-sea interface. The discretization of the coupling problem leads to inconsistencies in time. It splits the simulation into small time intervals (coupling periods) over which averaged-in-time boundary data are exchanged. Atmosphere computes the fluxes at the interface (heat, water and momentum), and ocean computes the sea surface properties (water and sea ice temperatures, sea ice fraction, albedos, surface current). Two main algorithms are used, the *parallel* and the *sequential atmosphere-first*. In the *parallel* algorithm, both models run a coupling period with the boundary conditions of the other model from the previous coupling period. In the *sequential atmosphere-first* algorithm, atmosphere runs with oceanic boundary conditions from the previous coupling time. Then ocean runs with atmospheric boundary conditions of the current coupling period. Both algorithms are lagged: there is a time lag (of one coupling period) between the model and its boundary conditions, for one (*sequential* algorithm) or both (*parallel* algorithm) models.

Schwarz algorithms are attractive iterative coupling methods to cure the temporal inconsistencies and provide tightly and mathematically consistent coupled solutions. As discussed in Lemarié (2008), the standard coupling methods correspond to one single iteration of a global-in-time iterative Schwarz method.

2 The Schwarz iterative method

A Schwarz iterative method has been implemented in the IPSLCM5A Earth System model (Sepulchre et al., 2020; Marti et al., 2010, 2021). The model repeats each coupling period, with the same initial state (the result from the previous coupling period). The boundary conditions are updated at each iteration, using the result of the previous iteration, until convergence. The solution is mathematically consistent: during a coupling

period, each model uses the boundary conditions from the other model for the same coupling period. There is no lag.

The method has a huge computational cost, as each coupling period is iterated. It is not affordable for production runs, but allows us to compare the legacy coupling algorithms used in state-of-the art GCM to a mathematically consistent and synchronous algorithm, which is used as a reference to compute the error made with legacy algorithms.

We run 5-day experiments with the Schwarz iterative method, using three algorithms (*parallel*, *sequential atmosphere-first* and *sequential ocean-first*), and two coupling periods $\Delta t = 1h$ and $\Delta t = 4h$. The Schwarz method can be used over all these algorithms. If the method is run until perfect convergence, the result is not dependant of the underlying coupling algorithm. In practice, the iterative method will be stopped before perfect convergence with an ad hoc criterion, and the different algorithms will give slightly different results.

3 Results

Fig. 1 shows the relative error in the change of sea surface temperature (SST) during one coupling period when the Schwarz method is not used. At each Schwarz iteration, the model computes an occurrence of the SST trend. At the first iteration, the trend is the one that the model would calculate with the legacy lagged coupling. It is compared with the trend obtained after convergence. The comparison is done on a unique trajectory of the model. The error is computed as the ratio between the correction due to the iterative procedure and the solution change with no Schwarz iteration. We consider each case of the iterative process, i.e. each point of the atmosphere grid, and for each coupling period.

In the *parallel*- $\Delta t = 1h$ experiment, the relative error is negligible (less than 0.01) in about 15 % of the cases. It is small (less than 0.1) in almost 50 % of the




Figure 1: Relative error of the change of sea surface temperature during a coupling period. The error is computed as the ratio between the correction due to the iterative procedure and the solution change during a coupling period with no Schwarz iteration. The ordinates show the number of cases in percentage of the total number of cases in $time \times space$.

cases. But it is larger than 0.1 for the other half. The relative error is even larger than 0.5 in 25 % of the cases. The *atmosphere-first* shows strongly improved results, with a negligible error for 97 % of the points. The results of the *ocean-first* experiment are close to the *atmosphere-first* experiments, with a slight deterioration. For the $\Delta t = 4h$ experiments, the errors are larger than in the $\Delta t = 1h$ case, but with the same hierarchy between the algorithms.

The error shows a strong diurnal cycle (not shown) with the lowest errors during the night. Errors are larger at noon than at midnight. The error is maximum after sunset and before sunrise, when the change of the insolation forcing evolves at the fastest pace.

We propose two hypotheses to explain the *atmosphere-first* algorithm performance compared to *ocean-first*. First, the atmosphere has shorter characteristic time scales than the ocean, with a more marked diurnal cycle. The atmospheric lower boundary conditions evolves slowly, and the atmospheric solution after the first half-iteration is then already quite close to its converged value, and provides a relevant and synchronized forcing to compute the oceanic solution in the second half-iteration. Second, the better performance of the *atmosphere-first* case can also be linked to the phasing of the solar radiation, which is the only external forcing and constrains the diurnal cycle. In the *ocean-first* case, the ocean is forced by fluxes, including solar radiation, calculated by the atmosphere at the previous coupling period. For *atmosphere-first*, the solar forcing is correctly phased.

Most current GCMs use the *parallel* algorithm which appears to have very large errors (see Marti et al.

(2021) for a short review). Our analysis shows that implementing *sequential* algorithms are simple ways to strongly reduce the error, with the *atmosphere-first* algorithm showing the best performance. The *sequential* algorithms, however, have a major drawback. The models do not run concurrently as, while one model is running, the other model waits for its boundary conditions. This eliminates a level of parallelism, and increases the time to solution of the coupled model.

4 Acknowledgements

This study is part of the ANR project COCOA (<https://anr.fr/Projet-ANR-16-CE01-0007>). This work was granted access to the HPC resources of TGCC under an allocation made by GENCI (Grand équipement National de Calcul Intensif, grant 2019-A0040100239). It benefits from the development of the common modelling IPSL infrastructure coordinated by the IPSL climate modeling center (<https://cmc.ipsl.fr>).

References

- Lemarié, F. (2008). *Algorithmes de Schwarz et couplage océan-atmosphère*. PhD thesis, Université Joseph Fourier, Grenoble.
- Marti, O., Braconnot, P., Dufresne, J.-L., Bellier, J., Benschila, R., Bony, S., Brockmann, P., Cadule, P., Caubel, A., Codron, F., de Noblet, N., Denvil, S., Fairhead, L., Fichefet, T., Foujols, M.-A., Friedlingstein, P., Goosse, H., Grandpeix, J.-Y., Guilyardi, E., Hourdin, F., Idelkadi, A., Kageyama, M., Krinner, G., Lévy, C., Madec, G., Mignot, J., Musat, I., Swingedouw, D., and Talandier, C. (2010). Key features of the IPSL ocean atmosphere model and its sensitivity to atmospheric resolution. *Clim. Dyn.*, 34(1):1–26.
- Marti, O., Nguyen, S., Braconnot, P., Valcke, S., Lemarié, F., and Blayo, E. (2021). A Schwarz iterative method to evaluate ocean-atmosphere coupling schemes. Implementation and diagnostics in IPSL-CM6-SW-VLR. *Geosci. Model Dev. Discuss.*, 2021:1–19.
- Sepulchre, P., Caubel, A., Ladant, J.-B., Bopp, L., Boucher, O., Braconnot, P., Brockmann, P., Cozic, A., Donnadiou, Y., Dufresne, J.-L., Estella-Perez, V., Ethé, C., Fluteau, F., Foujols, M.-A., Gastineau, G., Ghattas, J., Hauglustaine, D., Hourdin, F., Kageyama, M., Khodri, M., Marti, O., Meurdesoif, Y., Mignot, J., Sarr, A.-C., Servonnat, J., Swingedouw, D., Szopa, S., and Tardif, D. (2020). IPSL-CM5A2 – an Earth system model designed for multi-millennial climate simulations. *Geophys. Res. Lett.*, 13(7):3011–3053.