



FP09.05 Driver Oncogenic Alterations and Indoor Radon in NSCLC Patients From the IFCT Biomarker Cohort: Bioradon France Study

L. Mezquita, F. Barlesi, G. Ielsch, J. Merlio, D. Debieuvre, J. Mosser, C. Ricordel, L'Houcine Ouafik, I. Rouquette, I. Monnet, et al.

► To cite this version:

L. Mezquita, F. Barlesi, G. Ielsch, J. Merlio, D. Debieuvre, et al.. FP09.05 Driver Oncogenic Alterations and Indoor Radon in NSCLC Patients From the IFCT Biomarker Cohort: Bioradon France Study. *Journal of Thoracic Oncology*, 2021, 16 (3), pp.S214. 10.1016/j.jtho.2021.01.122 . hal-03603784

HAL Id: hal-03603784

<https://amu.hal.science/hal-03603784>

Submitted on 10 Mar 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

and 11+ drinks per week (Table). Those with FH of LC were more likely to be obese regardless of demographic factors, and less likely to exercise 2+ times per week (Table). Those with FH of LC were more likely to have received 10+ imaging scans and to have been exposed to hydrocarbons compared to those without FH (Table). There were no reported differences in any other occupational or home exposures, including asbestos and radon in those with v. without FH of LC (Table).

Conclusion: FH of LC is associated with higher-risk behaviors that increase the likelihood of developing both lung and other cancers. As such, FH can be used in cancer prevention and control programs to identify higher-risk patients and educate patients that FH of lung cancer is associated with risk-increasing behaviors that are nonetheless modifiable. **Keywords:** Family History, Risky Behaviors, Prevention

FP09.05

Driver Oncogenic Alterations and Indoor Radon in NSCLC Patients From the IFCT Biomarker Cohort: Bioradon France Study



L. Mezquita,¹ F. Barlesi,² G. Ielsch,³ J. Merlio,⁴ D. Debieuvre,⁵ J. Mosser,⁶ C. Ricordel,⁷ L. Ouafik,⁸ I. Rouquette,⁹ I. Monnet,¹⁰ F. Escande,¹¹ A. Langlais,¹² F. Morin,¹² C. Vignaud,³ E. Auclin,¹³ J.C. Benitez,² D. Planchard,² K. Leuraud,³ D. Laurier,³ B. Besse,² V. Westeel¹⁴
¹Hospital Clinic, Medical Oncology Department, Barcelona; ²Medical Oncology Department, Gustave Roussy Cancer Campus, Villejuif, France, Spain/ES; ³Medical Oncology Department, Gustave Roussy Cancer Campus, Villejuif, France, Villejuif/FR; ⁴Institute for Radiological Protection and Nuclear Safety – Irsn, France, Paris/FR; ⁵University Hospital and University Bordeaux, Bordeaux/france, Bordeaux/FR; ⁶Groupe Hospitalier Régional Mulhouse Sud Alsace (Ghrmsa), Hôpital Emile Muller, Mulhouse Cedex/france, Mulhouse Cedex/FR; ⁷CHU Rennes, Rennes/france, Rennes/FR; ⁸Centre Hospitalier Régional Et Universitaire de Rennes, Rennes/france, Rennes/FR; ⁹Aix Marseille Univ, Aphm, Cnrs, Inp, Inst Neurophysiopathol, Marseille, France, Faculté de Médecine Secteur Nord, Service de Transfert D'Oncologie Biologique, Marseille, France, Marseille/france, Marseille/FR; ¹⁰Oncopole - CHU Toulouse, Toulouse/france, Toulouse/FR; ¹¹Chi Creteil, Creteil/france, Creteil/FR; ¹²Chru Lille, Lille/france, Lille/FR; ¹³Ifct, Paris/france, Paris/FR; ¹⁴Hopital Européen Georges Pompidou, Paris/FR; ¹⁵University Hospital of Besançon, Besançon/france, Besançon/FR

Introduction: Radon is a radioactive gas, considered as the leading cause of lung cancer in non-smokers. In a previous work, we studied the correlation between the estimation of radon concentration from the French Indoor Radon Map (*Institut de Radioprotection et de Sûreté Nucléaire*, IRSN) and the regional prevalence of driver alterations in a cohort of 116,424 NSCLC patients in France (*Mezquita et al, WCLC 2018*). The prevalence of driver oncogene alterations was significantly

higher in high-radon areas but clinical data were not available. We aim to confirm this hypothesis in an annotated database of NSCLC patients with matched molecular data available for adjustment (*Barlesi et al, Lancet 2016*). **Methods:** Retrospective assessment of patients with NSCLC tested for *EGFR/BRAF/HER2/KRAS* mutations (m) and *ALK* fusion from the 28 Platform led by the National Cancer Institute between Apr.2012 and Apr.2013, and included in the Biomarkers France dataset. We studied the association between the prevalence of driver oncogenic alterations (*EGFR/ALK/BRAF/HER2/KRAS*) and the radon mean concentration in the area where the patient was born according to the IRSN Map. Adjustment on age, gender and smoking was performed. **Results:** Out of 17664 patients, we analyzed 3994 with birthplace available: 63% males, 82% smokers, with a median age of 64 years [18-94]. Lung cancer tumors were mostly adenocarcinoma (76%), followed by other histologies (18%) and squamous (6%). By molecular alterations: 468 tumors harbor *EGFRm* (12%), 129 *ALK* (3%), 89 *BRAFm* (2%), 32 *HER2m* (1%), 985 *KRASm* (25%); 2273 *wildtype* or harbor other non-driver alterations (control; 57%). Adenocarcinoma histology (83.7% vs. 80.2%, $p=0.0034$), and non-smoker habit (19.5% vs. 16.5%, $p=0.0251$) were more common in radon high-risk group (comparatively at the low risk group). The mean radon concentration by birthplace was $74.36 \text{ Bq/m}^3 \pm 53.28\text{SD}$ [range 16.6-622.3], and by molecular groups: *EGFRm* $72.49 \text{ Bq/m}^3 \pm 47.98\text{SD}$ [16.6-461.4], *ALK* $80.24 \text{ Bq/m}^3 \pm 55.22\text{SD}$ [19.3-384.7], *BRAFm* $73.22 \text{ Bq/m}^3 \pm 47.86\text{SD}$ [19.3-319.3], *HER2m* $72.74 \text{ Bq/m}^3 \pm 39.51\text{SD}$ [27.8-231.3], *KRASm* $71.79 \text{ Bq/m}^3 \pm 53.32\text{SD}$ [16.6-576.8] and control group $75.67 \text{ Bq/m}^3 \pm 54.55\text{SD}$ [16.6-622.3] ($p=0.20$). The prevalence of driver alterations was higher in high-radon areas (table 1; $p=0.0472$); but no significant difference was observed after adjustment on age, gender and smoking. **Conclusion:** We observed a higher prevalence of driver oncogenic alterations in NSCLC patients born in high radon areas; but no significant difference was observed after adjustment on age, gender and smoker. This study warrants further research on radon gas and driver oncogenes. **Keywords:** radon, lung cancer, driver oncogenic alterations

FP10 SMALL CELL LUNG CANCER/NET

FP10.01

Survival in Advanced SCLC: Projected Impact of Immuno-Oncology-Associated Durable Response on Population Health Gains in US



J. Gordon,¹ P. Mcewan,¹ R. Young,¹ P. Kirkness,¹ J. Penrod,² Y. Yuan^{2,4}
¹Health Economics and Outcomes Research Ltd., Cardiff/GB, ²Bristol-Myers Squibb, Princeton/NJ/US

			Low radon <50 Bq/m3	High radon ≥50 Bq/m3	P value	P value adjusted*
<i>EGFR</i>	<i>Mutation</i>	N (%)	155 (11.4%)	313 (13.2%)	0.1218	0.3024
	<i>Control (1)</i>	N (%)	1199 (88.6)	2059 (86.8)		
<i>ALK</i>	<i>Fusion</i>	N (%)	42 (3.4%)	87 (4.1%)	0.3272	0.4708
	<i>Control (1)</i>	N (%)	1199 (96.6)	2059 (95.9)		
<i>BRAF</i>	<i>Mutation</i>	N (%)	32 (2.6%)	57 (2.7%)	0.8708	0.9865
	<i>Control (1)</i>	N (%)	1199 (97.4)	2059 (97.3)		
<i>HER2</i>	<i>Mutation</i>	N (%)	7 (0.6%)	25 (1.2%)	0.0880	0.1781
	<i>Control (1)</i>	N (%)	1199 (99.4)	2059 (98.8)		
<i>KRAS</i>	<i>Mutation</i>	N (%)	375 (31.3%)	610 (29.6%)	0.3227	0.3478
	<i>Control (2)</i>	N (%)	824 (68.7)	1449 (70.4)		
<i>DRIVER</i>	<i>Positive</i>	N (%)	236 (16.4%)	482 (19%)	0.0472	0.2128
	<i>Control (1)</i>	N (%)	1199 (83.6)	2059 (81)		

(1) population no *EGFRm*, no *BRAFm*, no *HER2m* and no *ALKr*. (2) population no *EGFRm*, no *BRAFm*, no *HER2m*, no *ALKr* and no *KRASm*. *adjustment on age, gender and smoking