



HAL
open science

Functional connectivity predicts MI-based BCI learning

Marie-Constance Corsi, Mario Chavez, Denis Schwartz, Nathalie George,
Laurent Hugueville, Ari E. Kahn, Sophie Dupont, Danielle S. Bassett,
Fabrizio de Vico Fallani

► **To cite this version:**

Marie-Constance Corsi, Mario Chavez, Denis Schwartz, Nathalie George, Laurent Hugueville, et al..
Functional connectivity predicts MI-based BCI learning. BIOMAG 2022 - 22nd International Conference on Biomagnetism, Aug 2022, Birmingham, United Kingdom. hal-03773303

HAL Id: hal-03773303

<https://inria.hal.science/hal-03773303>

Submitted on 9 Sep 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Functional connectivity predicts MI-based BCI learning

Marie-Constance Corsi¹

Mario Chavez²

Denis Schwartz¹

Nathalie George¹

Laurent Hugueville¹

Ari E. Kahn²

Sophie Dupont¹

Danielle S. Bassett²

Fabrizio De Vico Fallani¹

¹ Sorbonne Université, Institut du Cerveau – Paris Brain Institute, ICM, CNRS, Inria, Inserm, AP-HP, Hôpital de la Pitié-Salpêtrière, F-75013, Paris, France

² University of Pennsylvania, Philadelphia, USA



marie.constance.corsi@gmail.com




@MConstanceCorsi

Non-invasive Brain-Computer Interfaces (BCIs) can exploit the ability of subjects to voluntarily modulate their brain activity through mental imagery. Despite its clinical applications, controlling a BCI appears to be a learned skill that requires several weeks to reach relatively high-performance in control, without being sufficient for 15 to 30 % of the users [1]. This gap has motivated a deeper understanding of mechanisms associated with motor imagery (MI) tasks. Here, we investigated dynamical changes in terms of cortical activations and network recruitment. We hypothesized that the associated characteristics would be able to predict the success of learning.

Longitudinal study


The EEG-based BCI task consisted of a standard 1D, two-target box task in which the subjects modulated their α and/or β activity. To hit the target-up, the subjects performed a sustained motor imagery of their right-hand grasping and to hit the target-down, they remained at rest. 20 healthy subjects (aged 27.45 ± 4.01 years, 12 men), all right-handed, participated in the study.

We found that the ability to control the BCI significantly increased across sessions (days) but not within sessions (hours). The session effect was also present when we averaged the BCI accuracy scores across the runs of each session (one-way ANOVA, $F_{3,57}=13.9$, $p=6.56 \cdot 10^{-7}$). Despite the expected high inter-subject variability ($> 8.95\%$), the group reached on average a reasonable level of control - above the chance level of 57 % by the end of the training.



Functional connectivity (FC) patterns


To evaluate the cortical changes at the network level, we considered the imaginary coherence [2] between the source reconstructed signals of each pair of regions of interest (ROI) corresponding to Destrieux atlas.



By statistically comparing the FC values between MI and Rest conditions at the group level, we found a progressive decrease of task-related connectivity in both α and β frequency ranges across sessions. In α_2 , the strongest decreases involved parieto-occipital interactions. For each subject, we also computed the relative node strength Δ_N . Middle occipital gyrus, cuneus, superior occipital gyrus, associated with visuo-spatial attentional tasks decreased significantly with the training.

Cortical changes


To evaluate the spatiotemporal cortical changes associated with the BCI training, we computed statistical differences among activations recorded in the MI and the rest conditions via a paired t-test, corrected for multiple comparisons using the cluster approach at the group level.



In session 3, these diffused decrements were particularly significant in the α_2 and β_1 frequency bands, and mainly spanned the primary sensorimotor cortex and in secondary higher-order premotor and somatosensory areas. At the end of training, these decrements were localized to the contralateral paracentral lobule, precentral gyrus, and superior parietal lobule.

Predictive markers of BCI scores

Higher BCI scores were associated with the recruitment of a higher number of task-related cortical sources and with larger decreases of relative power in α_2 . Better performance was rather associated with the decrease of Δ_N in associative areas and in premotor and primary visual areas. We observed significant correlations in regions known to be involved in specific cognitive aspects of human learning, such as decision making (fronto-marginal



gyrus), in visual attention task (occipital pole), and in both mental rotation and working memory (orbital part of the inferior frontal gyrus). With Δ_N , we found significant predictions for different ROIs involved during visual information processing (middle occipital gyrus), but also during motor imagery and working memory (precuneus).

Interested in this study?
Scan the QR code to get access to the associated paper!



References

- [1] B. Z. Allison and C. Neuper, "Could Anyone Use a BCI?," in *Brain-Computer Interfaces*, Eds. Springer London, 2010,
- [2] Sekihara K, Owen J, Trisno S, Nagarajan SS. Removal of spurious coherence in MEG source-space coherence analysis. *IEEE transactions on bio-medical engineering*. 2011.

Acknowledgements

This work was partially supported by the program "Investissements d'avenir" ANR-10-IAIHU-06; "ANR-NIH CRCNS" ANR-15-NEUC-0006-02 and by NICHD 1R01HD086888-01.