

A Stabilized Finite Element Method for the Oseen Equation with Dominating Reaction

Gabriel Raúl Barrenechea, Miguel Angel Fernández, Carolina Ivon Vidal

► **To cite this version:**

Gabriel Raúl Barrenechea, Miguel Angel Fernández, Carolina Ivon Vidal. A Stabilized Finite Element Method for the Oseen Equation with Dominating Reaction. [Research Report] RR-5213, INRIA. 2004, pp.21. inria-00070780

HAL Id: inria-00070780

<https://hal.inria.fr/inria-00070780>

Submitted on 19 May 2006

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



INSTITUT NATIONAL DE RECHERCHE EN INFORMATIQUE ET EN AUTOMATIQUE

*A Stabilized Finite Element Method for the Oseen
Equation with Dominating Reaction*

Gabriel Raúl Barrenechea — Miguel Ángel Fernández — Carolina Ivon Vidal

N° 5213

Mai 2004

Thème BIO



*R*apport
de recherche



A Stabilized Finite Element Method for the Oseen Equation with Dominating Reaction

Gabriel Raúl Barrenechea* , Miguel Ángel Fernández† , Carolina Ivon Vidal‡

Thème BIO — Systèmes biologiques
Projet REO

Rapport de recherche n° 5213 — Mai 2004 — 21 pages

Abstract: A new stabilized finite element method is introduced for the linearized version of the Navier-Stokes equation (or Oseen equation), containing a dominating zeroth order term. The method consists in subtracting a mesh dependent term from the formulation without compromising consistency. The design of this mesh dependent term, as well as the stabilization parameter involved, are suggested by bubble condensation. Stability is proved for any combination of velocity and pressure spaces, under the hypotheses of continuity for the pressure space. Optimal order error estimates are derived for the velocity and the pressure, using the standard norms for these unknowns. Numerical experiments confirming these theoretical results are presented.

Key-words: Oseen equation, reaction term, stabilized method

This research was partially supported by CONICYT Chile, through project FONDECYT No. 1030674, and FONDAF Program on Applied Mathematics, and by Universidad de Concepción through the Advanced Research Groups Program.

* GI²MA, Departamento de Ingeniería Matemática, Universidad de Concepción, Casilla 160-C Concepción, Chile. email: gbarrene@ing-mat.udec.cl

† INRIA projet REO. email: miguel.fernandez@inria.fr

‡ Departamento de Ingeniería Matemática, Universidad de Concepción, Casilla 160-C Concepción, Chile, email: cvidal@ing-mat.udec.cl

Une méthode d'éléments finis stabilisée pour l'équation d'Oseen avec réaction dominante

Résumé : On introduit une nouvelle méthode d'éléments finis stabilisée pour l'équation d'Oseen (Navier-Stokes linéarisée) avec un terme d'ordre zéro dominant. La méthode est obtenue après soustraction d'un terme dépendant du maillage à la formulation de Galerkin classique, sans compromettre la consistance de la méthode. L'expression de ce terme et celle du paramètre de stabilisation associé sont déterminées par une technique de type *condensation de la bulle*. La stabilité de la méthode est assurée pour tout couple d'espaces discrets, sous une hypothèse de continuité des pressions discrètes. On fournit des estimations d'erreur optimales pour la vitesse et la pression dans ses normes naturelles. Des expériences numériques confirment les résultats théoriques.

Mots-clés : Équation d'Oseen, terme de réaction, méthode stabilisée

1 Introduction

The Navier-Stokes equation constitutes a major challenge in applied mathematics. Specifically, its numerical solution presents two major difficulties, namely, the need for a compatibility condition (the inf-sup condition, see [17, 3]) relating the discrete spaces used to approximate the velocity field \mathbf{u} and the pressure p , and the treatment of the spurious modes generated by the convective term. For both these aspects, several solutions have been proposed in the last two decades. The convective terms have been treated by appropriate upwinding strategies (cf. [17, 21, 11], and the references therein), or stabilized finite element methods (cf. [6, 14], among others). On the other hand, the inf-sup condition may be treated directly (cf. [3, 17], and the references therein), or circumvented via stabilized finite element methods (cf. [20, 19, 5, 4, 15] in the context of a Stokes flow).

On the other hand, if we are dealing with the time discretization of the Navier-Stokes equation, and we choose the “classical” approach (i.e., discretizing in time by time-advancing finite differences) we have different choices for the scheme (for a resume of these techniques, see [21]). A common fact of all these techniques is the presence of a zeroth order term of type $\frac{1}{\Delta t}\mathbf{u}$, where Δt is the time step (usually very small), and \mathbf{u} is the unknown velocity field. In the late nineties, several works concerning stabilization procedures for problems with zeroth order terms (or reaction terms), were proposed (see, e.g., [18, 24], and the recent paper [7], where edge stabilization has been proposed for a scalar convection-diffusion-reaction problem). In particular, in [13, 16, 2], the connection between stabilized finite element methods and Galerkin methods enriched with bubble functions was used to derive a new family of stabilized finite element method, namely, the Unusual Stabilized Finite Element Method (USFEM), which are particularly suited for treating problems with dominating reaction.

The purpose of this work is to derive, analyze and test a new stabilized finite element method, analogous to the one presented in [2], for the Oseen problem with a dominating reaction term. The method is introduced in Section 2 where the stability of the method is proved and an error analysis is performed. The error estimates obtained in Section 2 are derived, using a suitable mesh dependent norm, for the standard norms for the unknowns, namely the $H^1(\Omega)$ norm for the velocity field, and the $L^2(\Omega)$ norm for the pressure. In Section 3 we report some numerical experiments that confirm our approximation results. Finally, some concluding remarks are made in Section 4.

2 The finite element method

First, we present the problem of interest. Let Ω be a bounded open subset of \mathbb{R}^2 , $\mathbf{f} \in [L^2(\Omega)]^2$, σ a positive real number (typically, $\sigma \approx \frac{1}{\Delta t}$ where Δt is the time step in a time discretization procedure), and $\mathbf{a} : \Omega \rightarrow \mathbb{R}^2$ a vectorial function such that $\nabla \cdot \mathbf{a} = 0$ in Ω (this function \mathbf{a} may be interpreted as the velocity field in the previous time step). Then, our

generalized Oseen equation reads: Find $(\mathbf{u}, p) \in [H_0^1(\Omega)]^2 \times L_0^2(\Omega)$ such that

$$\begin{aligned} \sigma \mathbf{u} - \nu \Delta \mathbf{u} + \mathbf{a} \cdot \nabla \mathbf{u} + \nabla p &= \mathbf{f} & \text{in } \Omega, \\ \nabla \cdot \mathbf{u} &= 0 & \text{in } \Omega, \\ \mathbf{u} &= 0 & \text{on } \partial\Omega, \end{aligned} \quad (1)$$

where $L_0^2(\Omega) \stackrel{\text{def}}{=} \{q \in L^2(\Omega) : (q, 1)_\Omega = 0\}$, and $(\cdot, \cdot)_D$ denotes the L^2 inner product in $L^2(D)$ (or in $L^2(D)^2, L^2(D)^{2 \times 2}$, when necessary). Also, by $\|\cdot\|_{l,D}$ and $|\cdot|_{l,D}$ we will denote the $H^l(D)$ norm and seminorm, respectively, with the usual convention $H^0(D) = L^2(D)$.

From now on, let us suppose that Ω is a polygonal domain in \mathbb{R}^2 , and let \mathcal{T}_h be a triangulation of Ω constituted by triangles (or quadrilaterals) which are shape regular. Let h_K be the usual element diameter, and denote $h \stackrel{\text{def}}{=} \max\{h_K : K \in \mathcal{T}_h\}$. We suppose from now on that $h \leq 1$. Now, for $k \geq 1$, let V_k be the space of piecewise polynomial functions given by

$$V_k \stackrel{\text{def}}{=} \{v \in \mathcal{C}^0(\bar{\Omega}) / v|_K \in R^k(K), \forall K \in \mathcal{T}_h\}.$$

Here, $R^k(K) = P^k(K)$ for triangular elements and $R^k(K) = \{p \circ F_K^{-1} / p \in Q^k(\hat{K})\}$ for quadrilateral elements, where F_K stands for the usual transformation mapping the reference element \hat{K} onto K .

Our stabilized finite element method reads: Find $(\mathbf{u}_h, p_h) \in \mathbf{V}_h \times Q_h$ such that:

$$\mathbf{B}((\mathbf{u}_h, p_h), (\mathbf{v}, q)) = \mathbf{F}(\mathbf{v}, q) \quad \forall (\mathbf{v}, q) \in \mathbf{V}_h \times Q_h, \quad (2)$$

where $\mathbf{V}_h \stackrel{\text{def}}{=} [V_k \cap H_0^1(\Omega)]^2$ and $Q_h \stackrel{\text{def}}{=} V_l \cap L_0^2(\Omega)$, $k, l \geq 1$, \mathbf{B} and \mathbf{F} are given by

$$\begin{aligned} \mathbf{B}((\mathbf{u}_h, p_h), (\mathbf{v}, q)) &\stackrel{\text{def}}{=} \sigma(\mathbf{u}_h, \mathbf{v})_\Omega + \nu(\nabla \mathbf{u}_h, \nabla \mathbf{v})_\Omega + (\mathbf{a} \cdot \nabla \mathbf{u}_h, \mathbf{v})_\Omega \\ &\quad - (p_h, \nabla \cdot \mathbf{v})_\Omega + (q, \nabla \cdot \mathbf{u}_h)_\Omega + \sum_{K \in \mathcal{T}_h} (\nabla \cdot \mathbf{u}_h, \delta_K \nabla \cdot \mathbf{v})_K \\ &\quad - \sum_{K \in \mathcal{T}_h} (\sigma \mathbf{u}_h - \nu \Delta \mathbf{u}_h + \mathbf{a} \cdot \nabla \mathbf{u}_h + \nabla p_h, \tau_K (\sigma \mathbf{v} - \nu \Delta \mathbf{v} - \mathbf{a} \cdot \nabla \mathbf{v} - \nabla q))_K, \end{aligned} \quad (3)$$

$$\mathbf{F}(\mathbf{v}, q) \stackrel{\text{def}}{=} (\mathbf{f}, \mathbf{v})_\Omega - \sum_{K \in \mathcal{T}_h} (\mathbf{f}, \tau_K (\sigma \mathbf{v} - \nu \Delta \mathbf{v} - \mathbf{a} \cdot \nabla \mathbf{v} - \nabla q))_K. \quad (4)$$

Here, the stabilization parameters τ_K and δ_K are given by

$$\tau_K \stackrel{\text{def}}{=} \frac{h_K^2}{\sigma h_K^2 \xi(Pe_K^1) + \frac{4\nu}{m_k} \xi(Pe_K^2)}, \quad (5)$$

$$\delta_K \stackrel{\text{def}}{=} \lambda |\mathbf{a}(x)|_p h_K \min\{1, Pe_K^2\}, \quad (6)$$

where $\lambda \geq 0$, $p \in [1, +\infty]$, and

$$Pe_K^1 = \frac{4\nu}{m_k \sigma h_K^2}, \quad (7)$$

$$Pe_K^2 = \frac{m_k |\mathbf{a}|_p h_K}{4\nu}, \quad (8)$$

$$|\mathbf{a}(x)|_p \stackrel{\text{def}}{=} (|a_1|^p + |a_2|^p)^{1/p}, \quad (9)$$

$$m_k = \min\left\{\frac{1}{3}, C_k\right\}, \quad (10)$$

$$C_k h_K^2 \|\Delta v\|_{0,K}^2 \leq \|\nabla v\|_{0,K}^2 \quad \forall v \in V_k, \quad (11)$$

$$\xi(\lambda) = \max\{\lambda, 1\}. \quad (12)$$

Remark 1 *The design of the stabilization parameter τ_K has been suggested by bubble condensation, following very closely the arguments given in [2, 1]. The least-squares parameter δ_K is the one from [14]. In the case of a generalized Stokes problem ($\mathbf{a} = \mathbf{0}$) we recover the method from [2]. Now, in the case of a pure Oseen equation $\sigma = 0$, we recover the “plus” formulation from [14] with a stabilization parameter which satisfies $\tau_{FF} \leq \tau_K \leq 2\tau_{FF}$, where τ_{FF} denotes the stabilization parameter proposed in [14], given by*

$$\tau_{FF} \stackrel{\text{def}}{=} \frac{h_K}{2|\mathbf{a}(x)|_p} \min\{1, Pe_K^2\}.$$

Remark 2 *In [10], the orthogonal subscales approach was applied to a related problem containing a Coriolis terms and a zeroth order term. The resulting formulation involves stabilization parameters with free constants to be set. The performance of the method depends on how these constants are chosen.*

2.1 The stability of the method

Throughout all this section (and the following one), C will denote a positive constant independent of h (but who may depend on the physical coefficients), whose value may vary whenever it is written in two different places.

The following lemma provides the positive-definiteness of the stiffness matrix associated with our method.

Lemma 2.1 *There exists a constant C_Ω , depending only on Ω , such that*

$$\mathbf{B}((\mathbf{v}, q), (\mathbf{v}, q)) \geq C_\Omega \nu \|\mathbf{v}\|_{1,\Omega}^2 + \sum_{K \in \mathcal{T}_h} \left\{ \|\tau_K^{1/2} (\mathbf{a} \cdot \nabla \mathbf{v} + \nabla q)\|_{0,K}^2 + \|\delta_K^{1/2} \nabla \cdot \mathbf{v}\|_{0,K}^2 \right\},$$

for all $(\mathbf{v}, q) \in \mathbf{V}_h \times Q_h$.

Proof. - From the definition of τ_K and the fact that $\xi(Pe_K^1), \xi(Pe_K^2) \geq 1$, we observe first that

$$\tau_K \leq \overline{\tau}_K \stackrel{\text{def}}{=} \frac{m_k h_K^2}{m_k \sigma h_K^2 + 4\nu},$$

for all $K \in \mathcal{T}_h$. Now, let $(\mathbf{v}, q) \in \mathbf{V}_h \times Q_h$, then

$$\begin{aligned} \mathbf{B}((\mathbf{v}, q), (\mathbf{v}, q)) &\geq \sigma \|\mathbf{v}\|_{0,\Omega}^2 + \nu \|\nabla \mathbf{v}\|_{0,\Omega}^2 + (\mathbf{a} \cdot \nabla \mathbf{v}, \mathbf{v})_\Omega + \sum_{K \in \mathcal{T}_h} \|\delta_K^{1/2} \nabla \cdot \mathbf{v}\|_{0,K}^2 \\ &\quad - \sum_{K \in \mathcal{T}_h} \left\{ \overline{\tau}_K [\sigma^2 \|\mathbf{v}\|_{0,K}^2 - 2\sigma\nu(\mathbf{v}, \Delta \mathbf{v})_K + \nu^2 \|\Delta \mathbf{v}\|_{0,K}^2] \right. \\ &\quad \left. + \|\tau_K^{1/2} (\mathbf{a} \cdot \nabla \mathbf{v} + \nabla q)\|_{0,K}^2 \right\}. \end{aligned}$$

Now, integrating by parts we obtain $(\mathbf{a} \cdot \nabla \mathbf{v}, \mathbf{v})_\Omega = 0$, and hence

$$\begin{aligned} \mathbf{B}((\mathbf{v}, q), (\mathbf{v}, q)) &\geq \sum_{K \in \mathcal{T}_h} \left\{ (\sigma - \sigma^2 \overline{\tau}_K) \|\mathbf{v}\|_{0,K}^2 + \nu \|\nabla \mathbf{v}\|_{0,K}^2 + 2\sigma\nu \overline{\tau}_K (\mathbf{v}, \Delta \mathbf{v})_K \right. \\ &\quad \left. - \nu^2 \overline{\tau}_K \|\Delta \mathbf{v}\|_{0,K}^2 + \|\tau_K^{1/2} (\mathbf{a} \cdot \nabla \mathbf{v} + \nabla q)\|_{0,K}^2 + \|\delta_K^{1/2} \nabla \cdot \mathbf{v}\|_{0,K}^2 \right\} \\ &\geq \sum_{K \in \mathcal{T}_h} \left\{ (\sigma - \sigma^2 \overline{\tau}_K) \|\mathbf{v}\|_{0,K}^2 + \nu \|\nabla \mathbf{v}\|_{0,K}^2 - 2\sigma\nu \overline{\tau}_K \|\mathbf{v}\|_{0,K} \|\Delta \mathbf{v}\|_{0,K} \right. \\ &\quad \left. - \frac{\nu^2 \overline{\tau}_K h_K^{-2}}{C_k} \|\nabla \mathbf{v}\|_{0,K}^2 + \|\tau_K^{1/2} (\mathbf{a} \cdot \nabla \mathbf{v} + \nabla q)\|_{0,K}^2 + \|\delta_K^{1/2} \nabla \cdot \mathbf{v}\|_{0,K}^2 \right\} \\ &\geq \sum_{K \in \mathcal{T}_h} \left\{ (\sigma - \sigma^2 \overline{\tau}_K) \|\mathbf{v}\|_{0,K}^2 + \left(\nu - \frac{\nu^2 \overline{\tau}_K h_K^{-2}}{C_k} \right) \|\nabla \mathbf{v}\|_{0,K}^2 - \frac{\sigma\nu \overline{\tau}_K}{\gamma} \|\mathbf{v}\|_{0,K}^2 \right. \\ &\quad \left. - \gamma\sigma\nu \overline{\tau}_K \|\Delta \mathbf{v}\|_{0,K} + \|\tau_K^{1/2} (\mathbf{a} \cdot \nabla \mathbf{v} + \nabla q)\|_{0,K}^2 + \|\delta_K^{1/2} \nabla \cdot \mathbf{v}\|_{0,K}^2 \right\} \\ &\geq \sum_{K \in \mathcal{T}_h} \left\{ \left(\sigma - \sigma^2 \overline{\tau}_K - \frac{\sigma\nu \overline{\tau}_K}{\gamma} \right) \|\mathbf{v}\|_{0,K}^2 \right. \\ &\quad \left. + \left(\nu - \frac{\nu^2 \overline{\tau}_K h_K^{-2}}{C_k} - \frac{\gamma\sigma\nu \overline{\tau}_K h_K^{-2}}{C_k} \right) \|\nabla \mathbf{v}\|_{0,K}^2 \right. \\ &\quad \left. + \|\tau_K^{1/2} (\mathbf{a} \cdot \nabla \mathbf{v} + \nabla q)\|_{0,K}^2 + \|\delta_K^{1/2} \nabla \cdot \mathbf{v}\|_{0,K}^2 \right\} \end{aligned}$$

$$\begin{aligned}
&\geq \sum_{K \in \mathcal{T}_h} \left\{ \left(\sigma - \frac{\sigma^2 m_k h_K^2}{m_k \sigma h_K^2 + 4\nu} - \frac{\sigma \nu m_k h_K^2}{\gamma(m_k \sigma h_K^2 + 4\nu)} \right) \|\mathbf{v}\|_{0,K}^2 \right. \\
&\quad + \left(\nu - \frac{m_k \nu^2}{C_k(m_k \sigma h_K^2 + 4\nu)} - \frac{m_k \gamma \sigma \nu}{C_k(m_k \sigma h_K^2 + 4\nu)} \right) \|\nabla \mathbf{v}\|_{0,K}^2 \\
&\quad \left. + \|\tau_K^{1/2}(\mathbf{a} \cdot \nabla \mathbf{v} + \nabla q)\|_{0,K}^2 + \|\delta_K^{1/2} \nabla \cdot \mathbf{v}\|_{0,K}^2 \right\} \\
&\geq \sum_{K \in \mathcal{T}_h} \left\{ \frac{\sigma \nu (4\gamma - m_k h_K^2)}{\gamma(m_k \sigma h_K^2 + 4\nu)} \|\mathbf{v}\|_{0,K}^2 + \frac{3\nu^2 + \sigma \nu (m_k h_K^2 - \gamma)}{m_k \sigma h_K^2 + 4\nu} \|\nabla \mathbf{v}\|_{0,K}^2 \right. \\
&\quad \left. + \|\tau_K^{1/2}(\mathbf{a} \cdot \nabla \mathbf{v} + \nabla q)\|_{0,K}^2 + \|\delta_K^{1/2} \nabla \cdot \mathbf{v}\|_{0,K}^2 \right\},
\end{aligned}$$

and it suffices to take $\gamma = \frac{m_k h_K^2}{4}$ and apply Poincaré's inequality to complete the proof. \square

Now, in order to prove our main result in stability, namely the inf-sup condition for \mathbf{B} , we define the following mesh-dependent norm:

$$\|(\mathbf{v}, q)\|_h \stackrel{\text{def}}{=} \left\{ \|\mathbf{v}\|_{1,\Omega}^2 + \sum_{K \in \mathcal{T}_h} \left[\|\tau_K^{1/2}(\mathbf{a} \cdot \nabla \mathbf{v} + \nabla q)\|_{0,K}^2 + \|\delta_K^{1/2} \nabla \cdot \mathbf{v}\|_{0,K}^2 \right] + \|q\|_{0,\Omega}^2 \right\}^{1/2}, \quad (13)$$

for all $(\mathbf{v}, q) \in \mathbf{V}_h \times Q_h$.

We now state the main stability result.

Theorem 2.2 *There exists a constant $\beta = \beta(\sigma, \mathbf{a}, \nu)$, independent of h , such that*

$$\sup_{\theta \neq (\mathbf{w}, t) \in \mathbf{V}_h \times Q_h} \frac{\mathbf{B}((\mathbf{u}, p), (\mathbf{w}, t))}{\|(\mathbf{w}, t)\|_h} \geq \beta \|(\mathbf{u}, p)\|_h,$$

for all $(\mathbf{u}, p) \in \mathbf{V}_h \times Q_h$.

Proof. - Let $(\mathbf{u}, p) \in \mathbf{V}_h \times Q_h$. Since $p \in L_0^2(\Omega)$, there exists $\mathbf{v} \in [H_0^1(\Omega)]^2$ such that $\nabla \cdot \mathbf{v} = -p$ in Ω and $\|\mathbf{v}\|_{1,\Omega} \leq C \|p\|_{0,\Omega}$. Now, let \mathbf{v}_h be the Clément interpolate of \mathbf{v} (cf. [17, 9]), which satisfies

$$\|\mathbf{v} - \mathbf{v}_h\|_{0,K} \leq C h_K \|\mathbf{v}\|_{1,V(K)}, \quad (14)$$

$$\|\mathbf{v}_h\|_{1,\Omega} \leq C \|\mathbf{v}\|_{1,\Omega}, \quad (15)$$

where $V(K)$ is the set of elements in \mathcal{T}_h who share at least one node with K . Now, since \mathbf{v} and \mathbf{v}_h vanish on $\partial\Omega$ and $Q_h \subseteq C^0(\bar{\Omega})$, we can integrate by parts, use the fact that

$\nabla \cdot \mathbf{v} = -p$, and obtain

$$\begin{aligned}
\mathbf{B}((\mathbf{u}, p), (\mathbf{v}_h, 0)) &= \sigma(\mathbf{u}, \mathbf{v}_h)_\Omega + \nu(\nabla \mathbf{u}, \nabla \mathbf{v}_h)_\Omega + (\mathbf{a} \cdot \nabla \mathbf{u}, \mathbf{v}_h)_\Omega \\
&\quad - (p, \nabla \cdot \mathbf{v}_h)_\Omega + \sum_{K \in \mathcal{T}_h} (\nabla \cdot \mathbf{u}, \delta_K \nabla \cdot \mathbf{v}_h)_K \\
&\quad - \sum_{K \in \mathcal{T}_h} (\sigma \mathbf{u} - \nu \Delta \mathbf{u} + \mathbf{a} \cdot \nabla \mathbf{u} + \nabla p, \tau_K (\sigma \mathbf{v}_h - \nu \Delta \mathbf{v}_h - \mathbf{a} \cdot \nabla \mathbf{v}_h))_K \\
&= \sigma(\mathbf{u}, \mathbf{v}_h)_\Omega + \nu(\nabla \mathbf{u}, \nabla \mathbf{v}_h)_\Omega + (\mathbf{a} \cdot \nabla \mathbf{u}, \mathbf{v})_\Omega - (\mathbf{a} \cdot \nabla \mathbf{u}, \mathbf{v} - \mathbf{v}_h)_\Omega \\
&\quad - \sum_{K \in \mathcal{T}_h} (\nabla p, \mathbf{v} - \mathbf{v}_h)_K + \|p\|_{0,\Omega}^2 + \sum_{K \in \mathcal{T}_h} (\nabla \cdot \mathbf{u}, \delta_K \nabla \cdot \mathbf{v}_h)_K \\
&\quad - \sum_{K \in \mathcal{T}_h} (\sigma \mathbf{u} - \nu \Delta \mathbf{u} + \mathbf{a} \cdot \nabla \mathbf{u} + \nabla p, \tau_K (\sigma \mathbf{v}_h - \nu \Delta \mathbf{v}_h - \mathbf{a} \cdot \nabla \mathbf{v}_h))_K.
\end{aligned}$$

Hence, using Cauchy-Schwartz one gets

$$\begin{aligned}
\mathbf{B}((\mathbf{u}, p), (\mathbf{v}_h, 0)) &\geq -\sigma \|\mathbf{u}\|_{0,\Omega} \|\mathbf{v}_h\|_{0,\Omega} - \nu |\mathbf{u}|_{1,\Omega} |\mathbf{v}_h|_{1,\Omega} - \|\mathbf{a} \cdot \nabla \mathbf{u}\|_{0,\Omega} \|\mathbf{v}\|_{0,\Omega} \\
&\quad - \sum_{K \in \mathcal{T}_h} \|\tau_K^{1/2} (\mathbf{a} \cdot \nabla \mathbf{u} + \nabla p)\|_{0,K} \|\tau_K^{-1/2} (\mathbf{v} - \mathbf{v}_h)\|_{0,K} + \|p\|_{0,\Omega}^2 \\
&\quad - \sum_{K \in \mathcal{T}_h} \|\delta_K^{1/2} \nabla \cdot \mathbf{u}\|_{0,K} \|\delta_K^{1/2} \nabla \cdot \mathbf{v}_h\|_{0,K} \\
&\quad - \sum_{K \in \mathcal{T}_h} \|\tau_K^{1/2} (\sigma \mathbf{u} - \nu \Delta \mathbf{u} + \mathbf{a} \cdot \nabla \mathbf{u} + \nabla p)\|_{0,K} \\
&\quad \cdot \|\tau_K^{1/2} (\sigma \mathbf{v}_h - \nu \Delta \mathbf{v}_h - \mathbf{a} \cdot \nabla \mathbf{v}_h)\|_{0,K}. \tag{16}
\end{aligned}$$

Now, in each $K \in \mathcal{T}_h$, we have

$$\frac{1}{\tau_K} = \begin{cases} \frac{\frac{4\nu}{m_k} + |\mathbf{a}(\mathbf{x})|_p h_K}{h_K^2} \leq \frac{C \|\mathbf{a}\|_{\infty,\Omega}}{h_K} & \text{if } Pe_K^1, Pe_K^2 > 1, \\ \frac{8\nu}{m_k h_K^2} & \text{if } Pe_K^1 > 1, Pe_K^2 < 1, \\ \frac{\sigma h_K^2 + |\mathbf{a}(\mathbf{x})|_p h_K}{h_K^2} \leq C \frac{\sigma h_K + \|\mathbf{a}\|_{\infty,\Omega}}{h_K} & \text{if } Pe_K^1 < 1, Pe_K^2 > 1, \\ \frac{\sigma h_K^2 + \frac{4\nu}{m_k}}{h_K^2} \leq 2\sigma & \text{if } Pe_K^1, Pe_K^2 < 1, \end{cases} \tag{17}$$

Thus, denoting generically by $\frac{1}{\tau_K^*}$ the right hand side of (17), and applying (14) we arrive at

$$\begin{aligned}
\mathbf{B}((\mathbf{u}, p), (\mathbf{v}_h, 0)) &\geq -\sigma \|\mathbf{u}\|_{0,\Omega} \|\mathbf{v}_h\|_{0,\Omega} - \nu |\mathbf{u}|_{1,\Omega} |\mathbf{v}_h|_{1,\Omega} - \|\mathbf{a} \cdot \nabla \mathbf{u}\|_{0,\Omega} \|\mathbf{v}\|_{0,\Omega} \\
&\quad - C \sum_{K \in \mathcal{T}_h} \|\tau_K^{1/2} (\mathbf{a} \cdot \nabla \mathbf{u} + \nabla p)\|_{0,K} \frac{h_K}{\sqrt{\tau_K^*}} \|\mathbf{v}\|_{1,V(K)} + \|p\|_{0,\Omega}^2 \\
&\quad - \sum_{K \in \mathcal{T}_h} \|\delta_K^{1/2} \nabla \cdot \mathbf{u}\|_{0,K} \|\delta_K^{1/2} \nabla \cdot \mathbf{v}_h\|_{0,K} \\
&\quad - \sum_{K \in \mathcal{T}_h} \left[\sigma \overline{\tau_K}^{1/2} \|\mathbf{u}\|_{0,\Omega} + \nu \overline{\tau_K}^{1/2} \|\Delta \mathbf{u}\|_{0,\Omega} + \|\tau_K^{1/2} (\mathbf{a} \cdot \nabla \mathbf{u} + \nabla p)\|_{0,K} \right] \\
&\quad \cdot \left[\sigma \overline{\tau_K}^{1/2} \|\mathbf{v}_h\|_{0,\Omega} + \nu \overline{\tau_K}^{1/2} \|\Delta \mathbf{v}_h\|_{0,\Omega} + \overline{\tau_K}^{1/2} \|\mathbf{a} \cdot \nabla \mathbf{v}_h\|_{0,K} \right] \\
&\geq -C \left\{ \sigma \|\mathbf{u}\|_{0,\Omega}^2 + \nu |\mathbf{u}|_{1,\Omega}^2 + \|\mathbf{a} \cdot \nabla \mathbf{u}\|_{0,\Omega}^2 \right. \\
&\quad \left. + \sum_{K \in \mathcal{T}_h} \|\tau_K^{1/2} (\mathbf{a} \cdot \nabla \mathbf{u} + \nabla p)\|_{0,K}^2 + \sum_{K \in \mathcal{T}_h} \|\delta_K^{1/2} \nabla \cdot \mathbf{u}\|_{0,K}^2 \right\}^{1/2} \\
&\quad \cdot \left\{ \sigma \|\mathbf{v}_h\|_{0,\Omega}^2 + \nu |\mathbf{v}_h|_{1,\Omega}^2 + \|\mathbf{v}\|_{0,\Omega}^2 + \sum_{K \in \mathcal{T}_h} \frac{h_K^2}{\tau_K^*} \|\mathbf{v}\|_{1,V(K)}^2 \right. \\
&\quad \left. + \sum_{K \in \mathcal{T}_h} \|\delta_K^{1/2} \nabla \cdot \mathbf{v}_h\|_{0,K}^2 \right\}^{1/2} + \|p\|_{0,\Omega}^2 \\
&\quad - C \left\{ \sum_{K \in \mathcal{T}_h} \left[\sigma^2 \overline{\tau_K} \|\mathbf{u}\|_{0,\Omega}^2 + \nu^2 \overline{\tau_K} \|\Delta \mathbf{u}\|_{0,\Omega}^2 + \|\tau_K^{1/2} (\mathbf{a} \cdot \nabla \mathbf{u} + \nabla p)\|_{0,K}^2 \right] \right\}^{1/2} \\
&\quad \cdot \left\{ \sum_{K \in \mathcal{T}_h} \left[\sigma^2 \overline{\tau_K} \|\mathbf{v}_h\|_{0,\Omega}^2 + \nu^2 \overline{\tau_K} \|\Delta \mathbf{v}_h\|_{0,\Omega}^2 + \overline{\tau_K} \|\mathbf{a} \cdot \nabla \mathbf{v}_h\|_{0,K}^2 \right] \right\}^{1/2}.
\end{aligned}$$

Now, from inverse inequality (11) a simple computation shows that

$$\nu^2 \overline{\tau_K} \|\Delta \mathbf{w}\|_{0,K}^2 \leq \nu |\mathbf{w}|_{1,K}^2 \quad \forall \mathbf{w} \in \mathbf{V}_h, \quad (18)$$

and hence, from the definition of δ_K and τ_K^* , the fact that $\sigma\overline{\tau_K} \leq 1$, and $\|\mathbf{v}_h\|_{1,\Omega} \leq C\|\mathbf{v}\|_{1,\Omega}$, inequality above becomes

$$\begin{aligned}
\mathbf{B}((\mathbf{u}, p), (\mathbf{v}_h, 0)) &\geq -C \left\{ \sigma \|\mathbf{u}\|_{0,\Omega}^2 + \nu |\mathbf{u}|_{1,\Omega}^2 + \|\mathbf{a}\|_{\infty,\Omega}^2 |\mathbf{u}|_{1,\Omega}^2 \right. \\
&\quad \left. + \sum_{K \in \mathcal{T}_h} \|\tau_K^{1/2} (\mathbf{a} \cdot \nabla \mathbf{u} + \nabla p)\|_{0,K}^2 + \sum_{K \in \mathcal{T}_h} \|\delta_K^{1/2} \nabla \cdot \mathbf{u}\|_{0,K}^2 \right\}^{1/2} \\
&\quad \cdot \left\{ \sigma \|\mathbf{v}_h\|_{0,\Omega}^2 + \nu |\mathbf{v}_h|_{1,\Omega}^2 + \|\mathbf{v}\|_{0,\Omega}^2 \right. \\
&\quad \left. + \max\left\{ \frac{8\nu}{m_k}, \|\mathbf{a}\|_{\infty,\Omega} h, \sigma h^2 \right\} \|\mathbf{v}\|_{1,\Omega}^2 + \lambda \|\mathbf{a}\|_{\infty,\Omega} h |\mathbf{v}_h|_{1,\Omega}^2 \right\}^{1/2} \\
&\quad + \|p\|_{0,\Omega}^2 \\
&\quad - C \left\{ \sum_{K \in \mathcal{T}_h} \left[\sigma \|\mathbf{u}\|_{0,K}^2 + \nu |\mathbf{u}|_{1,K}^2 + \|\tau_K^{1/2} (\mathbf{a} \cdot \nabla \mathbf{u} + \nabla p)\|_{0,K}^2 \right] \right\}^{1/2} \\
&\quad \cdot \left\{ \sum_{K \in \mathcal{T}_h} \left[\sigma \|\mathbf{v}_h\|_{0,K}^2 + \nu |\mathbf{v}_h|_{1,K}^2 + \|\mathbf{a}\|_{\infty,K\overline{\tau_K}}^2 |\mathbf{v}_h|_{1,K}^2 \right] \right\}^{1/2} \\
&\geq -C_* \left\{ \|\mathbf{u}\|_{1,\Omega}^2 + \sum_{K \in \mathcal{T}_h} \left[\|\tau_K^{1/2} (\mathbf{a} \cdot \nabla \mathbf{u} + \nabla p)\|_{0,K}^2 + \|\delta_K^{1/2} \nabla \cdot \mathbf{u}\|_{0,K}^2 \right] \right\}^{1/2} \cdot \|\mathbf{v}\|_{1,\Omega} \\
&\quad + \|p\|_{0,\Omega}^2 \\
&\quad - C_{**} \left\{ \|\mathbf{u}\|_{1,\Omega}^2 + \sum_{K \in \mathcal{T}_h} \|\tau_K^{1/2} (\mathbf{a} \cdot \nabla \mathbf{u} + \nabla p)\|_{0,K}^2 \right\}^{1/2} \cdot \|\mathbf{v}\|_{1,\Omega},
\end{aligned}$$

where C_* and C_{**} are positive constants depending on σ , ν , λ and $\|\mathbf{a}\|_{\infty,\Omega}$.

Now, applying the fact that $\|\mathbf{v}\|_{1,\Omega} \leq C\|p\|_{0,\Omega}$, and using the inequality $ab \leq \frac{1}{\gamma}a^2 + \gamma b^2$ in each expression involving a product of norms above, we obtain

$$\begin{aligned}
\mathbf{B}((\mathbf{u}, p), (\mathbf{v}_h, 0)) &\geq -\frac{C_*}{\gamma_1} \left\{ \|\mathbf{u}\|_{1,\Omega}^2 + \sum_{K \in \mathcal{T}_h} \left[\|\tau_K^{1/2} (\mathbf{a} \cdot \nabla \mathbf{u} + \nabla p)\|_{0,K}^2 + \|\delta_K^{1/2} \nabla \cdot \mathbf{u}\|_{0,K}^2 \right] \right\} \\
&\quad - \frac{C_{**}}{\gamma_2} \left\{ \|\mathbf{u}\|_{1,\Omega}^2 + \sum_{K \in \mathcal{T}_h} \left[\|\tau_K^{1/2} (\mathbf{a} \cdot \nabla \mathbf{u} + \nabla p)\|_{0,K}^2 \right] \right\} \\
&\quad + (1 - C_*\gamma_1 - C_{**}\gamma_2) \|p\|_{0,\Omega}^2 \\
&\geq -C^{**} \left\{ \|\mathbf{u}\|_{1,\Omega}^2 + \sum_{K \in \mathcal{T}_h} \left[\|\tau_K^{1/2} (\mathbf{a} \cdot \nabla \mathbf{u} + \nabla p)\|_{0,K}^2 + \|\delta_K^{1/2} \nabla \cdot \mathbf{u}\|_{0,K}^2 \right] \right\} \\
&\quad + C^* \|p\|_{0,\Omega}^2, \tag{19}
\end{aligned}$$

where C^* and C^{**} are positive constants, independent of h , if γ_1 and γ_2 are chosen small enough.

In this form, if we set $(\mathbf{z}, q) \stackrel{\text{def}}{=} (\mathbf{u} + \gamma \mathbf{v}_h, p)$, $\gamma > 0$, we have by the bilinearity of \mathbf{B} , Lemma 2.1 and (19) that

$$\begin{aligned} \mathbf{B}((\mathbf{u}, p), (\mathbf{z}, q)) &= \mathbf{B}((\mathbf{u}, p), (\mathbf{u}, p)) + \gamma \mathbf{B}((\mathbf{u}, p), (\mathbf{v}_h, 0)) \\ &\geq C_{\Omega} \nu \|\mathbf{u}\|_{1,\Omega}^2 + \sum_{K \in \mathcal{T}_h} \left[\|\tau_K^{1/2} (\mathbf{a} \cdot \nabla \mathbf{u} + \nabla p)\|_{0,K}^2 + \|\delta_K^{1/2} \nabla \cdot \mathbf{u}\|_{0,K}^2 \right] \\ &\quad - \gamma C^{**} \left\{ \|\mathbf{u}\|_{1,\Omega}^2 + \sum_{K \in \mathcal{T}_h} \left[\|\tau_K^{1/2} (\mathbf{a} \cdot \nabla \mathbf{u} + \nabla p)\|_{0,K}^2 + \|\delta_K^{1/2} \nabla \cdot \mathbf{u}\|_{0,K}^2 \right] \right\} \\ &\quad + \gamma C^* \|p\|_{0,\Omega}^2 \\ &\geq C \|(\mathbf{u}, p)\|_h^2, \end{aligned} \tag{20}$$

where $C > 0$ is independent of h if γ is chosen small enough.

Finally, using that $h \leq 1$ and $\|\mathbf{v}\|_{1,\Omega} \leq C \|p\|_{0,\Omega}$, we obtain

$$\begin{aligned} \|(\mathbf{z}, q)\|_h^2 &\leq 2 \|\mathbf{u}\|_{1,\Omega}^2 + 2 \sum_{K \in \mathcal{T}_h} \|\tau_K^{1/2} (\mathbf{a} \cdot \nabla \mathbf{u} + \nabla p)\|_{0,K}^2 + 2 \sum_{K \in \mathcal{T}_h} \|\delta_K^{1/2} \nabla \cdot \mathbf{u}\|_{0,K}^2 + \|p\|_{0,\Omega}^2 \\ &\quad + 2\gamma^2 \left[\|\mathbf{v}_h\|_{1,\Omega}^2 + \sum_{K \in \mathcal{T}_h} \tau_K^{-1/2} \|\mathbf{a} \cdot \nabla \mathbf{v}_h\|_{0,K}^2 + \sum_{K \in \mathcal{T}_h} \|\delta_K^{1/2} \nabla \cdot \mathbf{v}_h\|_{0,K}^2 \right] \\ &\leq C \|(\mathbf{u}, p)\|_h^2, \end{aligned}$$

which, together with (20), finish the proof. \square

2.2 Error Analysis

Let k, l be integers with $k, l \geq 1$. We use the Lagrange interpolation operator $\mathbf{I}_h^k : (C^0(\bar{\Omega}))^2 \rightarrow [V_k]^2$, we denote $\tilde{\mathbf{u}}_h \stackrel{\text{def}}{=} \mathbf{I}_h^k(\mathbf{u})$, define the interpolation error $\eta^{\mathbf{u}} \stackrel{\text{def}}{=} \mathbf{u} - \tilde{\mathbf{u}}_h$, and we have (cf. [8])

$$|\eta^{\mathbf{u}}|_{m,K} \leq Ch_K^{s-m} |\mathbf{u}|_{s,K}, \tag{21}$$

if $\mathbf{u} \in H^s(K)^2$ for all $K \in \mathcal{T}_h$, with $0 \leq m \leq 2$ and $\max\{m, 2\} \leq s \leq k+1$. Now, for the pressure we define \tilde{p}_h as being the Clément interpolate of p . Denoting now by $\eta^p \stackrel{\text{def}}{=} p - \tilde{p}_h$, where

$$\tilde{p}_h \stackrel{\text{def}}{=} \tilde{p}_h - \frac{1}{|\Omega|} (\tilde{p}_h, 1)_{\Omega} \in L_0^2(\Omega),$$

we have (cf. [17])

$$\|\eta^p\|_{0,\Omega} \leq Ch^t \|p\|_{t,\Omega}, \tag{22}$$

$$|\eta^p|_{1,K} \leq Ch^{t-1} \|p\|_{t,V(K)}, \tag{23}$$

if $p \in H^t(\Omega)$, with $1 \leq t \leq l+1$.

The main result concerning approximation is now stated.

Theorem 2.3 *Let us suppose that the solution (\mathbf{u}, p) of (1) belongs to $(H^{k+1}(\Omega) \cap H_0^1(\Omega))^2 \times (H^l(\Omega) \cap L_0^2(\Omega))$. Then, there exists $C > 0$, independent of h , such that the error $(\mathbf{e}^{\mathbf{u}}, e^p) \stackrel{\text{def}}{=} (\mathbf{u} - \mathbf{u}_h, p - p_h)$ satisfies*

$$\|(\mathbf{e}^{\mathbf{u}}, e^p)\|_h \leq C \left[h^k |\mathbf{u}|_{k+1, \Omega} + h^l \|p\|_{l, \Omega} \right].$$

Proof. - Let $\mathbf{e}_h^{\mathbf{u}} \stackrel{\text{def}}{=} \mathbf{u}_h - \tilde{\mathbf{u}}_h$ and $e_h^p \stackrel{\text{def}}{=} p_h - \bar{p}_h$. From the proof of previous theorem, we see that the supremum is attained, and then there exists $(\mathbf{v}, q) \in \mathbf{V}_h \times Q_h$ such that

$$\|(\mathbf{v}, q)\|_h \leq C,$$

and

$$\beta \|(\mathbf{e}_h^{\mathbf{u}}, e_h^p)\|_h \leq \mathbf{B}((\mathbf{e}_h^{\mathbf{u}}, e_h^p), (\mathbf{v}, q)).$$

Using the consistency of the method we obtain

$$\mathbf{B}((\mathbf{e}_h^{\mathbf{u}}, e_h^p), (\mathbf{v}, q)) = \mathbf{B}((\eta^{\mathbf{u}}, \eta^p), (\mathbf{v}, q)). \quad (24)$$

Now, for the right hand side of (24) we have by using Schwartz's inequality we obtain

$$\begin{aligned} \mathbf{B}((\eta^{\mathbf{u}}, \eta^p), (\mathbf{v}, q)) &= \sigma(\eta^{\mathbf{u}}, \mathbf{v})_{\Omega} + \nu(\nabla \eta^{\mathbf{u}}, \nabla \mathbf{v})_{\Omega} + (\mathbf{a} \cdot \nabla \eta^{\mathbf{u}}, \mathbf{v})_{\Omega} \\ &\quad - (\eta^p, \nabla \cdot \mathbf{v})_{\Omega} + (q, \nabla \cdot \eta^{\mathbf{u}})_{\Omega} + \sum_{K \in \mathcal{T}_h} (\nabla \cdot \eta^{\mathbf{u}}, \delta_K \nabla \cdot \mathbf{v})_K \\ &\quad - \sum_{K \in \mathcal{T}_h} (\sigma \eta^{\mathbf{u}} - \nu \Delta \eta^{\mathbf{u}} + \mathbf{a} \cdot \nabla \eta^{\mathbf{u}} + \nabla \eta^p, \tau_K (\sigma \mathbf{v} - \nu \Delta \mathbf{v} - \mathbf{a} \cdot \nabla \mathbf{v} - \nabla q))_K \\ &\leq \left\{ \sum_{K \in \mathcal{T}_h} \sigma \|\eta^{\mathbf{u}}\|_{0,K}^2 + \nu |\eta^{\mathbf{u}}|_{1,K}^2 + \|\mathbf{a} \cdot \nabla \eta^{\mathbf{u}}\|_{0,K}^2 + \|\eta^p\|_{0,K}^2 + \|\nabla \cdot \eta^{\mathbf{u}}\|_{0,K}^2 \right. \\ &\quad \left. + \|\delta_K^{1/2} \nabla \cdot \eta^{\mathbf{u}}\|_{0,K}^2 + \|\tau_K^{1/2} (\sigma \eta^{\mathbf{u}} - \nu \Delta \eta^{\mathbf{u}} + \mathbf{a} \cdot \nabla \eta^{\mathbf{u}} + \nabla \eta^p)\|_{0,K}^2 \right\}^{\frac{1}{2}} \\ &\quad \cdot \left\{ \sum_{K \in \mathcal{T}_h} \sigma \|\mathbf{v}\|_{0,K}^2 + \nu |\mathbf{v}|_{1,K}^2 + \|\mathbf{v}\|_{0,K}^2 + \|\nabla \cdot \mathbf{v}\|_{0,K}^2 + \|q\|_{0,K}^2 \right. \\ &\quad \left. + \|\delta_K^{1/2} \nabla \cdot \mathbf{v}\|_{0,K}^2 + \|\tau_K^{1/2} (\sigma \mathbf{v} - \nu \Delta \mathbf{v} - \mathbf{a} \cdot \nabla \mathbf{v} - \nabla q)\|_{0,K}^2 \right\}^{\frac{1}{2}}. \end{aligned}$$

Now, applying inequality (18) for the $-\nu^2 \tau_K \|\Delta \mathbf{v}\|_{0,K}^2$ term and $\sigma \overline{\tau_K} \leq 1$, previous inequality becomes

$$\begin{aligned}
\mathbf{B}((\eta^{\mathbf{u}}, \eta^p), (\mathbf{v}, q)) &\leq C \left\{ \sum_{K \in \mathcal{T}_h} \sigma \|\eta^{\mathbf{u}}\|_{0,K}^2 + \nu |\eta^{\mathbf{u}}|_{1,K}^2 + \|\mathbf{a} \cdot \nabla \eta^{\mathbf{u}}\|_{0,K}^2 + \|\eta^p\|_{0,K}^2 + \|\nabla \cdot \eta^{\mathbf{u}}\|_{0,K}^2 \right. \\
&\quad + \|\delta_K^{1/2} \nabla \cdot \eta^{\mathbf{u}}\|_{0,K}^2 + \sigma^2 \overline{\tau_K} \|\eta^{\mathbf{u}}\|_{0,K}^2 + \nu^2 \overline{\tau_K} \|\Delta \eta^{\mathbf{u}}\|_{0,K}^2 \\
&\quad \left. + \overline{\tau_K} \|\mathbf{a} \cdot \nabla \eta^{\mathbf{u}}\|_{0,K}^2 + \overline{\tau_K} \|\nabla \eta^p\|_{0,K}^2 \right\}^{\frac{1}{2}} \\
&\quad \cdot \left\{ \sum_{K \in \mathcal{T}_h} (\sigma + 1) \|\mathbf{v}\|_{0,K}^2 + (\nu + 1) |\mathbf{v}|_{1,K}^2 + \|q\|_{0,K}^2 + \|\delta_K^{1/2} \nabla \cdot \mathbf{v}\|_{0,K}^2 \right. \\
&\quad \left. + \sigma \|\mathbf{v}\|_{0,K}^2 + \nu |\mathbf{v}|_{1,K}^2 + \|\tau_K^{1/2} (\mathbf{a} \cdot \nabla \mathbf{v} + \nabla q)\|_{0,K}^2 \right\}^{\frac{1}{2}} \\
&\leq C \left\{ \max\{\sigma, \nu, \|\mathbf{a}\|_{\infty, \Omega}, \|\mathbf{a}\|_{\infty, \Omega} h, \frac{1}{\nu}\} \sum_{K \in \mathcal{T}_h} \left[\|\eta^{\mathbf{u}}\|_{0,K}^2 + |\eta^{\mathbf{u}}|_{1,K}^2 + \|\eta^p\|_{0,K}^2 \right. \right. \\
&\quad \left. \left. + h_K^2 \|\Delta \eta^{\mathbf{u}}\|_{0,K}^2 + h_K^2 |\eta^p|_{1,K}^2 \right] \right\}^{\frac{1}{2}} \\
&\quad \cdot \left\{ \max\{\sigma, \nu, 1\} \sum_{K \in \mathcal{T}_h} \left[\|\mathbf{v}\|_{0,K}^2 + |\mathbf{v}|_{1,K}^2 + \|q\|_{0,K}^2 + \|\delta_K^{1/2} \nabla \cdot \mathbf{v}\|_{0,K}^2 \right. \right. \\
&\quad \left. \left. + \|\mathbf{v}\|_{0,K}^2 + |\mathbf{v}|_{1,K}^2 + \|\tau_K^{1/2} (\mathbf{a} \cdot \nabla \mathbf{v} + \nabla q)\|_{0,K}^2 \right] \right\}^{\frac{1}{2}}.
\end{aligned}$$

Now, the second term in the product above is smaller than $C \|(\mathbf{v}, q)\|_h$ and hence it is bounded by a constant. Therefore, applying interpolation inequalities (21), (22) and (23) we arrive at

$$\begin{aligned}
\mathbf{B}((\eta^{\mathbf{u}}, \eta^p), (\mathbf{v}, q)) &\leq C(\sigma, \nu, \mathbf{a}) \left\{ \sum_{K \in \mathcal{T}_h} \left[\|\eta^{\mathbf{u}}\|_{0,K}^2 + |\eta^{\mathbf{u}}|_{1,K}^2 + h_K^2 \|\Delta \eta^{\mathbf{u}}\|_{0,K}^2 \right] \right. \\
&\quad \left. + \|\eta^p\|_{0,\Omega}^2 + h^2 |\eta^p|_{1,\Omega}^2 \right\}^{\frac{1}{2}} \\
&\leq C (h^k \|\mathbf{u}\|_{k+1,\Omega} + h^l \|p\|_{l,\Omega}). \tag{25}
\end{aligned}$$

Finally, applying triangle inequality and (25) we arrive at

$$\begin{aligned}
\|(\mathbf{e}^{\mathbf{u}}, e^p)\|_h &\leq \|(\mathbf{e}_h^{\mathbf{u}}, e_h^p)\|_h + \|(\eta^{\mathbf{u}}, \eta^p)\|_h \\
&\leq \frac{C}{\beta} (h^k |\mathbf{u}|_{k+1, \Omega} + h^l \|p\|_{L(\Omega)}) \\
&\quad + \left\{ \|\eta^{\mathbf{u}}\|_{1, \Omega}^2 + \|\eta^p\|_{0, \Omega}^2 + \sum_{K \in \mathcal{T}_h} \|\delta_K^{1/2} \nabla \cdot \eta^{\mathbf{u}}\|_{0, K}^2 + \|\tau_K^{1/2} (\mathbf{a} \cdot \nabla \eta^{\mathbf{u}} + \nabla \eta^p)\|_{0, K}^2 \right\}^{\frac{1}{2}} \\
&\leq \frac{C}{\beta} (h^k |\mathbf{u}|_{k+1, \Omega} + h^l \|p\|_{L(\Omega)}) \\
&\quad + C \left\{ \|\eta^{\mathbf{u}}\|_{1, \Omega}^2 + \|\eta^p\|_{0, \Omega}^2 + \|\mathbf{a}\|_{\infty, \Omega} h |\eta^{\mathbf{u}}|_{1, \Omega}^2 + \|\mathbf{a}\|_{\infty, \Omega}^2 \frac{h^2}{\nu} |\eta^{\mathbf{u}}|_{1, \Omega}^2 + \frac{h^2}{\nu} |\eta^p|_{1, \Omega}^2 \right\}^{\frac{1}{2}} \\
&\leq \frac{C}{\beta} (h^k |\mathbf{u}|_{k+1, \Omega} + h^l \|p\|_{L(\Omega)}) + C \{ \|\eta^{\mathbf{u}}\|_{1, \Omega}^2 + \|\eta^p\|_{0, \Omega}^2 + h^2 |\eta^p|_{1, \Omega}^2 \}^{\frac{1}{2}},
\end{aligned}$$

and the proof follows by applying interpolation inequalities (21), (22) and (23) once again. \square

Remark 3 *We observe that the estimate given by previous theorem implies an error estimate in the natural norms of \mathbf{u} and p , (the $H^1(\Omega)^2$ -norm for the velocity and the $L^2(\Omega)$ -norm for the pressure), that are optimal in order and regularity. In next section we will show that the errors are not too much affected by the physical coefficients. \square*

3 Numerical experiments

We use as domain the square $(0, 1) \times (0, 1)$, and we set \mathbf{f} to be such as the exact solution of our problem (1) is given by

$$\begin{aligned}
u_1(x_1, x_2) &= -256 x_1^2 (x_1 - 1)^2 x_2 (x_2 - 1) (2x_2 - 1), \\
u_2(x_1, x_2) &= -u_1(x_2, x_1), \\
p(x_1, x_2) &= 150 (x_1 - 0.5)(x_2 - 0.5).
\end{aligned}$$

First, we report the diffusive dominated case with $\sigma = 1, \nu = 1, \mathbf{a} = (1, 1)$ and $\lambda = 0$. The results are depicted in Figures 1 and 2, using P^1/P^1 and Q^1/Q^1 finite elements. We recover optimal orders of convergence for velocity and pressure.

Next, we have considered the reaction-convection dominated case with $\sigma = 10^2, \nu = 10^{-3}, \mathbf{a} = (1, 1)$ and $\lambda = 0$ and $\lambda = 0.5$. Figures 3 to 6 show the convergence history, and they agree with the theoretical results. We recover an optimal order of convergence for the pressure. However, a quasi-optimal $h^{3/2}$ order of convergence is observed for $\|\mathbf{u} - \mathbf{u}_h\|_{0, \Omega}$. This well known fact in SDFEM-like stabilized methods (cf. [22, 12, 25], specially refer to the introduction in [23]) is recovered by our method. On the other hand, we notice that the *div-div* term, $\sum_{K \in \mathcal{T}_h} (\nabla \cdot \mathbf{u}, \delta_K \nabla \cdot \mathbf{v})_K$, does not provide a significant improvement of




Figure 1: P^1 convergence history: $\sigma = 1, \nu = 1, \mathbf{a} = (1, 1), \lambda = 0$






Figure 2: Q^1 convergence history: $\sigma = 1, \nu = 1, \mathbf{a} = (1, 1), \lambda = 0$

the convergence rate. Because of this, in the sequel we will consider only $\lambda = 0$ in our computations.

Figure 3: P^1 convergence history: $\sigma = 100$, $\nu = 10^{-3}$, $\mathbf{a} = (1, 1)$, $\lambda = 0$ Figure 4: P^1 convergence history: $\sigma = 100$, $\nu = 10^{-3}$, $\mathbf{a} = (1, 1)$, $\lambda = 0.5$

Now, we address the study of the sensitivity of the error to the physical coefficients. To this purpose, we use a uniform 40×40 mesh ($= 3200 P^1/P^1$ elements), and we measure the errors in velocity and pressure.




Figure 5: Q^1 convergence history: $\sigma = 100$, $\nu = 10^{-3}$, $\mathbf{a} = (1, 1)$, $\lambda = 0$




Figure 6: Q^1 convergence history: $\sigma = 100$, $\nu = 10^{-3}$, $\mathbf{a} = (1, 1)$, $\lambda = 0.5$

First, we fix $\nu = 10^{-3}$ and $\mathbf{a} = (1, 1)$ and make σ grow. The results are shown in Table 1, where we see that the velocity error remains bounded while σ grows and that the pressure error presents a good behavior even for very large values of σ .

σ	$\ \mathbf{u} - \mathbf{u}_h\ _{0,\Omega}$	$\ \mathbf{u} - \mathbf{u}_h\ _{1,\Omega}$	$\ p - p_h\ _{0,\Omega}$
0.1	5.5269×10^{-3}	0.7425	9.9611×10^{-2}
1	5.1536×10^{-3}	0.7397	9.5769×10^{-2}
10	4.4729×10^{-3}	0.7322	0.1079
100	3.4800×10^{-3}	0.7238	0.1316
10^3	3.3633×10^{-3}	0.7245	0.1427
10^4	3.3666×10^{-3}	0.7252	0.1443

Table 1: Behavior of the Finite Element error when σ grows

ν	$\ \mathbf{u} - \mathbf{u}_h\ _{0,\Omega}$	$\ \mathbf{u} - \mathbf{u}_h\ _{1,\Omega}$	$\ p - p_h\ _{0,\Omega}$
1	5.7078×10^{-3}	0.7204	5.3583×10^{-2}
0.1	5.4621×10^{-3}	0.7207	5.8640×10^{-2}
0.01	9.2784×10^{-3}	0.7234	0.1095
10^{-3}	3.4800×10^{-3}	0.7238	0.1316
10^{-4}	2.6934×10^{-3}	0.7278	0.1320
10^{-5}	2.6781×10^{-3}	0.7285	0.1320
10^{-6}	2.6774×10^{-3}	0.7286	0.1320

Table 2: Behavior of the Finite Element error when ν decreases

\mathbf{a}	$\ \mathbf{u} - \mathbf{u}_h\ _{0,\Omega}$	$\ \mathbf{u} - \mathbf{u}_h\ _{1,\Omega}$	$\ p - p_h\ _{0,\Omega}$
(0.1,0.1)	9.3554×10^{-3}	0.7248	0.1383
(1,1)	3.4800×10^{-3}	0.7238	0.1316
(5,5)	2.8497×10^{-3}	0.7253	0.1150
(10,10)	3.0116×10^{-3}	0.7261	0.1108
(20,20)	3.2770×10^{-3}	0.7274	0.1190
(40,40)	3.5450×10^{-3}	0.7289	0.1737

Table 3: Behavior of the Finite Element error when $|\mathbf{a}|$ grows

Next, we fix $\sigma = 100$ and $\mathbf{a} = (1, 1)$, and make the viscosity ν decrease. The results are shown in Table 2, where we see that both errors are not significantly affected by the viscosity.

Finally, we fix $\sigma = 100$ and $\nu = 10^{-3}$, and let $|\mathbf{a}|$ grow. We observe, in Table 3, that the error in velocity remains bounded, while the error in pressure remains bounded for a quite large range of local Péclet numbers $\frac{m_\kappa |\mathbf{a}| h_K}{4\nu}$. Numerical experiences beyond that range of Péclet numbers, have shown that the pressure error grows. This is reasonable since we are already dealing with relatively high Reynolds number.

4 Conclusion

In this paper we have introduced and analyzed a new stabilized finite element method for the Oseen's equations containing dominating reaction terms. The method is obtained by subtracting a mesh dependent term to the standard Galerkin formulation without compromising consistency. The stabilization parameter is suggested by bubble condensation and it is completely determined from the data of the problem. The error estimates provide optimal convergence rate for velocity and pressure in its natural norms. The reported numerical experiments confirm the approximation results and point out that the performance of the method is not dramatically affected by the physical coefficients.

Acknowledgements

First author was partially supported by MACSI-net Program during his visit to the Modeling and Scientific Computing Chair of the EPFL (Switzerland) in February 2003. Second author acknowledges support by the Swiss National Science Foundation (contract number 20-65110.01) during his Post-Doctoral position at the EPFL.

References

- [1] C. Baiocchi, F. Brezzi, and L.P. Franca. Virtual bubbles and Galerkin-least-squares type methods (Ga. L. S.). *Comput. Methods Appl. Mech. Engrg.*, 105:125–141, 1993.
- [2] G. Barrenechea and F. Valentin. An unusual stabilized finite element method for a generalized Stokes problem. *Numer. Math.*, 92(4):653–677, 2002.
- [3] F. Brezzi and M. Fortin. *Mixed and Hybrid Finite Element Methods*. Springer-Verlag, 1991.
- [4] F. Brezzi and J. Douglas Jr. Stabilized mixed methods for the Stokes problem. *Numer. Math.*, 53:225–235, 1988.
- [5] F. Brezzi and J. Pitkaranta. On the stabilization of finite element approximations of the Stokes problem. In W. Hackbush, editor, *Efficient Solution of Elliptic Systems*, volume 10 of *Notes on Numerical Fluid Mechanics*, pages 11–19. Vieweg, 1984.
- [6] A.N. Brooks and T.J.R. Hughes. Streamline upwinding Petrov-Galerkin formulations for convective dominated flows with particular emphasis on the incompressible Navier-Stokes equations. *Comput. Methods Appl. Mech. Engrg.*, 32:199–259, 1982.
- [7] E. Burman and P. Hansbo. Edge stabilization for Galerkin approximations of convection-diffusion-reaction problems. *Comp. Methods Appl. Mech. Engrg.*, 193:1437–1453, 2004.

-
- [8] P.G. Ciarlet. *The Finite Element Method for Elliptic Problems*. North Holland, 1978.
- [9] P. Clément. Approximation by finite element functions using local regularization. *R.A.I.R.O. Anal. Numer.*, 9:77–84, 1975.
- [10] R. Codina. A stabilized finite element method for generalized stationary incompressible flows. *Comput. Methods Appl. Mech. Engrg.*, 190(20-21):2681–2706, 2001.
- [11] J. Donea and A. Huerta. *Finite Element Methods for Flow Problems*. Wiley, 2003.
- [12] A. Ern and J.L. Guermond. *Éléments finis: théorie, applications et mise en œuvre*. Springer-SMAI, 2002.
- [13] L.P. Franca and C. Farhat. Bubble functions prompt unusual stabilized finite element methods. *Comput. Methods Appl. Mech. Engrg.*, 123:299–308, 1995.
- [14] L.P. Franca and S.L. Frey. Stabilized finite element methods: II. the incompressible Navier-Stokes equations. *Comput. Methods Appl. Mech. Engrg.*, 99(2–3):209–233, 1992.
- [15] L.P. Franca and R. Stenberg. Error analysis of some Galerkin least squares methods for the elasticity equations. *SIAM J. Numer. Anal.*, 28(6):1680–1697, 1991.
- [16] L.P. Franca and F. Valentin. On an improved unusual stabilized finite element method for the advective-reactive-diffusive equation. *Comput. Methods Appl. Mech. Engrg.*, 190:1785–1800, 2000.
- [17] V. Girault and P.A. Raviart. *Finite Element Methods for the Navier-Stokes Equations*. Springer-Verlag, 1986.
- [18] I. Harari and T.J.R. Hughes. Stabilized finite element methods for steady advection-diffusion with production. *Comput. Methods Appl. Mech. Engrg.*, 115:165–191, 1994.
- [19] T.J.R. Hughes and L.P. Franca. A new finite element formulation for computational fluid dynamics: VII. the Stokes problem with various well-posed boundary conditions: Symmetric formulations that converge for all velocity/pressure spaces. *Comput. Methods Appl. Mech. Engrg.*, 65(1):85–96, 1987.
- [20] T.J.R. Hughes, L.P. Franca, and M. Balestra. A new finite element formulation for computational fluids dynamics: V. Circumventing the Babuska-Brezzi condition : A stable Petrov-Galerkin formulation of the Stokes problem accommodating equal-order interpolations. *Comput. Methods Appl. Mech. Engrg.*, 59(1):85–99, 1986.
- [21] A. Quarteroni and A. Valli. *Numerical Approximation of Partial Differential Equations*. Springer-Verlag, 1991.
- [22] H. Roos, M. Stynes, and L. Tobiska. *Numerical Methods for Singularly Perturbed Differential Equations. Convection-Diffusion and Flow Problems*. Springer-Verlag, 1996.

- [23] M. Stynes and L. Tobiska. The SDFEM for a convection-diffusion problem with a boundary layer: optimal error analysis and enhancement of accuracy. *SIAM J. Numer. Anal.*, 41(5):1620–1642, 2003.
- [24] L. Tobiska and R. Verfürth. Analysis of a streamline diffusion finite element method for the Stokes and Navier-Stokes equations. *SIAM J. Numer. Anal.*, 33(1):107–127, 1996.
- [25] G. Zhou. How accurate is the streamline diffusion finite element method? *Math. Comp.*, 66(217):31–44, 1997.



Unité de recherche INRIA Rocquencourt
Domaine de Voluceau - Rocquencourt - BP 105 - 78153 Le Chesnay Cedex (France)

Unité de recherche INRIA Futurs : Parc Club Orsay Université - ZAC des Vignes
4, rue Jacques Monod - 91893 ORSAY Cedex (France)

Unité de recherche INRIA Lorraine : LORIA, Technopôle de Nancy-Brabois - Campus scientifique
615, rue du Jardin Botanique - BP 101 - 54602 Villers-lès-Nancy Cedex (France)

Unité de recherche INRIA Rennes : IRISA, Campus universitaire de Beaulieu - 35042 Rennes Cedex (France)

Unité de recherche INRIA Rhône-Alpes : 655, avenue de l'Europe - 38334 Montbonnot Saint-Ismier (France)

Unité de recherche INRIA Sophia Antipolis : 2004, route des Lucioles - BP 93 - 06902 Sophia Antipolis Cedex (France)

Éditeur
INRIA - Domaine de Voluceau - Rocquencourt, BP 105 - 78153 Le Chesnay Cedex (France)
<http://www.inria.fr>
ISSN 0249-6399