

Analytical results on Connected dominating sets in mobile ad hoc networks

Philippe Jacquet

▶ **To cite this version:**

Philippe Jacquet. Analytical results on Connected dominating sets in mobile ad hoc networks. [Research Report] RR-5173, INRIA. 2004. inria-00071415

HAL Id: inria-00071415
https://inria.hal.science/inria-00071415

Submitted on 23 May 2006

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

INRIA

INSTITUT NATIONAL DE RECHERCHE EN INFORMATIQUE ET EN AUTOMATIQUE

Analytical results on Connected dominating sets in mobile ad hoc networks

Philippe Jacquet

N° 5173

THÈME 1



ISSN 0249-6399 ISRN INRIA/RR--5173--FR+ENG

INRIA
ROCQUENCOURT

Analytical results on Connected dominating sets in mobile ad hoc networks

Philippe Jacquet *

Thème 1 — Réseaux et systèmes
Projet Hipercom

Rapport de recherche n° 5173 — — 6 pages

Abstract: We provide analytical results about the performance of various Connected Dominating Set (CDS) algorithms: MultiPoint Relaying (MPR) flooding, MPR-CDS, Generalized Wu Li CDS (GWL-CDS). In particular we focuse on the 1D unit disk graph model.

Key-words: Mobile ad hoc, connected dominating set, unit disk model, asymptotic results.

* (philippe.jacquet@inria.fr)

Unité de recherche INRIA Rocquencourt
Domaine de Voluceau, Rocquencourt, BP 105, 78153 Le Chesnay Cedex (France)
Téléphone : +33 1 39 63 55 11 — Télécopie : +33 1 39 63 53 30

Résultats analytiques sur les ensembles dominants connectés dans les réseaux mobiles ad hoc

Résumé : Nous présentons des résultats analytiques sur les performances de différents algorithmes sur les ensemble dominants connectés (CDS): innondation par Relais MultiPoints (MPR), MPR-CDS, algorithmes Wu et Li généralisé (GWL-CDS). En particulier nous nous intéressons au modèle du graphe avec disque unité en dimension 1.

Mots-clés : Mobile ad hoc, ensemble dominants connectés, modèle du disque unité, résultats asymptotiques.

1 Introduction

In [2, 3, 1] we describe many connected dominating set built from Multipoint Relay method (MPR). In [4] there is a description of Genaralized Wu and Li algorithm for oriented graph that we simplified in [5] for the context of non oriented graphs. In this note we analyze the performance of these algorithm via analytical models. We investigate the case where the network is a unit disk graph of dimension 1. We assume that the nodes are randomly and uniformly dispatched on an infinite segment. The radio range is 1 and the density of nodes is $\nu/2$. This implies that the average node degree is ν.

In particular we will prove that the densities of MPR connecting sets tend to constant densities when $\nu \to \infty$. The density of CDS nodes of generalized Wu Li algorithm also tends to a constant when selection criterium is based on ID, but tends to be of order $\sqrt{\nu}$ when the criterium is the node degree. Therefore this confirms the poor performance of generalized Wu and Li algorithm with degree criterium.

2 MPR connected dominating sets

There are two kinds of MPR connected dominating set: the set obtained by the transmitter in an MPR flooding and the set obtained by the specific algorithm described in [3], called MPR-CDS. The former has the advantage to enable less transmitters than the latter but with the drawback that the connected dominating set is source dependent (*i.e.* the connected dominating set varied when the source of the initial broadcast varies). In fact the MPR flooding is last hop dependent.

Theorem 1 *The probability that a node belongs to the MPR flooding set tends to be equivalent to* $\frac{2}{\nu}$.

The proof is in [2, 1].

Theorem 2 *The probability that a node belongs to the MPR CDS set tends to be equivalent to* $\frac{3}{\nu}$ *when* $\nu \to \infty$.

It has been proven in [2] that a node has 2 MPR which are the extremal points of its coverage interval. Conversely the number of MPR selector neighbors is 2. A node belongs to the MPR CDS (i) if it has the shortest ID in its neighborhood, which occurs with probability $\frac{1}{\nu}$, or (ii) if one of its MPR selector has the shortest ID in the neighborhood, which occurs with probability $\frac{2}{\nu}$. These two events being exclusive the probabilities sum.

3 Generalized Wu Li algorithms

By ID criterium we assume that a node belongs to the CDS when the neighbors with higher ID don't form a connected component that covers the whole node neighborhood.

Theorem 3 *The probability that a node belongs to the generalized Wu and Li CDS with ID criterium tends to be equal to* $\frac{4}{\nu}$ *when* $\nu \to \infty$

In fact rule k is equivalent to rule 2 in dimension 1. It has been shown in [2] that the average number of CDS members in a segment of length x is exactly $2x-1$.

By degree criterium we assume that a node belongs to the CDS when the neighbors with higher degree don't form a connected component that covers the whole node neighborhood.

Theorem 4 *The probability that a node belongs to the generalized Wu and Li CDS with degree criterium tends to be equal to* $\sqrt{8}\pi\nu$ *when* $\nu \to \infty$.

Let $N(x)$ be the number of neighbor of a node at location x on the segment map. Let $I([a,b])$ be the number of nodes contained by interval $[a,b]$. Therefore $N(x) = I([x-1, x+1])$. Let $\Delta(x) = N(x) - N(0)$. We need the following lemma

Lemma 1 *For any* $x \in [-1,0]$ *and* $y \in [0,1]$ $\Delta(x)$ *and* $\Delta(y)$ *are independent variables.*

Proof: If $x \in [0,1]$, then we have $\Delta(x) = I([-1,-1+x]) - I([1,1+x])$. If $x \in [-1,0]$ then $\Delta(x) = I([-1+x,-1]) - I([1+x,1])$. Since the intervals don't overlap then $\Delta(x)$ and $\Delta(y)$ are independent when x and y have different signs.

Let R_r be the absolute value of the first $x \in [0,1]$ such that $\Delta(x) > 0$ ($\forall t \in [0,x] \Delta(x) \leq 0$). Let R_l be the absolute value position of the first $y \in [-1,0]$ such that $\Delta(x) > 0$ ($\forall t \in [y,0] \Delta(x) \leq 0$). We know that R_r and R_l are independent. A node at position 0 belongs to the CDS iff $R_r + R_l > 1$

Lemma 2 *Let* $x \in [0,1]$ *the probability* $P(x)$ *such that* $R_r = x$ *has Laplace transform*

$$\int_0^\infty P(x) e^{-\omega x} dx = \frac{2 + \frac{2\omega}{\nu} + \sqrt{(2+\frac{2\omega}{\nu})^2 - 4}}{2}$$

Having $\Delta(t) \leq 0$ for all $t \in [0,x]$ is equivalent that an M/M/1 system with service rate and arrival rate equal to $\frac{\nu}{2}$ starts with one customer and does not empty its queue during a time interval of length x. Let $f(\omega)$ be the Laplace transform of the distribution of the time T needed to empty the queue $f(\omega) = E[e^{-\omega T}]$. Let θ be the time needed for the exit of the first customer, we have from classic queueing theory:

$$T = \theta + N_\theta \times T \tag{1}$$

where N_θ is a Poisson random variable of mean θ and $N \times T$ means the addition of N independent copies of T (N i.i.d. variables distributed as T). Therefore

$$f(\omega) = \int_0^\infty P(\theta = x) e^{-x\omega} e^{x\frac{\nu}{2}(f(\omega)-1)} dx \tag{2}$$

$$= \frac{\nu/2}{\nu/2(1-f(\omega)) + \omega} \tag{3}$$

Consequently the Laplace transform of the distribution of $R_r + R_l$ is $f(\omega)^2$. Therefore the probability $P(\nu)$ that node at position 0 belongs to the CDS, that is $R_l + R_r > 1$, satisfies

$$P(\nu) = \frac{1}{2i\pi} \int_{-i\infty}^{i\infty} \frac{1 - (\omega + 1 + \sqrt{(\omega+1)^2 - 1})^2}{w} e^{\nu\omega} d\omega$$

Using the fact that $1 - (\omega + 1 + \sqrt{(\omega+1)^2 - 1})^2 \sim 2\sqrt{\omega} + O(\omega)$ when $\omega \to 0$ we have from Flajolet and Odlyzko theorem:

$$\frac{1}{2i\pi} \int_{-i\infty}^{i\infty} \frac{1 - (\omega + 1 + \sqrt{(\omega+1)^2 - 1})^2}{w} e^{\nu\omega} d\omega \sim \frac{2\Gamma(1/2)}{\pi} \nu^{1/2}.$$

We display in figure 1 the various node densities of the connected dominating set studied in this note.

References

[1] P. Jacquet, A. Laouiti, P. Minet, L. Viennot, "Performance of multipoint relaying in mobile ad hoc networks," Networking 2002, Pisa, 2002.

[2] P. Jacquet, A. Laouiti, P. Minet, L. Viennot, " Performance analysis of OLSR multipoint relaying in two ad hoc wireless network models, " INRIA Research Repport RR-4260, 2001.

[3] C. Adjih, P. Jacquet, L. Viennot, " Computing dominating set with multipoint relays, INRIA RR-4597, 2002.

[4] J. Wu, and F. Dai, "A generic distributed broadcast schem in ad hoc networks," in *ICDCS*, 2003

[5] P. Jacquet, "Performance of connected dominating set in OLSR protocol," INRIA Research Repport RR-5098, 2004.




Figure 1: Node density of various CDS algorithm from bottom to top: MPR flooding, MPR-CDS, GWL-CDS with ID, GWL-CDS with degree.



Unité de recherche INRIA Rocquencourt
Domaine de Voluceau - Rocquencourt - BP 105 - 78153 Le Chesnay Cedex (France)

Unité de recherche INRIA Lorraine : LORIA, Technopôle de Nancy-Brabois - Campus scientifique
615, rue du Jardin Botanique - BP 101 - 54602 Villers-lès-Nancy Cedex (France)
Unité de recherche INRIA Rennes : IRISA, Campus universitaire de Beaulieu - 35042 Rennes Cedex (France)
Unité de recherche INRIA Rhône-Alpes : 655, avenue de l'Europe - 38330 Montbonnot-St-Martin (France)
Unité de recherche INRIA Sophia Antipolis : 2004, route des Lucioles - BP 93 - 06902 Sophia Antipolis Cedex (France)

Éditeur
INRIA - Domaine de Voluceau - Rocquencourt, BP 105 - 78153 Le Chesnay Cedex (France)
http://www.inria.fr
ISSN 0249-6399