

## On complexity and motion planning for co-rank one sub-Riemannian metrics

C. Romero-Meléndez, J.P. Gauthier, Felipe Monroy-Pérez

► **To cite this version:**

C. Romero-Meléndez, J.P. Gauthier, Felipe Monroy-Pérez. On complexity and motion planning for co-rank one sub-Riemannian metrics. RR-4882, INRIA. 2003. inria-00071701

**HAL Id: inria-00071701**

**<https://hal.inria.fr/inria-00071701>**

Submitted on 23 May 2006

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

*On complexity and motion planning for co-rank one  
sub-Riemannian metrics*

Romero-Meléndez C. — Gauthier J.P. — Felipe Monroy-Pérez

**N° 4882**

Juillet 2003

THÈME 4



*Rapport  
de recherche*



## On complexity and motion planning for co-rank one sub-Riemannian metrics

Romero-Meléndez C. <sup>\*</sup>, Gauthier J.P. <sup>†</sup>, Felipe Monroy-Pérez<sup>‡</sup>

Thème 4 — Simulation et optimisation  
de systèmes complexes  
Projet MIAOU

Rapport de recherche n° 4882 — Juillet 2003 — 30 pages

**Abstract:** In this paper, we study the *motion planning problem* for generic sub-Riemannian metrics of co-rank one. We give explicit expressions for the metric complexity (in the sense of F. Jean [7],[8]), in terms of the elementary invariants of the problem. We construct asymptotic optimal syntheses, it turns out that among the results we show, the most complicated case is the 3-dimensional. Besides the generic  $C^\infty$  case, we study some non-generic generalizations in the analytic case.

**Key-words:** motion planning problem, metric complexity, normal forms, asymptotic optimal synthesis

<sup>\*</sup> Laboratoire d'Analyse Appliquée et Optimisation, Département de Mathématiques, Université de Bourgogne, 21078 Dijon, France.

<sup>†</sup> Laboratoire d'Analyse Appliquée et Optimisation, Département de Mathématiques, Université de Bourgogne, 21078 Dijon, France.

<sup>‡</sup> On sabbatical leave from the Basic Sciences Department of the Universidad Autonoma Metropolitana Azcapotzalco. Financially supported by the CONACYT, México.

## Sur la complexité et la planification de trajectoires pour métriques sous-Riemanniennes de co-rang un

**Résumé :** Dans cet article, nous étudions le *problème de planification de trajectoires* pour une métrique sous-Riemannienne générique de co-rang un. Nous donnons des expressions explicites pour la complexité métrique (au sens de F. Jean, [7],[8]), en terme des invariants élémentaires du problème. Nous construisons aussi la synthèse optimale asymptotique. Il se trouve que, parmi les résultats énoncés, le cas le plus compliqué est celui de dimension 3. Les résultats génériques sont obtenus dans la catégorie  $C^\infty$ , mais nous étudions aussi quelques généralisations non-génériques dans la catégorie analytique.

**Mots-clés :** problème de planification de trajectoires, complexité métrique, forme normale, synthèse optimale asymptotique

# 1 Introduction, notations and organization of the paper

## 1.1 Definitions

A *motion planning problem* is a triple  $\Sigma = (\Delta, g, \Gamma)$ , where  $(\Delta, g)$  is a sub-Riemannian metric over  $\Xi_N$ , a  $N$ -dimensional manifold,  $N \geq 3$ , and  $\Gamma : [0, 1] \rightarrow \Xi_N$  is a smooth parameterized curve (although there is no Riemannian structure assumed over  $\Xi_N$ , the curve  $\Gamma$  has a length parameter). As usual,  $\Delta$  is a distribution over  $\Xi_N$  and  $g$  is a Riemannian metric over  $\Delta$ . Everything is smooth, which means at least  $C^\infty$  and for technical reasons  $C^\omega$  when explicitly stated.

We shall consider co-rank 1 distributions only, that is, the dimension of  $\Delta_x$  is  $N - 1$  for all  $x \in \Xi_N$ . A convenient way to specify such a sub-Riemannian metric is to define it by means of  $N - 1$  vector fields over  $\Xi_N$  that are everywhere independent, generate the distribution  $\Delta$ , and provide at each point an orthonormal frame for  $g$ . It is not always possible to define globally a sub-Riemannian metric in this way, but we shall limit ourselves to this case. Hence in this paper a sub-Riemannian metric is a couple  $\Sigma = (F, \Gamma)$ , where  $F = (X_1, \dots, X_{N-1})$  is a frame of smooth vector fields everywhere defined.

We shall consider generic motion planning problems. Since  $\Gamma([0, 1]) \subset \Xi_n$  is a compact set, and since the problem is only local around the curve, the behavior of  $F$  at infinity is irrelevant. We shall denote as  $S^\infty$  ( $S^\omega$ ) the set of all *motion planning problems*, the topology over  $S^\infty$  ( $S^\omega$ ) is the  $C^\infty$  topology (not Whitney).

We assume that the curve  $\Gamma$  is well-parameterized, that is,  $\frac{d\Gamma}{dt}(t) \neq 0$  for all  $t \in [0, 1]$ . Since  $N \geq 3$ , it is clear that the set of motion planning problems, with  $\Delta$  *non integrable* (otherwise it does not define a sub-Riemannian metric), and with  $\Gamma$  having no double points, is an open-dense set in  $S^\infty$  ( $S^\omega$ ). We limit ourselves to this case, and we denote again by  $S^\infty$  ( $S^\omega$ ) the set of such problems.

Let  $d$  denote the sub-Riemannian distance. Following F. Jean's definition, see [7] and [8], we chose a small real parameter  $\varepsilon \geq 0$  and denote by  $T_\varepsilon$  the sub-Riemannian  $\varepsilon$ -tube

$$T_\varepsilon = \{x \in \Xi_N \mid d(x, \Gamma) \leq \varepsilon\},$$

we define then the *metric complexity*  $MC_\Sigma(\varepsilon)$  of  $\Sigma$  as follows:

$$MC_\Sigma(\varepsilon) = \frac{1}{\varepsilon} \min_\gamma \{\text{SR-length}(\gamma)\},$$

the minimum being taken among all (smooth) admissible (tangent to  $\Delta$ ) curves  $\gamma : [0, T_\gamma] \rightarrow \Xi_N$ , entirely contained in the tube  $T_\varepsilon$  and joining  $\Gamma(0)$  to  $\Gamma(1)$ , that is,  $\gamma(0) = \Gamma(0)$  and  $\gamma(T_\gamma) = \Gamma(1)$ . For our purposes, it would be equivalent to consider continuous, piecewise-smooth, or even absolutely continuous curves  $\gamma$ .

We are only interested with the asymptotic behavior of  $MC_\Sigma(\varepsilon)$  when  $\varepsilon$  tends to zero. In all the situations we consider,  $MC_\Sigma(\varepsilon)$  has an *equivalent* when  $\varepsilon$  tends to zero, we shall take both,  $MC_\Sigma(\varepsilon)$  and its equivalent indistinctly. In general, two functions  $\varphi_1(\varepsilon)$  and  $\varphi_2(\varepsilon)$  tending to  $\infty$  when  $\varepsilon \rightarrow 0$  are said to be *weakly equivalent* (denoted  $\varphi_1 \stackrel{w}{\sim} \varphi_2$ ), if

$$\frac{\varphi_2(\varepsilon)}{\varphi_1(\varepsilon)} \quad \text{and} \quad \frac{\varphi_1(\varepsilon)}{\varphi_2(\varepsilon)},$$

are bounded when  $\varepsilon \rightarrow 0$ . They said to be *strongly equivalent* (denoted  $\varphi_1 \stackrel{s}{\simeq} \varphi_2$ ), if

$$\lim_{\varepsilon \rightarrow 0} \frac{\varphi_1(\varepsilon)}{\varphi_2(\varepsilon)} = 1.$$

Contrarily to the results of F. Jean [7], [8] and [9], most of our expressions for the metric complexity in this paper, are in the sense of strong equivalence.

## 1.2 The fundamental 2-form.

Let  $\Sigma = (F, \Gamma)$  be given, we assume that  $\Gamma(t)$  is transversal to  $\Delta(\Gamma(t))$  for all  $t \in [0, 1]$ . In this case, we say that  $\Sigma$  is *relevant*. Let  $S_R^\infty$  and  $S_R^\omega$  denote the corresponding sets of relevant systems. They form open subsets of  $S^\infty$  and  $S^\omega$  respectively.

Now, let  $\omega$  be a one-form such that  $\Delta = \ker \omega$ , we can choose it in such a way that  $\omega(\frac{d\Gamma}{dt}(t)) = 1$ , for all  $t \in [0, 1]$ . This one-form  $\omega$  is defined modulo multiplication by a smooth function  $f$  that takes value 1 on  $\Gamma$ .

We take  $\alpha = d\omega$  and consider  $\tilde{\alpha} = \alpha|_\Delta$ . The map  $t \mapsto \tilde{\alpha}(\Gamma(t))$  determines along  $\Gamma$  a field 2-forms on  $\Delta$ . This field is well defined and it is independent of  $f$ . In fact, since

$$\tilde{\alpha} = d(f\omega)|_\Delta,$$

then for all  $G, H$  sections of  $\Delta$ , we have

$$\tilde{\alpha}(G, H)(\Gamma(t)) = d(f\omega)(G, H) = df \wedge \omega(G, H) + fd\omega(G, H) = 0 + fd\omega(G, H)$$

but  $f(\Gamma(t)) = 1$ , then  $\tilde{\alpha}(\Gamma(t))$  is independent of  $f$ .

**Definition 1.1** We call  $\tilde{\alpha}$  the *fundamental 2-form associated to  $\Sigma$* .

The form  $\tilde{\alpha}$  defines a skew symmetric (with respect to  $g$ ) linear map  $A(t) : \Delta(\Gamma(t)) \longrightarrow \Delta(\Gamma(t))$ , in the following way:

$$g(A(t)X, Y) = \tilde{\alpha}(X, Y), \quad \text{for } X \text{ and } Y \text{ in } \Delta(\Gamma(t)).$$

Then, once an orthonormal frame  $F_t$  is given in  $\Delta(\Gamma(t))$ , it defines an element of  $\mathfrak{so}(N-1)$  given by the matrix of  $A(t)$  in the basis  $F_t$ . This matrix shall be typically denoted by  $\tilde{A}(t)$ .

### 1.3 Statement of our results and comments

Let  $\kappa(t) = \sup\{|\text{eigenvalues of } A(t)|\}$ . Let us denote by  $MC_\Sigma(\varepsilon, T)$  the metric complexity of the curve  $\Gamma : [0, T] \rightarrow \Xi_N$ , for  $T < 1$ . The following theorems are our main results for the dimensions  $N \geq 4$  and  $N = 3$ .

**Theorem 1.2** ( $N \geq 4$ ) *There is an open-dense subset of  $S_R^\infty$  (and also of  $S^\infty$ , but in that case  $\kappa(t)$  is well defined except on a finite subset of  $[0, 1]$ ), such that*

$$MC_\Sigma(\varepsilon, T) \underset{s}{\simeq} \frac{2}{\varepsilon^2} \int_0^T \frac{dt}{\kappa(t)}. \quad (1)$$

*This integral is well defined and finite. Moreover, the function  $T \mapsto \int_0^T \frac{dt}{\kappa(t)}$  is smooth, for  $\Sigma \in S_R^\infty$ .*

**Theorem 1.3** ( $N = 3$ ) *There is an open-dense subset of  $S_R^\infty$  (and also of  $S^\infty$ ), such that*

$$MC_\Sigma(\varepsilon, T) \geq -\frac{\log \varepsilon}{\varepsilon^2} \sum_{i=1}^M C(t_i), \quad (2)$$

*where  $t_i \in [0, T]$ ,  $\Gamma(t_i)$  is a Martinet point of  $\Delta$ , (i.e.  $\Gamma(t_i)$  is a point at which  $\Delta$  has a Martinet singularity), and  $C(t_i)$  are constants. If there are no Martinet points, then the formula for  $MC_\Sigma(\varepsilon, T)$  is still the one in Theorem 1.2.*

In this paper we do not compute explicitly the constants  $C(t_i)$  in the general case. We provide an example that shows equality in (2), and for the example, we exhibit only rough estimations for the constants  $C(t_i)$ .

In this paper we show also that in the *analytic case*, the asymptotics (1) is always valid, except in very degenerate cases. For the analytic case under the assumption that the fundamental form  $\tilde{\alpha}(t)$  never vanishes, we prove the following theorem:

**Theorem 1.4** *If  $\tilde{\alpha}$  is never zero, and  $\Sigma$  is analytic,  $\Sigma \in S_R^\omega$  (or  $S^\omega$ ), then the asymptotics (1) of Theorem 1.2 for the metric complexity is still valid. But, in that case, the map  $T \mapsto \int_0^T \frac{dt}{\kappa(t)}$  is only  $C^1$ , piecewise analytic. Its second derivative has jumps at isolated points.*

**Definition 1.5** *A weak (strong) asymptotic optimal synthesis is a control strategy, depending on  $\varepsilon$ , that realizes a weak (strong) equivalent of the metric complexity, when  $\varepsilon \rightarrow 0$ , i.e., it is a family  $\{\gamma_\varepsilon\}$ , of admissible curves,  $\gamma_\varepsilon([0, \theta_\varepsilon]) \subset T_\varepsilon$  satisfying  $\gamma_\varepsilon(0) = \Gamma(0)$  and  $\gamma_\varepsilon(\theta_\varepsilon) = \Gamma(1)$ , such that,*

$$\frac{1}{\varepsilon} (\text{SR-length}(\gamma_\varepsilon)) \simeq MC_\Sigma(\varepsilon).$$



In this paper, we will describe explicit asymptotic optimal syntheses (strong), in the case of the Theorems 1.2, 1.4 above. We will also show a (weak) explicit optimal synthesis in the Martinet case.

**Remark 1.6 Non-relevant motion planning problems.** *Generically, for  $\Sigma \in S^\infty$  or  $S^\omega$ ,  $\Delta$  is tangent to  $\Gamma(t)$  at isolated points. Around these points, since the curve  $\Gamma$  is almost admissible, asymptotic optimal synthesis is rather trivial. In fact, if  $t_0$  is such a point on  $\Gamma$  we have that  $\lim_{t \rightarrow 0} \kappa(t) = +\infty$ , the integral in (1) is well defined and still yields the metric complexity.*

Our results have to be compared to those of F. Jean in papers [7] and [8], on which the asymptotics  $\frac{1}{\varepsilon^2}$  and  $-\frac{\log \varepsilon}{\varepsilon}$  already appear, however, with his technique, he only obtains weak equivalence. Also, in his papers, there is no explicit construction of an asymptotic optimal synthesis. On the other hand, the weakness of our paper with respect to F. Jean's results is the fact that we consider distributions of codimension one only.

## 1.4 Organization of the paper.

Apart from this introduction the paper contains five sections and one appendix. In section 2, we state and prove some genericity results about co-rank 1 sub-Riemannian metrics and motion-planning problems.

In Section 3, we present normal forms and normal coordinates for motion-planning problems. These tools are the key points in our treatment. The idea of normal coordinates and the resulting normal forms were introduced for the 3-dimensional case in the papers [4] and [5], the general *contact case* is discussed in [6]. For the *quasi-contact*, they were introduced in [10]. Here, we generalize these normal forms and normal coordinates to generic motion-planning problems, and also to degenerate ones, in the analytic case.

The normal coordinates are the sub-Riemannian analogs of the classical normal coordinates in Riemannian geometry. Normal forms in the Riemannian case, similar to those presented in this paper, were already well known by Darboux.

In Section 4, we use these normal forms for  $N \geq 4$  (or  $N = 3$ , but contact) to construct in a rather natural way an *asymptotic optimal synthesis*, then we obtain the asymptotics (1) for the metric complexity of Theorem 1.2 and Theorem 1.3.

Section 5 is devoted to the Martinet Case. We prove Theorem 1.3, and, for an example, we show in a constructive way, the equality in the asymptotics (2), i.e., we exhibit an asymptotic optimal synthesis (but weak).

In the Section 6, we consider non generic situations, but analytic. We show that our normal forms of Section 3, and formula (1) for the metric complexity still hold, as long as the fundamental 2-form does not vanish along  $\Gamma$ . This is Theorem 1.4. In fact, the results are the same, the only difference is that the metric complexity becomes non smooth: it has jumps on the second derivative. The strong asymptotic optimal synthesis turns out to be slightly more complicated.

In the Appendix (Section 7), we collect a few technical facts that are needed along the paper. Results similar to Lemma (7.1) can be found in the literature. But for the sake of completeness we provide a simple proof.

## 2 Genericity results

We consider the set  $S_R^\infty$  of *relevant motion-planning problems*  $\Sigma = (F, \Gamma)$  with the  $C^\infty$  topology, we consider also the field  $\tilde{\alpha}_\Sigma$  of fundamental 2-forms along  $\Gamma$  defined by  $\Sigma$ . As explained above, in the coordinates on  $\Delta$  defined by the frame  $F$ , to  $\tilde{\alpha}_\Sigma(\Gamma(t))$  corresponds a skew symmetric matrix  $\bar{A}(t)$ .

As customary,  $\mathfrak{so}(N-1)$  denotes the Lie algebra of square  $(N-1) \times (N-1)$  skew symmetric matrices, we have the following result for relevant motion-planning problems.

**Theorem 2.1** *The mapping  $S_R^\infty \times [0, 1] \longrightarrow \mathfrak{so}(N-1)$ ,  $(\Sigma, t) \longmapsto \bar{A}(t)$  is a submersion.*

**Proof.** Let us consider a fixed  $\Sigma = (F, \Gamma) \in S_R^\infty$ , and an arbitrary  $t_0 \in [0, 1]$ . It is always possible to choose a neighborhood  $V_{t_0}$  of  $\Gamma(t_0)$ , and coordinates  $(x, w)$  on  $V_{t_0}$  such that:

$$\Gamma(t) = (0, t - t_0), \quad (3)$$

$$\Delta(0, w) = \ker dw. \quad (4)$$

The frame  $F = (F_1, \dots, F_{N-1})$  with

$$F_j = \sum_{i=1}^n Q_{ij} \frac{\partial}{\partial x_i} + L_j \frac{\partial}{\partial w},$$

writes in abbreviated notation as follows

$$F = Q \frac{\partial}{\partial x} + L \frac{\partial}{\partial w},$$

with  $L(0, w) = 0$ .

The matrix  $Q(x, w)$  is invertible, because  $\Gamma$  is transversal to  $\Delta$  and  $F$  is an orthonormal frame. Let  $Q_{ij}^{-1}$  denote the  $ij$  entry of the inverse matrix  $Q^{-1}$ .

Let  $\Omega = (\omega_1, \dots, \omega_N)$  be the coframe dual to the frame  $(F_1, \dots, F_{N-1}, \frac{\partial}{\partial w})$ . Then, by definition we have

$$d\omega_N|_{\Delta(\Gamma(t))} = \tilde{\alpha}(\Gamma(t)).$$

Now, for each  $(x, w)$  we have that  $Q(x, w)$  is a  $(N-1) \times (N-1)$  matrix, and also that  $L(x, w) = (L_1(x, w), \dots, L_{N-1}(x, w))$  is a linear form. Furthermore,  $\Omega = (\tilde{\Omega}dx, \tilde{L}dx + dw)$ , with

$$\begin{aligned}\tilde{\Omega} &= Q^{-1}(x, w), \\ \omega_N &= \tilde{L}dx + dw = dw - LQ^{-1}dx.\end{aligned}\tag{5}$$

Therefore

$$d\omega_N(F_l, F_s) = \sum_{i,j,k=1}^{N-1} \frac{\partial}{\partial x_k} (L_i Q_{ij}^{-1}) (Q_{lj} Q_{sk} - Q_{sj} Q_{lk}) + \sum_{i,j=1}^{N-1} \frac{\partial}{\partial w} (L_i Q_{ij}^{-1}) (L_s Q_{lj} - L_i Q_{sj}),$$

and since  $L(0, w) = 0$ , we have:

$$d\omega_N(F_l, F_s)|_{\Gamma} = \sum_{i,j,k=1}^{N-1} \frac{\partial L_i}{\partial x_k} Q_{i,j}^{-1} (Q_{lj} Q_{sk} - Q_{sj} Q_{lk}).\tag{6}$$

Let us fix  $Q$ , and leave  $L$  free. By definition we know that

$$\bar{A}_{sl}(t_0) = d\omega_N(F_l, F_s)(\Gamma(t_0)).$$

Let us consider the  $(N-1) \times (N-1)$  matrices  $\bar{L}$  and  $\tilde{L}$ , the entries of which are given as follows

$$\begin{aligned}\bar{L}_{jk} &= \sum_{i=1}^{N-1} \frac{\partial L_i}{\partial x_k} Q_{i,j}^{-1}(\Gamma(t_0)), \\ \tilde{L}_{ik} &= \frac{\partial L_i}{\partial x_k}(\Gamma(t_0)).\end{aligned}$$

Then  $\bar{L} = (Q^{-1})' \tilde{L}(\Gamma(t_0))$ , and (6) can be rewritten:

$$\bar{A}(t_0) = Q \bar{L} Q' - Q \tilde{L}' Q' = Q(\bar{L} - \tilde{L}') Q',\tag{7}$$

where  $'$  denotes the corresponding transpose matrix. Then,  $\bar{A}(t_0) = 0$  implies  $\bar{L} = \tilde{L}'$ , which means that  $\bar{L}$  is symmetric, and that  $\tilde{L} = Q'(\Gamma(t_0))S$ , where  $S$  is any symmetric matrix.

Therefore, the kernel of the linear map

$$\mathcal{M}_{N-1 \times N-1} \longrightarrow \mathfrak{so}(N-1), \quad \tilde{L} \longmapsto \bar{A}(t_0),$$

has dimension  $\frac{1}{2}(N-1)N$ . Hence, its image has dimension  $\frac{1}{2}(N-1)(N-2)$ , and in consequence it is surjective. This implies the theorem.  $\diamond$

Now, let us consider the following map:

$$\rho : S_{\mathbb{R}}^{\infty} \times [0, 1] \longrightarrow \mathfrak{so}(N-1), \quad (\Sigma = (F, \Gamma), t) \longmapsto \bar{A}(t).$$

As a consequence of Theorem 2.1 we have that  $\rho$  is a submersion. We apply now standard transversality techniques, (see for instance [1], [2], and [3]), to the mapping  $\rho$ , and to the closed stratified subsets  $\mathcal{A}_1, \mathcal{A}_2, \mathcal{A}_3, \bar{\mathcal{A}}_4$  of  $\mathfrak{so}(N-1)$  which are defined in the Appendix 7, Section 7.2. There is an open-dense subset  $S_*^\infty \subset S_R^\infty$  such that, for any  $\Sigma \in S_*^\infty$  the mapping

$$ev_\rho(\Sigma) : [0, 1] \longrightarrow \mathfrak{so}(N-1), \quad t \longmapsto \rho(\Sigma, t),$$

is transversal to the aforementioned closed stratified subsets of  $\mathfrak{so}(N-1)$ . It is enough to evaluate the codimension of these sets to conclude that for  $\Sigma \in S_*^\infty$  one has that

1. For all  $t \in [0, 1]$ ,  $\bar{A}(t)$  is not an element of  $\bar{\mathcal{A}}_4 \cup \mathcal{A}_2 \cup \mathcal{A}_3$ , and
2. The set  $\{t \in [0, 1] \mid \bar{A}(t) \in \mathcal{A}_1\}$  is finite.

In conclusion we have the following result:

**Theorem 2.2** *There is an open-dense subset  $S_*^\infty \subset S_R^\infty$ , such that for all  $\Sigma \in S_*^\infty$ , the mapping  $t \mapsto A(t)$  has the following properties. For all  $t \in [0, 1]$ :*

1.  $A(t)$  has no double nonzero eigenvalues.
2. The kernel of  $A(t)$  has dimension 1 if  $N = 2n$ .
3. The kernel of  $A(t)$  has dimension 2 on a (may be empty) finite subset of  $[0, 1]$ , and it is zero elsewhere if  $N = 2n + 1$ .

In particular, and this will be a key point in the next sections, for  $\Sigma \in S_*^\infty$  and all  $t \in [0, 1]$ , if  $N \geq 4$  then  $A(t)$  has a nonzero eigenvalue.

### 3 Normal Coordinates and Normal Forms

The results in this section are very close to the results in the papers [4], [5] and [6]. We refer the reader to these references, and limit ourselves to explain in detail the points that are different.

#### 3.1 Normal Coordinates

**Theorem 3.1** *Let  $\Sigma = (F, \Gamma) \in S_R^\infty$ . Then there is a global coordinate system  $(\xi, w) \in \mathbb{R}^{N-1} \times \mathbb{R}$ , defined on a neighborhood of  $\Gamma([0, 1])$ , such that:*

1.  $\Gamma(t) = (0, t)$ ,  $\Delta(\Gamma(t)) = \ker dw$  and  $g|_{\Gamma(t)} = \sum_{i=1}^{N-1} d\xi_i^2$ .

2. Geodesics satisfying the Pontryagin Maximum Principle's transversality conditions with respect to  $\Gamma$  are the straight lines through  $\Gamma$  contained in the planes  $S_{w_0} = \{(\xi, w) \mid w = w_0\}$ .
3. For  $\varepsilon$  small enough, the sub-Riemannian cylinder  $C_\varepsilon = \{q \mid d(q, \Gamma) = \varepsilon\}$  is the Riemannian cylinder  $\{\|\xi\|_2 = \varepsilon\}$ , (here  $d$  denotes the sub-Riemannian distance associated to  $\Sigma$ ).

The coordinate system in the Theorem 3.1 is usually called a *pre-normal coordinate system*. The proof of this theorem is included in the aforementioned references as well as in [10]. The paper [4] discusses dimension 3, but in formal power series, whereas [5] does it in the smooth case. Reference [6] studies the contact case in any odd dimension whereas reference [10] treats the quasi-contact in any even dimension. In fact, there is no need for the contact or quasi-contact assumptions, since the only crucial point in those cases is that the curve  $\Gamma$  has to be everywhere transversal to the distribution  $\Delta$ , which is, by definition, the case in  $S_R^\infty$ .

Now, in a pre-normal coordinate system  $(\xi, w)$  along the curve  $\Gamma$ , the vector fields

$$\frac{\partial}{\partial \xi_1}, \dots, \frac{\partial}{\partial \xi_{N-1}}, \quad (8)$$

provide an orthonormal frame on  $\Delta(\Gamma(t))$ . Let  $\bar{A}(t)$  be the matrix of the linear map  $A(t)$  with respect to this frame.

**Theorem 3.2 (Normal Coordinates)** *Let  $\Sigma = (F, \Gamma) \in S_*^\infty$ . Then, there is a global coordinate system  $(\xi, w)$ , defined on a neighborhood of  $\Gamma([0, 1])$ , such that:*

1.  $(\xi, w)$  is a pre-normal coordinate system.
2. In the coordinates  $\xi$ , the matrices  $\bar{A}(t)$ ,  $t \in [0, 1]$  are (skew-symmetric)  $2 \times 2$  block-diagonal. (plus a  $1 \times 1$  block if  $N = 2n$ ).

**Proof.** By Lemma 7.1, Appendix 7, we can smoothly block-diagonalize the matrix  $\bar{A}(t)$  along the curve  $\Gamma$ , using a smooth orthogonal transformation  $U(t)$ . Now, setting  $(\tilde{\xi}, w) = (U(w)\xi, w)$  we obtain the desired coordinates, because such a change of coordinates does not affect the properties of pre-normal coordinates in Theorem 3.1.  $\diamond$

Normal coordinates are unique up to a change of coordinates of the form:

$$(\tilde{\xi}, w) = (T(w)\xi, w), \quad (9)$$

where  $T(w)$  is a smooth curve in the natural maximal-torus of the orthogonal group  $\text{SO}(N-1)$ .

In the contact and quasi-contact cases, we can do even more, because in those cases we can consider, in normal coordinates, the planes  $S_{w_0} = \{(\xi, w) \mid w = w_0\}$ . Since the curve  $\Gamma$  is transversal to  $\Delta$ , then the sub-Riemannian metric  $g(\xi, w)$  projects on a Riemannian metric  $g_w$  on the plane  $S_w$ . In fact, if  $\Pi_{w_0}$  denotes the projection

$$\Pi_{w_0} : \Xi_N \rightarrow S_{w_0}, \quad (\xi, w) \mapsto (\xi, w_0),$$

we set

$$g_{w_0}(X, Y) = g((d\Pi_{w_0}|_{\Delta})^{-1}X, (d\Pi_{w_0}|_{\Delta})^{-1}Y), \quad \text{for } X, Y \in T_{(\xi, w_0)}S_{w_0}.$$

**Theorem 3.3** *If  $\Delta$  is contact or quasi-contact, we can chose  $T(w)$  in such a way that for all points  $(0, w)$  on the curve  $\Gamma$ , the sectional curvatures of the metrics  $g_w$ , with respect to the 2-dimensional real eigenspaces of  $A(w)$  are zero.*

This theorem is proved in [5] (3-dimensional case), [6] (general contact case), and [10] (quasi-contact case). These normal coordinates are again unique, up to an element  $T(w_0)$  of a maximal torus of  $\text{SO}(N - 1)$ . (Once the  $T(w_0)$  in (9) is fixed for some  $w_0$ , then, it determines  $T(w)$  for all  $w$ , for the sectional curvatures being zero).

### 3.2 Normal Forms

Assume that  $\Sigma = (F, \Gamma)$  is given, we write the normal coordinates  $(\xi, w)$  along the curve  $\Gamma$  according to the following notation:  $(\xi, w) = (x, y, w)$ , for  $N = 2n + 1$ ; and  $(\xi, w) = (x, y, z, w)$ , for  $N = 2n$ . Clearly  $\xi = (x, y)$  and  $\xi = (x, y, z)$  respectively, and

$$\begin{aligned} (x, y) &= (x_1, y_1, \dots, x_n, y_n), & \text{for } N = 2n + 1, \text{ whereas} \\ (x, y) &= (x_1, y_1, \dots, x_{n-1}, y_{n-1}), & \text{for } N = 2n. \end{aligned}$$

Then, with an obvious abuse of notation, already used in section (2), we write in coordinates:

$$F = Q \frac{\partial}{\partial \xi} + L \frac{\partial}{\partial w}. \quad (10)$$

We expand  $Q(\xi, w)$  and  $L(\xi, w)$  in power series of the variables  $\xi_j$ , at  $\xi = 0$ , along  $\Gamma$ , as follows:

$$Q(\xi, w) = Q_0(\xi, w) + Q_1(\xi, w) + Q_2(\xi, w) + \dots, \quad (11)$$

$$L(\xi, w) = L_0(\xi, w) + L_1(\xi, w) + L_2(\xi, w) + \dots, \quad (12)$$

where  $Q_k$  and  $L_k$  are homogeneous polynomials of degree  $k$ , in the variables  $x_j$  and  $y_j$  (and  $z$ , if  $N = 2n$ ). Then, for  $\Sigma = (F, \Gamma) \in S_*^\infty$ , setting  $i = \sqrt{-1}$ , we know the following two facts:

- a). For  $N = 2n$ , the eigenvalues of  $A(w)$  are smooth functions:  $\pm i\alpha_1(w), \dots, \pm i\alpha_{n-1}(w), 0$ , where  $\alpha_j(w) > 0$  for all  $j$ . For all  $w$ , we have that  $\alpha_k(w) \neq \alpha_l(w)$ , provided  $k \neq l$ . Hence, for all  $w$ , we can order the eigenvalues as follows:

$$\alpha_1(w) > \dots > \alpha_{n-1}(w) > 0.$$

- b). For  $N = 2n + 1$ , the eigenvalues of  $A(w)$  are smooth functions:  $\pm i\alpha_1(w), \dots, \pm i\alpha_n(w)$ , where again for all  $w$  we have that  $\alpha_k(w) \neq \pm\alpha_l(w)$  for  $k \neq l$ , and one at most is zero, for isolated values of  $w$ . If none of the  $\alpha_j$ 's vanishes somewhere, we proceed as in a). If one vanishes for some  $w_0$ , then, we order the eigenvalues at  $w_0$ , as follows:

$$\alpha_1(w_0) > \dots > \alpha_{n-1}(w_0) > \alpha_n(w_0) = 0,$$

in any case, we can always conclude that for all  $w$  we have

$$\alpha_1(w) > \dots > \alpha_{n-1}(w) > \alpha_n(w),$$

and  $\alpha_{n-1}(w) > 0$  for all  $w$ . This last point ( $\alpha_{n-1}(w) > 0$ ) holds, because if it were not true, at some point  $w'$ , then  $\alpha_{n-1}(w') = \pm\alpha_n(w') \neq 0$ , which is not possible.

In the normal coordinates, it might be eventually necessary to consider permutations  $(x_j, y_j) \rightarrow (y_j, x_j)$ , to write:

$$\begin{aligned} d\omega|_{\Delta(\Gamma(t))} &= \alpha_1(t)dx_1 \wedge dy_1 + \dots + \alpha_n(t)dx_n \wedge dy_n, \quad \text{and} \\ d\omega|_{\Delta(\Gamma(t))} &= \alpha_1(t)dx_1 \wedge dy_1 + \dots + \alpha_{n-1}(t)dx_{n-1} \wedge dy_{n-1}, \end{aligned}$$

for  $N = 2n + 1$ , and  $N = 2n$  respectively.

Here the frame (8) along the curve  $\Gamma$  is orthonormal, as a frame on  $\Delta$ , and the matrices  $\bar{A}(t)$  of the linear map  $A(t)$ , with respect to this frame are the following:

$$\bar{A}(w) = \begin{pmatrix} 0 & -\alpha_1(w) & 0 & \dots & 0 & 0 \\ \alpha_1(w) & 0 & -\alpha_2(w) & \dots & 0 & 0 \\ 0 & \alpha_2(w) & 0 & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \dots & 0 & -\alpha_n(w) \\ 0 & 0 & 0 & \dots & \alpha_n(w) & 0 \end{pmatrix},$$

for  $N = 2n + 1$ , and

$$\bar{A}(w) = \begin{pmatrix} 0 & -\alpha_1(w) & 0 & \dots & 0 & 0 & 0 \\ \alpha_1(w) & 0 & -\alpha_2(w) & \dots & 0 & 0 & 0 \\ 0 & \alpha_2(w) & 0 & \dots & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & \dots & 0 & -\alpha_{n-1}(w) & 0 \\ 0 & 0 & 0 & \dots & \alpha_{n-1}(w) & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix},$$

for  $N = 2n$ .

**Theorem 3.4 (Normal Form)** *Let  $\Sigma = (F, \Gamma) \in S_*^\infty$ , and let  $(\xi, w) = (x, y, w)$  or  $(\xi, w) = (x, y, z, w)$ , be a fixed normal coordinate system. Then, there is a unique sub-Riemannian orthonormal frame  $\mathcal{F}$  (gauge equivalent to  $F$ ),  $\mathcal{F} = Q \frac{\partial}{\partial \xi} + L \frac{\partial}{\partial w}$ , with the following properties:*

- 1)  $Q$  is symmetric
- 2)  $Q_0(\xi, w) = Id$
- 3)  $Q(\xi, w)\xi = \xi$
- 4)  $L(\xi, w) \cdot \xi = 0$
- 5)  $Q_1 = 0$
- 6)  $L_0 = 0$
- 7) *The first order linear forms are given as follows:*

$$\begin{aligned} L_1 &= \left( \frac{\alpha_1(w)}{2} y_1, -\frac{\alpha_1(w)}{2} x_1, \dots, \frac{\alpha_n(w)}{2} y_n, -\frac{\alpha_n(w)}{2} x_n \right), \text{ and} \\ L_1 &= \left( -\frac{\alpha_1(w)}{2} y_1, -\frac{\alpha_1(w)}{2} x_1, \dots, \frac{\alpha_{n-1}(w)}{2} y_{n-1}, \frac{\alpha_{n-1}(w)}{2} x_{n-1}, 0 \right), \end{aligned}$$

for  $N = 2n + 1$  and  $N = 2n$  respectively.

**Proof.** The identities 1) to 6) are proved in [6] for the contact case and in [10] for the quasi-contact one. Equation 5) is an interesting sub-Riemannian analog of a Bianchi identity in Riemannian geometry. We prove 7), which is the only different point. We prove it for  $N = 2n + 1$  only, (the case  $N = 2n$  is similar).

We already know (Section 2, proof of Theorem 2.1) that, if we take  $\omega_N = dw - LQ^{-1}d\xi$ , then  $\tilde{\alpha}(\Gamma(w)) = d\omega_N|_{\Delta(\Gamma(w))}$  has  $\bar{A}(w)$  as matrix. Now, on one hand, by definition of our coordinates, we have that

$$d\omega_N|_{\Delta(\Gamma(w))} = \alpha_1(w)dx_1 \wedge dy_1 + \dots + \alpha_n(w)dx_n \wedge dy_n, \quad (13)$$

and on the other hand that

$$d\omega_N|_{\Delta(\Gamma(w))} = -d(LQ^{-1}d\xi)(\Gamma(w)).$$

Now, 6) implies

$$L = L_1(w, z) + O^2(\xi) = (L_1^1, M_1^1, \dots, L_1^n, M_1^n) + O^2(\xi),$$

and in consequence 4) yields,

$$L_1^1 x_1 + M_1^1 y_1 + \dots + L_1^n x_n + M_1^n y_n = 0.$$

Therefore, by writing for all  $k$ :



$$\begin{aligned} L_1^k &= \sum_j (L_1^{kj} x_j + \widehat{L_1^{kj}} y_j), \quad \text{and} \\ M_1^k &= \sum_j (M_1^{kj} x_j + \widehat{M_1^{kj}} y_j), \end{aligned}$$

we obtain

$$\sum_{j,k} (L_1^{kj} x_j x_k + \widehat{L_1^{kj}} y_j x_k + M_1^{kj} x_j y_k + \widehat{M_1^{kj}} y_j y_k) = 0,$$

which implies:

$$L_1^{kk} = 0, \quad \widehat{M_1^{kk}} = 0, \quad L_1^{kj} = -L_1^{jk}, \quad \widehat{L_1^{kj}} = -M_1^{jk}, \quad \text{and} \quad \widehat{M_1^{kj}} = -\widehat{M_1^{jk}}.$$

A straightforward calculation yields

$$d(L_1 d\xi)|_\Gamma = 2 \sum_{j < k} L_1^{kj} dx_j \wedge dx_k + 2 \sum_{j < k} \widehat{M_1^{kj}} dy_j \wedge dy_k + \sum_{j,k} \widehat{L_1^{kj}} dy_j \wedge dx_k + \sum_{j,k} M_1^{kj} dx_j \wedge dy_k,$$

or equivalently

$$d(L_1 d\xi)|_\Gamma = 2 \left( \sum_{j < k} (L_1^{kj} dx_j \wedge dx_k + \widehat{M_1^{kj}} dy_j \wedge dy_k) - \sum_{j,k} \widehat{L_1^{kj}} dx_k \wedge dy_j \right).$$

Now,

$$-d(LQ^{-1}d\xi)|_{\Delta(\Gamma(w))} = -d(Ld\xi)|_{\Delta(\Gamma(w))},$$

(because  $Q^{-1} = Id + O^2(\xi)$ ), and,

$$-d(Ld\xi)|_{\Delta(\Gamma(w))} = -d(L_1 d\xi)|_{\Delta(\Gamma(w))}.$$

By (13), we deduce that for all  $k$  and  $j$

$$\begin{aligned} L_1^{kj} &= 0, \\ \widehat{M_1^{kj}} &= 0, \quad \text{and} \\ 2 \sum_{j,k} \widehat{L_1^{kj}} dx_k \wedge dy_j &= - \sum_j \alpha_j(w) dx_j \wedge dy_j. \end{aligned}$$

Hence,

$$\begin{aligned}\widehat{L}_1^{kj} &= 0, \quad \text{for } k \neq j, \quad \text{and} \\ \widehat{L}_1^{kk} &= -M_1^{kk} = -\frac{\alpha_k}{2}.\end{aligned}$$

This gives 7), changing  $(x_i, y_i)$  for  $(y_i, x_i)$ . ◇

Conversely, if we have a coordinate neighborhood of  $\Gamma$  and a frame  $F$  meeting conditions 1), 2), 3), 4), 7), then 5), 6) are automatically satisfied, and the coordinates turn out to be normal.

### 3.3 Special Case: normal form in dimension 3

In dimension 3, the normal form is given by the frame  $F = \{F_1, G_1\}$

$$\begin{aligned}F_1 &= (1 + y^2\beta(x, y, w))\frac{\partial}{\partial x} - xy\beta(x, y, w)\frac{\partial}{\partial y} + \frac{y}{2}\gamma(x, y, w)\frac{\partial}{\partial w}, \\ G_1 &= -xy\beta(x, y, w)\frac{\partial}{\partial x} + (1 + x^2\beta(x, y, w))\frac{\partial}{\partial y} - \frac{x}{2}\gamma(x, y, w)\frac{\partial}{\partial w},\end{aligned}\quad (14)$$

where  $\beta$  and  $\gamma$  are smooth functions.

In the contact case,  $\gamma(0, 0, w)$  never vanishes, and we can take it positive. Furthermore, we can normalize the function  $\beta$  by the condition  $\beta(0, 0, w) = 0$ , which is tantamount of saying that the curvature or the metric  $g_w$ , (projection of  $g$  on the cross section  $S_w = \{w = \text{constant}\}$ ), is zero. This normal form has been first given in [4], for the formal power series case, and after in [5] for the smooth case.

## 4 Complexity and asymptotic optimal synthesis

We discuss now the metric complexity. The results of this section are valid for all  $N \geq 4$ , and in the absence of Martinet points along the curve  $\Gamma$ , they are also valid for  $N = 3$ .

Using Theorem 3.4 (Normal Form), and the notations of Section 3, we consider a normal coordinate neighborhood  $(\xi, w)$  of the curve  $\Gamma$ , and a normal frame along  $\Gamma$ , that we write as:

$$\begin{aligned}F &= (F_1, G_1, \dots, F_n, G_n), \quad \text{for } N = 2n + 1 \quad \text{or} \\ F &= (F_1, G_1, \dots, F_{n-1}, G_{n-1}, H_n), \quad \text{for } N = 2n,\end{aligned}$$

where

$$\begin{aligned}
F_i &= (1 + O^2(\xi)) \frac{\partial}{\partial x_i} + \left(\frac{y_i}{2} \alpha_i(w) + O^2(\xi)\right) \frac{\partial}{\partial w}, \\
G_i &= (1 + O^2(\xi)) \frac{\partial}{\partial y_i} + \left(-\frac{x_i}{2} \alpha_i(w) + O^2(\xi)\right) \frac{\partial}{\partial w}, \\
H_n &= (1 + O^2(\xi)) \frac{\partial}{\partial x_{N-1}} + O^2(\xi) \frac{\partial}{\partial w}.
\end{aligned} \tag{15}$$

All the functions denoted by  $O^i(\xi)$  are (different) smooth functions in the variables  $(\xi, w)$ , that are in the  $i^{\text{th}}$  power of the ideal generated by  $\xi_i$ ,  $i = 1, \dots, N-1$ .

Let us now consider any (smooth) admissible curve  $\delta : [0, T] \rightarrow T_\varepsilon$ ,  $t \mapsto \delta(t) = (\widehat{\xi}(t), \widehat{w}(t))$  entirely contained in the sub-Riemannian tube  $T_\varepsilon$ , (hence, in normal coordinates, contained in the Riemannian tube, because in normal coordinates, they are the same). Furthermore, let us assume that the following conditions hold:

$$\widehat{w}(0) = w_0, \quad \text{and} \quad \widehat{w}(T) = w_1.$$

Assume that the curve  $\delta$  is parameterized by arc length, assume also that the corresponding control functions are  $u_i(\cdot)$  with  $i = 1, \dots, N-1$ . The length of  $\delta$  is  $T$ , and

$$\sum_{i=1}^{N-1} u_i^2(t) = 1, \quad \text{for all } t \in [0, T],$$

and also

$$\frac{d\widehat{w}}{dt} = \sum_{i=1}^{N-1} (L_i^1(\widehat{\xi}, \widehat{w}) + O^2(\widehat{\xi})) u_i.$$

Let  $\kappa(w) = \sup_i \{\alpha_i(w)\}$ , we have that  $\kappa$  is strictly positive on the interval  $[w_0, w_1]$ . Furthermore, for  $N = 2n + 1$  we have that

$$\frac{d\widehat{w}}{dt} \leq \frac{\kappa(\widehat{w})}{2} (|\widehat{y}_1| |u_1| + |\widehat{x}_1| |u_2| + \dots + |\widehat{y}_n| |u_{N-2}| + |\widehat{x}_n| |u_{N-1}|) + |O^2(\widehat{\xi})|,$$

whereas for  $N = 2n$

$$\begin{aligned}
\frac{d\widehat{w}}{dt} &\leq \frac{\kappa(\widehat{w})}{2} (|\widehat{y}_1| |u_1| + |\widehat{x}_1| |u_2| + \dots + |\widehat{y}_{n-1}| |u_{N-3}| + |\widehat{x}_{n-1}| |u_{N-2}|) + |O^2(\widehat{\xi})| \\
&\leq \frac{\kappa(\widehat{w})}{2} \langle |\widehat{\xi}|, |u| \rangle_g + O^2(\widehat{\xi}),
\end{aligned}$$

where  $|\widehat{\xi}|$  and  $|u|$  are the vectors with entries  $|\widehat{\xi}_i|$  and  $|u_i|$  respectively. In any case, since  $\widehat{\xi}$  is in the tube  $T_\varepsilon$ , and  $\|u\|_2 = 1$ , we have that

$$\frac{d\hat{w}}{dt} \leq \frac{\kappa(\hat{w})}{2} \|\hat{\xi}\|_2 \|u\|_2 + |O^2(\hat{\xi})| \leq \varepsilon \frac{\kappa(\hat{w})}{2} + K\varepsilon^2,$$

for some constant  $K > 0$ . But then, there is a constant  $K'$  such that

$$\frac{d\hat{w}}{\kappa(\hat{w})} \leq \varepsilon \frac{dt}{2} (1 + \varepsilon K'),$$

and consequently, there is a constant  $K''$ , such that

$$T \geq \frac{2}{\varepsilon} \int_{w_0}^{w_1} \frac{dw}{\kappa(w)} (1 + \varepsilon K''). \quad (16)$$

**Remark 4.1** *In order to reduce the burden in the notation for the constants appearing in the estimates ( $K, K', \dots$ ), in the remaining of the paper, we shall abuse the notation using only one  $K$  for all the (different) constants.*

Now, we shall exhibit a (strong) *asymptotic optimal synthesis*. We know that, for  $\varepsilon$  small enough, the cylinder  $C_\varepsilon = \{\|\xi\| = \varepsilon\}$  is transversal to the curve  $\Gamma$ . Let us consider also the following 2-dimensional cylinder

$$C_\varepsilon^1 = \{\xi \in C_\varepsilon \mid x_1^2 + y_1^2 = \varepsilon^2\}.$$

Then, for  $\varepsilon$  small enough, we have a (never vanishing) vector field  $X_\varepsilon$  on  $C_\varepsilon^1$  defined by the following three conditions

- a)  $X_\varepsilon(\xi, w) \in \Delta(\xi, w) \cap T_{(\xi, w)}(C_\varepsilon^1)$ ,
- b)  $\|X_\varepsilon(\xi, w)\|_g = 1$ , and
- c)  $L_{X_\varepsilon}(w) > 0$ .

On the cylinder  $C_\varepsilon^1$ , the integral curves of  $X_\varepsilon$  satisfy:

$$\dot{w} = \frac{\alpha_1(w)}{2} \varepsilon + \varepsilon^2 F(\varepsilon, \theta, w),$$

where  $F$  is a bounded function; and  $x_1 = \varepsilon \cos \theta$ ,  $y_1 = \varepsilon \sin \theta$ . This can be shown by elementary but tedious calculations using the normal form. This shows that, for  $\varepsilon$  sufficiently small, trajectories of  $X_\varepsilon$  can join any  $w_0$  and  $w_1$  with  $w_0 < w_1$  and  $(0, w_0), (0, w_1) \in T_\varepsilon$ .

But then, since  $F$  is bounded on  $T_\varepsilon$  and  $|\alpha_1(w)| > A > 0$  for some constant  $A$ , there is a positive constant  $K$  such that

$$\frac{\dot{w}}{\alpha_1(w)} = \varepsilon \left( \frac{1}{2} + \varepsilon F(\varepsilon, \theta, w) \right) \geq \frac{\varepsilon}{2} - \frac{\varepsilon^2 K}{2}.$$

In consequence, if  $T(\varepsilon)$  is the time to join  $w_0$  to  $w_1$ , then:

$$\int_{w_0}^{w_1} \frac{dw}{\alpha_1(w)} \geq \frac{\varepsilon}{2} (1 - \varepsilon K) T(\varepsilon),$$

and furthermore

$$T(\varepsilon) \leq \frac{2}{\varepsilon} \frac{1}{(1 - \varepsilon K)} \int_{w_0}^{w_1} \frac{dw}{\kappa(w)} \leq \frac{2}{\varepsilon} \int_{w_0}^{w_1} \frac{dw}{\kappa(w)} (1 - \varepsilon A),$$

for some  $A > 0$ , if  $\varepsilon$  is small enough.

Now, we can apply this to  $w_0$  and  $w_1$ , such that:

$$\Gamma(0) = (0, w_0), \quad \Gamma(1) = (0, w_1).$$

Our trajectory does not join  $\Gamma(0)$  to  $\Gamma(1)$ , since it is entirely contained in  $C_\varepsilon$ . But then, any  $(\xi, w) \in C_\varepsilon$  can be joined to  $(0, w)$  in time  $\varepsilon$ , by definition of  $C_\varepsilon$ , and by the fact that the rays through  $\Gamma$  in the planes  $S_w$  are geodesics. Therefore, we can construct an admissible trajectory  $\delta$  from  $\Gamma(0)$  to  $\Gamma(1)$ , entirely contained in  $T_\varepsilon$ , such that the time  $T(\varepsilon)$  to join  $\Gamma(0)$  to  $\Gamma(1)$  verifies:

$$T(\varepsilon) \leq \frac{2(1 - \varepsilon A)}{\varepsilon} \int_{w_0}^{w_1} \frac{dw}{\kappa(w)} + 2\varepsilon, \quad (17)$$

Since both curves are parameterized by arc-length, then (17) together with (16), imply

$$\frac{2(1 + \varepsilon K)}{\varepsilon^2} \int_{w_0}^{w_1} \frac{dw}{\kappa(w)} \leq MC_\Sigma(\varepsilon) \leq \frac{2(1 - \varepsilon A)}{\varepsilon^2} \int_{w_0}^{w_1} \frac{dw}{\kappa(w)} + 2.$$

As a conclusion we have that

$$MC_\Sigma(\varepsilon) \underset{s}{\simeq} \frac{2}{\varepsilon^2} \int_{w_0}^{w_1} \frac{dw}{\kappa(w)},$$

which finishes the proof of Theorem 1.2 and also that of Theorem 1.3 in the absence of Martinet points. Even more, we have explicitly exhibited an asymptotic optimal synthesis that goes under the following steps

- a) Follow a horizontal geodesic, to go from  $\Gamma(0)$  to  $C_\varepsilon^1$ ,
- b) follow then the flow of  $X_\varepsilon$  on  $C_\varepsilon^1$  up to  $w_1$ ,
- c) connect the point reached on  $C_\varepsilon^1$  to  $\Gamma(1)$  by a horizontal geodesic.

## 5 Martinet Case

### 5.1 Proof of the Theorem 1.3

Now, we consider in  $\mathbb{R}^3$  the orthonormal frame  $\{F_1, G_1\}$ , in normal form (14),

$$\begin{aligned} F_1 &= (1 + y^2\beta(x, y, w))\frac{\partial}{\partial x} - xy\beta(x, y, w)\frac{\partial}{\partial y} + \frac{y}{2}\gamma(x, y, w)\frac{\partial}{\partial w}, \\ G_1 &= -xy\beta(x, y, w)\frac{\partial}{\partial x} + (1 + x^2\beta(x, y, w))\frac{\partial}{\partial y} - \frac{x}{2}\gamma(x, y, w)\frac{\partial}{\partial w}. \end{aligned}$$

Let us assume the existence of a single isolated Martinet point, at  $q = (x, y, w) = 0$  (we know, generically that Martinet points are isolated). Then:

$$\gamma(x, y, w) = \gamma_0(w) + (x, y) \cdot \gamma_1(x, y, w), \quad \text{and} \quad \gamma_0(0) = 0, \quad (18)$$

here and from now,  $(x, y) \cdot \gamma_1$  means the scalar product  $x\gamma_1^1 + y\gamma_1^2$ .

For a generic problem  $\Sigma = (F_1, G_1)$ , it is easy to check (we leave this to the reader), that  $\frac{\partial\gamma_0}{\partial w}$  is non zero at the Martinet point. In fact,  $\frac{\partial\gamma_0}{\partial w}(0)$  is an invariant of the problem.

We can change the parameterization of  $\Gamma$  using a smooth diffeomorphism  $\Phi$  in such a way that, setting  $\tilde{w} = \Phi(w)$ , we get:

$$\gamma_0(\tilde{w})\frac{\partial}{\partial\tilde{w}} = -\kappa\tilde{w}\frac{\partial}{\partial\tilde{w}}.$$

In what follows, we shall forget about  $\tilde{w}$ , and we shall take,

$$\gamma_0(w) = -\kappa w.$$

Let us assume  $\kappa > 0$  (the case  $\kappa < 0$  is similar). Since the admissible curves are parameterized by arc length, the controls  $u(t)$  and  $v(t)$  satisfy  $u^2(t) + v^2(t) = 1$ , and for a trajectory entirely contained in the tube  $T_\varepsilon$ , we have:

$$\frac{dw}{dt} = (-\kappa w + (x, y) \cdot \gamma_1(x, y, w))\left(\frac{y}{2}u - \frac{x}{2}v\right). \quad (19)$$

Let us take an admissible curve  $q(t) = (x(t), y(t), w(t))$ , such that  $w_0 = w(0) < 0 < w(T) = w_1$ . And let us consider  $t_1$  and  $t_2$  such that  $0 < t_2 \leq t_1 < T$ ,  $w(t_1)$  is zero for the first time, and  $w(t_2)$  is zero for the last time (continuity of  $q(t)$ .)

Let us consider the piece  $w(\cdot) > 0$  of the curve (the piece  $w(\cdot) < 0$  is similar) and assume that  $q(t)$  is contained in the  $\varepsilon$ -tube  $T_\varepsilon$ . Then, on  $[t_2, T]$ , we have:

$$\left|\frac{y}{2}u - \frac{x}{2}v\right| \leq \frac{\varepsilon}{2}, \quad \text{and} \quad \frac{dw}{dt} \leq (\kappa w + \varepsilon K)\frac{\varepsilon}{2},$$

since  $\gamma_1(x, y, w)$  is bounded on the compact tube  $T_\varepsilon$ .

Now, since  $w(\cdot)$  is at least absolutely continuous then,

$$\tilde{w}(t) = w(t)e^{-\frac{\kappa}{2}\varepsilon(t-t_2)},$$

is also absolutely continuous. A direct estimation yields

$$\frac{d\tilde{w}}{dt} = \frac{dw}{dt}e^{-\frac{\kappa}{2}\varepsilon(t-t_2)} - \frac{\kappa\varepsilon}{2}\tilde{w} \leq \frac{\kappa\varepsilon}{2}\tilde{w} + \frac{\varepsilon^2}{2}Ke^{-\frac{\kappa}{2}\varepsilon(t-t_2)} - \frac{\kappa\varepsilon}{2}\tilde{w},$$

but then

$$\frac{d\tilde{w}}{dt} \leq \frac{\varepsilon^2}{2}Ke^{-\frac{\kappa}{2}\varepsilon(t-t_2)}, \quad \text{and} \quad \tilde{w}(t) \leq \frac{\varepsilon}{\kappa}K \left(1 - e^{-\frac{\kappa}{2}\varepsilon(t-t_2)}\right),$$

therefore

$$w(t) \leq \frac{\varepsilon}{\kappa}K \left(e^{\frac{\kappa\varepsilon}{2}(t-t_2)} - 1\right) \leq \frac{\varepsilon}{\kappa}Ke^{\frac{\kappa\varepsilon}{2}(t-t_2)},$$

which means that:

$$(t - t_2) \geq \frac{2}{\varepsilon\kappa} \ln \left( \frac{\kappa w(t)}{\varepsilon K} \right) = -\frac{2}{\kappa} \left( \frac{\ln \varepsilon}{\varepsilon} - \frac{\ln \left( \frac{\kappa w(t)}{K} \right)}{\varepsilon} \right).$$

For the part of the curve corresponding to  $w(t) < 0$ , we get a similar estimation (changing  $w$  for  $-w$  in (19) above, the computations are similar). At the end, if  $T$  is the time to connect  $w_0$  to  $w_1$ , we obtain that:

$$T \geq -\frac{4}{\kappa} \frac{\ln \varepsilon}{\varepsilon} + \frac{2}{\varepsilon\kappa} \ln \left( \frac{\kappa^2 w_0 w_1}{K^2} \right),$$

which shows that:

$$MC(\varepsilon) \geq -\frac{4}{\kappa} \frac{\ln \varepsilon}{\varepsilon^2} + \frac{2}{\varepsilon^2\kappa} \ln \left( \frac{\kappa^2 w_0 w_1}{K^2} \right).$$

Now if we take any constant  $C$  such that  $0 < C < \frac{4}{\kappa}$ , for  $\varepsilon$  small enough then

$$MC(\varepsilon) \geq -C \frac{\ln \varepsilon}{\varepsilon^2}. \tag{20}$$

And the proof of Theorem 1.3 is complete.

**Remark 5.1** *In fact, it can be proved that in general,*

$$MC(\varepsilon) \underset{s}{\simeq} -C \frac{\ln \varepsilon}{\varepsilon^2}, \quad \text{where} \quad \frac{4}{\kappa} \leq C \leq \frac{12}{\kappa}, \tag{21}$$

*however this estimation requires tedious calculations, that shall be discussed elsewhere. We limit ourselves to analyze an example, for which the estimation (21) above holds.*




Figure 1: The limit cycle

## 5.2 An example.

Before starting with the example, we shall explain what should be a (weak) asymptotic optimal synthesis in the general Martinet case ( $\kappa \neq 0$  above). Starting with (19), we consider the cylinder  $C_\varepsilon$ . Since  $\Gamma$  is transversal to  $\Delta$ , and  $\Gamma$  is compact, then for  $\varepsilon$  small enough, as in the section 4,  $\Delta$  is transversal also to  $C_\varepsilon$ , and therefore,  $\Delta$  defines a vector field  $X_\varepsilon$  on  $C_\varepsilon$ , (see figure 1).

We see in equation (19) that, for  $|w|$  large, this vector field points in the direction of  $w = 0$ . Therefore, by the Poincaré-Bendixon theorem, there is a limit-cycle.

It is possible to check that, for generic problems, this limit-cycle is not horizontal (not contained in a plane  $\{w = \text{constant}\}$ ), has size  $\varepsilon^3$ , and is centered at a point  $w_c$  of order  $\varepsilon^2$ .

Then, a well defined synthesis goes under the following steps:

- Starting from  $w_0$ , follow any integral curve of  $X_\varepsilon$ , up to the first time it crosses the plane  $\{w = w_c\}$ .
- Following horizontal geodesic, use at most  $2\varepsilon$  units of time, to cross the plane  $\{w = w_c\}$ .
- Following (in reversed time) a suitably chosen trajectory of  $X_\varepsilon$ , reach the desired destination  $w_1$ .

As we shall show in the example, this strategy gives a complexity which is strongly equivalent to  $-12 \frac{\ln \varepsilon}{\varepsilon^2}$ .



**Example 5.2** Consider in  $\mathbb{R}^3$  the frame

$$\begin{aligned} F &= \frac{\partial}{\partial x} + \frac{y}{2}\gamma \frac{\partial}{\partial w}, \\ G &= \frac{\partial}{\partial y} - \frac{x}{2}\gamma \frac{\partial}{\partial w}, \end{aligned}$$

with  $\gamma = -\alpha w + ax$ .

The fact that  $a$  is nonzero (an invariant fact in the general normal form) makes the distribution non-integrable, and the limit-cycle non horizontal. In cylindrical coordinates,  $(w, \theta, \varepsilon)$ , it is to easy to see that:

$$X_\varepsilon = -\frac{1}{\varepsilon} \frac{\partial}{\partial \theta} + \frac{\varepsilon}{2}(-\alpha w + a\varepsilon \cos \theta) \frac{\partial}{\partial w}.$$

Then, we can explicitly compute the flow of  $X_\varepsilon$ , to obtain

$$\begin{aligned} \theta(t) &= -\frac{t}{\varepsilon} + \theta_0, \quad \text{and} \\ w(t) &= e^{-\frac{\varepsilon\alpha}{2}t} w_0 + \frac{\varepsilon^2 a}{2} \int_0^t e^{-\frac{\varepsilon\alpha}{2}(t-s)} \cos \theta(s) ds. \end{aligned}$$

After straightforward computations, we obtain

$$\begin{aligned} w(t) &= (4 + \alpha^2 \varepsilon^4)^{-1} \left( e^{-\frac{\varepsilon\alpha t}{2}} \left[ 4w_0 + \alpha^2 \varepsilon^4 w_0 + a\alpha e^{\frac{\varepsilon\alpha t}{2}} \varepsilon^5 \cos\left(\frac{t}{\varepsilon} - \theta_0\right) \right. \right. \\ &\quad \left. \left. - a\alpha \varepsilon^5 \cos \theta_0 + 2a e^{\frac{\varepsilon\alpha t}{2}} \varepsilon^3 \sin\left(\frac{t}{\varepsilon} - \theta_0\right) + 2a \varepsilon^3 \sin \theta_0 \right] \right). \end{aligned} \quad (22)$$

The expression for the limit cycle is the following

$$\begin{aligned} \theta(t) &= -\frac{t}{\varepsilon}, \\ w(t) &= \frac{a\varepsilon^3}{4 + \alpha^2 \varepsilon^4} \left( \alpha \varepsilon^2 \cos\left(\frac{t}{\varepsilon}\right) + 2 \sin\left(\frac{t}{\varepsilon}\right) \right) \end{aligned}$$

which shows that  $w_c = 0$ , and the size of the limit cycle is  $\frac{a}{2}\varepsilon^3$ .

Now, starting from  $w_0 > 0$ , we want to go to  $w_c = 0$ , using (22). We get

$$0 = e^{-\frac{\varepsilon\alpha}{2}t} \left( 4w_0 + 2a\varepsilon^3 (\sin \theta_0 + e^{\frac{\varepsilon\alpha}{2}t} \sin\left(\frac{t}{\varepsilon} - \theta_0\right)) \right) + O(\varepsilon^4),$$

(were  $\frac{O(\varepsilon^4)}{\varepsilon^4}$  is bounded). We rewrite this:

$$4w_0 e^{-\frac{\varepsilon\alpha}{2}t} = -2a\varepsilon^3 \left( \sin \theta_0 e^{-\frac{\varepsilon\alpha}{2}t} + \sin\left(\frac{t}{\varepsilon} - \theta_0\right) \right) + O(\varepsilon^4). \quad (23)$$

It is clear that this equation in  $t$  has a solution, for which:

$$e^{-\frac{\varepsilon\alpha}{2}t} \simeq \frac{a}{2w_0} \varepsilon^3.$$

Hence,

$$t \simeq -\frac{6}{\alpha\varepsilon} \ln \varepsilon.$$

Taking into account the part  $w(t) < 0$  of the trajectory, we get that the complexity  $C(\varepsilon)$  of our synthesis (complexity =  $\frac{1}{\varepsilon}$  length):

$$C(\varepsilon) \simeq -12 \frac{\ln \varepsilon}{\alpha\varepsilon^2}. \quad (24)$$

## 6 The non generic case.

### 6.1 Normal form

Here we shall restrict ourselves to  $C^\omega$  (analytic) motion-planning problems. In this case, prenormal coordinates of Theorem 3.1, Section 3 still exist, and they are  $C^\omega$  coordinates along the curve  $\Gamma$ .

Now, to obtain normal coordinates, (Theorem 3.2, Section 3), we need the analog of Lemma 7.1, Appendix 7, for the analytic case. This is Lemma 7.2 of the same Appendix, which claims that in fact, the *smooth diagonalization property* for the fundamental form  $\tilde{\alpha}$  always holds. Because of that, we have the following theorem:

**Theorem 6.1 Normal Coordinates in the Analytic Case.** *Let  $\Sigma = (F, G) \in S_R^\omega$ . Then, without any extra assumption, the conclusions of Theorem 3.2, Section 3, still hold, and the normal coordinate system is compatible with the analytic structure.*

We have also the analog of the Theorem 3.4, Section 3.2 in the analytic context. Let us repeat it precisely:

**Theorem 6.2 Analytic Normal Form** *Let  $\Sigma = (F, \Gamma) \in S_R^\omega$ , and let  $(\xi, w)$  be a fixed normal  $C^\omega$  coordinate system along the curve  $\Gamma$  (it does exist by Theorem 6.1). Then, there is a family of analytic functions  $\alpha_1(\cdot), \dots, \alpha_n(\cdot) : [0, 1] \rightarrow \mathbb{R}$ , such that the fundamental form  $\tilde{\alpha}$  along  $\Gamma$  is*

$$\tilde{\alpha}(w) = \alpha_1(w) dx_1 \wedge dy_1 + \dots + \alpha_n(w) dx_n \wedge dy_n, \quad N = 2n, \text{ or } 2n + 1.$$

Also, there is an orthonormal analytic sub-Riemannian frame  $\mathcal{F} = Q \frac{\partial}{\partial \xi} + L \frac{\partial}{\partial w}$ , with the same properties 1) to 7) as those of Theorem 3.4, Section 3.

The only difference is that these analytic functions  $\alpha_1(\cdot), \dots, \alpha_{n-1}(\cdot)$  are not necessarily positive everywhere and their graphs might cross each other.

## 6.2 Metric Complexity

Let us make just the following assumption

$$\kappa(w) = \sup_i |\alpha_i(w)| \neq 0 \quad \text{for all } w \in [0, 1]. \quad (25)$$

In fact this is equivalent to say that the fundamental form  $\tilde{\alpha}(w) \neq 0$ , for all  $w \in [0, 1]$ . Then, the statements of Theorem 1.4 hold, namely

Complexity:

- Under assumption (25), formula (1) for the metric complexity is still valid.
- The function  $T \mapsto 2 \int_0^T \frac{dt}{\kappa(t)}$ , is of class  $C^1$ , piecewise analytic, and its second derivative having jumps at a finite number of times.

## 6.3 Asymptotic optimal synthesis

We describe now the corresponding asymptotic optimal synthesis for the non-generic case. The function  $\kappa : [0, 1] \rightarrow \mathbb{R}$  is clearly continuous, piecewise analytic, positive, with jumps on its derivative at a finite number of points, say  $t_1, \dots, t_m$ . Between two successive times  $t_j$  and  $t_{j+1}$ , the supremum in assumption (25) is reached by one of the eigenvalues moduli  $|\alpha_{i_j}|$ . Also between the consecutive times  $t_j$  and  $t_{j+1}$ , we consider the cylinder  $C_\varepsilon^j$  in the normal coordinates:

$$C_\varepsilon^j = \{(\xi, w) : \|\xi\|^2 = \varepsilon^2, x_{i_j}^2 + y_{i_j}^2 = \varepsilon^2\}.$$

For  $\varepsilon$  small enough,  $\Delta$  is transversal to  $C_\varepsilon^j$ , and defines a vector field  $X_\varepsilon^j$  on  $C_\varepsilon^j$ .

Asymptotic optimal synthesis:

- between  $t_j$  and  $t_{j+1}$ : follow the flow (any trajectory) of  $X_\varepsilon^j$
- at  $t = t_j$ , or  $t = 0$ , or  $t = 1$ : follow horizontal geodesics in the planes  $\{w = \text{constant}\}$ , passing through  $\Gamma$ :
  - at  $t = 0$ , in time exactly  $\varepsilon$ , reach  $C_\varepsilon^1$  from  $\Gamma(0)$
  - at  $t = 1$ , in time exactly  $\varepsilon$ , go from  $C_\varepsilon^m$  to  $\Gamma(1)$

- at  $t = t^j$ , join  $C_\varepsilon^{j-1}$  to  $C_\varepsilon^j$  in time less than  $2\varepsilon$ .

The proof of the fact that this is a strong asymptotic optimal synthesis, and the proof of Theorem 1.4, are obvious modifications of the proof (Section 4) of Theorem 1.2: Theorem 6.2 allows us to repeat exactly the same arguments between two successive crossing points  $t_j$  and  $t_{j+1}$ .

## 7 Appendix

### 7.1 Diagonalization of 1-parameter families of skew-symmetric matrices

**Lemma 7.1** *Let  $[0, 1] \rightarrow \mathfrak{so}(N, \mathbb{R})$ ,  $t \mapsto A(t)$ , be a smooth ( $C^\infty$ ) curve of real skew-symmetric matrices satisfying that for each  $t \in [0, 1]$  :*

- a) *The kernel of  $A(t)$  is at most 2-dimensional, and*
- b)  *$A(t)$  has no double nonzero eigenvalue.*

*Then,  $A(t)$  is  $2 \times 2$  block-diagonalizable by a smooth curve*

$$[0, 1] \longrightarrow \text{SO}(N, \mathbb{R}) \quad t \longmapsto U(t).$$

*In the case  $N = 2n + 1$ , there is an extra  $1 \times 1$  block.*

**Proof.**

First, let us consider the map  $[0, 1] \rightarrow P_N(\mathbb{R})$ ,  $t \mapsto A^2(t)$ , where  $P_N(\mathbb{R})$  denotes the space of real  $N \times N$  symmetric matrices. By the assumptions a) and b), we have that for all  $t \in [0, 1]$ , the eigenvalues of  $A^2(t)$  are double (at most one is identically zero and it is simple, if  $N = 2n + 1$ ).

We take  $P_{A(t)}(\lambda)$  to be either the characteristic polynomial of  $A(t)$ , if  $N = 2n$ , or this polynomial divided by  $\lambda$ , if  $N = 2n + 1$ . Then,  $P_{A(t)}(\lambda)$  is a polynomial of degree  $n$  in  $X = \lambda^2$ .

All the roots of polynomial  $P_{A(t)}(X)$  are simple. Hence, for a root  $X_0$ , we can take a  $t_0 \in [0, 1]$ , such that in a neighborhood of  $t_0$ , there is a smooth real-valued function  $t \mapsto X(t)$  such that  $X(t_0) = X_0$ .

If  $X_0 < 0$ , then, in a neighborhood of  $t_0$  we have that  $\pm i\sqrt{X(t)}$ , is a smooth couple of distinct eigenvalue of  $A(t)$ . The only thing that might be unclear is that if  $X_0 = 0$  then  $\pm i\sqrt{X(t)}$  is smooth. In any case, we have at least  $n - 1$  positive smooth functions

$$[0, 1] \longrightarrow \mathbb{R}, \quad t \longmapsto \alpha_j(t),$$

such that  $\pm i\alpha_j(t)$  are simple eigenvalues of  $A(t)$ . These functions are smooth and *globally* defined on  $[0, 1]$ . Now fix one of these couples of eigenvalues  $\pm i\alpha_j(t)$  and consider the maps:

$$[0, 1] \longrightarrow P_N(\mathbb{R}), \quad t \longmapsto A^2(t) + \alpha_j^2(t)Id.$$

The kernels  $V_t$  of these matrices  $A^2(t) + \alpha_j^2(t)Id$ , are well defined, smooth and 2-dimensional. Then the set of the couples  $(t, V_t)$  defines a smooth sub-bundle of the trivial bundle

$$\Pi : [0, 1] \times \mathbb{R}^N \rightarrow [0, 1].$$

This sub-bundle is a vector bundle of 2-planes over  $[0, 1]$ . Then, it is trivial and possesses a smooth nonzero section  $t \rightarrow X_t$ . We can assume that  $\|X_t\| = 1$ . The restriction of the Euclidean metric  $g$  on  $\mathbb{R}^N$  to this bundle defines the following two smooth curves:

$$t \longmapsto \pm Y_t, \quad \text{such that} \quad \|Y_t\| = 1, \quad \langle X_t, Y_t \rangle_g = 0.$$

We choose one of these curves, say  $t \mapsto +Y_t$ , and then, we have a smooth map  $t \mapsto (X(t), Y(t))$ , defined on  $[0, 1]$ , such that for all  $t \in [0, 1]$ , the couple  $(X_t, Y_t)$  forms an orthonormal basis of the kernel:

$$\ker (A^2(t) + \alpha_j^2(t)Id).$$

For  $N = 2n + 1$ , we have also the mapping  $t \mapsto \bar{V}_t = \ker A^2(t)$ , which defines a bundle of lines over  $[0, 1]$ , that has also a nonzero smooth section  $t \rightarrow \bar{X}_t$ , with  $\|\bar{X}_t\| = 1$ . Now if  $N = 2n$ , it remains a single couple of eigenvalues,  $\pm i\alpha_n(t)$  such that  $\alpha_n^2(t)$  is smooth, but vanishes somewhere, and it is not clear that  $\alpha_n(t)$  is smooth. But then the space:

$$V_t = \ker (A^2(t) + \alpha_n^2(t)Id),$$

is still well defined and smooth (we may also get it as the one which is orthogonal to all the other smooth vector bundles).

Then, in the same way, we obtain a couple  $(X_t, Y_t)$  of smooth orthogonal vectors in this kernel. Now we have the mapping:

$$[0, 1] \longrightarrow \text{SO}(N), \quad t \longmapsto U(t),$$

which is smooth, and block-diagonalizes the curve  $A(t)$ . This shows moreover that  $t \mapsto \alpha_n(t)$  is smooth. ◇

In the analytic case, there is a much stronger result. Let us consider the real  $C^\omega$  map:

$$[0, 1] \longrightarrow \mathfrak{so}(N, \mathbb{R}), \quad t \longmapsto A(t).$$

It follows from known results in perturbation theory for linear operators, see for instance [12], that in the complex plane  $\mathbb{C}$ , there is a neighborhood  $V([0, 1]) \subset \mathbb{C}$  of the interval  $[0, 1]$ , and a holomorphic map,

$$V([0, 1]) \longrightarrow M_N(\mathbb{C}), \quad t \longmapsto U(t),$$

such that  $U(t)$  is unitary when restricted to  $[0, 1]$ , and diagonalizes  $A(t)$  for all  $t$ . The following lemma is an immediate consequence of these considerations.

**Lemma 7.2** *Given any  $C^\omega$  map*

$$[0, 1] \longrightarrow \mathfrak{so}(N), \quad t \longmapsto A(t).$$

*There is a  $C^\omega$  map*

$$[0, 1] \longrightarrow \text{SO}(N), \quad t \longmapsto U(t),$$

*that  $2 \times 2$  block-diagonalizes  $A(t)$  for all  $t$ . If  $N = 2n + 1$ , then there is an extra  $1 \times 1$  block.*

## 7.2 Distinguished sets of the Lie algebra of skew-symmetric matrices

We consider in this paragraph basic properties of certain distinguished sets of the Lie algebra  $\mathfrak{so}(N)$  of skew-symmetric matrices, that play a role in the proofs of our results. It is easy to prove the following general fact:

(F.) The subset  $\widehat{\mathcal{A}}_n \subset \mathfrak{so}(N)$  of matrices with kernel of codimension exactly  $n$ , is a sub-manifold of codimension  $\frac{n(n+1)}{2}$ , (not all  $n$  are possible for each  $N$ ).

As easy consequences of this result we have the following three properties.

- $F_1$ . The subset  $\mathcal{A}_1$  of  $\mathfrak{so}(2n)$  of matrices that have a nonzero kernel, is an algebraic closed set with codimension 1.
- $F_2$ . The subset  $\mathcal{A}_3$  of  $\mathfrak{so}(2n+1)$  of matrices that have a non-minimum kernel (i.e. not of dimension 1) is an algebraic closed set with codimension 3.
- $F_3$ . The subset  $\mathcal{A}_2$  of  $\mathfrak{so}(2N)$  of matrices that have a kernel of codimension  $> 2$  (i.e.  $\geq 4$ ) is algebraic, closed, and of codimension 6.

The following property is less straightforward and we shall provide a proof.

- $F_4$ . The subset  $\mathcal{A}_4$  of  $\mathfrak{so}(N)$  of matrices with a double nonzero eigenvalue is semi-algebraic, of codimension 3. Its closure  $\overline{\mathcal{A}}_4$  has the same codimension.

*Proof.* The fact that  $\mathcal{A}_4$  is semi-algebraic can be easily verified, we check only that the codimension of  $\mathcal{A}_4$  is  $\geq 3$ . In fact, this is the only result we use in the proofs. The set  $\mathcal{A}_4$  is the union of the orbits under conjugation in  $\text{SO}(N, R)$  of block-diagonal matrices of the form:

$$\begin{cases} \text{diag}(\alpha J, \alpha J, \alpha_3 J, \dots, \alpha_n J), & \text{if } N = 2n, \text{ and} \\ \text{diag}(\alpha J, \alpha J, \alpha_3 J, \dots, \alpha_n J, 0), & \text{if } N = 2n + 1, \end{cases}$$

with  $\alpha \neq 0$  and  $J = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$ .

Let  $\mathcal{D}$  be the set of such matrices, and let  $\widehat{\mathcal{A}}_4 \subset \mathcal{D}$  be the set of generic elements of  $\mathcal{D}$ , that is, matrices such that

$$\begin{cases} \text{diag}(\alpha_3 J, \dots, \alpha_n J), \\ \text{diag}(\alpha J, \alpha J, \alpha_3 J, \dots, \alpha_n J, 0), \end{cases}$$

is a regular element in  $\mathfrak{so}(N-4)$ , that further satisfies  $\alpha \neq 0$ ,  $\alpha \neq \alpha_j$  and  $\alpha_j \neq 0$ .

Then, for  $D \in \widehat{\mathcal{A}}_4$  an easy computation shows that  $\ker adD = \mathfrak{g} \times \mathfrak{c}_{N-4}$ , where  $\mathfrak{c}_{N-4}$  is a Cartan sub-algebra of  $\mathfrak{so}(N-4)$ , and  $\mathfrak{g}$  is the Lie algebra of matrices of the form:

$$\begin{pmatrix} 0 & a & c & d \\ -a & 0 & -d & c \\ -c & -d & 0 & e \\ d & -c & -e & 0 \end{pmatrix}.$$

Let now  $G$  be the 4-dimensional Lie group corresponding to  $\mathfrak{g}$ . The stabilizer  $S_1$  of  $D \in \widehat{\mathcal{A}}_4$  is the set  $S_1 = \mathbb{T}(N-4) \times G$ , where  $\mathbb{T}(N-4)$  is a maximal torus of  $\text{SO}(N-4)$ . Its dimension is  $\dim(S_1) = 4 + n - 2 = n + 2$ . Therefore, the orbit under conjugation of  $D$  has dimension  $M - n - 2$ , where

$$M = \frac{N(N-1)}{2}.$$

Stabilizers of non-generic points of  $\mathcal{D}$  are bigger, hence the orbits of these points have lower dimension. At the end, all of these orbits under conjugation of the elements of  $\mathcal{D}$  have dimension  $\leq M - n - 2$ . Since the dimension of  $\mathcal{D}$  is  $n - 1$ , it follows that the dimension of  $\mathcal{A}_4$  is smaller than  $M - n - 2 + n - 1 = M - 3$ . Hence:

$$\text{codim } \mathcal{A}_4 \geq 3.$$

◇

### Acknowledgments

The authors want to thank Frédéric Jean for the fruitful discussions and for his thorough observations about our results.

The present paper was written while the third author (Monroy-Pérez) was on sabbatical leave at INRIA, Unité de Recherche Sophia-Antipolis, France. The author thanks Prof. J.B. Pomet and the rest of the team MIAOU for their support. He was also partly supported by the CONACYT's grant for sabbatical periods abroad.

## References

- [1] Abraham R, Robbin, J., *Transversal mappings and flows*, W. A. Benjamin, Inc. , 1967.
- [2] Hirsch, M. W., *Differential Topology*, Springer Verlag, N.Y. , 1976.
- [3] Goreski, Mc Pherson, *Stratified Morse Theory*, Springer Verlag, N.Y., 1988.
- [4] Chakir, El-A., El-H., Gauthier, J. P., Kupka, I., Small Sub-Riemannian balls on  $R^3$ , in *Journal and Dynamical Control Systems*, Vol. 2, No. 3, pp. 359-421, 1996.
- [5] Agrachev, A., Chakir El-A., El-H.; Gauthier, J. P., Sub-Riemannian metrics on  $R^3$ , in *Geometric Control and non-holonomic mechanics, Mexico City, 1996*, pp. 29-76, Canadian Mathematical Society Conf. Proc., 25, Amer. Math. Soc., Providence, RI, 1998.
- [6] Agrachev, A., Gauthier, J. P., Sub-Riemannian Metrics and Isoperimetric Problems in the Contact case, L.S. Pontriaguine, 90<sup>th</sup> Birthday Commemoration, *Contemporary Mathematics*, Tome 64, pp. 5-48, 1999 (Russian). English version: *Journal of Mathematical Sciences*, Vol. 103, No. 6, pp. 639-663.
- [7] Jean, F., Complexity of nonholonomic motion planning, *International Journal of Control*, Vol. 74, No. 8, pp. 776-782, 2001.
- [8] Jean, F., Entropy and Complexity of a Path in Sub-Riemannian Geometry, to appear in *COCV*.
- [9] Jean, F., Falbel, E., Measures and transverse paths in Sub-Riemannian Geometry, to appear in *Journal d'Analyse mathématique*
- [10] Charlot, G., Quasi-Contact sub-Riemannian Metrics, Normal Form in  $R^{2n}$ , Front and Caustic in  $R^4$ , to appear in *Acta Applicandae Mathematicae*.
- [11] Gromov, M. Carnot-Caratheodory spaces seen from within, in *Sub-Riemannian geometry*, Eds. A. Bellaïche, J.J. Risler, Birkhauser, p.79-323, 1996.
- [12] Kato, T., *Perturbation theory for linear operators*, Springer Verlag, p.120-122, 1966.



## Contents

<b>1</b>	<b>Introduction, notations and organization of the paper</b>	<b>3</b>
1.1	Definitions . . . . .	3
1.2	The fundamental 2-form. . . . .	4
1.3	Statement of our results and comments . . . . .	5
1.4	Organization of the paper. . . . .	6
<b>2</b>	<b>Genericity results</b>	<b>7</b>
<b>3</b>	<b>Normal Coordinates and Normal Forms</b>	<b>9</b>
3.1	Normal Coordinates . . . . .	9
3.2	Normal Forms . . . . .	11
3.3	Special Case: normal form in dimension 3 . . . . .	15
<b>4</b>	<b>Complexity and asymptotic optimal synthesis</b>	<b>15</b>
<b>5</b>	<b>Martinet Case</b>	<b>19</b>
5.1	Proof of the Theorem 1.3 . . . . .	19
5.2	An example. . . . .	21
<b>6</b>	<b>The non generic case.</b>	<b>23</b>
6.1	Normal form . . . . .	23
6.2	Metric Complexity . . . . .	24
6.3	Asymptotic optimal synthesis . . . . .	24
<b>7</b>	<b>Appendix</b>	<b>25</b>
7.1	Diagonalization of 1-parameter families of skew-symmetric matrices . . . . .	25
7.2	Distinguished sets of the Lie algebra of skew-symmetric matrices . . . . .	27



---

Unité de recherche INRIA Sophia Antipolis  
2004, route des Lucioles - BP 93 - 06902 Sophia Antipolis Cedex (France)

Unité de recherche INRIA Futurs : Parc Club Orsay Université - ZAC des Vignes  
4, rue Jacques Monod - 91893 ORSAY Cedex (France)

Unité de recherche INRIA Lorraine : LORIA, Technopôle de Nancy-Brabois - Campus scientifique  
615, rue du Jardin Botanique - BP 101 - 54602 Villers-lès-Nancy Cedex (France)

Unité de recherche INRIA Rennes : IRISA, Campus universitaire de Beaulieu - 35042 Rennes Cedex (France)

Unité de recherche INRIA Rhône-Alpes : 655, avenue de l'Europe - 38334 Montbonnot Saint-Ismier (France)

Unité de recherche INRIA Rocquencourt : Domaine de Voluceau - Rocquencourt - BP 105 - 78153 Le Chesnay Cedex (France)

---

Éditeur  
INRIA - Domaine de Voluceau - Rocquencourt, BP 105 - 78153 Le Chesnay Cedex (France)  
<http://www.inria.fr>  
ISSN 0249-6399