

Comparing multifractal formalisms: the Neighboring Boxes Condition

Julien Barral, Fathi Ben Nasr, Jacques Peyrière

► **To cite this version:**

Julien Barral, Fathi Ben Nasr, Jacques Peyrière. Comparing multifractal formalisms: the Neighboring Boxes Condition. [Research Report] RR-4562, INRIA. 2002. inria-00072026

HAL Id: inria-00072026

<https://hal.inria.fr/inria-00072026>

Submitted on 23 May 2006

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Comparing multifractal formalisms: the Neighboring Boxes Condition

Julien Barral — Fathi Ben Nasr — Jacques Peyrière

N° 4562

Septembre 2002

THÈME 4



*Rapport
de recherche*

Comparing multifractal formalisms: the Neighboring Boxes Condition

Julien Barral ^{*}, Fathi Ben Nasr [†], Jacques Peyrière [‡]

Thème 4 — Simulation et optimisation
de systèmes complexes
Projet Fractales

Rapport de recherche n° 4562 — Septembre 2002 — 23 pages

Abstract: Physicists usually compute dimensions by using boxes and they also do so when dealing with multifractals. Also in the study of some dynamical systems and multiplicative processes, boxes naturally appear. On the other hand, in geometric measure theory, it is preferred to perform computations which do not depend on a grid.

This paper provides a bridge between the boxes and the grid-free approaches to the multifractal analysis of measures. Results for quasi-Bernoulli measures and statistically self-similar measures are obtained.

Key-words: Hausdorff dimension, Packing dimension, Multifractal formalism for measures, Statistically self-similar measures.

^{*} Projet Fractales, INRIA Rocquencourt, 78153 Le Chesnay Cedex, FRANCE. This article was completed while visiting Yale University. {Julien.Barral@inria.fr}

[†] Département de Mathématiques, Faculté des Sciences de Monastir, Monastir 5000, TUNISIE. {Fathi.BenNasr@fsm.rnu.tn}

[‡] Yale University, Department of Mathematics, PO Box 208283, New Haven, CT 06520-8283, USA; Permanent address: Université Paris-Sud, Mathématiques bât. 425, CNRS UMR 8628, 91405 Orsay Cedex, FRANCE. {Jacques.Peyriere@u-psud.fr}

Analyses multifractales: des “boîtes” au formalisme centré.

Résumé : Les physiciens utilisent les boîtes d’une grille fine pour estimer la dimension d’un ensemble, et plus généralement le spectre multifractal d’une mesure. Les structures dépendant d’une grille apparaissent également naturellement dans certains systèmes dynamiques et certains processus multiplicatifs. D’autre part, en théorie géométrique de la mesure, on préfère calculer un spectre multifractal défini sans avoir recours à une grille. C’est ce qui a amené Olsen à développer un formalisme multifractal centré.

Ce travail donne des conditions suffisantes sur une mesure pour que la validité du formalisme pour une grille entraîne celle du formalisme centré. On obtient notamment la validité du formalisme d’Olsen pour les mesures quasi-Bernoulli et les mesures statistiquement auto-similaires associées à une grille c -adique.

Mots-clés : Dimension de Hausdorff, Dimension de Packing, Formalisme multifractal pour les mesures, Mesures statistiquement auto-similaires.

Contents

1	Introduction	4
2	Box analysis	4
2.1	Trees and weighted trees	4
2.2	Homogeneous trees	6
3	A centered multifractal formalism	7
4	Comparing multifractal analyses	9
4.1	Stating the problem	9
4.2	The comparison theorems	10
5	Quasi-Bernoulli measures.	15
6	Statistically self-similar measures.	18

1 Introduction

The multifractal analysis of a measure μ aims at relating the dimension of the level sets of the pointwise Hölder exponent of μ to the Legendre transform of some kind of entropy or free energy function, a problem initially raised and studied for physical motivations ([19, 17, 18, 25, 26]).

To define these pointwise Hölder exponents one has two alternatives: to define $\alpha(x)$ as the limit, when it exists, of either the ratio $\frac{\log \mu(B(x, r))}{\log r}$ when r goes to 0, or the ratio

$\frac{\log \mu(Q_n(x))}{\log \text{diam}(Q_n(x))}$ when n goes to $+\infty$ (where $B(x, r)$ stands for the ball of radius r centered at x and $Q_n(x)$ stands for the c -adic box of size c^{-n} which contains x). Of course the partition function is defined in terms of covers or packings by balls in the former case, by boxes in the latter case. It is usual to observe connections between these two approaches when μ possesses self-similarity properties and μ is supported by a regular enough Cantor set ([4, 9, 13, 34, 15, 16, 31, 1]). But there is no a priori reason why these two approaches should be connected in full generality. In this work, we give a condition ensuring that if a measure obeys the “box formalism”, then it also obeys the other one. Our results apply on two families of measures supported by the full c -adic grid of $[0, 1]$, namely the quasi-Bernoulli measures and the Mandelbrot measures.

The so called “box formalism” is better explained in the abstract setting of trees. This is the matter of the next section.

In Section 3, the Olsen multifractal formalism is recalled for the reader’s convenience. The main comparison theorem is stated and proven in Section 4. This is explained in the one dimensional case, for the sake of simplicity, but at the end of this section it is said how to deal with higher dimensions. Section 5 deals with quasi-Bernoulli measures, and Section 6 with the Mandelbrot multiplicative measures.

2 Box analysis

2.1 Trees and weighted trees

Let \mathcal{T} be a locally finite rooted tree. If w is a node, we denote by $F(w)$ the set of nodes which immediately follow w . Let \mathcal{T}_n denote the set of nodes whose geodesic distance from the root equals n . There is a natural topology on the set $\partial\mathcal{T}$ of geodesics stemming from the root which makes of $\partial\mathcal{T}$ a metrizable totally disconnected compact space.

We identify a node of \mathcal{T} with the set of geodesics going through it and the root. These sets are open and closed and they are the balls of an ultrametric distance defining the topology. A Borel measure on \mathcal{T} can be identified with a function μ from the nodes of \mathcal{T} to $[0, +\infty)$

subject to the condition

$$\mu(v) = \sum_{w \in F(v)} \mu(w) \quad \text{for all } v \in \mathcal{T}.$$

From now on, we are given a continuous probability measure ξ on $\partial\mathcal{T}$ whose support is the whole $\partial\mathcal{T}$.

If μ is a measure on $\partial\mathcal{T}$, one defines

$$C_n^\mu(q, t) = \sum_{w \in \mathcal{T}_n}^* \mu(w)^q \xi(w)^t, \quad (1)$$

where the star means that the terms for which $\mu(w) = 0$ are removed from the sum — a convention valid throughout this article —, and

$$\tau_\mu(q) = \sup\{t \in \mathbb{R} \mid \limsup_{n \rightarrow \infty} C_n^\mu(q, t) = +\infty\}.$$

Since C_n^μ is a log-convex function, the function τ_μ is easily seen to be convex and non-increasing.

On the other hand, for any $\alpha \in \mathbb{R}$, one considers the set

$$E_\alpha = \left\{ x \in \partial\mathcal{T} \mid \lim_{\substack{\xi(w) \rightarrow 0 \\ x \in w}} \frac{\log \mu(w)}{\log \xi(w)} = \alpha \right\}.$$

Let \dim_ξ be the Hausdorff dimension defined by using $\xi(w)$ instead of the diameter of w (see [8]).

Then it is known (for instance, see [11]) that, for each q for which $\tau_\mu'(q)$ exists, one has

$$\dim_\xi E_{-\tau_\mu'(q)} \leq \tau_\mu^*(-\tau_\mu'(q)),$$

where τ_μ^* is the Legendre transform of τ_μ , *i.e.* $\tau_\mu^*(t) = \inf_{q \in \mathbb{R}} qt + \tau_\mu(q)$. Indeed, the stronger inequality [11]

$$\text{Dim}_\xi E_{-\tau_\mu'(q)} \leq \tau_\mu^*(-\tau_\mu'(q)),$$

where Dim_ξ stands for the packing dimension (defined in [36]), is almost an instance of the Chernoff formula [12]. Of course, if these inequalities lead to negative dimensions, this means that the corresponding set is empty.

The formula

$$\dim_\xi E_{-\tau_\mu'(q)} = \tau_\mu^*(-\tau_\mu'(q)) \quad (2)$$

is shown [11] to hold if there exists a measure μ_q , called a Gibbs measure, such that there exists $C > 0$ such that, for all $w \in \mathcal{T}$,

$$\frac{1}{C} \mu(w)^q \xi(w)^{\tau_\mu(q)} \leq \mu_q(w) \leq C \mu(w)^q \xi(w)^{\tau_\mu(q)} \quad (3)$$

Indeed, the equality (2) appeared, in a non rigorous mathematical form, in [18] and was proven to hold for some examples [9, 13].

It is noticed in [33] that these results still hold when the fonction C_n above is defined as the supremum of the corresponding sums on the sections of \mathcal{T} which lie below level n (in other terms, one considers packings of $\partial\mathcal{T}$ by ultrametric balls). Let us call the corresponding separator function φ instead of τ .

Indeed, as shown in [5], only the right hand side inequality in (3) is needed. Since this is reminiscent of the Frostman lemma, it was natural to consider generalized Hausdorff measures: for q and t in \mathbb{R} , and $A \subset \partial\mathcal{T}$, define

$$\mathcal{H}_\mu^{q,t}(A) = \liminf_{\delta \searrow 0} \left\{ \sum \mu(w_j)^q \xi(w_j)^t \mid w_j \in \mathcal{T}, A \subset \bigcup w_j, \xi(w_j) \leq \delta \right\}.$$

Lemma 1 ([5]). *If a compact set A is such that $\mathcal{H}_\mu^{q,t}(A) > 0$, then it carries a probability measure ν satisfying $\nu(w) \leq C \mu(w)^q \xi(w)^t$ for all w .*

This lemma implies the following theorem.

Theorem 1 ([5]). *Suppose that $\varphi'_\mu(q)$ exists and that $\mathcal{H}_\mu^{q,\varphi_\mu(q)}(\text{supp } \mu) > 0$. Then one has $\dim_\xi E_{-\varphi'_\mu(q)} = \varphi_\mu^*(-\varphi'_\mu(q))$.*

2.2 Homogeneous trees

In this section, we suppose that the tree \mathcal{T} is homogeneous of order c and that ξ is uniformly distributed. In this context, it is more convenient to see \mathcal{T} as the free monoid on a c -letter alphabet \mathcal{A} .

So, let $\mathcal{A}^* = \bigcup_{n \geq 0} \mathcal{A}^n$ be the free monoid consisting of words on \mathcal{A} endowed with the concatenation. The operation of concatenation will be simply denoted by juxtaposition, and by a dot, when this is necessary for a better understanding. The empty word ϵ is the identity element. The length of a word w is denoted by $|w|$. If a word v is a prefix of the word w , we write $v \prec w$. This defines an order on \mathcal{A}^* . Endowed with this order, \mathcal{A} is a tree the root of which is ϵ . If v and w are words, $v \wedge w$ stands for their largest common prefix. The quantity $d(v, w) = c^{-|v \wedge w|}$ defines an ultrametric distance on \mathcal{A}^* .

The completion $\widehat{\mathcal{A}^*}$ of (\mathcal{A}^*, d) is a compact space which is the disjoint union of \mathcal{A}^* and $\partial\mathcal{A}^*$. The elements of $\partial\mathcal{A}^*$ can be viewed as the infinite sequences of elements of \mathcal{A} . The notion of prefix extends in a natural way to elements of $\partial\mathcal{A}^*$.

We identify $w \in \mathcal{A}^*$ with the cylinder $\{x \in \partial\mathcal{A}^* \mid w \prec x\}$, so a Borel measure μ on $\partial\mathcal{A}^*$ is a mapping from \mathcal{A}^* to \mathbb{R}^+ fulfilling the following compatibility condition

$$\mu(w) = \sum_{a \in \mathcal{A}} \mu(wa) \quad \text{for all } w \in \mathcal{A}^*.$$

At last, the measure ξ is so defined: $\xi(w) = c^{-|w|}$.

In this setting, the τ_μ function can be defined as follows.

$$\tau_\mu(q) = \limsup_{n \rightarrow +\infty} \frac{1}{n} \log_c \sum_{w \in \mathcal{A}^n}^* \mu(w)^q,$$

where q is a real number and the star means, as already said, that the summation runs only on the w having a non-zero measure. This function τ_μ is convex and non-increasing as observed previously.

3 A centered multifractal formalism

Let (\mathcal{X}, d) be a metric space and μ a positive atomless Borel measure on \mathcal{X} . The support of μ is denoted by $\text{supp } \mu$.

According to Olsen [32], we define several measures and premeasures indexed by a couple (q, t) of real numbers. If E is a subset of \mathcal{X} and δ is a positive real number, we set

$$\overline{\mathcal{P}}_{\mu, \delta}^{q, t}(E) = \sup \sum^* \mu(B(x_j, r_j))^q r_j^t,$$

this supremum being taken over the collections $\{B(x_j, r_j)\}$ of mutually disjoint balls whose centers x_j belong to E and whose radii r_j are less than δ . The star means that we omit in the summation the terms which are obviously infinite (*i.e.* zero raised to a negative power). As previously said, this convention holds throughout this paper. Of course, as long as the measure μ has no punctual masses, it does not matter that balls be open or closed. In the sequel, we deal with such measures and, it will be convenient for our reasonings to consider only closed balls.

We consider the limit

$$\overline{\mathcal{P}}_\mu^{q, t}(E) = \lim_{\delta \searrow 0} \overline{\mathcal{P}}_{\mu, \delta}^{q, t}(E).$$

The function $\overline{\mathcal{P}}_\mu^{q, t}$ is called *packing pre-measure*. It lacks σ -subadditivity to be a Caratheodory outer measure. This is why one considers the following quantity

$$\mathcal{P}_\mu^{q, t}(E) = \inf_{E \subset \bigcup E_j} \sum \overline{\mathcal{P}}_\mu^{q, t}(E_j),$$

which, as a function of E , is an outer measure. (This is the same process as for defining packing measures, which were introduced in [36].)

In a similar way, one defines Hausdorff-like measures.

$$\overline{\mathcal{H}}_{\mu, \delta}^{q, t}(E) = \inf \sum^* \mu(B(x_j, r_j))^q r_j^t,$$

this infimum being taken over the coverings $\{B(x_j, r_j)\}$ of E by balls whose centers x_j belong to E and whose radii r_j are less than δ , and consider the limit

$$\overline{\mathcal{H}}_\mu^{q, t}(E) = \lim_{\delta \searrow 0} \overline{\mathcal{H}}_{\mu, \delta}^{q, t}(E).$$

In order to deal with an outer measure, one defines

$$\mathcal{H}_\mu^{q,t}(E) = \sup_{F \subset E} \overline{\mathcal{H}}_\mu^{q,t}(F).$$

These last measures are the multifractal counterparts of the centered Hausdorff measures introduced in [35].

For a fixed q , if, for some t one has $\overline{\mathcal{P}}_\mu^{q,t}(\text{supp } \mu) < +\infty$, then, for all $t' > t$, one has $\overline{\mathcal{P}}_\mu^{q,t'}(\text{supp } \mu) = 0$. Therefore, there exists a unique $\Lambda_\mu(q) \in \overline{\mathbb{R}}$ such that $\overline{\mathcal{P}}_\mu^{q,t}(\text{supp } \mu)$ is infinite if $t < \Lambda_\mu(q)$ and zero if $t > \Lambda_\mu(q)$.

In a similar way, two functions B_μ and b_μ are associated respectively to $\mathcal{P}_\mu^{q,t}$ and $\mathcal{H}_\mu^{q,t}$.

All these three functions are non-increasing; Λ_μ and B_μ are convex. It is also clear that $b_\mu \leq B_\mu \leq \Lambda_\mu$.

If α and β are two real numbers such that $\alpha \leq \beta$, one considers the following sets

$$X_\mu(\alpha, \beta) = \left\{ x \mid \alpha \leq \liminf_{r \searrow 0} \frac{\log \mu(B(x, r))}{\log r} \leq \limsup_{r \searrow 0} \frac{\log \mu(B(x, r))}{\log r} \leq \beta \right\}.$$

Instead of $X_\mu(\alpha, \alpha)$, we shall simply write $X_\mu(\alpha)$.

If the derivative of B_μ exists at point q , it is known [32] that the following inequalities hold, with the convention that a set of negative dimension¹ is empty,

$$\begin{aligned} \dim X_\mu(-B'_\mu(q)) &\leq b_\mu^*(-B'_\mu(q)) \\ \text{Dim } X_\mu(-B'_\mu(q)) &\leq B_\mu^*(-B'_\mu(q)), \end{aligned} \tag{4}$$

where, as previously, the star as an exponent denotes the Legendre transform —i.e. $f^*(\alpha) = \inf_{q \in \mathbb{R}} \alpha q + f(q)$ — and where \dim and Dim stand for the Hausdorff and packing dimensions.

Definition. If $B'_\mu(q)$ exists and if all the quantities in (4) are equal, one says that the measure μ obeys the multifractal formalism at point q .

Before recalling the following theorem [6], we need another definition.

Definition. A metric space (X, d) is said to have the Besicovitch covering property if there exists a positive integer β such that, given any collection $\{B(x_i, r_i)\}_{i \in I}$ of balls, one can extract from it β packings which altogether cover the set $\{x_i\}_{i \in I}$.

Any euclidean space has this property, as well as, of course, any ultrametric space.

Theorem 2. *If (X, d) has the Besicovitch covering property, if $\alpha = -B'_\mu(q)$ exists, and if $\mathcal{H}_\mu^{q, B_\mu(q)}(\text{supp } \mu) > 0$, then*

$$\dim X_\mu(\alpha) = \text{Dim } X_\mu(\alpha) = B_\mu^*(\alpha) \quad \text{and} \quad b_\mu(q) = B_\mu(q).$$

¹For a further interpretation of negative dimensions, see [26, 27].

4 Comparing multifractal analyses

4.1 Stating the problem

Lemma 2. *Let μ be a non-zero positive measure on $\partial\mathcal{A}^*$. Then one has $\tau_\mu = \Lambda_\mu$.*

Proof. We begin by noticing that, if $t > \tau_\mu(q)$, one has

$$\sum_{w \in \mathcal{A}^*}^* \mu(w)^q c^{-t|w|} < +\infty \quad (5)$$

It results that, if $t > \tau_\mu(q)$, $\overline{\mathcal{P}}_\mu^{q,t}(\text{supp } \mu)$ is finite, and therefore $t \geq \Lambda_\mu(q)$. This proves $\Lambda_\mu \leq \tau_\mu$. The converse inequality is obvious.

The equality asserted in this lemma is similar to the equivalence of two different definitions of the quantity called $\Delta(E)$ in [36].

The map γ

There is a natural map from $\partial\mathcal{A}^*$ onto \mathbb{R} : let us take $\mathcal{A} = \{0, 1, 2, \dots, c-1\}$ and consider the map γ which sends the element $x = a_1 a_2 \cdots a_n \cdots$ on the number $\sum_{n \geq 1} a_n c^{-n}$. This map sends cylinders onto c -adic intervals, and when a measure μ on $\partial\mathcal{A}^*$ has no point masses, it is equivalent in many problems to consider it or its image $\nu = \gamma_*(\mu)$ under γ .

This rises the natural question of deciding when the multifractal analysis of ν (in $(\mathbb{R}, |\cdot|)$) and μ (in $(\partial\mathcal{A}^*, d)$) are linked. Indeed, there are no reasons (even when ν is doubling) why B_μ and B_ν , for instance, should coincide. There are no reasons either that $X_\nu(\alpha, \beta)$ should be the image of $X_\mu(\alpha, \beta)$ under γ . Nevertheless, one has the following fact.

Lemma 3. *One has $\Lambda_\nu \leq \tau_\mu$.*

Proof. Consider first the case $q \leq 0$. Let $\{B(x_j, r_j)\}$ be a centered packing of $\text{supp } \nu$, and $t > \tau_\mu(q)$. Each $B(x_j, r_j)$ contains a c -adic interval I_j such that $x_j \in I_j$, $|I_j| > r_j/c$ (where $|I_j|$ stands for the length of I_j), and $\nu(I_j) > 0$ (do not forget that $x_j \in \text{supp } \nu$). Therefore

$$\begin{aligned} \sum \nu(B(x_j, r_j))^q r_j^t &\leq c^{t^+} \sum \nu(I_j)^q |I_j|^t \\ &\leq c^{t^+} \sum_{w \in \mathcal{A}^*}^* \mu(w)^q c^{-t|w|} < +\infty, \end{aligned}$$

where $t^+ = \max\{t, 0\}$. This yields $\mathcal{P}_\nu^{q,t}(\text{supp } \nu) < \infty$. Consequently $\Lambda_\nu(q) \leq \tau_\mu(q)$.

Now, consider the case $q > 0$. Let $\{B(x_j, r_j)\}$ be a centered packing of $\text{supp } \nu$, and $t > \tau_\mu(q)$. Each $B(x_j, r_j)$ is covered by at most $c+1$ c -adic intervals $I_{j,1}, I_{j,2}, \dots, I_{j,k_j}$ of non-zero ν -measure and of common length l satisfying $l \leq 2r_j < cl$. One has

$$\nu(B(x_j, r_j))^q \leq k_j^{(q-1)^+} \sum_{k=1}^{k_j} \nu(I_{j,k})^q$$

and $r_j^t \leq 2^{-t} c^{t^+} |I|^t$. Observe also that a c -adic interval can appear at most twice as an $I_{j,k}$, since $\{B(x_j, r_j)\}$ is a packing. Therefore

$$\begin{aligned} & \sum \nu(B(x_j, r_j))^q r_j^t \\ & \leq 2^{1-t} (c+1)^{(q-1)^+} c^{t^+} \sum_{w \in \mathcal{A}^*}^* \mu(w)^q c^{-t|w|} < +\infty. \end{aligned}$$

This proves again that $\Lambda_\nu(q) \leq \tau_\mu(q)$.

In the next section, we give a conditions which ensure that the multifractal analysis of μ and ν are equivalent. In the subsequent sections, we analyse some examples.

4.2 The comparison theorems

As recalled in Lemma 1 the inequality $\mathcal{H}_\mu^{q,t}(\text{supp } \mu) > 0$ (where μ is a measure on $\partial\mathcal{A}^*$) is equivalent to the existence of a measure $\mu_{q,t}$ with the property that, for a suitable $C > 0$, one has $\mu_{q,t}(B) \leq C\mu(B)^q |B|^t$, for any ball B of radius $|B|$ small enough. It is convenient to set the following definition.

Definition. Let μ be a measure on a metric space \mathcal{X} and q and t be two real numbers. A non-zero measure μ' having the property that there exists two positive numbers C and η such that, for all $x \in \text{supp } \mu$ and $r \leq \eta$, one has $\mu'(B(x, r)) \leq C\mu(B(x, r))^q r^t$ is called a *Frostman measure* for μ at (q, t) .

From now on we stick to the convention that μ (with or without subscript) is a measure on $\partial\mathcal{A}^*$ and that ν (with or without subscript) stands for the corresponding image measure under γ .

As, via γ , the elements of \mathcal{A}^n correspond to the c -adic intervals of length c^{-n} contained in $[0, 1]$, one can assign to each $w \in \mathcal{A}^n$ a number $i(w)$ so as to have $\gamma(w) = [i(w)c^{-n}, (i(w)+1)c^{-n}]$. Finally, if the words v and w have the same length, we set $\delta(v, w) = |i(v) - i(w)|$.

If $\nu = \gamma_*(\mu)$ has no masses at the endpoints of $\gamma(w)$, then $\nu(\gamma(w)) = \mu(w)$. But, one always has
$$\nu(w) \leq \sum_{\substack{v \in \mathcal{A}^* \\ \delta(v,w) \leq 1}} \mu(v).$$

From now on, in this section, we deal with an atomless measure μ on $\partial\mathcal{A}^*$ and its image ν under γ .

Lemma 4. *Let $q \in \mathbb{R}$ and $\eta > 0$ be two numbers. Assume there exists a Frostman measure μ_q for μ at $(q, \tau_\mu(q))$. Let $\nu_q = \gamma_*(\mu_q)$ stand for the image of μ_q under γ . Then we have the following facts.*

1. When $q < 0$, if $c^{-(n+1)} < 2r \leq c^{-n}$, one has

$$\int_{\text{supp } \nu} \left[\nu(B(x, r))^{-q} \nu_q(B(x, r)) \right]^\eta d\nu_q(x) \leq C c^{-n(1+\eta)\tau_\mu(q)} \sum_{\substack{v, w \in \mathcal{A}^n \\ \delta(v, w) \leq 3}}^* \mu(v)^{(1+\eta)q} \mu(w)^{-\eta q}.$$

2. When $q \geq 0$, if $c^{-n} \leq r < c^{1-n}$, one has

$$\int_{\text{supp } \nu} \left[\nu(B(x, r))^{-q} \nu_q(B(x, (1 + \frac{1}{c})r)) \right]^\eta d\nu_q(x) \leq C c^{-n(1+\eta)\tau_\mu(q)} \sum_{\substack{v, w \in \mathcal{A}^n \\ \delta(v, w) \leq 4c+2}}^* \mu(v)^{(1+\eta)q} \mu(w)^{-\eta q}.$$

In both cases the constant C depends only on q , η , and c .

Proof. We begin by assuming $q < 0$. By the Besicovitch covering lemma, we extract from the collection $\{B(x, r)\}_{x \in \text{supp } \nu}$ of balls β (a universal constant) packings which altogether cover $\text{supp } \nu$.

One has

$$\int_{B(a, r) \cap \text{supp } \nu} \left[\nu(B(x, r))^{-q} \nu_q(B(x, r)) \right]^\eta d\nu_q(x) \leq \nu(B(a, 2r))^{-\eta q} \nu_q(B(a, 2r))^{1+\eta}.$$

The interval $B(a, 2r)$ is contained in at most two c -adic consecutive intervals I_1 and maybe I_2 of common length l such that $l/c < 2r \leq l$. We add to this collection of one or two intervals the two c -adic intervals of the same length which are contiguous. We keep only the ones having a non-zero ν -measure. Then we have intervals J_1, J_2, J_3 , and maybe J_4 such that, if I is any of the intervals I_1 or I_2 , $\nu_q(I)$ is majorized by the sum $\sum \mu_q(J_i)$. We had to adjunct the right and left intervals to cope with the case where ν_q has masses at some c -adic points. In these conditions, we have

$$\begin{aligned} \nu(B(a, 2r))^{-\eta q} \nu_q(B(a, 2r))^{1+\eta} &\leq C_1 \left(\sum \nu(J_j)^{-q\eta} \right) \left(\sum \nu_q(J_j)^{1+\eta} \right) \\ &\leq C_1 \left(\sum \nu(J_j)^{-q\eta} \right) \left(\sum \nu(J_j)^{(1+\eta)q} |J_j|^{(1+\eta)\tau_\mu(q)} \right) \\ &\leq C_2 c^{-n(1+\eta)\tau_\mu(q)} \sum \nu(J_i)^{-q\eta} \nu(J_j)^{(1+\eta)q}, \end{aligned}$$

where the constants C_1 and C_2 depend only on c , q , and η .

Now, consider the case $q \geq 0$. For every $x \in \text{supp } \nu$, $B(x, r)$ contains at least one c -adic interval I_x of length c^{-n} such that $\mu(I_x) > 0$ and $c^{-n} \leq r < c^{1-n}$.

Fix $a \in \text{supp } (\mu)$ and consider the set of closed c -adic intervals of length c^{-n} the interior of which intersect $B(a, (2 + \frac{1}{c})r)$, and, as previously, add to them two intervals of the same generation, one on the right, the other on the left. The number of elements of the set S of these intervals is bounded by $4c + 3$. Therefore

$$\begin{aligned}
& \int_{B(a,r) \cap \text{supp } \nu} \left[\nu(B(x,r))^{-q} \nu_q \left(B \left(x, \frac{(c+1)r}{c} \right) \right) \right]^\eta d\nu_q(x) \\
& \leq \int_{B(a,r) \cap \text{supp } \nu} \left[\nu^{-q}(I_x) \nu_q \left(B \left(x, \frac{(c+1)r}{c} \right) \right) \right]^\eta d\nu_q(x) \\
& \leq \left[\sum_{I \in S} \nu^{-q}(I) \right]^\eta \int_{B(a,r) \cap \text{supp } \nu} \nu_q \left(B \left(x, \frac{(c+1)r}{c} \right) \right)^\eta d\nu_q \\
& \leq \left[\sum_{I \in S} \nu^{-q}(I) \right]^\eta \nu_q \left(B \left(a, \left(2 + \frac{1}{c} \right) r \right) \right)^{1+\eta} \\
& \leq C \left[\sum_{I \in S} \nu^{-q\eta}(I) \right] \left[\sum_{I \in S} \nu_q(I)^{1+\eta} \right].
\end{aligned}$$

The proof ends as previously.

Definition (NBC). A measure μ on $\partial\mathcal{A}^*$ is said to satisfy the Neighboring Boxes Condition (NBC) for $q \in \mathbb{R}$ if, for all $\varepsilon > 0$, there exists $\eta > 0$ such that

$$\sum_{n>0} c^{-n(\tau_\mu(q)+\varepsilon\eta)} \sum_{\substack{v,w \in \mathcal{A}^n \\ \delta(v,w) \leq c'}} \mu(v)^{-\eta q} \mu(w)^{(1+\eta)q} < +\infty, \quad (6)$$

where $c' = 3$ if $q < 0$, $c' = 4c + 2$ otherwise.

Lemma 5. Let $q \in \mathbb{R}$. Assume there exists a Frostman measure μ_q for μ at $(q, \tau_\mu(q))$ and that μ fulfills the NBC for q . Then, we have $b_\nu(q) = \tau_\mu(q)$.

Also, there exists a constant $C > 0$ such that, for any $\varepsilon > 0$, for ν_q -almost every x , and for r small enough, $\nu_q(B(x,r)) \leq C \nu(B(x,r))^q r^{\tau_\mu(q)-\varepsilon}$.

Proof. First consider the case $q < 0$. Set

$$F(r) = \int_{\text{supp } \nu} \left[\nu(B(x,r))^{-q} \nu_q(B(x,r)) \right]^\eta d\nu_q(x).$$

For any $\varepsilon > 0$, we have

$$\sum_{n \geq 0} F(c^{-n}) c^{n\eta(\tau_\mu(q) - \varepsilon)} \leq C \sum_{n \geq 0} c^{-n(\tau_\mu(q) + \eta\varepsilon)} \sum_{\substack{v, w \in \mathcal{A}^n \\ \delta(v, w) \leq c'}} \mu(v)^{-\eta q} \mu(w)^{(1+\eta)q},$$

due Lemma 4. The right hand side quantity is finite for a suitable choice of η , because of the NBC.

As a consequence of the Borel-Cantelli lemma, for ν_q -almost every x , one has

$$\nu_q(B(x, c^{-n})) \leq \nu(B(x, c^{-n}))^q c^{-n(\tau_\mu(q) - \varepsilon)}$$

for n large enough. Consequently for ν_q -almost every x , one has

$$\nu_q(B(x, r)) \leq C \nu(B(x, r))^q r^{\tau_\mu(q) - \varepsilon}$$

for r small enough.

Similarly, when $q \geq 0$, for ν_q -almost every x , one has

$$\nu_q(B(x, \kappa^{1-n})) \leq C \nu(B(x, \kappa^{-n}))^q \kappa^{-n(\tau_\mu(q) - \varepsilon)}$$

for n large enough, where $\kappa = 1 + 1/c$. This implies that, for ν_q -almost every x , one has

$$\nu_q(B(x, r)) \leq C \nu(B(x, r))^q r^{\tau_\mu(q) - \varepsilon}$$

for r small enough.

It results from these estimates that $\mathcal{H}_\nu^{q, \tau_\mu(q) - \varepsilon}(\text{supp } \nu) > 0$, and therefore $b_\nu(q) \geq \tau_\mu(q)$.

Lemma 6. *Under the same assumptions as in Lemma 5, one has*

$$\nu_q(\mathbb{R} \setminus X_\nu(-B'_\nu(q+), -B'_\nu(q-))) = 0.$$

Proof. Due to Lemmas 3 and 5, $B_\nu(q) = \tau_\mu(q)$. Take $\alpha < -B'_\nu(q+)$ and set

$$E_n = \{x \in \text{supp } \nu \mid \nu(B(x, r)) > r^\alpha \text{ for } r < 1/n\}.$$

Choose $\varepsilon > 0$ and $t > 0$ such that $\tau_\mu(q) - \varepsilon - \alpha t > B_\nu(q + t)$. This can be done for positive, but arbitrarily small, ε . Once ε is chosen, consider the following sets

$$G_m = \left\{ x \in \text{supp } \nu \mid \nu_q(B(x, r)) \leq C \nu(B(x, r))^q r^{\tau_\mu(q) - \varepsilon} \text{ for } r \leq 1/m \right\},$$

where C is the constant in Lemma 5. Due to this lemma, $\bigcup_{m \geq 1} G_m$ has full ν_q -measure.

Let F be a subset of E_n . For any $\theta < \min(1/n, 1/m)$, consider a centered θ -cover $\{B(x_j, r_j)\}$ of $F \cap G_m$ which splits in β packings.

We have, if ν_q^* stands for the outer measure associated with ν_q ,

$$\begin{aligned} \nu_q^*(F \cap G_m) &\leq \sum \nu_q(B(x_j, r_j)) \leq C \sum \nu(B(x_j, r_j))^q r_j^{\tau_\mu(q) - \varepsilon} \\ &\leq C \sum \nu(B(x_j, r_j))^{q+t} r_j^{\tau_\mu(q) - \varepsilon - \alpha t} \nu(B(x_j, r_j))^{-t} r_j^{\alpha t} \\ &\leq C \sum \nu(B(x_j, r_j))^{q+t} r_j^{\tau_\mu(q) - \varepsilon - \alpha t} \\ &\leq \beta C \overline{\mathcal{P}}_{\nu, \theta}^{q+t, \tau_\mu(q) - \varepsilon - \alpha t}(F). \end{aligned}$$

This means that $\nu_q^*(F) \leq C \overline{\mathcal{P}}_{\nu}^{q+t, \tau_\mu(q) - \varepsilon - \alpha t}(F)$, and

$$\nu_q(E_n) \leq \mathcal{P}_{\nu}^{q+t, \tau_\mu(q) - \varepsilon - \alpha t}(E_n) \leq \mathcal{P}_{\nu}^{q+t, \tau_\mu(q) - \varepsilon - \alpha t}(\text{supp } \nu).$$

But this last quantity equals 0 because $\tau_\mu(q) - \varepsilon - \alpha t \geq B_\nu(q+t)$. Therefore

$$\nu_q \left(\left\{ x \in \text{supp } \nu \mid \limsup_{r \searrow 0} \frac{\log \nu(B(x, r))}{\log r} < \alpha \right\} \right) = \nu_q \left(\bigcup E_n \right) = 0.$$

Thus we proved the equality

$$\nu_q \left(\left\{ x \in \text{supp } \nu \mid \limsup_{r \searrow 0} \frac{\log \nu(B(x, r))}{\log r} < -B'_\nu(q+) \right\} \right) = 0.$$

The equality

$$\nu_q \left(\left\{ x \in \text{supp } \nu \mid \limsup_{r \searrow 0} \frac{\log \nu(B(x, r))}{\log r} > -B'_\nu(q-) \right\} \right) = 0.$$

is proven in a similar way.

Theorem 3. *Under the same hypotheses as in Lemma 5, we have*

$$\dim X_\nu(-B'_\nu(q+), -B'_\nu(q-)) \geq \begin{cases} \tau_\mu^*(-B'_\nu(q+)) & \text{if } q \geq 0, \\ \tau_\mu^*(-B'_\nu(q-)) & \text{if } q \leq 0. \end{cases}$$

Proof. Due to Lemma 6, ν_q is carried by $X_\nu(-B'_\nu(q+), -B'_\nu(q-))$. On the other hand, due to Lemma 5, for ν_q -almost every x , for $\varepsilon > 0$, and for r small enough, one has

$$\frac{\log \nu_q(B(x, r))}{\log r} \geq \frac{\log C}{\log r} + q \frac{\log \nu(B(x, r))}{\log r} + \tau_\mu(q) - \varepsilon.$$

One concludes by using the Billingsley lemma [8].

Corollary 1. *Let $q \in \mathbb{R}$. Assume there exists a Frostman measure μ_q for μ at $(q, \tau_\mu(q))$. If $\tau'(q)$ exists and if μ fulfills the NBC for q , then both measures μ and $\nu = \gamma_*(\mu)$ satisfy the multifractal formalism at q and one has $b_\nu(q) = B_\nu(q) = \Lambda_\nu(q) = b_\mu(q) = B_\mu(q) = \Lambda_\mu(q) = \tau_\mu(q)$.*

By careful analysis of the steps which lead to the previous theorem and its corollary, and a few modification, one can prove the following result.

Theorem 4. *Let $q \in \mathbb{R}$. Assume there exists a non-zero measure μ_q on $\partial\mathcal{A}^*$ and a function C_q on \mathcal{A}^* such that $\mu_q(w) \leq C_q(w) \mu(w)^q c^{-|w|\tau_\mu(q)}$ for all $w \in \mathcal{A}^*$ such that $\mu(w) > 0$. If $\tau'(q)$ exists and if, for all $\varepsilon > 0$, there exists $\eta > 0$ such that, we have*

$$\sum_{n>0} c^{-n(\tau_\mu(q)+\eta\varepsilon)} \sum_{\substack{v,w \in \mathcal{A}^n \\ \delta(v,w) \leq c'}} \mu(v)^{-\eta q} C_q(w)^{1+\eta} \mu(w)^{(1+\eta)q} < +\infty$$

(where $c' = 3$ if $q < 0$, $c' = 4c + 2$ otherwise), then both measures μ and $\nu = \gamma_*(\mu)$ obey the multifractal formalism at q , and $b_\nu(q) = B_\nu(q) = \Lambda_\nu(q) = b_\mu(q) = B_\mu(q) = \Lambda_\mu(q) = \tau_\mu(q)$.

One can remark that the conjunction of the existence of a Gibbs measure and of the NBC is stronger than the hypothesis of this last theorem, which therefore can be called weak NBC.

Remark (The case of higher dimension). We stated and proved these comparison theorems in a one dimensional setting in order not to deal with too complicated notations. But these results hold for measures on \mathbb{R}^d . Here are the few modifications to be made to accomodate this case.

This time, the alphabet is of the form \mathcal{A}^d . So each node w of the tree $(\mathcal{A}^d)^*$ can be viewed as a collection (w_1, w_2, \dots, w_d) of words on \mathcal{A} . An element x of $\partial(\mathcal{A}^d)^*$ can be identified to $(x_1, x_2, \dots, x_d) \in (\partial\mathcal{A}^*)^d$ and a mapping, which we again call γ , from $\partial(\mathcal{A}^d)^*$ to \mathbb{R}^d is defined by $\gamma(x) = (\gamma(x_1), \gamma(x_2), \dots, \gamma(x_d))$. Similarly, the distance of two nodes $v = (v_1, v_2, \dots, v_d)$ and $w = (w_1, w_2, \dots, w_d)$ of the same generation, again denoted by δ , is $\delta(v, w) = \max \{\delta(v_1, w_1), \delta(v_2, w_2), \dots, \delta(v_d, w_d)\}$. At last, the value of c' is unchanged. Then, as already said, the previous results hold, with the same proofs, if μ is a measure on $\partial(\mathcal{A}^d)^*$ and $\nu = \gamma_*(\mu)$.

5 Quasi-Bernoulli measures.

A probability measure on $\partial\mathcal{A}^*$ is said to be quasi-Bernoulli if there exists $C > 0$ such that, for any v and w in \mathcal{A}^* , one has

$$\frac{1}{C} \mu(v) \mu(w) \leq \mu(vw) \leq C \mu(v) \mu(w). \quad (7)$$

The multinomial measures constitute a paradigm of such measures. They are very special elements of a larger family. Indeed, denote by S the shift operation on $\partial\mathcal{A}^*$. Due

to its construction [10], any Gibbs measure with Hölder potential on the dynamical system $(\partial\mathcal{A}^*, S)$ is quasi-Bernoulli. Moreover, any quasi-Bernoulli measure is equivalent to an ergodic quasi-Bernoulli measure (see [20]).

It turns out from [28, 29, 11] that for every $q \in \mathbb{R}$ there exists a probability measure μ_q and a constant $C_q > 0$, such that, for every $v \in \mathcal{A}^*$ such that $\mu(v) \neq 0$, one has

$$\frac{1}{C_q} \mu(v)^q c^{-\tau_\mu(q)|v|} \leq \mu_q(v) \leq C_q \mu(v)^q c^{-\tau_\mu(q)|v|}. \quad (8)$$

Moreover, the function τ_μ is differentiable [20]. As a consequence, the first two assumptions in Corollary 1 are fulfilled for every $q \in \mathbb{R}$.

Clearly, a quasi-Bernoulli probability measure μ on $\partial\mathcal{A}^*$ has an atom if and only if $\mu(j) = 1$ for some $j \in \mathcal{A}^1$, in which case $\mu = \delta_{jj\dots j\dots}$. We discard this case and prove the following result.

Theorem 5. *Let μ be a continuous quasi-Bernoulli measure on $\partial\mathcal{A}^*$. Then both measures μ and $\nu = \gamma_*(\mu)$ obey the multifractal formalism everywhere and one has $b_\nu = B_\nu = \Lambda_\nu = b_\mu = B_\mu = \Lambda_\mu = \tau_\mu$.*

This result is established without using the NBC in [7] under the strong hypothesis: for all $n \geq 1$, if $a, b \in \mathcal{A}^n$ and $\delta(a, b) = 1$ then $\mu(a)\mu(b) = 0$. In particular, the case $c = 2$ is excluded and in the case $c = 3$ μ is a Dirac mass.

Proof. We begin by a preliminary remark: if v and w are words of length n , and if \dot{v} and \dot{w} stand for their prefixes of length $n - 1$, then $\delta(\dot{v}, \dot{w}) > k$ implies $\delta(v, w) > ck$. It results that, given two integers $n \geq m > 0$ and two words v and w in \mathcal{A}^n such that $c^{m-1} < \delta(v, w) \leq c^m$, there exists two prefixes \bar{v} and \bar{w} of v and w respectively of common length $n - m$ such that $\delta(\bar{v}, \bar{w}) \leq 1$.

It results from (7) and the above remark that, for a quasi-Bernoulli measure to fulfill the NBC it is enough that for all $\varepsilon > 0$, there exists $\eta > 0$ such that

$$\sum_{n>0} c^{-n(\tau_\mu(q)+\eta\varepsilon)} \sum_{\substack{v, w \in \mathcal{A}^n \\ \delta(v, w) \leq 1}} \mu(v)^{-\eta q} \mu(w)^{(1+\eta)q} < +\infty. \quad (9)$$

Due to the existence of the Gibbs measure μ_q this reduces to

$$\sum_{n>0} c^{-n(\tau_\mu(q)+\eta\varepsilon)} \sum_{\substack{v, w \in \mathcal{A}^n \\ \delta(v, w) = 1}} \mu(v)^{-\eta q} \mu(w)^{(1+\eta)q} < +\infty.$$

Define $\tilde{\mathcal{A}} = \{0 \leq j \leq c - 2 \mid \mu(j)\mu(j + 1) \neq 0\}$.

Define ρ_k to be the word consisting of k consecutive zeros and λ_k to be the word consisting of k consecutive $c - 1$ (considered as a letter from the alphabet $\{0, 1, 2, \dots, c - 1\}$).

By (7), for $n \geq 1$, a representation of the set of pairs (v, w) in \mathcal{A}^n such that $\mu(v)\mu(w) \neq 0$ and $\iota(w) = \iota(v) + 1$ is as follows:

$$\bigcup_{k=0}^{n-1} \bigcup_{\substack{u \in \mathcal{A}^{n-1-k} \\ \mu(u) \neq 0}} \{(u.j.\lambda_k, u.(j+1).\rho_k) \mid j \in \tilde{\mathcal{A}}\} \quad (10)$$

if $\mu(0)\mu(c-1) \neq 0$ and

$$\bigcup_{\substack{u \in \mathcal{A}^{n-1} \\ \mu(u) \neq 0}} \{(u.j, u.(j+1)) \mid j \in \tilde{\mathcal{A}}\}$$

otherwise.

We end the proof when $\mu(0)\mu(c-1) \neq 0$. The other case is simpler. We have to prove that for every $\varepsilon > 0$, there exists $\eta > 0$ such that

$$S_{q,\varepsilon}(\eta) = \sum_{n \geq 1} c^{-n(\tau_\mu(q) + \eta\varepsilon)} \sum_{k=0}^{n-1} \sum_{\substack{u \in \mathcal{A}^{n-1-k} \\ \mu(u) \neq 0}} \sum_{j \in \tilde{\mathcal{A}}} \mu(u.j.\lambda_k)^q \frac{\mu(u.j.\lambda_k)^{q\eta}}{\mu(u.(j+1).\rho_k)^{q\eta}} + \mu(u.(j+1).\rho_k)^q \frac{\mu(u.(j+1).\rho_k)^{q\eta}}{\mu(u.j.\lambda_k)^{q\eta}}$$

is finite.

Fix $\varepsilon > 0$. Let η be a positive number to be chosen later on. By (7),

$$\begin{aligned} \mu(u.j.\lambda_k)^q \frac{\mu(u.j.\lambda_k)^{q\eta}}{\mu(u.(j+1).\rho_k)^{q\eta}} + \mu(u.(j+1).\rho_k)^q \frac{\mu(u.(j+1).\rho_k)^{q\eta}}{\mu(u.j.\lambda_k)^{q\eta}} \leq \\ C(q, \eta) \mu(u)^q \left[\mu(\lambda_k)^q \frac{\mu(\lambda_k)^{q\eta}}{\mu(\rho_k)^{q\eta}} + \mu(\rho_k)^q \frac{\mu(\rho_k)^{q\eta}}{\mu(\lambda_k)^{q\eta}} \right] \end{aligned} \quad (11)$$

for some constant $C(q, \eta)$. Therefore

$$S_{q,\varepsilon}(\eta) \leq (c-1) C(q, \eta) \sum_{n \geq 1} c^{-n(\tau_\mu(q) + \eta\varepsilon)} f_n(q, \eta) \quad (12)$$

with

$$f_n(q, \eta) = \sum_{k=0}^{n-1} c^{(n-1-k)\tau_{\mu, n-1-k}(q)} \left[\mu(\lambda_k)^q \frac{\mu(\lambda_k)^{q\eta}}{\mu(\rho_k)^{q\eta}} + \mu(\rho_k)^q \frac{\mu(\rho_k)^{q\eta}}{\mu(\lambda_k)^{q\eta}} \right],$$

where

$$\tau_{\mu, k}(q) = \frac{1}{k} \log_c \sum_{u \in \mathcal{A}^k}^* \mu(u)^q.$$

By using (8) we get

$$\frac{1}{C_q} c^{k\tau_\mu(q)} \leq c^{k\tau_{\mu,k}(q)} \leq C_q c^{k\tau_\mu(q)}. \quad (13)$$

Then (8) and (13) yield

$$f_n(q, \eta) \leq C_q^{2+2\eta} c^{(n-1)\tau_\mu(q)} \sum_{k=0}^{n-1} \left[\mu_q(\lambda_k) \frac{\mu_q(\lambda_k)^\eta}{\mu_q(\rho_k)^\eta} + \mu_q(\rho_k) \frac{\mu_q(\rho_k)^\eta}{\mu_q(\lambda_k)^\eta} \right]. \quad (14)$$

By construction μ_q has no atoms and is quasi-Bernoulli. Hence, by the sub-multiplicativity property (up to a multiplicative constant) of μ_q , $\lim_{n \rightarrow \infty} \frac{1}{n} \log \mu_q(\rho_n)$ exists and is negative, as well as $\lim_{n \rightarrow \infty} \frac{1}{n} \log \mu_q(\lambda_n)$. Therefore if η is small enough (notice that η does not depend on ε), one has

$$M_q(\eta) = \sup_{k \in \mathbb{N}} \left[\mu_q(\rho_k) \frac{\mu_q(\rho_k)^\eta}{\mu_q(\lambda_k)^\eta} + \mu_q(\lambda_k) \frac{\mu_q(\lambda_k)^\eta}{\mu_q(\rho_k)^\eta} \right] < \infty$$

and, by (14),

$$f_n(q, \eta) \leq C_q^{2+2\eta} M_q(\eta) n c^{(n-1)\tau_\mu(q)}.$$

Finally by (12)

$$S_{q,\varepsilon}(\eta) \leq (c-1) C(q, \eta) C_q^{2+2\eta} M_q(\eta) c^{-\tau_\mu(q)} \sum_{n \geq 1} n c^{-n\eta\varepsilon} < \infty.$$

6 Statistically self-similar measures.

In this section, we consider the random measures introduced by B. Mandelbrot in [24]. Up to now, their multifractal analysis, as the one of quasi-Bernoulli measures, was performed only in the setting of the tree of c -adic intervals (see [22, 21, 14, 30, 2, 3]). Moreover, these measures, which can be viewed as random Bernoulli measures, are not quasi-Bernoulli. This motivated the consideration of the weak NBC (Theorem 4).

For the sake of simplicity, we only deal with the so-called canonical multiplicative cascades. Let us recall a construction of these measures.

Fix W a non-negative random variable. Assume that W is not almost surely constant and that $\mathbb{E}(W) = 1/c$.

Define the function $\tilde{\tau}(q) = 1 + \log_c \mathbb{E}(\mathbf{1}_{\{W>0\}} W^q)$ for $q \in \mathbb{R}$.

In order to avoid technicalities, unessential to our purpose, we assume that W is positive and that $\tilde{\tau}(q)$ is finite for any $q \in \mathbb{R}$.

Let $(W_w)_{w \in \mathcal{A}^*}$ be a sequence of independent copies of W . For every $n \geq 1$, consider the random measure μ_n whose density with respect to the uniform probability measure on $\partial \mathcal{A}^*$ is locally constant and equals

$$c^n W_{w_1} W_{w_1 w_2} \cdots W_{w_1 w_2 \dots w_n}$$

on the cylinder $w = w_1 w_2 \dots w_n$. With probability one, the sequence μ_n converges weakly to a measure μ as n goes to infinity. Moreover, if $\tilde{\tau}(1) < 0$, one has $\mu \neq 0$ (see [23]).

Then, define $J = \{q \in \mathbb{R}; \tilde{\tau}^*(-\tilde{\tau}'(q)) > 0\}$. It follows from Theorem 8(*iv*) in [3] that $\tau_\mu = \tilde{\tau}$ on J .

Theorem 6. *With probability one, both measures μ and $\nu = \gamma_*(\mu)$ satisfy the multifractal formalism on J and $b_\nu = B_\nu = \Lambda_\nu = b_\mu = B_\mu = \Lambda_\mu = \tau_\mu = \tilde{\tau}$ on J .*

Proof. For every $q \in J$, $v \in \mathcal{A}^*$ and $n \geq 1$, define

$$Y_{q,n}(v) = c^{-n\tilde{\tau}(q)} \sum_{w_1 \dots w_n \in \mathcal{A}^n} W_{vw_1}^q W_{vw_1 w_2}^q \cdots W_{vw_1 w_2 \dots w_n}^q.$$

It follows from Corollary 5 in [3] that, with probability one, for all $v \in \mathcal{A}^*$ and all $q \in J$, the limit $Y_q(v) = \lim_{n \rightarrow \infty} Y_{q,n}(v)$ exists. Moreover, with probability one, for all $q \in J$, the mapping μ_q defined on $\partial \mathcal{A}^*$ by

$$\mu_q(v) = c^{-|v|\tilde{\tau}(q)} Y_q(v) \prod_{j=1}^{|v|} W_{v_1 \dots v_j}^q \quad (15)$$

defines a measure (notice that $\mu_1 = \mu$); all the measures μ_q have $\partial \mathcal{A}^*$ as support and for all $v \in \mathcal{A}^*$ and $q \in J$,

$$\mu_q(v) = C_q(v) \mu(v)^q c^{-|v|\tilde{\tau}(q)} \quad (16)$$

with

$$C_q(v) = \frac{Y_q(v)}{Y_1^q(v)}.$$

Since $\tau_\mu = \tilde{\tau}$ on J , the result will be a consequence of Theorem 4 if we show that for every non trivial compact subinterval K of J , with probability one, for all $q \in K$, for all $\varepsilon > 0$, there exists $\eta > 0$ such that

$$\sum_{n \geq 1} c^{-n(\tilde{\tau}(q) + \eta\varepsilon)} \sum_{\substack{v, w \in \mathcal{A}^n \\ \delta(v, w) \leq c'}} \mu(v)^{-\eta q} C_q(w)^{1+\eta} \mu(w)^{(1+\eta)q} < +\infty, \quad (17)$$

i.e.

$$\sum_{n \geq 1} c^{-n(\tilde{\tau}(q) + \eta\varepsilon)} f_{n, \varepsilon, \eta}(q) < \infty,$$

where

$$f_{n,\varepsilon,\eta}(q) = \sum_{\substack{v,w \in \mathcal{A}^n \\ \delta(v,w) \leq c'}} Y_1(v)^{-\eta q} Y_q(w)^{1+\eta} \prod_{k=1}^n W_{v_1 \dots v_k}^{-\eta q} W_{w_1 \dots w_k}^{(1+\eta)q}.$$

Fix such a compact K . It turns out that it suffices to show that for every $\varepsilon > 0$, if $\eta > 0$ is small enough, then

$$\begin{cases} \sum_{n \geq 1} \sup_{q \in K} n c^{-n(\bar{\tau}(q) + \eta\varepsilon)} \mathbb{E}(f_{n,\varepsilon,\eta}(q)) < \infty \\ \sum_{n \geq 1} \sup_{q \in K} c^{-n(\bar{\tau}(q) + \eta\varepsilon)} \mathbb{E}(|f'_{n,\varepsilon,\eta}(q)|) < \infty \end{cases} \quad (18)$$

(see the proof of Corollary 1 in [3] for a more detailed similar argument).

It follows from Lemma 6 in [3] that for η small enough

$$C_K(\eta) = \sup_{\substack{q \in K, \\ n \geq 1, \\ v, w \in \mathcal{A}^n}} \mathbb{E} \left(\left| \frac{d}{dq} (Y_1(v)^{-\eta q} Y_q(w)^{1+\eta}) \right| \right) + \mathbb{E} (Y_1(v)^{-\eta q} Y_q(w)^{1+\eta}) < \infty$$

and

$$C'_K(\eta) = \sup_{\substack{q \in K, \\ n \geq 1, \\ v, w \in \mathcal{A}^n}} \frac{\mathbb{E} \left(\left| \frac{d}{dq} W_v^{-\eta q} W_w^{(1+\eta)q} \right| \right)}{\mathbb{E} \left(W_v^{-\eta q} W_w^{(1+\eta)q} \right)} < \infty.$$

By taking into account the fact that the W s are mutually independent, we get

$$\mathbb{E}(|f'_{n,\varepsilon,\eta}(q)|) \leq C_K(\eta)(1 + nC'_K(\eta))g_{n,\varepsilon,\eta}(q)$$

where

$$g_{n,\varepsilon,\eta}(q) = \sum_{\substack{v,w \in \mathcal{A}^n \\ \delta(v,w) \leq c'}} \prod_{k=1}^n \mathbb{E} \left(W_{v_1 \dots v_k}^{-\eta q} W_{w_1 \dots w_k}^{(1+\eta)q} \right).$$

Also, we have

$$\mathbb{E}(f_{n,\varepsilon,\eta}(q)) \leq C_K(\eta)g_{n,\varepsilon,\eta}(q).$$

By using a reduction similar to the one made in the beginning of Section 5, we can assume without loss of generality that in the sums over $\{v, w \in \mathcal{A}^n; \delta(v, w) \leq c'\}$, there are only pairs (v, w) for which $\delta(v, w) \leq 1$. Then, by using (10), we get

$$g_{n,\varepsilon,\eta}(q) = c^{n\bar{\tau}(q)} + h_{n,\varepsilon,\eta}(q),$$

where

$$\begin{aligned}
h_{n,\varepsilon,\eta}(q) &= (c-1) \sum_{k=0}^{n-1} c^k (\mathbb{E}(W^q))^k \left(\mathbb{E}(W^{-\eta q}) \mathbb{E}(W^{(1+\eta)q}) \right)^{n-k} \\
&= (c-1) \left(\mathbb{E}(W^{-\eta q}) \mathbb{E}(W^{(1+\eta)q}) \right)^n \sum_{k=0}^{n-1} \left[\frac{c \mathbb{E}(W^q)}{\mathbb{E}(W^{-\eta q}) \mathbb{E}(W^{(1+\eta)q})} \right]^k \\
&\leq c^{1+n} \left(-2 + \bar{\tau}(q) + \bar{\tau}(0) + \eta q (\bar{\tau}'(q) - \bar{\tau}'(0)) + \eta O_q(\eta) \right) \\
&\quad \times \sum_{k=0}^{n-1} c^k \left(2 - \bar{\tau}(0) - \eta q (\bar{\tau}'(q) - \bar{\tau}'(0)) + \eta O_q(\eta) \right) \\
&\leq \frac{c^{1+n} (\bar{\tau}(q) + \eta O_q(\eta))}{c^{2 - \bar{\tau}(0) - \eta q (\bar{\tau}'(q) - \bar{\tau}'(0)) + \eta O_q(\eta)} - 1,
\end{aligned}$$

with $o_q(\eta) \rightarrow 0$ uniformly on K when $\eta \rightarrow 0$. Then, it is easily seen that (18) holds if η is small enough.

References

- [1] M. Arbeiter and N. Patzschke, *Random self-similar multifractals*. Math. Nachr. **181** (1996), 5–42.
- [2] J. Barral, *Moments, continuité et analyse multifractale des martingales de Mandelbrot*. Probab. Theory Relat. Fields **113** (1999), 535–569.
- [3] J. Barral, *Continuity of the multifractal spectrum of a random statistically self-similar measures*. J. Theor. Probab., **13** (2000), 1027–1060.
- [4] T. Bedford, *Applications of dynamical systems theory to fractals: a study of cookie-cutter Cantor sets*. In Fractal Geometry and Analysis. Eds. Belair J. and Dubuc S., Kluwer Academic Publishers, 1991.
- [5] F. Ben Nasr, *Analyse multifractale de mesures*. C. R. Acad. Sci. Paris, **319**, Série I (1994), 807–810.
- [6] F. Ben Nasr, I. Bhour, and Y. Heurteaux, *The validity of the multifractal formalism: results and examples*. Adv. in Math. **165** (2002), 264–284.
- [7] I. Bhour, *Une condition de validité du formalisme multifractal pour les mesures*. Thèse de Doctorat. Tunis, 1999.
- [8] P. Billingsley, *Ergodic Theory and Information*. J. Wiley, New York, 1965.

-
- [9] T. Bohr and D. Rand, *The entropy function for characteristic exponents*. *Physica* **25D** (1987), 387–398.
- [10] R. Bowen, *Equilibrium states and the ergodic theory of Anosov Diffeomorphisms*. Lectures Notes in Mathematics, **470**, Dold A. and Eckmann B. Eds., Springer.
- [11] G. Brown, G. Michon, and J. Peyrière, *On the multifractal analysis of measures*. *J. Stat. Phys.*, **66** (1992), No 3/4, 775–790.
- [12] H. Chernoff, *A measure of asymptotic efficiency for tests of a hypothesis based on the sum of observations*. *Ann. Math. Statist.* **23** (1952), 493–507.
- [13] P. Collet, J.L. Lebowitz, and A. Porzio, *The dimension spectrum of some dynamical systems*. *J. Stat. Phys.* **47** (1987), 609–644.
- [14] P. Collet, F. Koukiou, *Large deviations for multiplicative chaos*. *Commun. Math. Phys.* **147** (1992), 329–342.
- [15] G.A. Edgar and R.D. Mauldin, *Multifractal decomposition of digraph recursive fractals*. *Proc. London Math. Soc.* **65** (1992), 604–628.
- [16] K.J. Falconer, *The multifractal spectrum of statistically self-similar measures*. *J. Theor. Prob.* **7** (1994), 681–702.
- [17] U. Frisch and G. Parisi, *Fully developed turbulence and intermittency in turbulence, and predictability in geophysical fluid dynamics and climate dynamics*. International school of physics "Enrico Fermi", course 88, edited by M. Ghil, North Holland (1985), p. 84.
- [18] T.C. Halsey, M.H. Jensen, L.P. Kadanoff, I. Procaccia, and B.I. Shraiman, *Fractal measures and their singularities: the characterisation of strange sets*. *Phys. Rev. A* **33** (1986), p. 1141.
- [19] H.G.E. Hentschel and I. Procaccia, *The infinite number of generalized dimensions of fractals and strange attractors*. *Physica* **8D** (1983), p. 435.
- [20] Y. Heurteaux, *Estimations de la dimension inférieure et de la dimension supérieure des mesures*. *Ann. Inst. Henri Poincaré*, **34** (1998), 309–338.
- [21] R. Holley and E.C. Waymire, *Multifractal dimensions and scaling exponents for strongly bounded random cascades*. *Ann. Appl. Probab.*, **2** (1992), 819–845.
- [22] J.-P. Kahane, *Produits de poids aléatoires et indépendants et applications*. in *Fractal Geometry and Analysis*, J. Bélair and S. Dubuc (eds.), 1991, 277–324.
- [23] J.-P. Kahane et J. Peyrière, *Sur certaines martingales de Benoît Mandelbrot*. *Adv. in Math.*, **22** (1976), 131–145.

-
- [24] B.B. Mandelbrot, *Intermittent turbulence in self-similar cascades: divergence of high moments and dimension of the carrier*. J. Fluid Mech. **62** (1974), 331–358.
- [25] B.B. Mandelbrot, *Multifractal measures, especially for the geophysicist*. Annual reviews of materials sciences **19**, 1989, 514–516.
- [26] B.B. Mandelbrot, *A class of multifractal measures with negative (latent) value for the dimension $f(\alpha)$* . Fractals: physical origin and properties (Erice, 1988). Edited by Luciano Pietronero, New York: Plenum, 1989.
- [27] B.B. Mandelbrot, *Two meanings of multifractality, and the notion of negative fractal dimension*. Soviet-American chaos meeting (Woods Hole, 1989). Edited by Kenneth Ford and David Campbell, American Institute of Physics, 1990.
- [28] G. Michon, *Une construction des mesures de Gibbs sur certains ensembles de Cantor*. C. R. Acad. Sc. Paris **308** (1989), 315–318.
- [29] G. Michon, *Mesures de Gibbs sur les Cantor Réguliers*. Ann. Inst. Henri Poincaré, Phys. Théor. **58**, No.3, (1993), 267–285.
- [30] G.M. Molchan, *Scaling exponents and multifractal dimensions for independent random cascades*. Commun. Math. Phys., **179** (1996), 681–702.
- [31] L. Olsen, *Random geometrically graph directed self-similar multifractals*. Pitman Res. Notes Math. Ser., **307** (1994).
- [32] L. Olsen, *A multifractal formalism*. Adv. Math. **116** (1995), 92–195.
- [33] J. Peyrière, *Multifractal measures*. In Probabilistic and Stochastic Methods in Analysis (Proceedings of the NATO ASI, Il Ciocco 1991). Ed. J. Byrnes, Kluwer Academic Publishers, 1992.
- [34] D.A. Rand, *The singularity spectrum $f(\alpha)$ for cookie-cutters*. Ergod. Th. & Dynam. Sys. **9** (1989), 527–541.
- [35] X. Saint Raymond and C. Tricot, *Packing regularity of sets in n -space*. Math. Proc. Cambridge Philos. Soc. **288** (1985), 679–699.
- [36] C. Tricot, *Two definitions of fractional dimension*. Math. Proc. Cambridge Philos. Soc. **91** (1982), 57–74.



Unité de recherche INRIA Rocquencourt

Domaine de Voluceau - Rocquencourt - BP 105 - 78153 Le Chesnay Cedex (France)

Unité de recherche INRIA Lorraine : LORIA, Technopôle de Nancy-Brabois - Campus scientifique
615, rue du Jardin Botanique - BP 101 - 54602 Villers-lès-Nancy Cedex (France)

Unité de recherche INRIA Rennes : IRISA, Campus universitaire de Beaulieu - 35042 Rennes Cedex (France)

Unité de recherche INRIA Rhône-Alpes : 655, avenue de l'Europe - 38330 Montbonnot-St-Martin (France)

Unité de recherche INRIA Sophia Antipolis : 2004, route des Lucioles - BP 93 - 06902 Sophia Antipolis Cedex (France)

Éditeur

INRIA - Domaine de Voluceau - Rocquencourt, BP 105 - 78153 Le Chesnay Cedex (France)

<http://www.inria.fr>

ISSN 0249-6399