



Proximal Convexification Procedures in Combinatorial Optimization

Aris Daniilidis, Claude Lemaréchal

► **To cite this version:**

Aris Daniilidis, Claude Lemaréchal. Proximal Convexification Procedures in Combinatorial Optimization. [Research Report] RR-4550, INRIA. 2002. inria-00072038

HAL Id: inria-00072038

<https://hal.inria.fr/inria-00072038>

Submitted on 23 May 2006

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

*Proximal Convexification Procedures in
Combinatorial Optimization*

Aris Daniilidis, Claude Lemaréchal

N° 4550

Septembre 2002

THÈME 4



*Rapport
de recherche*

Proximal Convexification Procedures in Combinatorial Optimization

Aris Daniilidis, Claude Lemaréchal

Thème 4 — Simulation et optimisation
de systèmes complexes
Projet Numopt

Rapport de recherche n° 4550 — Septembre 2002 — 18 pages

Abstract: Lagrangian relaxation is useful to bound the optimal value of a given optimization problem, and also to obtain relaxed solutions. To obtain primal solutions, it is conceivable to use a convexification procedure suggested by D.P. Bertsekas in 1979, based on the proximal algorithm in the primal space.

The present paper studies the theory assessing the approach in the framework of combinatorial optimization. Our results indicate that very little can be expected in theory, even though fairly good practical results have been obtained for the unit-commitment problem.

Key-words: Proximal algorithm, Lagrangian relaxation, duality gap, primal-dual heuristics

Procédures Primales de Convexification en Optimisation Combinatoire

Résumé : La relaxation lagrangienne est utile pour borner la valeur optimale d'un problème d'optimisation, et aussi pour obtenir des solutions relaxées. Pour obtenir des solutions primales, on peut concevoir d'utiliser une procédure de convexification proposée par Bertsekas en 1979, fondée sur l'algorithme proximal dans l'espace primal.

Dans cet article, nous étudions la théorie sous-jacente à cette approche, dans le cadre de l'optimisation combinatoire. Nos résultats suggèrent que bien peu de justifications théoriques peuvent être attendues, même si de très bons résultats pratiques ont été obtenus, pour des problèmes d'optimisation de la production.

Mots-clés : Algorithme proximal, relaxation lagrangienne, saut dual, heuristiques primales-duales

1 Introduction, Motivation

This paper is motivated by a practical application: the unit-commitment problem, more precisely to optimize the generation schedules of the set of electrical power plants in France. Such a problem is usually solved through duality ([2, 11], see also [5, 12] for additional references). After solving the dual, comes the question of recovering a primal feasible solution, possibly suboptimal. An idea is to add in the production cost a quadratic term penalizing the deviation from the relaxed solution, obtained by dual means. This can give fairly good practical results, to be published elsewhere: [7].

Our aim here is to study theoretically this approach, with emphasis on combinatorial problems lending themselves to Lagrangian relaxation. To this aim, we consider first the general optimization problem

$$\inf f(x), \quad x \in \mathbb{R}^n. \quad (1)$$

In this simplified notation, possible constraints are incorporated via the indicator function (0 on the feasible set, $+\infty$ outside). We have particularly in mind *0-1 linear programming problems*, say

$$\min b^\top x, \quad Ax \geq c \in \mathbb{R}^m, \quad x^i \in \{0, 1\}, \quad i = 1, \dots, n; \quad (2)$$

in this case the objective function is

$$f(x) := \begin{cases} b^\top x & \text{if } Ax \geq c \text{ and } x \in \{0, 1\}^n, \\ +\infty & \text{otherwise.} \end{cases} \quad (3)$$

Our aim is to study for such problems a convexification procedure introduced by D.P. Bertsekas in [1], based on the (primal) *proximal algorithm*.

1.1 The General Idea: Moreau Envelope

The basic idea of this procedure is to add some convexity into f , which is replaced by

$$f_\rho(x, y) := f(x) + \rho \|x - y\|^2, \quad (4)$$

where $\rho > 0$ and $y \in \mathbb{R}^n$ is an additional variable. Then f_ρ is minimized hierarchically: one sets

$$\varphi_\rho(y) := \inf \{f_\rho(x, y) : x \in \mathbb{R}^n\} \quad (5)$$

and one minimizes φ_ρ *on the whole space* (observe that φ_ρ is $+\infty$ nowhere – unless $f \equiv +\infty$). The function φ_ρ thus defined is fairly known in nonlinear analysis. Introduced originally for a convex function f , it is usually called the Moreau-Yosida regularization; for more general situations see [13], where it is called the Moreau envelope.

Lemma 1.1 *The following general properties hold.*

- (i) For all $y \in \mathbb{R}^n$, the function $\rho \mapsto \varphi_\rho(y)$ is nondecreasing.
- (ii) $\varphi_\rho(y) \leq f(y)$ for all $y \in \mathbb{R}^n$.

Assume that f is lower semicontinuous and bounded from below. Then:

- (iii) if y_0 is a local minimum of φ_ρ then $x = y_0$ is the unique minimum point of $f_\rho(\cdot, y_0)$ in (5).

In particular $\varphi_\rho(y_0) = f(y_0)$;

- (iv) local minima of φ_ρ are also local minima of $\varphi_{\rho'}$ for $\rho' \geq \rho$.

Proof. (i) Obviously, $f_\rho(x, y)$ in (4) is a nondecreasing function of ρ and this property is transmitted to the infima.

(ii) Just observe that $\varphi_\rho(y) \leq f_\rho(y, y) = f(y)$.

(iii) Let $y = y_0$ in (5). Then our assumption implies that the infimum is attained at some x_0 . Now for any y close to y_0 , we can write

$$f(x_0) + \rho\|y_0 - x_0\|^2 = \varphi_\rho(y_0) \leq \varphi_\rho(y) \leq f(x_0) + \rho\|y - x_0\|^2,$$

hence $\|y_0 - x_0\|^2 \leq \|y - x_0\|^2$. Take in particular $y = y_0 - t(y_0 - x_0)$, so that $y - x_0 = (1 - t)(y_0 - x_0)$. Then $\|y_0 - x_0\|^2 \leq (1 - t)^2\|y_0 - x_0\|^2$. Taking $t > 0$ small enough shows that $y_0 - x_0$ must be 0.

(iv) Using (i), we have for a local minimum y_0 of φ_ρ :

$$\varphi_{\rho'}(y) \geq \varphi_\rho(y) \geq \varphi_\rho(y_0) \quad \text{for } y \text{ close to } y_0;$$

but from (iii) and (ii), $\varphi_\rho(y_0) = f(y_0) \geq \varphi_{\rho'}(y_0)$, which completes the proof. \square

Remark 1.2 *In these results, (iii) is the most important and its proof deserves comment. Being a min-function, φ_ρ is usually not differentiable: the concept of derivative, or gradient, is then replaced by that of directional derivatives:*

$$\varphi'_\rho(y, d) := \lim_{t \downarrow 0} \frac{\varphi_\rho(y + td) - \varphi_\rho(y)}{t}, \quad \text{for given } d \in \mathbb{R}^n.$$

Now a well-known formula (due to J.M. Danskin in [3]) says that, under appropriate assumptions on f_ρ , the directional derivative of functions given by (5) exists and has the expression

$$\varphi'_\rho(y, d) = \min \{d^\top \nabla_y f_\rho(x, y) : x \text{ minimizes } f_\rho(\cdot, y)\}. \quad (6)$$

Here $\nabla_y f_\rho(x, y) = 2\rho(y - x)$ is the partial derivative of f_ρ with respect to y . In plain words: when moving from y to $y + td$ ($t > 0$ small), the marginal change of φ_ρ is the smallest scalar product of d with the partial derivatives of the minimand, computed at all the minimizing x 's. For a local minimum, this change must be nonnegative: $\varphi'_\rho(y, d) \geq 0$ for any $d \in \mathbb{R}^n$, i.e. $d^\top \nabla_y f_\rho(x, y) \geq 0$ for any minimizing x and any $d \in \mathbb{R}^n$. This just means $\nabla_y f_\rho(x, y) = 2\rho(y - x) = 0$, i.e. $x = y$ for any x minimizing (4). \square

1.2 Minimizing f via φ_ρ

Intuitively, minimizing φ_ρ in (5) is equivalent to minimizing f ; this can be made precise:

Theorem 1.3 *The minimization of f and of φ_ρ are related as follows:*

- (i) $\inf \{f(x) : x \in \mathbb{R}^n\} = \inf \{\varphi_\rho(y) : y \in \mathbb{R}^n\}$.
- (ii) If x^* minimizes f , then x^* minimizes φ_ρ .
- (iii) Assume f is lower semicontinuous and bounded from below. If y^* minimizes φ_ρ then y^* minimizes f .

Proof. (i) For any x and y in \mathbb{R}^n , $f_\rho(x, y) \geq f(x)$; hence $\varphi_\rho(y) \geq \inf f$ and $\inf \varphi_\rho \geq \inf f$. On the other hand, Lemma 1.1 (ii) gives $\inf \varphi_\rho \leq \inf f$.

(ii) In view of (i) and of Lemma 1.1 (ii), $\inf \varphi_\rho = \inf f = f(x^*) \geq \varphi_\rho(x^*)$.

(iii) It follows from Lemma 1.1 (ii), (iii) that $f(y^*) = \varphi_\rho(y^*) \leq \varphi_\rho(y) \leq f(y)$ for all $y \in \mathbb{R}^n$. \square

Thus, the approach replaces a single minimization (of f) by:

- the minimization of $f_\rho(\cdot, y)$, a function which is “more convex” than f ,
- the minimization of φ_ρ .

Then a natural question is whether φ_ρ has a chance of being convex; this essentially corresponds to f being convex:

Theorem 1.4 *The following holds, relating the convexity of f with that of φ_ρ :*

- (i) *Assume f is bounded from below. Then φ_ρ is convex whenever f_ρ is convex (jointly with respect to x and y);*
- (ii) *f_ρ is convex (jointly) if and only if f is convex.*

Proof. (i) This is a classical result, see for example [9, Corollary B.2.4.5].

(ii) If f is convex, then f_ρ is obviously convex (jointly). If f_ρ is convex, then in particular $x \mapsto f(x) = f_\rho(x, x)$ is convex. \square

In view of Theorem 1.3, better convexification properties could hardly be expected from this procedure. Indeed (1) contains just about any optimization problem; in particular, there are instances of (1) which are difficult, but for which f_ρ is convex in x and computing φ_ρ is “easy” – an example will be given in §3.4. In these cases, minimizing f could not be equivalent to minimizing φ_ρ , if the latter were convex!

On the other hand, a classical convexification scheme is augmented Lagrangian. Applied to a problem such as (2), for example, it would add the term $\rho \|Ax - c\|^2$. This also corresponds to introducing a Moreau envelope, but in the *dual* space; it does result in a convex optimization problem for ρ large enough but is hardly implementable; see [8, § XII.5.2] for example. As mentioned in [1], the present *primal* approach has the advantage of preserving separability of f , if any. The crucial point is that the quadratic term in (4) is a sum over the coordinates of the primal variable x ; as such, it is not too complicating. In fact, we are interested in instances of (1) amenable to Lagrangian relaxation – such is the case of (2). Our aim will then be to reduce the duality gap, and/or to produce heuristic primal solutions, generated by the algorithm minimizing φ_ρ .

1.3 The Proximal Algorithm

The algorithm suggested in [1] to minimize φ_ρ is essentially

$$y_{k+1} \in \operatorname{Argmin} \{ f_\rho(x, y_k) : x \in \mathbb{R}^n \}, \quad (7)$$

where Argmin denotes the set of global minimizers, assumed nonempty (which is the case when f is lower semicontinuous and bounded from below); naturally, the algorithm stops when $y_{k+1} = y_k$. This is called the *proximal algorithm*, whose convergence properties rely on the following result:

Lemma 1.5 Assume y_{k+1} exists in (7). There holds at each iteration:

$$f(y_{k+1}) \leq f(y_k) - \rho \|y_k - y_{k+1}\|^2.$$

As a result:

- either $f(y_k) \rightarrow -\infty$,
- or $\sum_k \|y_{k+1} - y_k\|^2 < +\infty$.

Proof. By definition of y_{k+1} and from Lemma 1.1 (ii),

$$f(y_{k+1}) + \rho \|y_{k+1} - y_k\|^2 = \varphi_\rho(y_k) \leq f(y_k),$$

which is the required inequality. In particular, the sequence $\{f(y_k)\}$ is decreasing:

- either $f(y_k) \rightarrow -\infty$,
- or $f(y_k)$ is bounded below; we obtain by summation

$$f(y_{K+1}) - f(y_1) \leq -\rho \sum_{k=1}^K \|y_{k+1} - y_k\|^2,$$

which shows that the series $\sum_k \|y_{k+1} - y_k\|^2$ converges. \square

We explain the motivation of the proximal algorithm in the light of Remark 1.2. To avoid excessive generality, assume that φ_ρ is a smooth function, namely that it has a gradient $\nabla \varphi_\rho(y)$ at every $y \in \mathbb{R}^n$. Then its directional derivatives are $\varphi'_\rho(y, d) = d^\top \nabla \varphi_\rho(y)$ for all $d \in \mathbb{R}^n$; with Danskin's formula (6), this clearly implies that $f_\rho(\cdot, y)$ must have a *unique* minimizer¹ $x(y)$, and that $\nabla \varphi_\rho(y) = 2\rho[y - x(y)]$. In particular, each next iterate y_{k+1} in (7) exists and is defined without ambiguity. Besides, since

$$y_{k+1} = x(y_k) = y_k + \frac{1}{2\rho} 2\rho(x(y_k) - y_k) = y_k - \frac{1}{2\rho} \nabla \varphi_\rho(y_k),$$

we see that the proximal algorithm is just the minimization of φ_ρ by a standard gradient method. Now, from Lemma 1.5:

- either $f(y_k) \rightarrow -\infty$ (then we are certainly minimizing f successfully!)
- or $2\rho(y_k - y_{k+1}) = \nabla \varphi_\rho(y_k) \rightarrow 0$.

In the latter situation, assume that the sequence $\{y_k\}$ has some cluster point y^* . If φ_ρ is actually continuously differentiable (this means that the mapping $y \mapsto x(y)$ is *continuous*), then we see that $\nabla \varphi_\rho(y^*) = 2\rho[y^* - x(y^*)] = 0$: the proximal algorithm can only produce stationary points of φ_ρ , which have a chance to be local minimizers.

2 Discrete Optimization Problems: Conceptual Forms

In this section we focus on the particular case where (1) is actually an optimization problem on a *finite* set: we consider

$$\min g(x), \quad x \in F = \{x_1, \dots, x_K\}. \quad (8)$$

¹To see this, observe that the directional derivative is linear in d , hence symmetric; with several minimizers, we would have $\varphi'_\rho(y, d) \neq -\varphi'_\rho(y, -d)$ for some direction d , a contradiction.

The function g is left unspecified for the moment; it is completely characterized by its (finitely many) values $\{g(x_k)\}_{k=1}^K$. With respect to our previous notation,

$$f(x) = \begin{cases} g(x_k) & \text{if } x = x_k \text{ for some } k = 1, \dots, K, \\ +\infty & \text{otherwise.} \end{cases} \quad (9)$$

Needless to say, this function is bounded and lower semi-continuous: the assumptions made in §1 are trivially satisfied. As for the “outer” objective function, it becomes

$$\varphi_\rho(y) = \min_{x \in F} \{g(x) + \rho\|x - y\|^2\}. \quad (10)$$

First of all, local minima of φ_ρ can be characterized in this particular situation:

Proposition 2.1 *A point y_0 is a local minimum of φ_ρ if and only if $x = y_0$ is the unique optimal solution of (10) for $y = y_0$. In particular, every local minimum is a strict local minimum.*

Proof. In view of Lemma 1.1 (iii), we have only to prove that, if $x = y_0$ is the unique solution of (10) for $y = y_0$, then y_0 is a strict local minimum of φ_ρ . Note first that, since F is a finite set, there is $\varepsilon > 0$ such that, for all $x \in F$ different from y_0 ,

$$g(x) + \rho\|x - y_0\|^2 \geq g(y_0) + 2\varepsilon = \varphi_\rho(y_0) + 2\varepsilon.$$

Now take y close enough to y_0 so that, for all $x \in F$,

$$\rho\|x - y\|^2 \geq \rho\|x - y_0\|^2 - \varepsilon.$$

Summing these two inequalities, we obtain

$$g(x) + \rho\|x - y\|^2 \geq \varphi_\rho(y_0) + \varepsilon$$

and hence

$$\min \{g(x) + \rho\|x - y\|^2 : x \in F \setminus \{y_0\}\} \geq \varphi_\rho(y_0) + \varepsilon.$$

As a result: for y close enough to y_0 ,

$$\varphi_\rho(y) \geq \min \{\varphi_\rho(y_0) + \varepsilon, \varphi_\rho(y_0) + \rho\|y_0 - y\|^2\} \geq \varphi_\rho(y_0),$$

and the second inequality is strict if $y \neq y_0$. □

Note that φ_ρ is the minimum of finitely many quadratic functions. The y -space is divided into regions inside which the minimum in (10) is attained at some point $x \in F$, call it x_k (k depending on the region in question). The corresponding quadratic portion has the equation $g(x_k) + \rho\|y - x_k\|^2$, with gradient $2\rho(y - x_k)$. Altogether, φ_ρ looks as indicated in Fig. 1. Local minima can be points such as y_1 or y_2 ; but at y_3 , there are two minimum points in (10): $x = y_3$ and some other $x \in F$; in view of Lemma 1.1 (iii), y_3 cannot be a local minimum of φ_ρ . It is instructive to look at this picture with Remark 1.2 in mind.

We already know from Theorem 1.3 (iii) that local minima of φ_ρ lie in F . Our next result specifies which feasible points can be thus obtained.

Proposition 2.2 *For any $\rho > 0$, any local minimum of φ_ρ lies in F . Moreover:*

(i) *For ρ large enough, the local minima of φ_ρ are exactly the points in F .*




Figure 1: A piecewise quadratic function.

(ii) For ρ small enough, the local minima of φ_ρ are exactly the optimal solutions of (8).

Proof. If y_0 is a local minimum of φ_ρ , Proposition 2.1 shows that y_0 minimizes $f_\rho(x, y_0) = g(x) + \rho\|x - y_0\|^2$ over $x \in F$; then in particular $y_0 \in F$.

(i) We have to prove that an arbitrary $y_0 \in F$ is a local minimum of φ_ρ if ρ is large enough. Define the diameter of $g(F)$

$$\Gamma := \max \{g(x) - g(x') : x \in F, x' \in F\},$$

the “discreteness” of F

$$\varepsilon := \min \{\|x - x'\| : x \in F, x' \in F, x \neq x'\},$$

and take $\rho > \Gamma/\varepsilon^2$.

For y close to y_0 , namely for $\|y - y_0\| \leq \delta := \varepsilon - \sqrt{\Gamma/\rho} > 0$, there holds for all $x \in F$ different from y_0 :

$$\|x - y\| \geq \|x - y_0\| - \|y_0 - y\| \geq \varepsilon - \delta.$$

Then we write

$$\begin{aligned} g(x) + \rho\|y - x\|^2 &\geq g(y_0) + g(x) - g(y_0) + \rho(\varepsilon - \delta)^2 \\ &\geq g(y_0) - \Gamma + \rho(\varepsilon - \delta)^2 \\ &= g(y_0) \geq \varphi_\rho(y_0), \end{aligned}$$

where the last inequality is Lemma 1.1 (ii). The conclusion follows by taking the infimum over x (knowing that the inequality holds trivially for $x = y_0$).

(ii) Call v^* the optimal value in (8), V^* the set of optimal solutions,

$$v^+ := \min \{g(x) : x \in F \setminus V^*\}$$

the “next to optimal” value of g over F , and finally

$$D := \max \{\|x - x'\| : x, x' \in F\}$$

the diameter of F . Note that $v^+ > v^*$ and take $0 < \rho < (v^+ - v^*)/D^2$.

Take $x^* \in V^*$ and let y be a local minimizer of φ_ρ . We already know that $y \in F$ and that $\varphi_\rho(y) = g(y)$ (Proposition 2.1); then write

$$g(y) = \varphi_\rho(y) \leq g(x^*) + \rho\|x^* - y\|^2 \leq g(x^*) + \rho D^2 < g(x^*) + v^+ - v^* = v^+.$$

By definition of v^+ , $g(y)$ has to be equal to v^* ; from Theorem 1.3 (i), this means $y \in V^*$. \square

The proximal algorithm of §1.3 can then be described in the present context:

Algorithm 2.3 (Conceptual proximal algorithm) Choose $\rho > 0$.

STEP 0. Take y_1 arbitrary in \mathbb{R}^n ; set $k = 1$.

STEP 1. Let x_k realize the smallest of the numbers $g(x_{k'}) + \rho\|y_k - x_{k'}\|^2$, for $k' = 1, \dots, K$.

STEP 2. If $x_k = y_k$ then stop.

STEP 3. Set $y_{k+1} = x_k$, replace k by $k + 1$ and go to Step 1. \square

Convergence is easy to establish:

Theorem 2.4 The above proximal algorithm terminates at some k with a $y_k \in F$ which is a local minimum of $\varphi_{\rho'}$ for any $\rho' > \rho$.

Proof. The y_k 's can take on finitely many values; in view of Lemma 1.5, $y_{k+1} - y_k$ tends to 0; hence $y_{k+1} - y_k = 0$ at some k . By construction, this means $x_k = y_k$ minimizes the function $x \mapsto g(x) + \rho\|x - y_k\|^2$:

$$\varphi_{\rho}(y_k) = g(y_k) \leq g(x) + \rho\|x - y_k\|^2, \quad \text{for all } x \in F;$$

the inequality becomes strict if $x \neq y_k$ and ρ is replaced by $\rho' > \rho$. To finish, apply Proposition 2.1. \square

Let us sum up this §2.

- (i) First remember Lemma 1.1 (iv): when ρ grows from 0 to $+\infty$, the local minima of φ_{ρ} form nested sets, growing from the “ideal” set of optimal solutions of (8), to the “worst” whole feasible set. It is therefore advantageous to take a “small” ρ (whatever this means).
- (ii) The proximal algorithm produces such a local minimum – a feasible point for (8). In terms of the objective function g , the quality of this point depends on
 - the initialization: in view of Lemma 1.5, the objective function is improved by at least $\rho\|y_{k+1} - y_k\|^2$ at each iteration; accordingly, a “good” point will be obtained if y_1 is itself “good”;
 - the value of ρ : in view of Proposition 2.2, only an optimal solution can be produced if ρ is small enough.
 - Note that, if we were able to guarantee a *global* minimum of φ_{ρ} , instead of local, then we would for sure have an optimal solution (Theorem 1.3).

Unfortunately, this algorithm is only conceptual anyway: φ_{ρ} cannot be computed exactly – a fortiori minimized globally.

3 Discrete Optimization Problems: Implementable Forms

From now on, we assume the feasible set F of (8) to be some structured set of 0-1 points; for convenience, we also assume g to be linear. Thus we consider a frequent situation in combinatorial optimization:

$$\min b^{\top} x, \quad x \in F := \{x \in G : Ax \geq c \in \mathbb{R}^m\}, \quad G \subset \{0, 1\}^n. \quad (11)$$

We may have $G = \{0, 1\}^n$ but common instances have

$$G = \{x \in \{0, 1\}^n : Bx \geq d \in \mathbb{R}^p\}. \quad (12)$$

The important feature is that G is assumed to be an “easy polyhedron”, in the sense that linear functions can “easily” be minimized over it; then φ_{ρ} can be underestimated via Lagrangian relaxation.

3.1 Introducing Lagrangian Relaxation

Here, with the extra variable $\lambda \in \mathbb{R}_+^m$, we introduce the Lagrangian associated with (11) and the corresponding relaxation $\tilde{\varphi}_\rho$ of φ_ρ :

$$\begin{aligned}\tilde{\varphi}_\rho(y) &:= \sup_{\lambda \geq 0} \theta_\rho(\lambda, y), \quad \text{where} \\ \theta_\rho(\lambda, y) &:= \min_{x \in G} \{b^\top x + \lambda^\top (c - Ax) + \rho \|x - y\|^2\}.\end{aligned}\tag{13}$$

The function $\theta_\rho(\cdot, y)$ defined above is piecewise linear and its maximization is a standard convex optimization problem. On the other hand, computing $\theta_\rho(\lambda, y)$ (for given λ and y) is “easy” because, using a standard trick in combinatorial optimization, the quadratic term $\|x - y\|^2$ can be “linearized”; see (18) below.

A crucial object in this approach is the bounded polyhedron obtained by convexifying G : we set

$$P := \{x \in \text{co } G : Ax \geq c\}, \quad \tilde{F} := \text{ext } P\tag{14}$$

(co denotes the convex hull, ext the set of extreme points). It is useful to understand that, being a set of 0-1 points, F in (11) is also the set of extreme points in its own convex hull; \tilde{F} is made up of F , plus some “parasitic” extreme points, which are all fractional. For more on Lagrangian relaxation, see for example [10] and the references therein.

We will need some more notation:

$$\ell_\rho(x, y) := b^\top x + \rho(e - 2y)^\top x, \quad g_\rho(x) := (b + \rho e)^\top x - \rho \|x\|^2,\tag{15}$$

where $e := (1, \dots, 1)$ is the vector of all ones;

$$v^* := \min_{x \in F} b^\top x, \quad v_c := \sup_{\lambda \geq 0} \min_{x \in G} \{b^\top x + \lambda^\top (c - Ax)\}\tag{16}$$

are respectively the optimal value of (11) and of its relaxation (naturally $v^* = \varphi_0(y)$ and $v_c = \tilde{\varphi}_0(y)$ for all y).

Lemma 3.1 *The function $\rho \mapsto \tilde{\varphi}_\rho(y)$ is nondecreasing and $v_c \leq \tilde{\varphi}_\rho(y) \leq \varphi_\rho(y)$ for all y . It follows that*

$$v_c \leq \inf_{y \in \mathbb{R}^n} \tilde{\varphi}_\rho(y) \leq v^*.$$

Proof. Weak duality (see [10] for example) guarantees $\tilde{\varphi}_\rho(y) \leq \varphi_\rho(y)$. Furthermore, the Lagrangian $b^\top x + \lambda^\top (c - Ax) + \rho \|x - y\|^2$ in (13) is obviously a nondecreasing function of ρ ; and this property is transmitted to the infima and suprema. The rest follows easily, use in particular (16) and Theorem 1.3 (i). \square

The form (13) of $\tilde{\varphi}_\rho$, as well as the expression of v_c in (16), are not easy to deal with, we express them via *minimization* problems:

Proposition 3.2 *Use the notation (15), (16). There holds*

$$v_c = \min_{x \in P} b^\top x = \min_{x \in \tilde{F}} b^\top x\tag{17}$$

and the function $\tilde{\varphi}_\rho$ has any of the following expressions:

$$\begin{aligned}\tilde{\varphi}_\rho(y) &= \min_{x \in P} \ell_\rho(x, y) + \rho \|y\|^2 \\ &= \min_{x \in \tilde{F}} \ell_\rho(x, y) + \rho \|y\|^2,\end{aligned}\tag{18}$$

$$\begin{aligned}\tilde{\varphi}_\rho(y) &= \min_{x \in P} [g_\rho(x) + \rho \|x - y\|^2] \\ &= \min_{x \in \tilde{F}} [g_\rho(x) + \rho \|x - y\|^2].\end{aligned}\tag{19}$$

Proof. We prove (18) for $\rho \geq 0$; this will prove (19): in fact both problems have the same minimand. Then (17) will follow for $\rho = 0$.

Observe that $\|x\|^2 = e^\top x$ for $x \in G \subset \{0, 1\}^n$. Then develop $\|y - x\|^2$ in (13) to realize that

$$\tilde{\varphi}_\rho(y) = \sup_{\lambda \geq 0} \min_{x \in G} \{ \ell_\rho(x, y) + \lambda^\top (c - Ax) \} + \rho \|y\|^2.$$

Then recognize duality applied to the problem

$$\min \{ \ell_\rho(x, y) : x \in G, Ax \geq c \}.$$

This is classically equivalent to convexifying G (see for example [4, 6, 12]). Finally, minimizing the linear function $\ell_\rho(\cdot, y)$ over the bounded polyhedron P gives the same values if the minimization is restricted to the extreme points of P . \square

Remark 3.3 *Thanks to the extreme simplicity of the minimization problem in (18) (the set \tilde{F} is finite), the function $\tilde{\varphi}_\rho$ is continuous; besides it increases at infinity (as fast as the squared norm). As a result, it does have some minimum point y^* .* \square

To reproduce the results of Sections 1 and 2, we will use either the form (18) – which is simple enough –, or (19) – which has the general form (10), but with the substantial difference that the function g depends on ρ .

Proposition 3.4 *The point y_0 is a local minimum of $\tilde{\varphi}_\rho$ if and only if $x = y_0$ is the unique optimal solution in (18), (19) with $y = y_0$. This implies in particular that y_0 lies in \tilde{F} , is a strict local minimum of $\tilde{\varphi}_\rho$, and that*

$$\tilde{\varphi}_\rho(y_0) = \ell_\rho(y_0, y_0) + \rho \|y_0\|^2 = g_\rho(y_0).$$

Proof. Apply Proposition 2.1 to the form (19) of $\tilde{\varphi}_\rho$. \square

The lower bound obtained by minimizing $\tilde{\varphi}_\rho$ does improve the relaxed value v_c of (16). We can even prove slightly more:

Proposition 3.5 *For $\rho > 0$ small enough, any global minimizer y_ρ of $\tilde{\varphi}_\rho$ also minimizes $x \mapsto b^\top x$ over \tilde{F} , and therefore satisfies $\tilde{\varphi}_\rho(y_\rho) = v_c + \rho(e^\top y_\rho - \|y_\rho\|^2)$.*

As a result, if there is a duality gap, then

$$v_c < \inf \tilde{\varphi}_\rho \leq v^*, \quad \text{for all } \rho > 0.$$

Proof. For any $y \in \tilde{F}$, define the number

$$\varepsilon(y) := \inf\{\rho > 0 : y \text{ minimizes (globally) } \tilde{\varphi}_\rho\}.$$

This is a nonnegative number, possibly $+\infty$. Then define $\hat{F} := \{y \in \tilde{F} : \varepsilon(y) > 0\}$ and set

$$\varepsilon := \min\{\varepsilon(y) : y \in \hat{F}\};$$

note that $\varepsilon > 0$ because \hat{F} is a finite set.

Fix $\rho < \varepsilon$ and let y_ρ be an arbitrary global minimizer of $\tilde{\varphi}_\rho$; clearly $y \in \tilde{F} \setminus \hat{F}$, so $\varepsilon(y_\rho) = 0$. This means that there exists a sequence $\rho_k \downarrow 0$ such that y_ρ is a global minimizer of $\tilde{\varphi}_{\rho_k}$. In view of Proposition 3.4, $x = y_\rho$ minimizes $\ell_{\rho_k}(\cdot, y_\rho)$:

$$b^\top x + \rho_k(e - 2y_\rho)^\top x \geq b^\top y_\rho + \rho_k(e - 2y_\rho)^\top y_\rho \quad \text{for all } x \in \tilde{F} \text{ and } k = 1, 2, \dots$$

Letting $\rho_k \downarrow 0$ shows that y_ρ minimizes $x \mapsto b^\top x$ over \tilde{F} : $b^\top y_\rho = v_c$.

As a result, $\inf \tilde{\varphi}_\rho = \tilde{\varphi}_\rho(y_\rho) = v_c + \rho(e^\top y_\rho - \|y_\rho\|^2) \geq v_c$. If $\tilde{\varphi}_\rho(y_\rho)$ were equal to v_c , y_ρ would be a 0-1 point, lying in F and there would be no duality gap. The rest follows from monotonicity (Lemma 3.1). \square

Proposition 3.6 *For ρ large enough, the local minima of $\tilde{\varphi}_\rho$ contain the whole feasible set F of (11).*

Proof. Take $\rho > |b|_\infty := \max_i |b_i|$ and let $y_0 \in F$; in particular, $y_0 \in \{0, 1\}^n$. For any $x \in \tilde{F} \subset [0, 1]^n$ and $i = 1, \dots, n$, consider two cases:

- if $y_0^i = 0$, then $[b + \rho(e - 2y_0)]^i (x - y_0)^i = (b^i + \rho)(x^i) \geq 0$,
the inequality being strict if $x^i > 0 = y_0^i$;
- if $y_0^i = 1$, then $[b + \rho(e - 2y_0)]^i (x - y_0)^i = (b^i - \rho)(x^i - 1) \geq 0$,
the inequality being strict if $x^i < 1 = y_0^i$.

Thus, we see by summation that

$$\ell_\rho(x, y_0) - \ell_\rho(y_0, y_0) = [b + \rho(e - 2y_0)]^\top (x - y_0) \geq 0,$$

the inequality being strict if $x \neq y_0$. Because $y_0 \in F \subset \tilde{F}$, the only optimal solution in (18) is clearly $x = y_0$. The result follows from Proposition 3.4. \square

Finally we show that the upper bound v^* is attained in Proposition 3.5.

Proposition 3.7 *For ρ large enough, the global minima of $\tilde{\varphi}_\rho$ are exactly the optimal solutions of (11).*

Proof. Let ρ be so large that the local minima of $\tilde{\varphi}_\rho$ contain the whole of F (Proposition 3.6). Consider first the local minima y_0 of $\tilde{\varphi}_\rho$ that lie in F . They satisfy (Proposition 3.4) $\tilde{\varphi}_\rho(y_0) = b^\top y_0 \geq v^*$, where equality holds exactly for those y_0 's solving (11).

Now set $\alpha := \min\{e^\top y - \|y\|^2 : y \in \tilde{F} \setminus F\}$; this is a positive number because $\tilde{F} \setminus F$ is made up of finitely many fractional points. Increase ρ if necessary so that $\rho\alpha > v^* - v_c$ and let y_0 be a local minimum of $\tilde{\varphi}_\rho$ lying in $\tilde{F} \setminus F$. It satisfies

$$\tilde{\varphi}_\rho(y_0) = b^\top y_0 + \rho(e^\top y_0 - \|y_0\|^2) \geq b^\top y_0 + \rho\alpha > b^\top y_0 + v^* - v_c \geq v^*,$$

where the last inequality comes from (17). In view of Lemma 3.1, y_0 cannot be a global minimum of $\tilde{\varphi}_\rho$. \square

3.2 An Example

As an illustrative example, consider the problem

$$\min_{x \in F} \{x^1 + 4x^2\}, \quad \text{where } F := \{x \in \{0, 1\}^2 : x^1 + 2x^2 \geq 2\}. \quad (20)$$

The feasible points are $x^* := (0, 1)$ (the optimal solution) and $e = (1, 1)$, so that

$$\varphi_\rho(y) = \min \{4 + \rho\|y - x^*\|^2, 5 + \rho\|y - e\|^2\}.$$

Working out the calculations shows that

$$\varphi_\rho(y) = \begin{cases} 4 + \rho\|y - x^*\|^2 & \text{if } (e - x^*)^\top y \leq \frac{1+\rho}{2\rho}, \\ 5 + \rho\|y - e\|^2 & \text{otherwise.} \end{cases}$$

First observe that $(e - x^*)^\top x^* = 0$ is always smaller than $(1 + \rho)/2\rho$, hence $y = x^*$ is always a local minimum. Now consider three cases:

- If $(1 + \rho)/2\rho < 1$ (i.e. $\rho > 1$) then $y = e$ is another local minimum – Proposition 2.2 (i).
- If $(1 + \rho)/2\rho > 1$ (i.e. $\rho < 1$), this latter local minimum vanishes – Proposition 2.2 (ii).
- We leave it to the reader to check the case $(1 + 2\rho)/2\rho = 1$, and in particular to see why $y = e$ is not a local minimum.

Now we study $\tilde{\varphi}_\rho$. The relaxed solution is $x_c = (1, 1/2)$ (see (16)), with objective value $b^\top x_c = 3$; P is the triangle whose vertices make up $\tilde{F} = \{x^*, e, x_c\}$. Then

$$\tilde{\varphi}_\rho(y) = \min \{\ell_\rho(x^*, y), \ell_\rho(e, y), \ell_\rho(x_c, y)\} + \rho\|y\|^2,$$

where

$$\begin{aligned} \ell_\rho(x^*, y) &= 4 + \rho - 2\rho y^2, \\ \ell_\rho(e, y) &= 5 + 2\rho - 2\rho(y^1 + y^2), \\ \ell_\rho(x_c, y) &= 3 + 3\rho/2 - \rho(2y^1 + y^2). \end{aligned}$$

Calculations are left to the reader. The final results are as indicated in Table 1. Note: x^* becomes the global minimum when $3 + \rho/4$ becomes larger than 4, i.e. when $\rho \geq 4$; compare with the theoretical results of Proposition 3.6, 3.7.

ρ	0	1/3	4	$+\infty$
local minima	x_c	x_c	x^*	x_c x^* e
$\tilde{\varphi}_\rho$ values	$3 + \rho/4$	$3 + \rho/4$	4	$3 + \rho/4$ 4 5

Table 1: Local minima of $\tilde{\varphi}_\rho$.

3.3 The Relaxed Proximal Algorithm

Consider now the proximal algorithm (7) to minimize $\tilde{\varphi}_\rho$. It needs a black box to compute $\tilde{\varphi}_\rho(y)$ for a given $y \in \mathbb{R}^n$. Of course this is done by some optimization process, which produces an $x(y)$ solving one of the “equivalent” problems (13), (18) or (19).

Then we do the following.

Algorithm 3.8 (Implementable proximal algorithm I) *A black box is assumed available to compute $x(y)$ for given y . Choose $\rho > 0$.*

STEP 0. Take y_1 arbitrary in \mathbb{R}^n ; set $k = 1$.

STEP 1. Call the black box to obtain $x(y_k)$.

STEP 2. If $x(y_k) = y_k$ then stop.

STEP 3. Set $y_{k+1} = x(y_k)$, replace k by $k + 1$ and go to Step 1. \square

Theorem 3.9 Assume that $x(y_k) \in \tilde{F}$ at each iteration k . Then the stop in Step 2 occurs at some finite iteration K .

Proof. By construction, the whole sequence $\{y_k\}$ lies in \tilde{F} , a finite set. Applying Lemma 1.5 to the form (19), we see that the sequence has to be finite. \square

Remark 3.10 This result outlines some weaknesses of Algorithm 3.8.

First, little can be said about the output y_K : all we know is that $y_K \in \text{Argmin} \ell_\rho(\cdot, y_K)$ – which does not imply that y_K is a local minimum of $\tilde{\varphi}_\rho$.

Besides, the assumption $x(y_k) \in \tilde{F}$ is rather dary:

- either we solve (13) by some dual algorithm; this produces a point in P , which has no reason to lie in \tilde{F} ;
- or we solve (18) by the simplex algorithm; but this supposes a close description of the polyhedron P – which, in practice, amounts to assuming $G = \{0, 1\}^n$.

\square

As a conclusion, let us compare with the situation in §2.

- (i) Replacing φ_ρ by $\tilde{\varphi}_\rho$ has a substantial price. Instead of producing a feasible point for sure (Theorem 2.4), we may land at some parasitic point in $\tilde{F} \setminus F$, including the relaxed solution x_c from (16).
- (ii) Taking a small ρ , which was recommended, is now unwise, since this will probably produce the relaxed solution x_c (Proposition 3.5). Note at this point that $y_1 = x_c$ is probably the most natural initialization in Algorithm 3.8.
- (iii) The only way to escape from x_c is to increase ρ . However this is dangerous, since the algorithm might produce a bad feasible point of (11) (Proposition 3.6); besides, it is not sure that larger values of ρ will eventually eliminate x_c from the set of local minima.

Note of course that rounding x_c may not produce any optimal point; it may not even produce any feasible point (replace in §3.2 the inequality constraint by the equality $x^1 + 2x^2 = 2$).

- (iv) Once again, global minimization of $\tilde{\varphi}_\rho$ would solve the problem; yet we should also take ρ large enough (Proposition 3.7).

3.4 Penalizing the 0-1 Constraints

The trouble with the previous convexification procedure is that the quadratic term can by no means convexify the objective function f of (9). In fact, the original motivation for [1] was to treat “ordinary” (continuous) nonlinear programming problems, whose Lagrangian had a Hessian which could be made positive definite. To put (11) into this mould, we introduce the penalty factor

$$p(x) := e^\top x - \|x\|^2, \quad (21)$$

we take a large parameter $\kappa > 0$, and we consider problems of the type

$$\min [b^\top x + \kappa p(x)], \quad x \in Q \subset [0, 1]^n, \quad (22)$$

Q being a convex polyhedron. To define Q , we may for example just change $\{0, 1\}$ to $[0, 1]$ in the definition of F ; it is generally accepted that this does not change (11). We give a proof of a slightly more general result:

Theorem 3.11 *Assume that the convex polyhedron Q in (22) contains the feasible set F of (11), but does not contain any other 0-1 point than those in F .*

(i) *Any optimal solution of (22) lying in F is also optimal for (11).*

(ii) *For κ large enough, the sets of optimal solutions in (11) and (22) coincide.*

Proof. (i) Just observe that (22) is a *relaxation* of (11): their objective functions coincide on the feasible set $F \subset \{0, 1\}^n$ of the latter.

(ii) Use the notation (16). Because $p \geq 0$ on $Q \subset [0, 1]^n$, v^* is an upper bound for the optimal value of (22). Because p is strictly concave on $Q \subset [0, 1]^n$, the feasible set in (22) can be restricted to the set $\text{ext } Q$ of its extreme points. Denote by $\bar{F} := \text{ext } Q \setminus F$ the set of parasitic points. Set $\delta := \min_{x \in \bar{F}} p(x)$ and note that $\delta > 0$ because $p > 0$ on the finite set \bar{F} .

Take $\kappa > \frac{(v^* - v_c)}{\delta}$, let $x^* \in \text{ext } Q$ solve (22) and assume $x^* \in \bar{F}$:

$$b^\top x^* + \kappa p(x^*) \geq v_c + \kappa \delta > v^*,$$

a contradiction. Thus, any x^* optimal in (22) lies in F ; in view of (i), x^* solves (11). Besides, the optimal value of (22) is v^* . Conversely, any \hat{x} optimal in (11) is feasible in (22) and has $b^\top \hat{x} + \kappa p(\hat{x}) = b^\top \hat{x} = v^*$: \hat{x} is optimal in (22). \square

Thus, the proximal convexification procedure can be applied to (22) as well: we define

$$\psi_{\rho\kappa}(y) := \min_{x \in Q} [b^\top x + \kappa p(x) + \rho \|y - x\|^2] \quad (23)$$

and all the results of §1 apply, $\psi_{\rho\kappa}$ playing the role of φ_ρ . Observe that the minimand $f_\rho(\cdot, y)$ in (23) is convex for $\rho \geq \kappa$: in contrast with φ_ρ , the computation of $\psi_{\rho\kappa}$ is straightforward. Its global minimization is not straightforward, though: with relation to Theorem 1.4, the Hessian of $f_\rho(\cdot, \cdot)$ is $2 \begin{pmatrix} (\rho - \kappa)I & -\rho I \\ -\rho I & \rho I \end{pmatrix}$ (I is the identity in \mathbb{R}^n), which is never positive semidefinite.

Algorithm 3.12 (Implementable proximal algorithm II) *A black box is assumed available to solve (23) for a given y . Take $\rho > \kappa$.*

STEP 0. *Take y_1 arbitrary in \mathbb{R}^n ; set $k = 1$.*

STEP 1. *Call the black box to obtain $x(y_k) \in Q$.*

STEP 2. *If $x(y_k) = y_k$ then stop.*

STEP 3. *Set $y_{k+1} = x(y_k)$, replace k by $k + 1$ and go to Step 1.* \square

Of course, the choice $\rho > \kappa$ implies that the minimum in (23) is unique: $x(y_k)$ is well-defined in Step 2. Another reason for this choice will become apparent in Proposition 3.15 below.

Theorem 3.13 *The sequence y_k generated by the above algorithm has some cluster point, and any such cluster point is a local minimum y^* of $\psi_{\rho\kappa}$ on Q .*

Proof. The first statement holds because y_k varies in Q , a compact set. Take a subsequence – also denoted by y_k – such that $y_k \rightarrow y^*$. Because of Lemma 1.5, $x(y_k) - y_k \rightarrow 0$, hence $x(y_k) \rightarrow y^*$. Then pass to the limit in the inequality

$$b^\top x + \kappa p(x) + \rho \|x - y_k\|^2 \geq b^\top x(y_k) + \kappa p(x(y_k)) + \rho \|x(y_k) - y_k\|^2 \quad \text{for all } x \in Q$$

to see that $x = y^*$ is the unique minimum point in (23) for $y = y^*$. The result follows from Proposition 2.1. \square

Remark 3.14 *Theorem 3.11 allows a large choice for the feasible set Q in (22).*

- *The most natural is $Q = P$ of (14); however, this requires a description of $\text{co } G$, which may not be easy numerically.*
- *When G has the form (12), one may also take the ordinary LP relaxation:*

$$Q = \{x \in [0, 1]^n : Ax \geq c, Bx \geq d\};$$

then (23) is an ordinary linear-quadratic program. However, this Q is then larger than P and may introduce more parasitic local minima. \square

The minimand of (23) is linear in x when $\rho = \kappa$, and this turns out to reproduce §3.1:

Proposition 3.15 *Suppose Q in (22) is P of (14).*

Then $\psi_{\rho\rho}(y) = \tilde{\varphi}_\rho(y)$ for all $y \in \mathbb{R}^n$ and $\rho > 0$. It follows that the local minima of $\tilde{\varphi}_\rho$ are also local minima of $\psi_{\rho'\rho}$, for any $\rho' \geq \rho$.

Proof. For all x, y ,

$$b^\top x + \rho p(x) + \rho \|x - y\|^2 = b^\top x + \rho(e - 2y)^\top x + \rho \|y\|^2,$$

in which we recognize the respective minimands in (23) (with $\kappa = \rho$) and in (18). Minimizing them gives the same value.

Now invoke Lemma 1.1 (iv): the set of local minima of $\psi_{\rho\kappa}$ can only increase when ρ increases. \square

Thus, the additional flexibility yielded by $\rho > \kappa$ improves nothing: this can only enlarge the set of local minima. The proximal algorithm has more chances to produce a parasitic point. A small bonus is obtained with respect to Algorithm 3.8, though: if the algorithm stops at Step 2, the corresponding y_k is *for sure* a local minimum of $\psi_{\rho\kappa}$ (Theorem 3.13).

General Conclusion We have studied in the framework of combinatorial optimization a general convexification procedure (the primal proximal algorithm), assessed by a useful heuristic for some special problem (unit-commitment). This procedure gives birth to various conceptual and implementable heuristic algorithms to generate primal solutions, and our results suggest that little is to be expected in theory from the approach.

First it should be mentioned that the assumptions allowing our study of implementable algorithms (§3) do not fit easily with the actual form of the unit-commitment problem: thermal plants and hydro valleys involve fairly more sophisticated models than (11).

More importantly, the ability itself of the approach in producing feasible suboptimal solutions is controversial. From our results in §3.1, such an ability implies a delicate tuning of the proximity parameter, to avoid

Charybdis: staying with a small ρ at the relaxed solution x_c ,
as well as

Scylla: jumping with a large ρ to an uncontrollable feasible point in (8).

Yet, even if these two dangers are avoided, the mere production of a feasible point is never guaranteed; see again our discussion at the end of §3.1.

References

- [1] D.P. Bertsekas. Convexification procedures and decomposition methods for nonconvex optimization problems. *Journal of Optimization Theory and Applications*, 29:169–197, 1979.
- [2] D.P. Bertsekas, G.S. Lauer, N.R. Sandell, and T.A. Posberg. Optimal short-term scheduling of large-scale power systems. *IEEE Transactions on Automatic Control*, AC-28:1–11, 1983.
- [3] J.M. Danskin. The theory of max-min with applications. *SIAM Journal on Applied Mathematics*, 14(4):641–655, 1966.
- [4] J.E. Falk. Lagrange multipliers and nonconvex programs. *SIAM Journal on Control*, 7(4):534–545, 1969.
- [5] S. Feltenmark and K. C. Kiwiel. Dual applications of proximal bundle methods, including lagrangian relaxation of nonconvex problems. *SIAM Journal on Optimization*, 10(3):697–721, 2000.
- [6] A.M. Geoffrion. Lagrangean relaxation for integer programming. *Mathematical Programming Study*, 2:82–114, 1974.
- [7] R. Gonzalez and C. Lemaréchal. Unit-commitment problem: from the dual to the primal. In preparation.
- [8] J.-B. Hiriart-Urruty and C. Lemaréchal. *Convex Analysis and Minimization Algorithms*. Springer Verlag, Heidelberg, 1993. Two volumes.
- [9] J.-B. Hiriart-Urruty and C. Lemaréchal. *Fundamentals of Convex Analysis*. Springer Verlag, Heidelberg, 2001.
- [10] C. Lemaréchal. Lagrangian relaxation. In M. Jünger and D. Naddef, editors, *Computational Combinatorial Optimization*, pages 115–160. Springer Verlag, Heidelberg, 2001.
- [11] C. Lemaréchal, F. Pellegrino, A. Renaud, and C. Sagastizábal. Bundle methods applied to the unit-commitment problem. In J. Doležal and J. Fidler, editors, *System Modelling and Optimization*, pages 395–402, 1996.

- [12] C. Lemaréchal and A. Renaud. A geometric study of duality gaps, with applications. *Mathematical Programming*, 90(3):399–427, 2001.
- [13] R.T. Rockafellar and R.J.-B. Wets. *Variational Analysis*. Springer Verlag, Heidelberg, 1998.



Unité de recherche INRIA Rhône-Alpes

655, avenue de l'Europe - 38330 Montbonnot-St-Martin (France)

Unité de recherche INRIA Lorraine : LORIA, Technopôle de Nancy-Brabois - Campus scientifique

615, rue du Jardin Botanique - BP 101 - 54602 Villers-lès-Nancy Cedex (France)

Unité de recherche INRIA Rennes : IRISA, Campus universitaire de Beaulieu - 35042 Rennes Cedex (France)

Unité de recherche INRIA Rocquencourt : Domaine de Voluceau - Rocquencourt - BP 105 - 78153 Le Chesnay Cedex (France)

Unité de recherche INRIA Sophia Antipolis : 2004, route des Lucioles - BP 93 - 06902 Sophia Antipolis Cedex (France)

Éditeur

INRIA - Domaine de Voluceau - Rocquencourt, BP 105 - 78153 Le Chesnay Cedex (France)

<http://www.inria.fr>

ISSN 0249-6399