

Limit theorems for extreme value estimates of point processes boundaries

Stéphane Girard, Ludovic Menneteau

► **To cite this version:**

Stéphane Girard, Ludovic Menneteau. Limit theorems for extreme value estimates of point processes boundaries. [Research Report] RR-4366, INRIA. 2002. inria-00072222

HAL Id: inria-00072222

<https://hal.inria.fr/inria-00072222>

Submitted on 23 May 2006

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

*Limit theorems for extreme value estimates of point
processes boundaries*

Stéphane Girard — Ludovic Menneteau

N° 4366

Février 2002

THÈME 4



*Rapport
de recherche*

Limit theorems for extreme value estimates of point processes boundaries

Stéphane Girard , Ludovic Menneteau *

Thème 4 — Simulation et optimisation
de systèmes complexes
Projet is2

Rapport de recherche n° 4366 — Février 2002 — 36 pages

Abstract: We give sufficient conditions to establish central limit theorems and moderate deviation principles for support estimates of Poisson point processes. The considered estimates write as linear combinations of extreme values of the point process. Our results are illustrated on four particular cases: Haar and trigonometric series estimates, Faber-Shauder estimate and kernel estimate. A hierarchy between these estimators is proposed by comparing their optimal convergence rates.

Key-words: Functional estimate, Central limit theorem, Moderate deviation principle, Extreme values, Poisson process, Shape estimation.

* Laboratoire de Probabilités et Statistique, Université Montpellier 2, Place Eugène Bataillon, 34095 Montpellier Cedex 5, France. mennet@stat.math.univ-montp2.fr

Théorèmes limites pour l'estimation à base de valeurs extrêmes du contour de processus ponctuels

Résumé : Nous donnons des conditions suffisantes permettant d'établir un théorème central limite ainsi qu'un principe de déviations modérées pour des estimateurs du support de processus de Poisson. Les estimateurs considérés s'écrivent comme une combinaison linéaire de valeurs extrêmes du processus ponctuel. Nos résultats sont illustrés dans quatre cas : méthode des fonctions orthogonales (base de Haar et base trigonométrique), estimateur utilisant la base de Faber-Shauder et méthode du noyau. Nous proposons une hiérarchie parmi cette famille d'estimateurs basée sur la comparaison des vitesses optimales de convergence.

Mots-clés : estimation fonctionnelle, théorème central limite, principe de déviations modérées, valeurs extrêmes, processus de Poisson, estimation de contour.

1 Introduction

Many proposals are given in the literature for estimating a set S given a finite random set N of points drawn from the interior. This problem of edge or support estimation arises in classification (HARDY & RASSON [20]), clustering problems (HARTIGAN [21]), discriminant analysis (BAUFAYS & RASSON [2]), and outliers detection (DEVROYE & WISE [6]). Applications are also found in image analysis (KOROSTELEV & TSYBAKOV [26]). For instance, the segmentation problem can be considered under the support estimation point of view, where the support is a connex bounded set in \mathbb{R}^2 . We also point out some applications in econometrics (e.g. DEPRINS, *et al* [5]). In such cases, the unknown support can be written

$$S = \{(x, y) : 0 \leq x \leq 1 ; 0 \leq y \leq f(x)\}, \quad (1.1)$$

where f is an unknown function. Here, the problem reduces to estimating f , called the production frontier (see for instance HÄRDLE *et al* [17]). The data consist of pair (X, Y) where X represents the input (labor, energy or capital) used to produce an output Y in a given firm. In such a framework, the value $f(x)$ can be interpreted as the maximum level of output which is attainable for the level of input x . KOROSTELEV *et al* [25] suppose f to be increasing and concave, from economical considerations, which suggests an adapted estimator, called the DEA (Data Envelopment Analysis) estimator. Its asymptotic distribution is established by GIJBELS *et al* [11].

Here N is a Poisson point process, with observed points belonging to a subset S defined as in (1.1) where f is an unknown function which needs not to be monotone. An early paper was written by GEFFROY [9] for independent identically distributed observations from a density ϕ . The proposed estimator is a kind of histogram based on the extreme values of the sample. This work was extended in two main directions.

- (a) On the one hand, piecewise polynomials were introduced and their optimality in an asymptotic minimax sense is proved under weak assumptions on the rate of decrease α of the density ϕ towards 0 by KOROSTELEV & TSYBAKOV [26] and by HÄRDLE *et al* [18]. Extreme values methods are then proposed by HALL *et al* [16] and by GIJBELS & PENG [10] to estimate the parameter α .
- (b) On the other hand, different propositions for smoothing Geffroy's estimate were made. GIRARD & JACOB [14] introduced estimates based on kernel regressions and orthogonal series method [12, 13]. In the same spirit, GARDES [8] proposed a Faber-Shauder estimate. In each case, the consistency and the limit distribution of the estimator are established.

We also refer to ABBAR [1] and JACOB & SUQUET [23] who used a similar smoothing approach, although their estimates are not based on the extreme values of the Poisson process.

The work presented here offers a general framework for studying the estimates of the family (b). We consider estimates writing as linear combinations of the extreme values of the Poisson process. We establish general central limit theorems and moderate deviation principles

for our estimates and we apply them to the particular cases quoted in (b). Such an approach allows to propose a hierarchy within this family of estimates according to their convergence rates.

2 The boundary estimate

Let $S \subset \mathbb{R}^2$ and consider a sequence of Poisson point processes

$$N_n = \{N_n(D) : D \in \mathcal{B}(S)\}, \quad n \geq 1,$$

with intensity measure $nc\lambda$, where $c > 0$, and λ is the Lebesgue measure on $\mathcal{B}(S)$, the Borel σ -algebra of S . Let $\{(X_{n,i}, Y_{n,i}), 1 \leq i \leq N_n\}$ be the set of points associated to the point process. In the following, we assume that there exists a function $f : [0, 1] \rightarrow \mathbb{R}^+$, such that S can be written as (1.1). Our aim is then to estimate S via an estimation of f . Let $k_n \uparrow \infty$ and define by $\{I_{n,r} : 1 \leq r \leq k_n\}$ a measurable equidistant partition of $[0, 1]$. For all $1 \leq r \leq k_n$, set

$$D_{n,r} = \{(x, y) : x \in I_{n,r}, 0 \leq y \leq f(x)\},$$

the cell of S built on $I_{n,r}$ and $N_{n,r} = N_n(D_{n,r})$. We introduce

$$Y_{n,r}^* = \max\{Y_{n,i} : (X_{n,i}, Y_{n,i}) \in D_{n,r}\},$$

if $D_{n,r} \neq \emptyset$ and $Y_{n,r}^* = 0$ otherwise. We use the convention $0 \times \infty = 0$. Our estimator of f is

$$\hat{f}_n : x \in [0, 1] \mapsto \frac{1}{k_n} \sum_{r=1}^{k_n} \kappa_{n,r}(x) \left(1 + \frac{1}{N_{n,r}}\right) Y_{n,r}^*, \quad (2.1)$$

where $\kappa_{n,r} : [0, 1] \rightarrow \mathbb{R}$ is an arbitrary weighting function determining the nature of the estimate. In the next section, conditions are imposed on $\kappa_{n,r}$ and examples are provided in Section 5. It is well-known that $Y_{n,r}^*$ is an estimator of the maximum of f on $I_{n,r}$ with negative bias. The use of the random variable $(1 + N_{n,r}^{-1})Y_{n,r}^*$ allows to reduce this bias. Therefore, \hat{f}_n appears as a linear combination of extreme value estimates of sampled values of f . The asymptotic properties of \hat{f}_n are established in Section 3, and proved in Section 4. Illustrations are presented in Section 5 on the estimates considered in family (b).

3 Main results

Define $m = \min\{f(x) : x \in [0, 1]\}$, $M = \max\{f(x) : x \in [0, 1]\}$ and

$$\kappa_n(x) = \left(\sum_{r=1}^{k_n} \kappa_{n,r}^2(x) \right)^{1/2}.$$

For all $1 \leq r \leq k_n$, set $m_{n,r} = \min\{f(x) : x \in I_{n,r}\}$, $M_{n,r} = \max\{f(x) : x \in I_{n,r}\}$ and

$$w_{n,r}(x) = \kappa_{n,r}(x)/\kappa_n(x). \quad (3.1)$$

We consider the following series of assumptions:

(H.1) $k_n \uparrow \infty$ and $k_n/n \rightarrow 0$ as $n \rightarrow \infty$.

(H.2) $0 < m \leq M < +\infty$ and

$$\Delta_n := \max_{1 \leq r \leq k_n} |M_{n,r} - m_{n,r}| = o(k_n/n) \text{ as } n \rightarrow \infty.$$

(H.3) For each $(x_1, \dots, x_d) \subset [0, 1]$, there exists a covariance matrix in \mathbb{R}^d

$$\Sigma_{(x_1, \dots, x_d)} = [\sigma_{i,j}]_{1 \leq i, j \leq d}$$

such that for all $1 \leq i, j \leq d$,

$$\sum_{r=1}^{k_n} w_{n,r}(x_i)w_{n,r}(x_j) \rightarrow \sigma_{i,j} \text{ as } n \rightarrow \infty.$$

(H.4) For all $x \in [0, 1]$,

$$\max_{1 \leq r \leq k_n} |w_{n,r}(x)| \rightarrow 0 \text{ as } n \rightarrow \infty.$$

(H.5) For all $x \in [0, 1]$,

$$\left| \frac{1}{k_n} \sum_{r=1}^{k_n} \kappa_{n,r}(x) - 1 \right| = o\left(\frac{\kappa_n(x)}{n}\right) \text{ as } n \rightarrow \infty.$$

(H.6) For all $x \in [0, 1]$,

$$\frac{1}{k_n} \sum_{r=1}^{k_n} \kappa_{n,r}(x)(f(x) - m_{n,r}) = o\left(\frac{\kappa_n(x)}{n}\right) \text{ as } n \rightarrow \infty.$$

(H.7) For all $x \in [0, 1]$,

$$\frac{1}{k_n} \sum_{r=1}^{k_n} |\kappa_{n,r}(x)| \max\left(\Delta_n, \exp\left(-m \frac{nc}{k_n}\right)\right) = o\left(\frac{\kappa_n(x)}{n}\right) \text{ as } n \rightarrow \infty.$$

Before proceeding, let us comment the assumptions. **(H.1)**–**(H.4)** are devoted to the control of the centered estimator $(\hat{f}_n - E \hat{f}_n)$. Assumption **(H.1)** imposes that the mean number of points in each $D_{n,r}$ goes to infinity. **(H.2)** controls the variations of the function f on $I_{n,r}$. More precisely it imposes that the mean number of points in $D_{n,r}$ above $m_{n,r}$ converges to 0. **(H.3)** is devoted to the multivariate aspects of the limit theorems. **(H.4)** imposes to all

the weight functions $\kappa_{n,r}(x)$ in the linear combination (2.1) to be approximatively of the same order. This is a natural condition to obtain an asymptotic Gaussian behavior. These assumptions are easy to verify in practice since they involve either f or $\kappa_{n,r}$ without mixing these two quantities. Assumptions **(H.5)**–**(H.7)** are devoted to the control of the bias term ($E \hat{f}_n - f$). They prevent it to be too important with respect to the variance of the estimate (which will reveal to be of order κ_n/n). Consequently, these three assumptions involve both the unknown function f and the weight functions $\kappa_{n,r}$. **(H.5)** imposes the mean of the coefficients in (2.1) to be approximatively 1. So, the estimate (2.1) can roughly be interpreted as a convex combination of extreme values. **(H.6)** is a localization condition: The weight function $\kappa_{n,r}$ should be concentrated on $I_{n,r}$. Then, the choice of proper weight functions requires to balance the antagonistic conditions **(H.4)** and **(H.6)**. Finally, **(H.7)** can be seen as a stronger version of **(H.2)**.

Our first result gives the multivariate central limit theorem for (\hat{f}_n) .

Theorem 1 *Under assumptions **(H.1)**–**(H.7)**, and for all $(x_1, \dots, x_d) \subset [0, 1]$,*

$$\left\{ \frac{nc}{\kappa_n(x_j)} (\hat{f}_n(x_j) - f(x_j)) : 1 \leq j \leq d \right\} \xrightarrow{d} N(0, \Sigma_{(x_1, \dots, x_d)}).$$

We present now some large deviation properties of (\hat{f}_n) (see Definition 1 of the appendix for the definition of the large deviation principle and see e.g. DEMBO & ZEITOUNI [4] for a general account on that topic).

For all $(x_1, \dots, x_d) \subset [0, 1]$ such that $\Sigma_{(x_1, \dots, x_d)}$ is regular, define

$$I_{(x_1, \dots, x_d)} : u \in \mathbb{R}^d \mapsto \frac{1}{2} {}^t u \Sigma_{(x_1, \dots, x_d)}^{-1} u.$$

The following family of large deviation principles is sometimes referenced in the literature as a moderate deviation principle. It could be used to measure the asymptotic performance of \hat{f}_n , see e.g. KALLENBERG [24] and references therein.

Theorem 2 *Under assumptions **(H.1)**–**(H.7)** and for all $(x_1, \dots, x_d) \subset [0, 1]$ and each $\varepsilon_n \downarrow 0$ such that $\Sigma_{(x_1, \dots, x_d)}$ is regular and*

$$\max_{1 \leq r \leq k_n} \max_{1 \leq j \leq d} |w_{n,r}(x_j)| = o(\sqrt{\varepsilon_n}),$$

the sequence of random vectors

$$\left\{ \sqrt{\varepsilon_n} \frac{nc}{\kappa_n(x_j)} (\hat{f}_n(x_j) - f(x_j)) : 1 \leq j \leq d \right\}$$

follows the large deviation principle in \mathbb{R}^d with speed (ε_n) and good rate function $I_{(x_1, \dots, x_d)}$.

In practice, c is not known and has to be estimated. In this aim, we introduce

$$\hat{c}_n = \frac{N_n}{n\hat{a}_n},$$

where

$$\hat{a}_n = \frac{1}{k_n} \sum_{r=1}^{k_n} \left(1 + \frac{1}{N_{n,r}}\right) Y_{n,r}^*$$

is an estimator of $a = \lambda(S)$. We then have the following corollary:

Corollary 1 *Theorem 1 and Theorem 2 still hold when c is replaced by \hat{c}_n .*

4 Proofs

The proofs of our main results are built as follow. First, we establish a multivariate central limit theorem and a moderate deviation principle for the finite dimensional projection of the centered process

$$\frac{nc}{\kappa_n(x)} \left(\hat{f}_n(x) - \mathbb{E} \left(\hat{f}_n(x) \right) \right), \quad x \in [0, 1]$$

(see Proposition 1 below). To this aim, by the general framework of the appendix (Theorem 3 and Theorem 4) it is sufficient to control the centered moments of

$$\xi_{n,r} = \left(1 + \frac{1}{N_{n,r}}\right) Z_{n,r}^* = \left(1 + \frac{1}{N_{n,r}}\right) \left(\frac{nc}{k_n}\right) Y_{n,r}^*.$$

This is achieved in Lemma 2 and Lemma 3. In a second time, we establish that the bias term

$$\frac{nc}{\kappa_n(x)} \left(\mathbb{E} \left(\hat{f}_n(x) \right) - f(x) \right)$$

vanishes when $n \uparrow \infty$ (see Proposition 2). Finally, we prove in Lemma 5 and Lemma 6 that c can be replaced by \hat{c}_n in the multivariate central limit theorem and in the moderate deviation principle. Before that, we introduce some new notations and definitions needed for our proofs. For all $1 \leq r \leq k_n$, set

$$D_{n,r}^- = \{(x, y) : x \in I_{n,r}, 0 \leq y \leq m_{n,r}\},$$

$$D_{n,r}^+ = \{(x, y) : x \in I_{n,r}, m_{n,r} < y \leq f(x)\},$$

$$N_{n,r}^- \text{ (resp. } N_{n,r}^+) = N_n(D_{n,r}^-) \text{ (resp. } N_n(D_{n,r}^+)),$$

$$Z_{n,r}^- \text{ (resp. } Z_{n,r}^+) = \left(\frac{nc}{k_n} \right) \max \{ Y_{n,i} : (X_{n,i}, Y_{n,i}) \in D_{n,r}^- \text{ (resp. } D_{n,r}^+) \},$$

if $D_{n,r}^-$ (resp. $D_{n,r}^+$) $\neq \emptyset$, and $Z_{n,r}^-$ (resp. $Z_{n,r}^+$) = 0 otherwise. Note that $\xi_{n,r}$ can be expanded as

$$\xi_{n,r} = Z_{n,r}^- + \gamma_{n,r}.$$

where

$$\gamma_{n,r} = \gamma_{n,r,1} + \gamma_{n,r,2} + \gamma_{n,r,3},$$

with

$$\gamma_{n,r,1} = (Z_{n,r}^+ - \mathbb{E}(Z_{n,r}^-)) \mathbb{I}_{\{N_{n,r}^+ > 0\}},$$

$$\gamma_{n,r,2} = (\mathbb{E}(Z_{n,r}^-) - Z_{n,r}^-) \mathbb{I}_{\{N_{n,r}^+ > 0\}},$$

and

$$\gamma_{n,r,3} = Z_{n,r}^* / N_{n,r}.$$

Some technical results are collected in the next lemma.

Lemma 1 *Under assumptions (H.1) and (H.2) we have*

i)

$$\max_{1 \leq r \leq k_n} \mathbb{P}(N_{n,r}^+ > 0) = O\left(\frac{n}{k_n} \Delta_n\right) = o(1). \quad (4.1)$$

ii) *For all $1 \leq r \leq k_n$, and any $t \in [0, m_{n,r}]$,*

$$\mathbb{P}(Z_{n,r}^- \leq t) = \exp(t - \lambda_{n,r}), \quad (4.2)$$

with

$$\lambda_{n,r} = \frac{nc}{k_n} m_{n,r}. \quad (4.3)$$

iii) *For all $1 \leq r \leq k_n$,*

$$\mathbb{E}(Z_{n,r}^-) = \lambda_{n,r} - (1 - e^{-\lambda_{n,r}}). \quad (4.4)$$

iv) *For all $1 \leq r \leq k_n$,*

$$\mathbb{V}(Z_{n,r}^-) = 1 - 2\lambda_{n,r}e^{-\lambda_{n,r}} - e^{-2\lambda_{n,r}}. \quad (4.5)$$

v) For all $1 \leq r \leq k_n$,

$$\mathbb{E} \left(\frac{Z_{n,r}^-}{N_{n,r}^-} \right) = 1 - e^{-\lambda_{n,r}} (1 + \lambda_{n,r}). \quad (4.6)$$

vi) For all $\ell \geq 1$ and $1 \leq r \leq k_n$,

$$\mathbb{E} \left(|Z_{n,r}^- - \mathbb{E}(Z_{n,r}^-)|^\ell \right) \leq 1 + \ell!. \quad (4.7)$$

vii) For all $\ell \geq 1$,

$$\max_{1 \leq r \leq k_n} \mathbb{E} \left(|\gamma_{n,r,1}|^\ell \right) \leq \left(1 + \frac{nc}{k_n} \Delta_n \right)^\ell \max_{1 \leq r \leq k_n} \mathbb{P}(N_{n,r}^+ > 0). \quad (4.8)$$

viii) For all $\ell \geq 1$,

$$\max_{1 \leq r \leq k_n} \mathbb{E} \left(|\gamma_{n,r,2}|^\ell \right) \leq (1 + \ell!) \max_{1 \leq r \leq k_n} \mathbb{P}(N_{n,r}^+ > 0). \quad (4.9)$$

ix) For all $\ell \geq 1$,

$$\max_{1 \leq r \leq k_n} \mathbb{E} \left(|\gamma_{n,r,3}|^\ell \right) \leq \ell! \left(1 + \left(1 + \frac{\Delta_n}{m} \right)^\ell \max_{1 \leq r \leq k_n} \mathbb{P}(N_{n,r}^+ > 0) \right). \quad (4.10)$$

Proof : i), ii), iii) and iv) are obtained by easy calculations.

v) By a well known property of poisson processes, $Z_{n,r}^- | N_{n,r}^-$ has the same distribution as the maximum of $N_{n,r}^-$ independent variables uniformly distributed on $[0, \lambda_{n,r}]$. Hence,

$$\begin{aligned} \mathbb{E} \left(\frac{Z_{n,r}^-}{N_{n,r}^-} \right) &= \mathbb{E} \left(\frac{\mathbb{I}_{\{N_{n,r}^- > 0\}}}{N_{n,r}^-} \mathbb{E}(Z_{n,r}^- | N_{n,r}^-) \right) = \lambda_{n,r} \mathbb{E} \left(\frac{\mathbb{I}_{\{N_{n,r}^- > 0\}}}{N_{n,r}^- + 1} \right) = \sum_{q=1}^{\infty} \frac{\lambda_{n,r}^{q+1}}{(q+1)!} e^{-\lambda_{n,r}} \\ &= 1 - e^{-\lambda_{n,r}} (1 + \lambda_{n,r}). \end{aligned}$$

vi) Note that

$$\mathbb{E} \left(|Z_{n,r}^- - \mathbb{E}(Z_{n,r}^-)|^\ell \right) = \ell \int_0^{+\infty} t^{\ell-1} \mathbb{P}(|Z_{n,r}^- - \mathbb{E}(Z_{n,r}^-)| > t) dt.$$

Moreover, by (4.2) and (4.4),

$$\begin{aligned} \mathbb{P}(|Z_{n,r}^- - \mathbb{E}(Z_{n,r}^-)| > t) &= \mathbb{P}(Z_{n,r}^- > \mathbb{E}(Z_{n,r}^-) + t) + \mathbb{P}(Z_{n,r}^- < \mathbb{E}(Z_{n,r}^-) - t) \\ &= [1 - \exp(t - \lambda_{n,r} + \mathbb{E}(Z_{n,r}^-))] \mathbb{I}_{[0, \lambda_{n,r} - \mathbb{E}(Z_{n,r}^-)]}(t) \\ &\quad + \exp(-t - \lambda_{n,r} + \mathbb{E}(Z_{n,r}^-)) \mathbb{I}_{[0, \mathbb{E}(Z_{n,r}^-)]}(t) \\ &\leq \mathbb{I}_{[0,1]}(t) + \exp(e^{-\lambda_{n,r}} - 1) e^{-t}. \end{aligned}$$

Hence,

$$\begin{aligned} \mathbb{E} \left(|Z_{n,r}^- - \mathbb{E}(Z_{n,r}^-)|^\ell \right) &\leq \ell \int_0^1 t^{\ell-1} dt + \exp(e^{-\lambda_{n,r}} - 1) \ell \int_0^{+\infty} t^{\ell-1} e^{-t} dt \\ &\leq 1 + \exp(e^{-\lambda_{n,r}} - 1) \ell!. \end{aligned}$$

vii) It is straightforward from (4.4) that

$$|\gamma_{n,r,1}| = |Z_{n,r}^+ - \lambda_{n,r} + \lambda_{n,r} - \mathbb{E}(Z_{n,r}^-)| \mathbb{I}_{\{N_{n,r}^+ > 0\}} \leq \left(\frac{nc}{k_n} \Delta_n + 1 \right) \mathbb{I}_{\{N_{n,r}^+ > 0\}}. \quad (4.11)$$

viii) Since $Z_{n,r}^-$ and $N_{n,r}^+$ are independent,

$$\max_{1 \leq r \leq k_n} \mathbb{E} \left(|\gamma_{n,r,2}|^\ell \right) \leq \max_{1 \leq r \leq k_n} \mathbb{E} \left(|Z_{n,r}^- - \mathbb{E}(Z_{n,r}^-)|^\ell \right) \max_{1 \leq r \leq k_n} \mathbb{P}(N_{n,r}^+ > 0)$$

and the result holds by (4.7).

ix) Since

$$\frac{Z_{n,r}^*}{N_{n,r}} \leq \frac{Z_{n,r}^-}{N_{n,r}^-} \mathbb{I}_{\{N_{n,r}^+ = 0\}} + \frac{nc}{k_n} \frac{M_{n,r}}{N_{n,r}^- + 1} \mathbb{I}_{\{N_{n,r}^+ \geq 1\}},$$

we have,

$$\mathbb{E} \left(\left(\frac{Z_{n,r}^*}{N_{n,r}} \right)^\ell \right) \leq \mathbb{E} \left(\left(\frac{Z_{n,r}^-}{N_{n,r}^-} \right)^\ell \right) + \mathbb{E} \left(\left(\frac{nc}{k_n} \frac{M_{n,r}}{N_{n,r}^- + 1} \right)^\ell \right) \mathbb{P}(N_{n,r}^+ \geq 1).$$

Moreover,

$$\begin{aligned} \mathbb{E} \left(\left(\frac{Z_{n,r}^-}{N_{n,r}^-} \right)^\ell \right) &= \mathbb{E} \left(\left(\frac{1}{N_{n,r}^-} \right)^\ell \mathbb{E} \left((Z_{n,r}^-)^\ell \mid N_{n,r}^- \right) \right) = \lambda_{n,r}^\ell \mathbb{E} \left(\frac{\mathbb{I}_{\{N_{n,r}^- \geq 1\}}}{(N_{n,r}^-)^{\ell-1} (\ell + N_{n,r}^-)} \right) \\ &= \ell! \sum_{q=1}^{\infty} \frac{(q + \ell - 1)!}{\ell! q^{\ell-1} q!} \frac{\lambda_{n,r}^{q+\ell}}{(q + \ell)!} e^{-\lambda_{n,r}} \\ &\leq \ell!, \end{aligned} \quad (4.12)$$

since for all ℓ and q ,

$$\frac{(q + \ell - 1)!}{\ell! q^{\ell-1} q!} \leq 1.$$

Besides,

$$\begin{aligned} \mathbb{E} \left(\left(\frac{nc}{k_n} \frac{M_{n,r}}{N_{n,r}^- + 1} \right)^\ell \right) &= \left(\frac{M_{n,r}}{m_{n,r}} \right)^\ell \ell! \sum_{q=0}^{\infty} \frac{(q + \ell)!}{\ell! (q + 1)^\ell q!} \frac{\lambda_{n,r}^{q+\ell}}{(q + \ell)!} e^{-\lambda_{n,r}} \leq \left(\frac{M_{n,r}}{m_{n,r}} \right)^\ell \ell! \\ &\leq \left(1 + \frac{\Delta_n}{m} \right)^\ell \ell!, \end{aligned} \quad (4.13)$$

since for all ℓ and q ,

$$\frac{(q + \ell)!}{\ell! (q + 1)^\ell q!} \leq 1.$$

Hence,

$$\max_{1 \leq r \leq k_n} \mathbb{E} \left(|\gamma_{n,r,3}|^\ell \right) \leq \ell! \left(1 + \left(1 + \frac{\Delta_n}{m} \right)^\ell \max_{1 \leq r \leq k_n} \mathbb{P} (N_{n,r}^+ > 0) \right).$$

■

In the next lemma we give an uniform upper bound on the centered moments of $(\xi_{n,r})$.

Lemma 2 *Under assumptions (H.1) and (H.2) we have*

$$\limsup_{n \rightarrow \infty} \max_{\ell \geq 2} \frac{1}{(12)^\ell \ell!} \max_{1 \leq r \leq k_n} \mathbb{E} \left(|\xi_{n,r} - \mathbb{E}(\xi_{n,r})|^\ell \right) < 1.$$

Proof : By Lemma 7 (see the appendix), we get that for all $1 \leq r \leq k_n$,

$$\max_{1 \leq r \leq k_n} \mathbb{E} \left(|\xi_{n,r} - \mathbb{E}(\xi_{n,r})|^\ell \right) \leq 2^\ell \max \left(\max_{1 \leq r \leq k_n} \mathbb{E} \left(|Z_{n,r}^- - \mathbb{E}(Z_{n,r}^-)|^\ell \right), 2^\ell \max_{1 \leq r \leq k_n} \mathbb{E} \left(|\gamma_{n,r}|^\ell \right) \right). \quad (4.14)$$

From Lemma 1 vi), we have

$$\max_{1 \leq r \leq k_n} \mathbb{E} \left(|Z_{n,r}^- - \mathbb{E}(Z_{n,r}^-)|^\ell \right) < 2\ell!. \quad (4.15)$$

Moreover, Lemma 7 and Lemma 1 vii)–ix) entail, for all large n , and all $\ell \geq 2$,

$$\max_{1 \leq r \leq k_n} \mathbb{E} \left(|\gamma_{n,r}|^\ell \right) \leq 3^\ell \max_{1 \leq j \leq 3} \max_{1 \leq r \leq k_n} \mathbb{E} \left(|\gamma_{n,r,j}|^\ell \right) \leq 6^\ell \ell!,$$

which, combined with (4.14) and (4.15) give the result. ■

The next lemma provides an exact uniform control of the variances of $(\xi_{n,r})_{1 \leq r \leq k_n}$.

Lemma 3 *Under assumptions (H.1) and (H.2) we have*

$$\max_{1 \leq r \leq k_n} |\mathbb{V}(\xi_{n,r}) - 1| = o(1).$$

Proof : Since

$$\mathbb{V}(\xi_{n,r}) = \mathbb{V}(Z_{n,r}^-) + 2\text{Cov}(Z_{n,r}^-, \gamma_{n,r}) + \mathbb{V}(\gamma_{n,r}),$$

we have,

$$\max_{1 \leq r \leq k_n} |\mathbb{V}(\xi_{n,r}) - 1| \leq \max_{1 \leq r \leq k_n} |\mathbb{V}(Z_{n,r}^-) - 1| + \max_{1 \leq r \leq k_n} \mathbb{V}(\gamma_{n,r}) + 2 \left[\max_{1 \leq r \leq k_n} \mathbb{V}(\gamma_{n,r}) \mathbb{V}(Z_{n,r}^-) \right]^{1/2}.$$

But, by Lemma 1 iv),

$$\max_{1 \leq r \leq k_n} |\mathbb{V}(Z_{n,r}^-) - 1| = \max_{1 \leq r \leq k_n} |2\lambda_{n,r}e^{-\lambda_{n,r}} + e^{-2\lambda_{n,r}}| = o(1).$$

Hence, it remains to show that

$$\max_{1 \leq r \leq k_n} \mathbb{V}(\gamma_{n,r}) = o(1),$$

and even, by Lemma 7, that

$$\max_{1 \leq j \leq 3} \max_{1 \leq r \leq k_n} \mathbb{V}(\gamma_{n,r,j}) = o(1).$$

Now, Lemma 1 vii) and viii) with $\ell = 2$ clearly imply that

$$\max_{1 \leq j \leq 2} \max_{1 \leq r \leq k_n} \mathbb{V}(\gamma_{n,r,j}) = o(1),$$

so it remains to prove that

$$\max_{1 \leq r \leq k_n} \mathbb{V}(\gamma_{n,r,3}) = o(1).$$

By (4.12) with $\ell = 2$,

$$\begin{aligned} \mathbb{E} \left(\left(\frac{Z_{n,r}^-}{N_{n,r}^-} \right)^2 \right) &= \sum_{q=1}^{\infty} \left(1 + \frac{1}{q} \right) \frac{\lambda_{n,r}^{q+2}}{(q+2)!} e^{-\lambda_{n,r}} \\ &= 1 - e^{-\lambda_{n,r}} \left(1 + \lambda_{n,r} + \frac{\lambda_{n,r}^2}{2} \right) + \sum_{q=1}^{\infty} \frac{1}{q} \frac{\lambda_{n,r}^{q+2}}{(q+2)!} e^{-\lambda_{n,r}}. \end{aligned}$$

Now, since,

$$\max_{1 \leq r \leq k_n} e^{-\lambda_{n,r}} \left(1 + \lambda_{n,r} + \frac{\lambda_{n,r}^2}{2} \right) \leq \max_{\lambda \geq mnc/k_n} e^{-\lambda} \left(1 + \lambda + \frac{\lambda^2}{2} \right) = o(1),$$

and for all $Q \geq 1$,

$$\begin{aligned} \max_{1 \leq r \leq k_n} \sum_{q=1}^{\infty} \frac{1}{q} \frac{\lambda_{n,r}^{q+2}}{(q+2)!} e^{-\lambda_{n,r}} &\leq \max_{1 \leq r \leq k_n} \sum_{q=1}^Q \frac{\lambda_{n,r}^{q+2}}{(q+2)!} e^{-\lambda_{n,r}} + \frac{1}{Q} \max_{1 \leq r \leq k_n} \sum_{q=1}^{\infty} \frac{\lambda_{n,r}^{q+2}}{(q+2)!} e^{-\lambda_{n,r}} \\ &\leq o(1) + \frac{1}{Q}, \end{aligned}$$

we get,

$$\max_{1 \leq r \leq k_n} \left| \mathbb{E} \left(\left(\frac{Z_{n,r}^-}{N_{n,r}^-} \right)^2 \right) - 1 \right| = o(1).$$

Furthermore, by Lemma 1 v),

$$\max_{1 \leq r \leq k_n} \left| \mathbb{E}^2 \left(\frac{Z_{n,r}^-}{N_{n,r}^-} \right) - 1 \right| = \max_{1 \leq r \leq k_n} \left[1 - (1 - e^{-\lambda_{n,r}} (1 + \lambda_{n,r}))^2 \right] = o(1).$$

Therefore,

$$\max_{1 \leq r \leq k_n} \mathbb{V}(\gamma_{n,r,3}) \leq \max_{1 \leq r \leq k_n} \left| \mathbb{E} \left(\left(\frac{Z_{n,r}^-}{N_{n,r}^-} \right)^2 \right) - 1 \right| + \max_{1 \leq r \leq k_n} \left| \mathbb{E}^2 \left(\frac{Z_{n,r}^-}{N_{n,r}^-} \right) - 1 \right| = o(1). \quad \blacksquare$$

Proposition 1 Under assumptions (H.1)–(H.4),

i) for all $(x_1, \dots, x_d) \subset [0, 1]$,

$$\left\{ \frac{nc}{\kappa_n(x_j)} \left(\hat{f}_n(x_j) - \mathbb{E} \left(\hat{f}_n(x_j) \right) \right) : 1 \leq j \leq d \right\} \xrightarrow{d} N(0, \Sigma_{(x_1, \dots, x_d)}),$$

ii) for all $(x_1, \dots, x_d) \subset [0, 1]$, and all $\varepsilon_n \downarrow 0$ such that $\Sigma_{(x_1, \dots, x_d)}$ is regular and

$$\max_{1 \leq r \leq k_n} \max_{1 \leq j \leq d} |w_{n,r}(x_j)| = o(\sqrt{\varepsilon_n}),$$

$$\left\{ \sqrt{\varepsilon_n} \frac{nc}{\kappa_n(x_j)} \left(\hat{f}_n(x_j) - \mathbb{E} \left(\hat{f}_n(x_j) \right) \right) : 1 \leq j \leq d \right\},$$

follows the large deviation principle in \mathbb{R}^d with speed (ε_n) and good rate function $I_{(x_1, \dots, x_d)}$.

Proof : For all $1 \leq r \leq k_n$, set

$$\zeta_{n,r} = \xi_{n,r} - \mathbb{E}(\xi_{n,r})$$

and

$$w_{n,r} = {}^t(w_{n,r}(x_1), \dots, w_{n,r}(x_d)).$$

Then, it is easily seen that

$$\left\{ \frac{nc}{\kappa_n(x_j)} \left(\hat{f}_n(x_j) - \mathbb{E} \left(\hat{f}_n(x_j) \right) \right) : 1 \leq j \leq d \right\} = \sum_{r=1}^{k_n} w_{n,r} \zeta_{n,r}.$$

Moreover, note that $(\zeta_{n,r})_{1 \leq r \leq k_n}$ are independent. Thus, by (H.3), (H.4), Lemma 3 and Lemma 2 we may apply Theorem 3 and Theorem 4 to get the intended results. \blacksquare

In order to control the bias term, we need information on the expectations of $(\xi_{n,r})_{1 \leq r \leq k_n}$.

Lemma 4 *Under assumptions (H.1) and (H.2) we have*

$$\max_{1 \leq r \leq k_n} |\mathbb{E}(\xi_{n,r}) - \lambda_{n,r}| = O\left(\frac{n}{k_n} \max\left(\Delta_n, \exp\left(-m \frac{nc}{k_n}\right)\right)\right).$$

Proof : Lemma 1 iii) yields

$$\mathbb{E}(\xi_{n,r}) - \lambda_{n,r} = e^{-\lambda_{n,r}} - 1 + \mathbb{E}(\gamma_{n,r,1}) + \mathbb{E}(\gamma_{n,r,3}), \quad (4.16)$$

with

$$\begin{aligned} \mathbb{E}(\gamma_{n,r,1}) &= \mathbb{E}\left((Z_{n,r}^+ - \lambda_{n,r})\mathbb{I}_{\{N_{n,r}^+ > 0\}}\right) + \mathbb{E}\left((\lambda_{n,r} - Z_{n,r}^-)\mathbb{I}_{\{N_{n,r}^+ > 0\}}\right) \\ &= \mathbb{E}\left((Z_{n,r}^+ - \lambda_{n,r})\mathbb{I}_{\{N_{n,r}^+ > 0\}}\right) + (1 - e^{-\lambda_{n,r}})\mathbb{P}(N_{n,r}^+ > 0), \end{aligned} \quad (4.17)$$

and, in view of Lemma 1 v),

$$\begin{aligned} \mathbb{E}(\gamma_{n,r,3}) &= \mathbb{E}\left(\frac{Z_{n,r}^-}{N_{n,r}^-}\right)\mathbb{P}(N_{n,r}^+ = 0) + \mathbb{E}\left(\frac{Z_{n,r}^+}{N_{n,r}^+}\right)\mathbb{P}(N_{n,r}^+ > 0) \\ &= (1 - e^{-\lambda_{n,r}}(1 + \lambda_{n,r}))\mathbb{P}(N_{n,r}^+ = 0) + \mathbb{E}\left(\frac{Z_{n,r}^+}{N_{n,r}^+}\right)\mathbb{P}(N_{n,r}^+ > 0). \end{aligned} \quad (4.18)$$

From (4.16)–(4.18), it follows that

$$\mathbb{E}(\xi_{n,r}) - \lambda_{n,r} = \mathbb{E}\left((Z_{n,r}^+ - \lambda_{n,r})\mathbb{I}_{\{N_{n,r}^+ > 0\}}\right) + \mathbb{E}\left(\frac{Z_{n,r}^+}{N_{n,r}^+}\right)\mathbb{P}(N_{n,r}^+ > 0) - \lambda_{n,r}e^{-\lambda_{n,r}}.$$

Now, by Lemma 1 i) and (4.13) with $\ell = 1$, we obtain

$$\begin{aligned} \max_{1 \leq r \leq k_n} |\mathbb{E}(\xi_{n,r}) - \lambda_{n,r}| &\leq \frac{nc}{k_n}\Delta_n + \left(1 + \frac{\Delta_n}{m}\right) \max_{1 \leq r \leq k_n} \mathbb{P}(N_{n,r}^+ > 0) + \frac{Mnc}{k_n} \exp\left(-m \frac{nc}{k_n}\right) \\ &= O\left(\frac{n}{k_n} \max\left(\Delta_n, \exp\left(-m \frac{nc}{k_n}\right)\right)\right), \end{aligned}$$

under (H.2). ■

Proposition 2 *Under assumptions (H.1), (H.2), (H.5)–(H.7), we have for all $x \in [0, 1]$,*

$$\frac{nc}{\kappa_n(x)} \left(\mathbb{E}\left(\hat{f}_n(x)\right) - f(x)\right) \rightarrow 0.$$

Proof : For all $x \in [0, 1]$, we get, by the triangle inequality and assumptions **(H.5)** and **(H.6)**,

$$\begin{aligned}
\frac{nc}{\kappa_n(x)} \left| \mathbb{E} \left(\hat{f}_n(x) \right) - f(x) \right| &= \left| \sum_{r=1}^{k_n} w_{n,r}(x) \mathbb{E}(\xi_{n,r}) - \frac{nc}{\kappa_n(x)} f(x) \right| \\
&\leq \left| \sum_{r=1}^{k_n} w_{n,r}(x) (\mathbb{E}(\xi_{n,r}) - \lambda_{n,r}) \right| \\
&\quad + \left| \frac{nc}{k_n} \sum_{r=1}^{k_n} w_{n,r}(x) (m_{n,r} - f(x)) \right| \\
&\quad + \left| \frac{1}{k_n} \sum_{r=1}^{k_n} \kappa_{n,r}(x) - 1 \right| \left| \frac{ncf(x)}{\kappa_n(x)} \right| \\
&\leq \left(\sum_{r=1}^{k_n} |w_{n,r}(x)| \right) \max_{1 \leq r \leq k_n} |\mathbb{E}(\xi_{n,r}) - \lambda_{n,r}| + o(1). \quad (4.19)
\end{aligned}$$

Lemma 4 and condition **(H.7)** give the result. \blacksquare

Theorem 1 and Theorem 2 are straightforward consequences of Proposition 1 and Proposition 2. The two following lemmas are dedicated to the replacement of c by \hat{c}_n in both theorems.

Lemma 5 *Under assumptions **(H.1)** and **(H.2)**, for all $\delta > 0$, there exist $\alpha_\delta > 0$ and $n_\delta > 0$ such that*

$$\forall n \geq n_\delta, \mathbb{P}(|\hat{c}_n - c| \geq \delta) \leq 3 \exp(-n\alpha_\delta).$$

Proof : We have

$$|\hat{c}_n - c| \leq \frac{1}{n\hat{a}_n} |N_n - nac| + ac \left| \frac{1}{\hat{a}_n} - \frac{1}{a} \right|.$$

Let $\delta > 0$ and $\eta_\delta = \min(a/2, a\delta/(4c))$. Then,

$$|\hat{c}_n - c| \leq |\hat{c}_n - c| \mathbb{I}_{\{|\hat{a}_n - a| > \eta_\delta\}} + \left(\frac{2}{na} |N_n - nac| + \frac{2c\eta_\delta}{a} \right) \mathbb{I}_{\{|\hat{a}_n - a| \leq \eta_\delta\}},$$

and therefore

$$\begin{aligned}
\mathbb{P}(|\hat{c}_n - c| \geq \delta) &\leq \mathbb{P} \left(\frac{1}{n} |N_n - nac| \geq \frac{a}{2} \left(\delta - \frac{2c\eta_\delta}{a} \right) \right) + \mathbb{P}(|\hat{a}_n - a| > \eta_\delta) \\
&\leq \mathbb{P} \left(\frac{N_n}{na} \notin \left[c - \frac{\delta}{4}, c + \frac{\delta}{4} \right] \right) + \mathbb{P}(|\hat{a}_n - a| > \eta_\delta). \quad (4.20)
\end{aligned}$$

Let us consider the first term of (4.20). Since N_n has a Poisson distribution, $N_n \rightsquigarrow \mathcal{P}(nac)$, it can be expanded as $N_n = \sum_{k=1}^n \pi_k$, where the random variables π_k are i.i.d. $\mathcal{P}(ac)$. Introducing

$$\Lambda_\pi(s) = \log \mathbb{E}(e^{s\pi}) = ac(e^s - 1)$$

and denoting

$$\Lambda_\pi^*(t) = \sup_{t \in \mathbb{R}} (st - \Lambda_\pi(s)) = \begin{cases} t \log(t/ac) - t + ac & \text{if } t > 0 \\ +\infty & \text{if } t \leq 0, \end{cases}$$

Cramer's theorem (see DEMBO & ZEITOUNI [4], Theorem 2.2.3) yields

$$\limsup_{n \rightarrow \infty} \frac{1}{n} \log \mathbb{P} \left(\frac{N_n}{na} \notin \left[c - \frac{\delta}{4}, c + \frac{\delta}{4} \right] \right) \leq -\inf \left\{ \Lambda_\pi^*(t), t \notin \left[c - \frac{\delta}{4}, c + \frac{\delta}{4} \right] \right\} < 0.$$

Consequently, there exists $\alpha'_\delta > 0$ such that

$$\forall n \geq 1, \mathbb{P} \left(\frac{N_n}{na} \notin \left[c - \frac{\delta}{4}, c + \frac{\delta}{4} \right] \right) \leq \exp(-n\alpha'_\delta). \quad (4.21)$$

Consider now the second term of (4.20). Lemma 4 yields

$$|\mathbb{E}(\hat{a}_n) - a| = O \left(\max \left(\Delta_n, \exp \left(-m \frac{nc}{k_n} \right) \right) \right)$$

which converges to 0 under **(H.1)** and **(H.2)**. Therefore, there exists $n_\delta > 0$ such that

$$\forall n \geq n_\delta, \mathbb{P}(|\hat{a}_n - a| \geq \eta_\delta) \leq \mathbb{P}(|\hat{a}_n - \mathbb{E}(\hat{a}_n)| \geq \eta_\delta/2) = \mathbb{P} \left(\left| \sum_{r=1}^{k_n} (\xi_{n,r} - \mathbb{E}(\xi_{n,r})) \right| \geq \eta_\delta nc/2 \right),$$

and in view of Lemma 2, applying Bernstein's inequality (see BOSQ [3], Theorem 2.6) yields

$$\forall n \geq n_\delta, \mathbb{P}(|\hat{a}_n - a| \geq \eta_\delta) \leq 2 \exp \left(-\frac{\eta_\delta^2 c^2 n^2}{2304 k_n + 48 \eta_\delta c n} \right) \leq 2 \exp \left(-\frac{\eta_\delta^2 c^2}{2304 + 48 \eta_\delta c} n \right). \quad (4.22)$$

Defining $\alpha_\delta = \min(\alpha'_\delta, \eta_\delta^2 c^2 / (2304 + 48 \eta_\delta c))$ and collecting (4.20)–(4.22) give the result. ■

The last lemma proves that the difference

$$D_n(x) = \frac{n}{\kappa_n(x)} (\hat{c}_n - c) (\hat{f}_n(x) - f(x))$$

can be neglected both in the central limit theorem and in the moderate deviation principle.

Lemma 6 *Let $(x_1, \dots, x_d) \subset [0, 1]$.*

i) Under assumptions of Theorem 1,

$$\{D_n(x_j), 1 \leq j \leq d\} \xrightarrow{\mathbb{P}} 0.$$

ii) Under assumptions of Theorem 2, for all $\eta > 0$,

$$\limsup_{n \rightarrow \infty} \varepsilon_n \log \mathbb{P} \left(\sqrt{\varepsilon_n} \max_{1 \leq j \leq d} |D_n(x_j)| \geq \eta \right) = -\infty.$$

Proof : i) For all $\eta > 0$ and $\delta > 0$, we have

$$\mathbb{P} \left(\max_{1 \leq j \leq d} |D_n(x_j)| \geq \eta \right) \leq \mathbb{P}(|\hat{c}_n - c| \geq \delta) + \mathbb{P} \left(\max_{1 \leq j \leq d} \frac{nc}{\kappa_n(x_j)} |\hat{f}_n(x_j) - f(x_j)| \geq \eta c / \delta \right),$$

where the first term converges to 0 as $n \rightarrow \infty$ in view of Lemma 5. Thus,

$$\begin{aligned} \limsup_{n \rightarrow \infty} \mathbb{P} \left(\max_{1 \leq j \leq d} |D_n(x_j)| \geq \eta \right) &\leq \limsup_{n \rightarrow \infty} \mathbb{P} \left(\max_{1 \leq j \leq d} \frac{nc}{\kappa_n(x_j)} |\hat{f}_n(x_j) - f(x_j)| \geq \eta c / \delta \right) \\ &= \mathbb{P} \left(\max_{1 \leq j \leq d} |G_j| \geq \eta c / \delta \right), \end{aligned} \quad (4.23)$$

where (G_1, \dots, G_d) follows the distribution $N(0, \Sigma_{(x_1, \dots, x_d)})$ (see Theorem 1). Letting $\delta \rightarrow 0$ in (4.23) concludes the proof:

$$\limsup_{n \rightarrow \infty} \mathbb{P} \left(\max_{1 \leq j \leq d} |D_n(x_j)| \geq \eta \right) = 0.$$

ii) For all $\eta > 0$ and $\delta > 0$, we have the well-known inequality (see DEMBO & ZEITOUNI [4], Lemma 1.2.15)

$$\begin{aligned} &\limsup_{n \rightarrow \infty} \varepsilon_n \log \mathbb{P} \left(\sqrt{\varepsilon_n} \max_{1 \leq j \leq d} |D_n(x_j)| \geq \eta \right) \leq \\ &\max \left\{ \limsup_{n \rightarrow \infty} \varepsilon_n \log \mathbb{P} \left(\sqrt{\varepsilon_n} \max_{1 \leq j \leq d} \frac{nc}{\kappa_n(x_j)} |\hat{f}_n(x_j) - f(x_j)| \geq \eta c / \delta \right), \limsup_{n \rightarrow \infty} \varepsilon_n \log \mathbb{P}(|\hat{c}_n - c| \geq \delta) \right\}. \end{aligned}$$

First, from Theorem 2,

$$\limsup_{n \rightarrow \infty} \varepsilon_n \log \mathbb{P} \left(\sqrt{\varepsilon_n} \max_{1 \leq j \leq d} \frac{nc}{\kappa_n(x_j)} |\hat{f}_n(x_j) - f(x_j)| \geq \eta c / \delta \right) \leq - \inf_{\|u\|_{\mathbb{R}^d} \geq \eta c / \delta} I_{(x_1, \dots, x_d)}(u) \rightarrow -\infty$$

when $\delta \rightarrow 0$. Second, Lemma 5 yields for $n \geq n_\delta$,

$$\varepsilon_n \log \mathbb{P}(|\hat{c}_n - c| \geq \delta) \leq -\alpha_\delta n \varepsilon_n + \varepsilon_n \log(3) \rightarrow -\infty$$

as $n \rightarrow \infty$, since for n large enough,

$$n \varepsilon_n \geq n \max_{1 \leq r \leq k_n} w_{n,r}^2(x) \geq n/k_n \rightarrow \infty.$$

As a conclusion,

$$\limsup_{n \rightarrow \infty} \varepsilon_n \log \mathbb{P} \left(\sqrt{\varepsilon_n} \max_{1 \leq j \leq d} |D_n(x_j)| \geq \eta \right) = -\infty.$$

■

5 Applications

We provide four different illustrations of the general results established in the previous section. We examine a kernel estimator, a particular case of wavelet estimator (the Haar estimate), an orthogonal series method (the trigonometric series method) as well as a non-orthogonal series method (the Faber-Schauder series method). First, the definition and the basic properties of each estimate are recalled. Second, we rewrite the sufficient conditions (H.1)–(H.7) in a more explicit way in these particular cases. Finally, we propose an illustration of the behavior of each estimate on simulated data.

5.1 Four boundary estimates

In the sequel (b_n) is a sequence of positive real numbers tending to infinity and $x_{n,r}$ is the center of the interval $I_{n,r}$, $r = 1, \dots, k_n$. The unknown function f is supposed to be twice continuously differentiable.

5.1.1 Kernel estimate

The kernel estimate is based on a nonparametric regression on the set of the bias corrected extreme values $(1 + N_{n,r}^{-1})Y_{n,r}^*$, $r = 1, \dots, k_n$. For $x \in [0, 1]$, it writes

$$\hat{f}_n(x) = \frac{1}{h_n k_n} \sum_{r=1}^{k_n} K\left(\frac{x - x_{n,r}}{h_n}\right) \left(1 + \frac{1}{N_{n,r}}\right) Y_{n,r}^*, \quad (5.1)$$

where K is a Parzen-Rosenblatt kernel, supposed to be bounded, positive, even, twice continuously differentiable and such that $x \rightarrow x^2 K(x)$ is integrable. (h_n) is a sequence of positive real numbers tending to zero called the window. It tunes the smoothing introduced by the kernel. For a review on nonparametric regression, see HÄRDLE [19]. It is apparent that (5.1) is a particular case of (2.1) in which $\kappa_{n,r}(x) = b_n K(b_n(x - x_{n,r}))$, with $b_n = 1/h_n$. The estimate \hat{f}_n is at least twice differentiable since it inherits from the regularity of the kernel K .

5.1.2 Haar estimate

Introduce a dyadic subdivision $\{J_\ell, \ell = b_n + 1, \dots, 2b_n + 1\}$ of $[0, 1]$ defined by $J_\ell = [c_\ell, d_\ell]$, where $c_\ell = p_\ell/2^{q_\ell-1}$, $d_\ell = (p_\ell + 1)/2^{q_\ell-1}$, p_ℓ and q_ℓ are the integers uniquely determined by $\ell = 2^{q_\ell-1} + p_\ell$ and $0 \leq p_\ell < 2^{q_\ell-1}$. The Haar basis (HAAR [15]) is the orthogonal basis defined by:

$$e_0 = \mathbb{I}_{[0,1]}, \quad e_\ell = 2^{\frac{q_\ell-1}{2}} (\mathbb{I}_{J_{2\ell}} - \mathbb{I}_{J_{2\ell+1}}), \quad \ell \geq 1.$$

We note $S_n(f)$ the expansion of f on the Haar basis truncated to the $(b_n + 1)$ first terms

$$S_n(f)(x) = \sum_{\ell=0}^{b_n} a_\ell e_\ell(x), \quad x \in [0, 1].$$

Each coefficient a_ℓ , $\ell = 0, \dots, b_n$ is estimated by the random Riemann sum

$$\hat{a}_{\ell, k_n} = \sum_{r=1}^{k_n} e_\ell(x_{n,r}) \left(1 + \frac{1}{N_{n,r}}\right) \frac{Y_{n,r}^*}{k_n},$$

leading to the estimator of f :

$$\hat{f}_n(x) = \sum_{\ell=0}^{b_n} \hat{a}_{\ell, k_n} e_\ell(x) = \frac{1}{k_n} \sum_{r=1}^{k_n} K_n^H(x_{n,r}, x) \left(1 + \frac{1}{N_{n,r}}\right) Y_{n,r}^*, \quad (5.2)$$

where K_n^H is the Dirichlet's kernel associated to the Haar basis:

$$K_n^H(x, y) = \sum_{\ell=0}^{b_n} e_\ell(x) e_\ell(y), \quad (x, y) \in [0, 1]^2.$$

It appears that (5.2) is a particular case of (2.1) with $\kappa_{n,r}(x) = K_n^H(x_{n,r}, x)$. Let us note that, since the functions of the Haar basis are not continuous, the estimator of f is also not continuous in general.

5.1.3 Trigonometric estimate

Formally, the trigonometric estimate writes as the Haar estimate:

$$\hat{f}_n(x) = \frac{1}{k_n} \sum_{r=1}^{k_n} K_n^T(x_{n,r}, x) \left(1 + \frac{1}{N_{n,r}}\right) Y_{n,r}^*, \quad (5.3)$$

except that K_n^T is the Dirichlet's kernel associated to the trigonometric basis which is defined by

$$e_0(x) = 1, \quad e_{2k-1}(x) = \sqrt{2} \cos(2k\pi x), \quad e_{2k}(x) = \sqrt{2} \sin(2k\pi x), \quad k \geq 1.$$

As previously, we have $\kappa_{n,r}(x) = K_n^T(x_{n,r}, x)$. In this case, the estimator \hat{f}_n is C^∞ .

5.1.4 Faber-Schauder estimate

Consider the dyadic subdivision introduced in Subsection 5.1.2. The Faber-Schauder is the non-orthogonal basis of continuous functions defined by

$$e_{-1}(x) = \mathbb{I}_{[0,1]}(x), \quad e_0(x) = x \mathbb{I}_{[0,1]}(x), \quad e_\ell(x) = 2^{q_\ell} ((x - c_\ell) \mathbb{I}_{J_{2^\ell}} + (d_\ell - x) \mathbb{I}_{J_{2^\ell+1}}), \quad \ell \geq 1.$$

Note $S_n(f)$ the expansion of f on the Faber-Schauder basis truncated to the $(b_n + 2)$ first terms

$$S_n(f)(x) = \sum_{\ell=-1}^{b_n} a_\ell e_\ell(x), \quad x \in [0, 1].$$

Each coefficient a_ℓ , $\ell = -1, \dots, k_n$ is estimated by

$$\hat{a}_{\ell, k_n} = (b_n + 1) \sum_{r=1}^{k_n} g_\ell(x_{n,r}) \left(1 + \frac{1}{N_{n,r}}\right) \frac{Y_{n,r}^*}{k_n},$$

where g_ℓ is a piecewise linear function on $[0, 1]$ that we do not precise here. Finally, the estimator can be written as:

$$\hat{f}_n(x) = \frac{1}{k_n} \sum_{r=1}^{k_n} K_n^F(x_{n,r}, x) \left(1 + \frac{1}{N_{n,r}}\right) Y_{n,r}^*, \quad (5.4)$$

where K_n^F is the kernel defined by:

$$K_n^F(x, y) = (b_n + 1) \sum_{\ell=-1}^{b_n} g_\ell(x) e_\ell(y), \quad (x, y) \in [0, 1]^2.$$

It appears that (5.4) is a particular case of (2.1) with $\kappa_{n,r}(x) = K_n^F(x_{n,r}, x)$. \hat{f}_n is a continuous estimate of f .

5.2 Basic properties

In this paragraph, we provide bounds on the three following quantities:

$$\max_{1 \leq r \leq k_n} |\kappa_{n,r}(x)|, \quad \frac{1}{k_n} \sum_{r=1}^{k_n} \kappa_{n,r}(x), \quad \kappa_n^2(x) = \sum_{r=1}^{k_n} \kappa_{n,r}^2(x), \quad x \in [0, 1]$$

that will reveal useful for rewriting the conditions **(H.1)**–**(H.7)**. The results which are collected here are classical ones so we omit their proofs.

5.2.1 Kernel estimate

It is clear that,

$$\max_{1 \leq r \leq k_n} |\kappa_{n,r}(x)| = O(b_n), \quad \forall x \in [0, 1].$$

Besides, under the condition $b_n^{3/2} = o(k_n)$, Proposition 1 in GIRARD & JACOB [14] yields

$$\left| \frac{1}{k_n} \sum_{r=1}^{k_n} \kappa_{n,r}(x) - 1 \right| = O(1/b_n^2) + O(b_n^3/k_n^2), \quad \forall x \in]0, 1[,$$

and

$$\kappa_n^2(x) \sim k_n b_n \int_0^1 K^2(t) dt, \quad \forall x \in]0, 1[. \quad (5.5)$$

5.2.2 Haar estimate

We introduce the following assumptions on the sequences (b_n) and (k_n) :

$$b_n + 1 = 2^{b'_n}, \quad b'_n \in \mathbb{N}, \quad k_n = d_n(b_n + 1), \quad d_n \in \mathbb{N}^*. \quad (5.6)$$

The first part of (5.6) is classical. It imposes that the number of terms in the expansion $S_n(x)$ is a power of two. The consequence of the second part of the condition is that for each $\ell \in \{b_n + 1, \dots, 2b_n + 1\}$, J_ℓ is exactly the union of d_n subintervals $I_{n,r}$. Now, let $x \in [0, 1]$ and $\ell(x)$ such that $x \in J_{\ell(x)}$. Under (5.6), the Dirichlet's kernel reduces to

$$K_n^H(x, x_{n,r}) = (b_n + 1)\mathbb{I}_{\{x_n, r \in J_{\ell(x)}\}}. \quad (5.7)$$

We thus obtain, for all $x \in [0, 1]$:

$$\begin{aligned} \max_{1 \leq r \leq k_n} |\kappa_{n,r}(x)| &= 1 + b_n, \\ \frac{1}{k_n} \sum_{r=1}^{k_n} \kappa_{n,r}(x) &= 1, \\ \kappa_n^2(x) &= k_n(b_n + 1). \end{aligned} \quad (5.8)$$

5.2.3 Trigonometric estimate

Suppose for convenience that b_n is even. Then, the Dirichlet's kernel can be explicitly computed:

$$K_n^T(x, y) = \begin{cases} \frac{\sin[(1 + b_n)\pi(x - y)]}{\sin[\pi(x - y)]} & x \neq y, \\ 1 + b_n. & x = y, \end{cases}$$

and it follows that, for all $x \in [0, 1]$,

$$\begin{aligned} \max_{1 \leq r \leq k_n} |\kappa_{n,r}(x)| &= 1 + b_n, \\ \left| \frac{1}{k_n} \sum_{r=1}^{k_n} \kappa_{n,r}(x) - 1 \right| &= O(b_n \ln b_n / k_n). \end{aligned}$$

Besides, under the condition $b_n \ln b_n = o(k_n)$, we have

$$\kappa_n^2(x) \sim k_n(b_n + 1), \quad x \in [0, 1].$$

For a proof, see for instance JACOB & SUQUET [23], equation (4.14).

5.2.4 Faber-Schauder estimate

Similarly to the Haar case, we suppose that the sequences (h_n) and (k_n) verify (5.6) where d_n is even. The kernel K_n^F can be explicitly computed (see Proposition 1 in GARDES [8]) although its expression is too heavy to be given here. The following results

$$\max_{1 \leq r \leq k_n} |\kappa_{n,r}(x)| = O(b_n),$$

$$\frac{1}{k_n} \sum_{r=1}^{k_n} \kappa_{n,r}(x) = 1,$$

$$\frac{1}{2} \leq \frac{\kappa_n^2(x)}{k_n(b_n + 1)} \leq 2,$$

($x \in [0, 1]$) can be found in GARDES [8], Lemma 2.

5.3 Convergence results

We first study the convergence of the centered estimates. Proposition 1 can be rewritten as:

Corollary 2 *Suppose f is C^2 . Under assumptions of Table 1, the Haar, Faber-Schauder, trigonometric and kernel estimates verify:*

i) for all $(x_1, \dots, x_d) \subset [0, 1]$,

$$\left\{ \frac{nc}{\kappa_n(x_j)} (\hat{f}_n(x_j) - E(\hat{f}_n(x_j))) : 1 \leq j \leq d \right\} \xrightarrow{d} N(0, I),$$

with $\kappa_n(x_j) \asymp (b_n k_n)^{1/2}$, $j = 1, \dots, d$.

ii) for all $(x_1, \dots, x_d) \subset [0, 1]$, and all $\varepsilon_n \downarrow 0$ such that $b_n/k_n = o(\varepsilon_n)$,

$$\left\{ \sqrt{\varepsilon_n} \frac{nc}{\kappa_n(x_j)} \left(\hat{f}_n(x_j) - E(\hat{f}_n(x_j)) \right) : 1 \leq j \leq d \right\}.$$

follows the large deviation principle in \mathbb{R}^d with speed (ε_n) and good rate function $I(u) = \|u\|_{\mathbb{R}^d}^2/2$.

The notation $\kappa_n(x_j) \asymp (b_n k_n)^{1/2}$ means that

$$0 < \liminf \kappa_n(x_j)/(b_n k_n)^{1/2} \leq \limsup \kappa_n(x_j)/(b_n k_n)^{1/2} < \infty.$$

In the case of the Haar, kernel and trigonometric estimate, this reduces to $\kappa_n(x_j) \sim (b_n k_n)^{1/2}$. The first line of assumptions of Table 1 ensures that the mean number of points in each cell $D_{n,r}$ converges to infinity. The second line of assumptions determines the minimal amount of smoothing required on each interval $I_{n,r}$ for asymptotic normality. Finally,

the third line of assumptions imposes that the mean number of points above $m_{n,r}$ in each cell $D_{n,r}$ converges to 0, so that the function f could be approximated by a constant on $I_{n,r}$.

Proof : Assumptions **(H.1)**, **(H.2)** and **(H.4)** are easily verified thanks to the results of the previous paragraph. Let us focus on assumption **(H.3)** and note

$$\sigma_{i,j}^{(n)} = \sum_{r=1}^{k_n} w_{n,r}(x_i)w_{n,r}(x_j), \quad 1 \leq i, j \leq d.$$

It is clear that one always have $\sigma_{i,i}^{(n)} = 1$, $1 \leq i \leq d$. The computation of $\sigma_{i,j} = \lim_{n \rightarrow \infty} \sigma_{i,j}^{(n)}$, $i \neq j$ is done separately for each estimate.

- Haar estimate: Let $\ell(x_i)$ and $\ell(x_j)$ such that $x_i \in J_{\ell(x_i)}$ and $x_j \in J_{\ell(x_j)}$. Then, in view of (5.7),

$$K_n^H(x_i, x_{n,r})K_n^H(x_j, x_{n,r}) = (b_n + 1)^2 \mathbb{1}_{\{x_{n,r} \in J_{\ell(x_i)}\}} \mathbb{1}_{\{x_{n,r} \in J_{\ell(x_j)}\}}.$$

Now, for n large enough, $J_{\ell(x_i)} \cap J_{\ell(x_j)} = \emptyset$ and then $\sigma_{i,j}^{(n)} = 0$. Thus $\sigma_{i,j} = 1$ if $i = j$, and $\sigma_{i,j} = 0$ otherwise. The case of the Faber-Schauder estimate is similar.

- Trigonometric estimate: Equation (4.35) of JACOB & SUQUET [23] entails $\sigma_{i,j} = 0$ for $i \neq j$ when $b_n \ln b_n = o(k_n)$.
- Kernel estimate: GIRARD & JACOB [14], Corollary 1 shows that $\sigma_{i,j} = 0$ for $i \neq j$ when $b_n^{3/2} = o(k_n)$. ■

The convergence of the estimates centered on the true function f is now derived from Theorem 1 and Theorem 2 at the expense on additional conditions on (b_n) and (k_n) .

Corollary 3 *Suppose f is C^2 . Under assumptions of Table 2, the Haar, Faber-Schauder, trigonometric and kernel estimates verify:*

i) for all $(x_1, \dots, x_d) \subset [0, 1]$,

$$\left\{ \frac{nc}{\kappa_n(x_j)} (\hat{f}_n(x_j) - f(x_j)) : 1 \leq j \leq d \right\} \xrightarrow{d} N(0, I),$$

with $\kappa_n(x_j) \asymp (b_n k_n)^{1/2}$, $j = 1, \dots, d$.

ii) for all $(x_1, \dots, x_d) \subset [0, 1]$, and all $\varepsilon_n \downarrow 0$ such that $b_n/k_n = o(\varepsilon_n)$,

$$\left\{ \sqrt{\varepsilon_n} \frac{nc}{\kappa_n(x_j)} (\hat{f}_n(x_j) - f(x_j)) : 1 \leq j \leq d \right\}.$$

follows the large deviation principle in \mathbb{R}^d with speed (ε_n) and good rate function $I(u) = \|u\|_{\mathbb{R}^d}^2/2$.

Proof : Conditions **(H.5)** and **(H.7)** are straightforwardly verified thanks to the results of paragraph 5.2. Let us focus on condition **(H.6)** and consider the different estimates separately.

- Haar estimate: Let $\ell(x)$ such that $x \in J_{\ell(x)}$. In view of (5.7),

$$\begin{aligned} \frac{1}{k_n} \sum_{r=1}^{k_n} \kappa_{n,r}(x)(f(x) - m_{n,r}) &= \frac{1}{d_n} \sum_{r=1}^{k_n} (f(x) - m_{n,r}) \mathbb{I}_{\{x_{n,r} \in J_{\ell(x)}\}} \\ &\leq \max_r \sup_{y \in J_{\ell(x)}} |f(y) - m_{n,r}| \\ &= O(1/b_n), \end{aligned}$$

and (5.8) concludes the proof. The case of the Faber-Schauder basis is similar.

- Trigonometric estimate: Consider the expansion

$$\begin{aligned} \frac{1}{k_n} \sum_{r=1}^{k_n} \kappa_{n,r}(x)(f(x) - m_{n,r}) &= \left(f(x) - \sum_{r=1}^{k_n} \kappa_{n,r}(x) \lambda_{n,r} \right) \\ &+ f(x) \left(\frac{1}{k_n} \sum_{r=1}^{k_n} \kappa_{n,r}(x) - 1 \right) \\ &+ \frac{1}{k_n} \sum_{r=1}^{k_n} \kappa_{n,r}(x) (k_n \lambda_{n,r} - m_{n,r}). \end{aligned}$$

The first term is bounded by GIRARD & JACOB [13], Proposition 2:

$$\left| f(x) - \sum_{r=1}^{k_n} \kappa_{n,r}(x) \lambda_{n,r} \right| = O(b_n \ln b_n / k_n) + O(1/b_n).$$

The second term is controlled with **(H.5)**, and the third one by the Cauchy-Schwarz inequality

$$\frac{1}{k_n} \sum_{r=1}^{k_n} \kappa_{n,r}(x) (k_n \lambda_{n,r} - m_{n,r}) = O(\kappa_n(x) / k_n^{3/2}).$$

- Kernel estimate: In a similar way,

$$\begin{aligned} \frac{1}{k_n} \sum_{r=1}^{k_n} \kappa_{n,r}(x)(f(x) - m_{n,r}) &= \left(f(x) - \frac{1}{k_n} \sum_{r=1}^{k_n} \kappa_{n,r}(x) f(x_{n,r}) \right) \\ &+ f(x) \left(\frac{1}{k_n} \sum_{r=1}^{k_n} \kappa_{n,r}(x) - 1 \right) \\ &+ \frac{1}{k_n} \sum_{r=1}^{k_n} \kappa_{n,r}(x) (f(x_{n,r}) - m_{n,r}). \end{aligned}$$

The first term is bounded by GIRARD & JACOB [14], Proposition 1:

$$\left| f(x) - \sum_{r=1}^{k_n} \kappa_{n,r}(x) f(x_{n,r}) \right| = O(b_n^3/k_n^2) + O(1/b_n^2).$$

The second term is controlled with **(H.5)**, and the third one by

$$\frac{1}{k_n} \sum_{r=1}^{k_n} \kappa_{n,r}(x) (f(x_{n,r}) - m_{n,r}) = \left(\frac{1}{k_n} \sum_{r=1}^{k_n} \kappa_{n,r}(x) \right) O(1/k_n) = O(1/k_n),$$

and (5.5) concludes the proof. ■

We conclude this investigation of the asymptotic properties of the estimates by providing the optimal choices of the (b_n) and (k_n) sequences in Table 3 and the resulting convergence rate of the estimate. From the asymptotic point of view, the kernel estimate is the better one, and the trigonometric estimate is the worst. The Haar and Faber-Shauder estimates share the same intermediary convergence rate. However, the convergence rate should not be the only criterion for choosing the estimator. For instance, prior information on the regularity of f can be a good indication for the choice of the estimator. Haar and Faber-Shauder estimates are well-adapted for the estimation of non smooth boundaries whereas, the trigonometric and kernel estimates are designed for regular boundaries. Besides, the trigonometric estimate is a good candidate for the estimation of periodic functions f . This kind of problem arises when a star shaped support has been reduced to the kind of domain (1.1) studied here by the transformation from polar to cartesian coordinates (see JACOB & ABBAR [22]). Let us also recall that the kernel estimate suffers from a boundary effects leading to poor estimations of f at the boundaries of the interval. Nevertheless, simple and efficient corrections exist, see e.g. GIRARD & JACOB [12]. We now illustrate the behaviour of the estimates on simulations.

5.4 Numerical experiments

In this illustration, the sum of $n = 200$ independent Poisson processes with the same intensity rate $c = 4$ is simulated on the set S defined by (1.1) with

$$f(x) = [0.1 + \sin(\pi x)] [1.1 - 0.5 \exp(-64(x - 0.5)^2)],$$

for $x \in [0, 1]$. The unit interval is divided into $k_n = 32$ equidistant intervals. In the case of the kernel estimate, we choose $h_n = 0.025$ for the smoothing parameter (leading to $b_n = 40$). In the case of the Haar, Faber-Shauder and trigonometric estimates, we consider an expansion of order $b_n = 15$ of f in the corresponding basis. The results are presented in Figure 1 for comparison. In each case, the estimate \hat{f}_n is superimposed to the true function f and the simulated point process. We also take profit of Theorem 1 to draw 90% confidence intervals

for $f(x)$ at 50 different locations. The qualitative differences between the four estimates appear clearly. The Haar estimate is piecewise constant (Figure 1(a)) and the Faber-Shauder estimate is piecewise linear (Figure 1(b)). Also note on Figure 1(c) the oscillations of the trigonometric estimate due to the periodic nature of the associated Dirichlet's kernel. It is apparent that the first peak is under-estimated by all the estimates. The smoothing induced by the different kernels is not sufficient to overcome the missing of simulated points in the peak. Nevertheless, the unknown function f is always included in the confidence intervals.

Haar & Faber-Shauder	Trigonometric	Kernel
$k_n = o(n)$	$k_n = o(n)$	$k_n = o(n)$
$b_n = o(k_n)$	$b_n \ln b_n = o(k_n)$	$b_n^{3/2} = o(k_n)$
$n = o(k_n^2)$	$n = o(k_n^2)$	$n = o(k_n^2)$

Table 1: Sufficient conditions **(H.1)**–**(H.4)** for the different estimators.

Haar & Faber-Shauder	Trigonometric	Kernel
$k_n = o(n/\ln n)$	$k_n = o(n/\ln n)$	$k_n = o(n/\ln n)$
$b_n = o(k_n)$	$b_n \ln b_n = o(k_n)$	$b_n^{3/2} = o(k_n)$ $k_n = o(b_n^{5/2})$
$n = o(k_n^{1/2} b_n^{3/2})$	$n = o(k_n^{1/2} b_n^{3/2})$ $n = o(k_n^{3/2} / b_n^{1/2} \ln b_n)$	$n = o(k_n^{5/2} / b_n^{5/2})$

Table 2: Sufficient conditions (H.1)–(H.7) for the different estimators.

Estimate	Haar & Faber-Shauder	Trigonometric	Kernel
k_n	$n^{1/2} \rho_n^2$	$n^{4/5} (\ln n)^{2/3} \rho_n^2$	$n^{2/3} \rho_n^2$
b_n	$n^{1/2}$	$n^{2/5}$	$n^{4/15}$
κ_n/n	$n^{-1/2} \rho_n$	$n^{-2/5} (\ln n)^{1/3} \rho_n$	$n^{-8/15} \rho_n$

Table 3: Optimal choices of (b_n) and (k_n) sequences, and convergence rate of the estimators. (ρ_n) is an arbitrary positive sequence such that $\rho_n \rightarrow \infty$ and $\rho_n = o(\ln n)$.




Figure 1: Superimposition of the simulated point process, the function f to estimate (continuous line) and the four estimates (dashed lines) obtained with the Haar (a), Faber-Schauder (b), trigonometric (c) and kernel (d) methods. In each case, the vertical lines represent the 90% confidence intervals.

6 Appendix

6.1 Multivariate central limit theorem and moderate deviations

In this part, we provide general theorems about the central limit property and the moderate deviation principle of a sequence of random \mathbb{R}^d valued vectors

$$\theta_n = \sum_{r=1}^{k_n} w_{n,r} \zeta_{n,r}, \quad n \geq 1,$$

where $(w_{n,r})_{1 \leq r \leq k_n} \subset \mathbb{R}^d$ and $(\zeta_{n,r})_{1 \leq r \leq k_n}$ are random variables such that:

(A.1) $(\zeta_{n,r})_{1 \leq r \leq k_n}$ are centered and independent random variables.

(A.2) $\max_{1 \leq r \leq k_n} |\mathbb{E}(\zeta_{n,r}^2) - 1| \rightarrow 0$.

(A.3) There exists a covariance matrix Σ in \mathbb{R}^d such that for all $\lambda \in \mathbb{R}^d$,

$$\sum_{r=1}^{k_n} \langle w_{n,r}, \lambda \rangle_{\mathbb{R}^d}^2 \rightarrow {}^t \lambda \Sigma \lambda.$$

(A.4) $\max_{1 \leq r \leq k_n} \|w_{n,r}\|_{\mathbb{R}^d} = o(1)$.

(A.5) $\limsup_{\alpha \rightarrow \infty} \limsup_{n \rightarrow \infty} \max_{1 \leq r \leq k_n} \mathbb{E}(\zeta_{n,r}^2 \mathbb{I}_{\{|\zeta_{n,r}| > \alpha\}}) = 0$.

(A.6) There exists $K > 0$ such that

$$\limsup_{n \rightarrow \infty} \max_{\ell \geq 3} \max_{1 \leq r \leq k_n} \frac{1}{K^\ell \ell!} \mathbb{E}(|\zeta_{n,r}|^\ell) < 1.$$

Let us note that (A.6) implies (A.5).

Theorem 3 Under assumptions (A.1) – (A.5),

$$\theta_n \xrightarrow[d]{} N(0, \Sigma).$$

Proof : We have to show that, for all $\lambda \in \mathbb{R}^d$,

$$\langle \theta_n, \lambda \rangle_{\mathbb{R}^d} \xrightarrow[d]{} N(0, {}^t \lambda \Sigma \lambda). \quad (6.1)$$

Now, observe that (A.1)–(A.3) entail

$$\mathbb{V}(\langle \theta_n, \lambda \rangle_{\mathbb{R}^d}) = \sum_{r=1}^{k_n} \langle w_{n,r}, \lambda \rangle_{\mathbb{R}^d}^2 \mathbb{E}(\zeta_{n,r}^2) = {}^t \lambda \Sigma \lambda + o(1).$$

Hence, by Lindeberg Theorem (see e.g : DUDLEY [7], p. 248), it is easy to see that (6.1) holds whenever for all $\varepsilon > 0$,

$$\limsup_{n \rightarrow \infty} \sum_{r=1}^{k_n} \langle w_{n,r}, \lambda \rangle_{\mathbb{R}^d}^2 \mathbb{E} \left(\zeta_{n,r}^2 \mathbb{I}_{\{|\zeta_{n,r} \langle w_{n,r}, \lambda \rangle_{\mathbb{R}^d}| > \varepsilon\}} \right) = 0.$$

Fix $\lambda \in \mathbb{R}^d$, $\varepsilon > 0$ and $\alpha > 0$. Using (A.4), we get for all n large enough and all $1 \leq r \leq k_n$ that

$$\mathbb{I}_{\{|\zeta_{n,r} \langle w_{n,r}, \lambda \rangle_{\mathbb{R}^d}| > \varepsilon\}} \leq \mathbb{I}_{\{|\zeta_{n,r}| > \alpha\}}.$$

Hence,

$$\begin{aligned} & \limsup_{n \rightarrow \infty} \sum_{r=1}^{k_n} \langle w_{n,r}, \lambda \rangle_{\mathbb{R}^d}^2 \mathbb{E} \left(\zeta_{n,r}^2 \mathbb{I}_{\{|\zeta_{n,r} \langle w_{n,r}, \lambda \rangle_{\mathbb{R}^d}| > \varepsilon\}} \right) \\ & \leq \limsup_{n \rightarrow \infty} \left(\sum_{r=1}^{k_n} \langle w_{n,r}, \lambda \rangle_{\mathbb{R}^d}^2 \right) \max_{1 \leq r \leq k_n} \mathbb{E} \left(\zeta_{n,r}^2 \mathbb{I}_{\{|\zeta_{n,r}| > \alpha\}} \right) \\ & \leq^t \lambda \Sigma \lambda \limsup_{n \rightarrow \infty} \max_{1 \leq r \leq k_n} \mathbb{E} \left(\zeta_{n,r}^2 \mathbb{I}_{\{|\zeta_{n,r}| > \alpha\}} \right). \end{aligned}$$

and we get the result by (A.5) when $\alpha \uparrow \infty$. ■

Let $\varepsilon_n \downarrow 0$, $I : \mathbb{R}^d \rightarrow [0, +\infty]$ such that for all $\alpha > 0$, $\{I \leq \alpha\}$ is compact and recall that :

Definition 1 A sequence of \mathbb{R}^d valued random variable (Θ_n) is said to follow the large deviation principle in \mathbb{R}^d with speed (ε_n) and good rate function I whenever for every set $A \in \mathcal{B}(\mathbb{R}^d)$,

$$\begin{aligned} - \inf \left\{ I(u) : u \in \overset{\circ}{A} \right\} & \leq \liminf_{n \rightarrow \infty} \varepsilon_n \log \mathbb{P}(\Theta_n \in A) \\ & \leq \limsup_{n \rightarrow \infty} \varepsilon_n \log \mathbb{P}(\Theta_n \in A) \\ & \leq - \inf \{ I(u) : u \in \overline{A} \}, \end{aligned}$$

where $\overset{\circ}{A}$ (resp. \overline{A}) denotes the interior (resp. closure) of A in \mathbb{R}^d .

Assume that Σ is regular and define

$$I_\Sigma : u \in \mathbb{R}^d \mapsto \frac{1}{2} {}^t u \Sigma^{-1} u.$$

Theorem 4 Assume that (A.1) – (A.4) and (A.6) hold, where the matrix Σ in (A.3) is regular, then, for all sequences $\varepsilon_n \downarrow 0$ such that

$$\max_{1 \leq r \leq k_n} \|w_{n,r}\|_{\mathbb{R}^d} = o(\sqrt{\varepsilon_n}), \quad (6.2)$$

$(\sqrt{\varepsilon_n}\theta_n)$ follows the large deviation principle in \mathbb{R}^d with speed (ε_n) and good rate function I_Σ .

Proof : Denote by

$$\Lambda_n : \lambda \in \mathbb{R}^d \mapsto \varepsilon_n \log \mathbb{E} \left(\exp \left(\varepsilon_n^{-1} \langle \lambda, \sqrt{\varepsilon_n} \theta_n \rangle_{\mathbb{R}^d} \right) \right),$$

and set

$$\Lambda : \lambda \in \mathbb{R}^d \mapsto \frac{1}{2} {}^t \lambda \Sigma \lambda.$$

Then,

$$\Lambda_n(\lambda) = \varepsilon_n \sum_{r=1}^{k_n} \log (1 + \alpha_{n,r}(\lambda) (1 + \beta_{n,r}(\lambda))),$$

with

$$\alpha_{n,r}(\lambda) = \frac{\langle \lambda, w_{n,r} \rangle_{\mathbb{R}^d}^2}{2\varepsilon_n}$$

and

$$\beta_{n,r}(\lambda) = \mathbb{E}(\zeta_{n,r}^2) - 1 + 2\mathbb{E} \left(\sum_{\ell=3}^{+\infty} \left(\frac{\langle \lambda, w_{n,r} \rangle_{\mathbb{R}^d}}{\sqrt{\varepsilon_n}} \right)^{\ell-2} \frac{\zeta_{n,r}^\ell}{\ell!} \right).$$

Now, by (6.2),

$$\max_{1 \leq r \leq k_n} \alpha_{n,r}(\lambda) \rightarrow 0 \text{ as } n \uparrow \infty.$$

Moreover, by (A.2), (A.6) and (6.2), for all large n ,

$$\begin{aligned} \max_{1 \leq r \leq k_n} |\beta_{n,r}(\lambda)| &\leq \max_{1 \leq r \leq k_n} |\mathbb{E}(\zeta_{n,r}^2) - 1| + 2 \max_{1 \leq r \leq k_n} \sum_{\ell=3}^{+\infty} \left| \varepsilon_n^{-1/2} \langle \lambda, w_{n,r} \rangle_{\mathbb{R}^d} \right|^{\ell-2} \mathbb{E} \left(\left| \frac{\zeta_{n,r}^\ell}{\ell!} \right| \right) \\ &\leq o(1) + 2K^2 \sum_{\ell=1}^{+\infty} \left(K \|\lambda\|_{\mathbb{R}^d} \varepsilon_n^{-1/2} \max_{1 \leq r \leq k_n} \|w_{n,r}\|_{\mathbb{R}^d} \right)^\ell \\ &\leq o(1) + 4K^3 \|\lambda\|_{\mathbb{R}^d} \varepsilon_n^{-1/2} \max_{1 \leq r \leq k_n} \|w_{n,r}\|_{\mathbb{R}^d} = o(1). \end{aligned}$$

Therefore, by Lemma 8 below and (A.3),

$$\begin{aligned}\Lambda_n(\lambda) &= \varepsilon_n \sum_{r=1}^{k_n} \alpha_{n,r}(\lambda) + o\left(\varepsilon_n \sum_{r=1}^{k_n} \alpha_{n,r}(\lambda)\right) = \frac{1}{2} \sum_{r=1}^{k_n} \langle \lambda, w_{n,r} \rangle_{\mathbb{R}^d}^2 + o\left(\sum_{r=1}^{k_n} \langle \lambda, w_{n,r} \rangle_{\mathbb{R}^d}^2\right) \\ &= \Lambda(\lambda) + o(1).\end{aligned}$$

Hence, by the Gartner-Ellis Theorem (see Theorem 2-3-6 of DEMBO & ZEITOUNI [4], p.44), $(\sqrt{\varepsilon_n} \theta_n)$ follows the large deviation principle in \mathbb{R}^d with speed (ε_n) and good rate function

$$\Lambda^*(u) = \sup \{ \langle \lambda, u \rangle_{\mathbb{R}^d} - \Lambda(\lambda) : \lambda \in \mathbb{R}^d \}$$

and it is easily seen that $\Lambda^* = \mathbb{I}_\Sigma$. ■

6.2 Technical lemmas

The following facts have been useful in our proofs :

Lemma 7 For every $\ell \geq 1$ and every random variables χ_1, \dots, χ_p in $L^\ell(\mathbb{P})$,

$$\mathbb{E}\left(|\chi_1 + \dots + \chi_p|^\ell\right) \leq p^\ell \max_{1 \leq j \leq p} \mathbb{E}\left(|\chi_j|^\ell\right).$$

In particular,

$$\mathbb{E}\left(|\chi_1 - \mathbb{E}(\chi_1)|^\ell\right) \leq 2^\ell \mathbb{E}\left(|\chi_1|^\ell\right).$$

The proof is a straightforward consequence of the $L^\ell(\mathbb{P})$ norm convexity.

Lemma 8 Consider triangular arrays $(\alpha_{n,r})_{1 \leq r \leq k_n} \subset \mathbb{R}^+$ and $(\beta_{n,r})_{1 \leq r \leq k_n} \subset \mathbb{R}$ such that :

$$\alpha_n := \max_{1 \leq r \leq k_n} \alpha_{n,r} \rightarrow 0 \text{ as } n \uparrow \infty,$$

and

$$\beta_n := \max_{1 \leq r \leq k_n} |\beta_{n,r}| \rightarrow 0 \text{ as } n \uparrow \infty.$$

Then,

$$\sum_{r=1}^{k_n} \log(1 + \alpha_{n,r}(1 + \beta_{n,r})) = \sum_{r=1}^{k_n} \alpha_{n,r} + o\left(\sum_{r=1}^{k_n} \alpha_{n,r}\right).$$

Proof : Since for all $u \geq 0$,

$$0 \leq u - \log(1 + u) \leq u^2,$$

we get, for all large n ,

$$\begin{aligned} \left| \sum_{r=1}^{k_n} \log(1 + \alpha_{n,r}(1 + \beta_{n,r})) - \sum_{r=1}^{k_n} \alpha_{n,r} \right| &\leq \sum_{r=1}^{k_n} [\alpha_{n,r}(1 + \beta_{n,r})]^2 + \left| \sum_{r=1}^{k_n} \alpha_{n,r} \beta_{n,r} \right| \\ &\leq (2\alpha_n + \beta_n) \sum_{r=1}^{k_n} \alpha_{n,r}, \end{aligned}$$

which gives the result. ■

References

- [1] Abbar, H. (1990) Un estimateur spline du contour d'une répartition ponctuelle aléatoire. *Statistique et analyse des données*, **15**(3), 1–19.
- [2] Baufays, P. and Rasson, J.P. (1985) A new geometric discriminant rule. *Computational Statistics Quarterly*, **2**, 15–30.
- [3] Bosq, D. (2000) Linear processes in function spaces. Theory and applications. in *Lecture Notes in Statistics*, **149**, Springer-Verlag, New York.
- [4] Dembo, A. and Zeitouni, O. (1991) *Large Deviations Techniques and Applications*, Jones and Bartlett.
- [5] Deprins, D., Simar, L. and Tulkens, H. (1984) Measuring Labor Efficiency in Post Offices. in *The Performance of Public Enterprises: Concepts and Measurements* by M. Marchand, P. Pestieau and H. Tulkens, North Holland ed, Amsterdam.
- [6] Devroye, L.P. and Wise, G.L. (1980) Detection of abnormal behavior via non parametric estimation of the support. *SIAM J. Applied Math.*, **38**, 448–480.
- [7] Dudley, R.M. (1989) *Real Analysis and Probability*, Wadsworth and Brooks / Cole.
- [8] Gardes, L. (2001) Estimating the support of a poisson process via the Faber-Schauder basis and extreme values. *Technical report ENSAM-INRA-UM2*, **01-04**.
- [9] Geffroy, J. (1964) Sur un problème d'estimation géométrique. *Publications de l'Institut de Statistique de l'Université de Paris*, **XIII**, 191–200.
- [10] Gijbels, I. and Peng, L. (1999). Estimation of a support curve via order statistics. *Discussion Paper 9905*, Institut de Statistique, Université Catholique de Louvain.
- [11] Gijbels, I., Mammen, E., Park, B.U. and Simar, L. (1999). On estimation of monotone and concave frontier functions. *Journal of the American Statistical Association*, **94**, 220–228.
- [12] Girard, S. and Jacob, P. (1999) Extreme values and Haar series estimates of point processes boundaries. *Technical report ENSAM-INRA-UM2*, **99-08**.
- [13] Girard, S. and Jacob, P. (2000) Extreme values and projection estimates of point processes boundaries. *Technical report ENSAM-INRA-UM2*, **00-03**.
- [14] Girard, S. and Jacob, P. (2001) Extreme values and kernel estimates of point processes boundaries. *Technical report ENSAM-INRA-UM2*, **01-02**.
- [15] Haar, A. (1910) Zur Theorie der orthogonalen Funktionensysteme. *Math. Ann.* **69**, 331–371.

-
- [16] Hall, P., Nussbaum, M. and Stern, S.E. (1997) On the estimation of a support curve of indeterminate sharpness. *J. Multivariate Anal.*, **62**, 204–232.
- [17] Härdle, W., Hall, P. and Simar, L. (1995) Iterated bootstrap with application to frontier models. *J. Productivity Anal.*, **6**, 63–76.
- [18] Härdle, W., Park, B. U. and Tsybakov, A. B. (1995) Estimation of a non sharp support boundaries. *J. Multiv. Analysis*, **43**, 205–218.
- [19] Härdle, W. (1990) *Applied nonparametric regression*, Cambridge University Press, Cambridge.
- [20] Hardy, A. and Rasson, J.P. (1982) Une nouvelle approche des problèmes de classification automatique. *Statistique et Analyse des données*, **7**, 41–56.
- [21] Hartigan, J.A. (1975) *Clustering Algorithms*. Wiley, Chichester.
- [22] Jacob, P. and Abbar, H. (1989) Estimating the edge of Cox process area. *Cahiers du Centre d'Etudes de Recherche Opérationnelle*, **31**, 215–226.
- [23] Jacob, P. and Suquet, P. (1995) Estimating the edge of a Poisson process by orthogonal series. *Journal of Statistical Planning and Inference*, **46**, 215–234.
- [24] Kallenberg, W. (1983) On moderate deviation theory in estimation. *The Annals of Statistics*, **11**(2), 498–504.
- [25] Korostelev, A., Simar, L. and Tsybakov, A. B. (1995) Efficient estimation of monotone boundaries. *The Annals of Statistics*, **23**, 476–489.
- [26] Korostelev, A.P. and Tsybakov, A.B. (1993) Minimax theory of image reconstruction. in *Lecture Notes in Statistics*, **82**, Springer-Verlag, New York.

Contents

1	Introduction	3
2	The boundary estimate	4
3	Main results	4
4	Proofs	7
5	Applications	18
5.1	Four boundary estimates	18
5.1.1	Kernel estimate	18
5.1.2	Haar estimate	18
5.1.3	Trigonometric estimate	19
5.1.4	Faber-Shauder estimate	19
5.2	Basic properties	20
5.2.1	Kernel estimate	20
5.2.2	Haar estimate	21
5.2.3	Trigonometric estimate	21
5.2.4	Faber-Shauder estimate	22
5.3	Convergence results	22
5.4	Numerical experiments	25
6	Appendix	29
6.1	Multivariate central limit theorem and moderate deviations	29
6.2	Technical lemmas	32



Unité de recherche INRIA Rhône-Alpes
655, avenue de l'Europe - 38330 Montbonnot-St-Martin (France)

Unité de recherche INRIA Lorraine : LORIA, Technopôle de Nancy-Brabois - Campus scientifique
615, rue du Jardin Botanique - BP 101 - 54602 Villers-lès-Nancy Cedex (France)

Unité de recherche INRIA Rennes : IRISA, Campus universitaire de Beaulieu - 35042 Rennes Cedex (France)

Unité de recherche INRIA Rocquencourt : Domaine de Voluceau - Rocquencourt - BP 105 - 78153 Le Chesnay Cedex (France)

Unité de recherche INRIA Sophia Antipolis : 2004, route des Lucioles - BP 93 - 06902 Sophia Antipolis Cedex (France)

Éditeur
INRIA - Domaine de Voluceau - Rocquencourt, BP 105 - 78153 Le Chesnay Cedex (France)
<http://www.inria.fr>
ISSN 0249-6399