



# On Solvability of Boundary Value Problems in Elastoplasticity

Alexander Khludnev, Jan Sokolowski

► **To cite this version:**

Alexander Khludnev, Jan Sokolowski. On Solvability of Boundary Value Problems in Elastoplasticity. [Research Report] RR-3163, INRIA. 1997, pp.18. <inria-00073525>

**HAL Id: inria-00073525**

**<https://hal.inria.fr/inria-00073525>**

Submitted on 24 May 2006

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

INSTITUT NATIONAL DE RECHERCHE EN INFORMATIQUE ET EN AUTOMATIQUE

***ON SOLVABILITY OF BOUNDARY VALUE  
PROBLEMS IN ELASTOPLASTICITY***

Alexander M. Khludnev , Jan Sokołowski

**N° 3163**

Mai 1997

\_\_\_\_\_ THÈME 4 \_\_\_\_\_



*R*apport  
de recherche







## ON SOLVABILITY OF BOUNDARY VALUE PROBLEMS IN ELASTOPLASTICITY

Alexander M. Khludnev , Jan Sokołowski

Thème 4 — Simulation et optimisation  
de systèmes complexes  
Projet Numath

Rapport de recherche n3163 — Mai 1997 — 18 pages

**Abstract:** In the paper the existence of a solution to the three dimensional elastoplastic problem with the Prandtl-Reuss constitutive law and the Neumann boundary conditions is obtained. The proof is based on a suitable combination of the parabolic regularization of equations and the penalty method for the elastoplastic yield condition. The method is applied in the case of the domain with smooth boundary as well as in the case of an interior crack. It is shown that the weak solutions to the elastoplastic problem satisfying the variational inequality meet all boundary conditions.

**Key-words:** elastoplastic problem, Prandtl-Reuss constitutive law, parabolic regularization, elastoplastic yield condition, variational inequality

*(Résumé : tsvp)*

\* Lavrentyev Institute of Hydrodynamics of the Russian Academy of Sciences, Novosibirsk 630090, Russia

† Institut Elie Cartan, Laboratoire de Mathématiques, Université Henri Poincaré Nancy I, B.P. 239, 54506 Vandœuvre lès Nancy Cedex, France and Systems Research Institute of the Polish Academy of Sciences, ul. Newelska 6, 01-447 Warszawa, Poland; e-mail: sokolowski@iecn.u-nancy.fr

## Sur les équations de l'élastoplasticité : existence des solutions

**Résumé :** Dans cet article, nous prouvons l'existence d'une solution pour le problème élastoplastique en dimension 3 avec la loi constitutive de Prandtl-Reuss et des conditions au bord de type Neumann.

La preuve repose sur la régularisation parabolique des équations et sur la méthode de pénalisation pour la condition  $\sigma \in K$  de la loi de comportement.

Cette méthode est utilisée aussi bien dans le cas d'un domaine à frontière régulière que dans le cas d'une fissure intérieure.

Nous montrons que les solutions faibles du problème élastoplastique vérifiant l'inégalité variationnelle vérifient les conditions au bord.

**Mots-clé :** loi de Prandtl-Reuss, élastoplasticité, fissure intérieure, inégalité variationnelle

## 1. Introduction

By standard approach to the elastoplastic problems the weak solutions are introduced in the form of solutions to the variational inequalities. We refer the reader to [1–9] for the related results. The variational inequalities are derived under assumption that solutions to the elastoplastic problem are sufficiently smooth and the originally prescribed boundary conditions are satisfied. On the other hand, the original boundary conditions are not easily recovered for the solutions to the variational inequalities, even if such a weak solution is regular since the set of admissible stresses is not any linear space. As the result at least a part of boundary conditions prescribed for the elastoplastic problem is not obtained for the weak solutions. The problem has been solved in a satisfactory way only in one dimensional cases; it has been shown that solutions of variational inequalities satisfy all boundary conditions one expects to obtain for the elastoplastic problem [2, 10–13].

In this paper we consider the three dimensional elastoplastic Prandtl-Reuss model with the Neumann boundary conditions and prove that all original boundary conditions hold. The proof is based on a suitable combination of the parabolic regularization and the penalty method for the constraints imposed upon the stresses. In section 2 domains with smooth boundaries are considered, and in section 3 the case of a domain obtained by removing a smooth two dimensional surface from its interior is analysed. The latter corresponds to elastoplastic problems for bodies with interior cracks.

We start with notations and preliminary remarks. Let  $\Omega \subset \mathbb{R}^3$  be a bounded and connected domain with a smooth boundary  $\Gamma$  having the exterior unit normal  $n = (n_1, n_2, n_3)$ . Denote  $v = (v_1, v_2, v_3)$ ,  $v_{i,j} = \frac{\partial v_i}{\partial x_j}$ ,  $x = (x_1, x_2, x_3) \in \Omega$ ,

$$\varepsilon_{ij}(v) = \frac{1}{2}(v_{i,j} + v_{j,i}), \quad i, j = 1, 2, 3.$$

It is well known that if  $\varepsilon_{ij}(v) = 0$  in  $\Omega$ ,  $i, j = 1, 2, 3$ , then  $v_i(x) = c_i + b_{ij}x_j$ ,  $i = 1, 2, 3$ , where  $c_i, b_{ij} \in \mathbb{R}$ ,  $b_{ij} = -b_{ji}$ ,  $i, j = 1, 2, 3$ . Linear space of all vectors  $v = (v_1, v_2, v_3)$ ,  $v_i(x) = c_i + b_{ij}x_j$ , is called the space of rigid displacements and denoted by  $R(\Omega)$ .

Summation convention over repeated indices  $i, j, k, l = 1, 2, 3$  is used throughout the paper. All functions with two lower indices are assumed to be symmetric in those indices, eg.  $\sigma_{ij} = \sigma_{ji}$ ,  $i, j = 1, 2, 3$ .

To simplify the formulae below the inclusion  $v \in L^2(\Omega)$  for the vector function  $v = (v_1, v_2, v_3)$  will mean  $v_i \in L^2(\Omega)$ ,  $i = 1, 2, 3$ , and the following notation is adopted for the scalar product of the vector functions  $u, v$ ,

$$\langle v, u \rangle = \langle v_i, u_i \rangle = \int_{\Omega} v_i(x) u_i(x) dx .$$

The same convention is used for tensor functions

$$\sigma \in L^2(\Omega) \text{ for } \sigma_{ij} \in L^2(\Omega), \quad i, j = 1, 2, 3,$$

$$\langle \sigma, \varepsilon \rangle = \langle \sigma_{ij}, \varepsilon_{ij} \rangle = \int_{\Omega} \sigma_{ij}(x) \varepsilon_{ij}(x) dx .$$

In view of the convention we use the notation  $v \in L^2(\Omega)$  for the space of vector functions as well as for the space of tensor functions, ie. we write

$$v \in L^2(\Omega) \text{ and } \sigma \in L^2(\Omega)$$

instead of  $v \in [L^2(\Omega)]^3$  and  $\sigma \in [L^2(\Omega)]^9$ .

Consider all vector functions with components from  $L^2(\Omega)$  satisfying the following conditions

$$\int_{\Omega} v = 0, \quad \int_{\Omega} (v_i x_j - v_j x_i) = 0, \quad i, j = 1, 2, 3, \quad v = (v_1, v_2, v_3). \quad (1)$$

It is clear that if  $\rho \in R(\Omega)$  satisfies (1) then  $\rho \equiv 0$ . In fact, we have  $L^2(\Omega) = R(\Omega) \oplus R(\Omega)^\perp$ , and  $R(\Omega)^\perp$  coincides with the subspace of  $L^2(\Omega)$  of vector functions with components from  $L^2(\Omega)$  satisfying (1).

Let

$$H^1(\Omega) = \{v = (v_1, v_2, v_3) \mid v_i \in L^2(\Omega), \quad i = 1, 2, 3, \quad v_{i,j} \in L^2(\Omega), \quad i, j = 1, 2, 3\}.$$

In the space  $H^1(\Omega)$  we consider the following equivalent norms,

$$\|v\|_1^2 = \|v\|_0^2 + \sum_{i,j=1}^3 \|\varepsilon_{ij}(v)\|_0^2, \quad (2)$$

$$|v|_1 = \left| \int_{\Omega} v \right| + \sum_{i,j=1}^3 \left| \int_{\Omega} (v_i x_j - v_j x_i) \right| + \sum_{i,j=1}^3 \|\varepsilon_{ij}(v)\|_0. \quad (3)$$

Here,  $\|\cdot\|_0$  is the norm in  $L^2(\Omega)$ . It can be shown by an application of the second Korn inequality that the norm (2) is equivalent to the usual norm in  $H^1(\Omega)$ . For the norm (3) it is easy to see that the first two terms in the right-hand side of (3) define the seminorm in  $H^1(\Omega)$  being the norm in  $R(\Omega)$ , and the statement follows from [1].

We can consider the scalar product in  $H^1(\Omega)$  inducing the norm (2),

$$(u, v) = \langle u, v \rangle + \langle \varepsilon_{ij}(u), \varepsilon_{ij}(v) \rangle, \quad u, v \in H^1(\Omega). \quad (4)$$

In this case  $H^1(\Omega) = R(\Omega) \oplus H_N^1(\Omega)$ , where

$$H_N^1(\Omega) = \{v = (v_1, v_2, v_3) \in H^1(\Omega) \mid v \text{ satisfies (1)}\}.$$

This means that in  $H_N^1(\Omega)$  the following equivalent norm is defined

$$|v|_1 = \sum_{i,j=1}^3 \|\varepsilon_{ij}(v)\|_0.$$

Furthermore, the following notation is used. Consider the space

$$LD(\Omega) = \{v = (v_1, v_2, v_3) \mid v_i \in L^1(\Omega), \quad i = 1, 2, 3, \quad \varepsilon_{ij}(v) \in L^1(\Omega), \quad i, j = 1, 2, 3\}$$

equipped with the norm

$$\|v\|_{LD(\Omega)} = \|v\|_{L^1(\Omega)} + \sum_{i,j=1}^3 \|\varepsilon_{ij}(v)\|_{L^1(\Omega)}. \quad (5)$$

Let

$$LD_N(\Omega) = \{v \in LD(\Omega) \mid v \text{ satisfies (1)}\}.$$

Along with the usual norm (5) we shall consider the following norm in  $LD(\Omega)$  (see [1])

$$|v|_{LD(\Omega)} = \left| \int_{\Omega} v \right| + \sum_{i,j=1}^3 \left| \int_{\Omega} (v_i x_j - v_j x_i) \right| + \sum_{i,j=1}^3 \|\varepsilon_{ij}(v)\|_{L^1(\Omega)}.$$

Consequently, in the linear subspace  $LD_N(\Omega)$  of the space  $LD(\Omega)$  the norm is defined as follows

$$\|v\|_{LD_N(\Omega)} = \sum_{i,j=1}^3 \|\varepsilon_{ij}(v)\|_{L^1(\Omega)}.$$

The space of bounded measures on  $\Omega$  is denoted by  $\mathcal{M}^1(\Omega)$ ;  $\mathcal{M}^1(\Omega) = (C_0(\Omega))^*$  is the dual space of the normed space  $C_0(\Omega)$  of continuous functions with compact supports, equipped with the uniform convergence topology. It is known that any bounded set in  $\mathcal{M}^1(\Omega)$  is relatively compact in the  $(*)$ -weak topology, ie. every bounded sequence in  $\mathcal{M}^1(\Omega)$  contains a subsequence which is  $(*)$ -weak convergent. We recall that  $h_m \in \mathcal{M}^1(\Omega)$  is  $(*)$ -weak convergent to an element  $h \in \mathcal{M}^1(\Omega)$  as  $m \rightarrow \infty$  if

$$h_m(\varphi) \rightarrow h(\varphi) \quad \forall \varphi \in C_0(\Omega).$$

Introduce the Banach space of vector functions of bounded deformations

$$BD(\Omega) = \{v = (v_1, v_2, v_3) \mid v_i \in L^1(\Omega), i = 1, 2, 3, \varepsilon_{ij}(v) \in \mathcal{M}^1(\Omega), i, j = 1, 2, 3\}$$

endowed with the norm

$$\|v\|_{BD(\Omega)} = \|v\|_{L^1(\Omega)} + \sum_{i,j=1}^3 \|\varepsilon_{ij}(v)\|_{\mathcal{M}^1(\Omega)}.$$

In the same way as above,

$$BD_N(\Omega) = \{v \in BD(\Omega) \mid v \text{ satisfies (1)}\}.$$

In the sequel we need the appropriate Green formula. Let  $\sigma_{ij} \in L^2(\Omega)$ ,  $i, j = 1, 2, 3$ ,  $\sigma_{ij,j} \in L^2(\Omega)$ ,  $i = 1, 2, 3$ . Then  $\sigma_{ij}n_j$ ,  $i = 1, 2, 3$ , can be considered as elements of the space  $H^{-1/2}(\Gamma)$  on the boundary  $\Gamma$  and, moreover, the following formula holds [1]

$$-\langle \sigma_{ij,j}, \theta \rangle = \langle \sigma_{ij}, \theta_{,j} \rangle - \langle \sigma_{ij}n_j, \theta \rangle_{1/2,\Gamma} \quad \forall \theta \in H^1(\Omega), i = 1, 2, 3, \quad (6)$$

where  $\langle \cdot, \cdot \rangle_{1/2,\Gamma}$  denotes a duality pairing between the spaces  $H^{-1/2}(\Gamma)$  and  $H^{1/2}(\Gamma)$ .

## 2. Domain with a smooth boundary

Let  $\Omega \subset \mathbb{R}^3$  be a bounded and connected domain with smooth boundary  $\Gamma$ ,  $Q = \Omega \times (0, T)$ ,  $x = (x_1, x_2, x_3) \in \Omega$ ,  $t \in (0, T)$ . The elastoplastic problem for a body occupying the domain  $\Omega$  in the nondeformable state is formulated as follows.

Find the functions  $v = (v_1, v_2, v_3)$ ,  $\sigma = \{\sigma_{ij}\}$ ,  $\eta_{ij}$ ,  $i, j = 1, 2, 3$ , defined in  $Q$  and satisfying the following equations and inequalities

$$-\sigma_{ij,j} = f_i, \quad i = 1, 2, 3, \quad (7)$$

$$\varepsilon_{ij}(v) = c_{ijkl}\dot{\sigma}_{kl} + \eta_{ij}, \quad i, j = 1, 2, 3, \quad (8)$$

$$\Phi(\sigma) \leq 0, \quad \eta_{ij}(\bar{\sigma}_{ij} - \sigma_{ij}) \leq 0 \quad \forall \bar{\sigma}, \Phi(\bar{\sigma}) \leq 0, \quad (9)$$

$$\sigma_{ij}n_j = 0, \quad i = 1, 2, 3, \quad \text{on } \Gamma \times (0, T), \quad (10)$$

$$\sigma = 0, \quad t = 0. \quad (11)$$

The functions  $v$ ,  $\sigma_{ij}$ ,  $\varepsilon_{ij}(v)$  represent the velocity, components of the stress tensor and components of the strain tensor velocity, respectively. The dot denotes the derivate with respect to  $t$ , eg.  $\dot{\sigma}_{kl} = \frac{d\sigma_{kl}}{dt}$ .



The convex and continuous function  $\Phi$  defines the plasticity yield condition. It is assumed that the set

$$\tilde{K} = \{\sigma = \{\sigma_{ij}\} \in \mathbb{R}^6 \mid \Phi(\sigma) \leq 0\} \quad (12)$$

contains zero as its interior point,  $0 \in \text{int}\tilde{K}$ . We assume that the symmetry conditions  $c_{ijkl}(x) = c_{jikl}(x) = c_{klij}(x)$  for  $i, j, k, l = 1, 2, 3$  are satisfied and there exist two positive constants  $c_1, c_2$  such that

$$c_1|\sigma|^2 \leq c_{ijkl}\sigma_{kl}\sigma_{ij} \leq c_2|\sigma|^2 \quad \forall \sigma = \{\sigma_{ij}\}. \quad (13)$$

To simplify the formulae below we assume that  $c_{ijkl} = \delta_k^i \delta_l^j$ ,  $\delta_j^i$  is the Kronecker symbol. Nevertheless, all the results obtained in the paper are valid in general case (13).

The functions  $\eta_{ij}$ ,  $i, j = 1, 2, 3$ , can be eliminated from (8), (9). In fact, multiply (8) by  $\bar{\sigma}_{ij} - \sigma_{ij}$  and sum up in  $i, j$ . This leads to

$$\Phi(\sigma) \leq 0, \quad (\dot{\sigma}_{ij} - \varepsilon_{ij}(v))(\bar{\sigma}_{ij} - \sigma_{ij}) \geq 0 \quad \forall \bar{\sigma}, \Phi(\bar{\sigma}) \leq 0. \quad (14)$$

Inequality (14) is used as the definition of a weak solution to the problem (7)–(11).

Consider the set of admissible stresses

$$K = \{\sigma = \{\sigma_{ij}\} \mid \sigma_{ij} \in L^2(\Omega), i, j = 1, 2, 3, \Phi(\sigma(x)) \leq 0 \text{ a.e. in } \Omega\}$$

and denote by  $p$  the penalty operator related to the set  $K$ . The operator is given by the formula  $p(\sigma) = \sigma - \pi\sigma$ , where  $\pi : L^2(\Omega) \rightarrow K \subset L^2(\Omega)$  is the operator of orthogonal projection with respect to the scalar product in  $L^2(\Omega)$ . Recall that the operator  $p : L^2(\Omega) \rightarrow L^2(\Omega)$  is bounded, monotone and continuous, and we use the convention to denote  $L^2(\Omega)$  for  $[L^2(\Omega)]^9$ .

Let the brackets  $(\cdot, \cdot)$  denote the scalar product in  $L^2(Q)$ ,

$$(u, v) = \int_0^T \langle u(t), v(t) \rangle dt$$

for vector functions  $u, v \in L^2(Q)$ ,  $f = (f_1, f_2, f_3)$ .

The space of statically admissible stresses is denoted by

$$V_0(\Omega) = \{\sigma = \{\sigma_{ij}\} \mid \sigma_{ij} \in L^2(\Omega), i, j = 1, 2, 3; \sigma_{ij,j} \in L^3(\Omega), i = 1, 2, 3;$$

$$\sigma_{ij}n_j = 0, i = 1, 2, 3, \text{ on } \Gamma\}.$$

Suppose that there exists a function  $\xi = \{\xi_{ij}\}$ ,  $i, j = 1, 2, 3$ , satisfying the equation (7) such that  $\xi \in C^2(\bar{Q})$ ,  $\xi(0) = \dot{\xi}(0) = 0$  and  $(1 + \kappa)\xi(t) \in K \cap V_0(\Omega)$ , where  $\kappa \in \mathbb{R}$ ,  $\kappa > 0$ ,  $t \in [0, T]$ .

Now we can prove the following existence theorem for the problem (7)–(11).

**Theorem 1.** Assume that  $f \in L^3(Q)$ ,  $\dot{f} \in L^2(Q)$ ,  $f(0) = 0$ ,  $\langle f(t), \rho \rangle = 0 \quad \forall \rho \in R(\Omega), t \in [0, T]$ , and the above assumption on  $\xi$  is satisfied.

Then there exist functions  $v = (v_1, v_2, v_3)$ ,  $\sigma = \{\sigma_{ij}\}$  such that

$$v \in L^2(0, T; BD_N(\Omega)), \quad \sigma \in L^2(0, T; V_0(\Omega)), \quad \dot{\sigma} \in L^2(Q), \quad \sigma(t) \in K, \quad t \in (0, T),$$

$$(\sigma_{ij}, \varepsilon_{ij}(\bar{v})) = (f, \bar{v}) \quad \forall \bar{v} \in L^2(0, T; H^1(\Omega)), \quad (15)$$

$$(\dot{\sigma}_{ij}, \bar{\sigma}_{ij} - \sigma_{ij}) + (v_i, \bar{\sigma}_{ij,j} - \sigma_{ij,j}) \geq 0 \quad \forall \bar{\sigma} \in L^2(0, T; V_0(\Omega)), \quad (16)$$

$$\text{with } \bar{\sigma}(t) \in K \quad \text{a.e. in } (0, T),$$

$$\sigma = 0, \quad t = 0. \quad (17)$$

**Proof.** First, an approximation of the problem (7)–(11) by penalization of the plastic yield condition is introduced. Then, the parabolic regularization of the penalized problem is considered. The auxiliary boundary value problem constructed in such a way depends on two positive parameters  $\alpha, \delta$ . The parameter  $\alpha$  is used for the parabolic regularization, the parameter  $\delta$  is used for the penalty term. We prove the existence of solutions for the fixed parameters  $\alpha > 0, \delta > 0$  and justify the passage to limits as  $\alpha, \delta \rightarrow 0$ . A priori estimates uniform with respect to  $\alpha, \delta$  are needed to perform the passage to the limits. All necessary estimates (29), (31), (32) are derived below.

We consider in the domain  $Q$  the following auxiliary boundary value problem. Find functions  $v = (v_1, v_2, v_3), \sigma = \{\sigma_{ij}\}$  such that

$$\alpha \dot{v}_i - \alpha \varepsilon_{ij}(v)_{,j} - \sigma_{ij,j} = f_i, \quad i = 1, 2, 3, \quad (18)$$

$$\dot{\sigma}_{ij} + \alpha \sigma_{ij} - \varepsilon_{ij}(v) + \frac{1}{\delta} p(\sigma)_{ij} = 0, \quad i, j = 1, 2, 3, \quad (19)$$

$$\sigma_{ij} n_j + \alpha \varepsilon_{ij}(v) n_j = 0, \quad i = 1, 2, 3, \quad \text{on } \Gamma \times (0, T), \quad (20)$$

$$v = 0, \quad \sigma = 0, \quad t = 0. \quad (21)$$

The dependence of solutions to (18)–(21) on the parameters  $\alpha, \delta$  is not indicated in (18)–(21) in order to simplify the formulae. Note that boundary conditions (20) do not coincide with (10); the conditions (20) can be viewed as the regularization of (10) connected with the proposed regularization of the equilibrium equations (7). Also, the artificial initial condition for  $v$  is introduced.

A solution to the problem (18)–(21) is defined in the following sense

$$v \in L^2(0, T; H^1(\Omega)), \quad \dot{v}, \sigma, \dot{\sigma} \in L^2(Q), \quad (22)$$

$$\alpha(\dot{v}, \bar{v}) + \alpha(\varepsilon_{ij}(v), \varepsilon_{ij}(\bar{v})) + (\sigma_{ij}, \varepsilon_{ij}(\bar{v})) = (f, \bar{v}) \quad \forall \bar{v} \in L^2(0, T; H^1(\Omega)), \quad (23)$$

$$\dot{\sigma}_{ij} + \alpha \sigma_{ij} - \varepsilon_{ij}(v) + \frac{1}{\delta} p(\sigma)_{ij} = 0, \quad i, j = 1, 2, 3, \quad (24)$$

$$v = 0, \quad \sigma = 0, \quad t = 0. \quad (25)$$

In this case the boundary conditions (20) are included in (23). At the first step of the proof we derive a priori estimates. To this end it is assumed that the solutions to (18)–(21) are sufficiently smooth. Multiply (18), (19) by  $v_i, \sigma_{ij} - \xi_{ij}$ , respectively, and integrate over  $\Omega$ . Taking into account that the penalty term is nonnegative this provides the inequality

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \{ \alpha \|v\|_0^2 + \|\sigma\|_0^2 \} + \alpha \|\sigma\|_0^2 - \langle \alpha \varepsilon_{ij}(v)_{,j} + \sigma_{ij,j}, v_i \rangle - \\ & - \langle \varepsilon_{ij}(v), \sigma_{ij} - \xi_{ij} \rangle - \langle f, v \rangle \leq \langle \alpha \sigma + \dot{\sigma}, \xi \rangle. \end{aligned} \quad (26)$$

For the sake of simplicity we do not show in (26) the dependence of  $v, \sigma, \xi, f$  on  $t$ . The integration by parts in the third term of the left-hand side in (26) can be performed. Recall that  $\xi$  satisfies the equation (7). As a result the following inequality is obtained

$$\begin{aligned} & \frac{1}{2} \{ \alpha \|v(t)\|_0^2 + \|\sigma(t)\|_0^2 \} + \alpha \sum_{i,j=1}^3 \int_0^t \|\varepsilon_{ij}(v)\|_0^2 d\tau \leq \\ & \leq \langle \sigma(t), \xi(t) \rangle - \int_0^t \langle \sigma(\tau), \dot{\xi}(\tau) \rangle d\tau + \frac{\alpha}{2} \int_0^t \|\xi(\tau)\|_0^2 d\tau. \end{aligned} \quad (27)$$

Since  $\langle \sigma, \xi \rangle \leq \frac{1}{4} \|\sigma\|_0^2 + \|\xi\|_0^2$ , the integration of (27) leads to the estimate

$$\sup_{0 \leq t \leq T} \|\sigma(t)\|_0^2 + \alpha \|v\|_{L^2(Q)}^2 + \alpha \sum_{i,j=1}^3 \int_0^T \|\varepsilon_{ij}(v)\|_0^2 d\tau \leq c \quad (28)$$

with the constant  $c$  being uniform in  $\alpha, \delta$ , for  $\alpha \leq \alpha_0$ . Hence

$$\sup_{0 \leq t \leq T} \|\sigma(t)\|_0^2 + \alpha \|v\|_{L^2(0,T;H^1(\Omega))}^2 \leq c. \quad (29)$$

The derivation of the following estimate requires the  $(\alpha, \delta)$ -uniform boundedness of  $\frac{1}{\delta} p(\sigma)$  in the  $L^1(Q)$  norm. By (29) it is easy to see that

$$\frac{1}{\delta} \int_0^T \langle p(\sigma), \sigma - \xi \rangle dt \leq c$$

uniformly in  $\alpha, \delta$ , provided that the penalty term is not neglected when deriving (26). Due to the monotonicity of  $p$

$$\frac{1}{\delta} \int_0^T \langle p(\sigma), \bar{\sigma} - \sigma \rangle dt \leq 0 \quad \forall \bar{\sigma} \in L^2(Q), \bar{\sigma}(t) \in K.$$

Summing up the two last inequalities we have

$$\frac{1}{\delta} \int_0^T \langle p(\sigma), \bar{\sigma} - \xi \rangle dt \leq c.$$

We can take here  $\bar{\sigma} = \xi + \bar{\xi}$ ,  $\|\bar{\xi}\|_{L^\infty(Q)} \leq \mu$ ,  $\mu > 0$  is sufficiently small. By the hypothesis imposed on  $\xi$  the inclusions  $\bar{\sigma}(t) \in K, t \in (0, T)$ , hold, hence

$$\frac{1}{\delta} \int_0^T \langle p(\sigma), \bar{\xi} \rangle dt \leq c \quad \forall \bar{\xi}, \|\bar{\xi}\|_{L^\infty(Q)} \leq \mu$$

and, consequently,

$$\frac{1}{\delta} \|p(\sigma)\|_{L^1(Q)} \leq c. \quad (30)$$

In the sequel this estimate is improved, namely, we show that  $\frac{1}{\delta} p(\sigma)$  is, in fact, bounded in  $L^2(0, T; L^1(\Omega))$ .

We derive a priori estimate for the time derivatives  $\dot{v}, \dot{\sigma}$ . It follows from (18), (19), (21) that we have the homogenous initial conditions for the derivatives,

$$\dot{v}_i(0) = 0, \quad i = 1, 2, 3, \quad \dot{\sigma}_{ij}(0) = 0, \quad i, j = 1, 2, 3.$$

Differentiate with respect to  $t$  the equations (18), (19) and multiply by  $\dot{v}_i, \dot{\sigma}_{ij} - \dot{\xi}_{ij}$ , respectively. Since the term

$$\frac{1}{\delta} \left\langle \frac{d}{dt} p(\sigma(t)), \dot{\sigma}(t) \right\rangle$$

is nonnegative for almost all  $t \in (0, T)$  (see [14], page 399) the above multiplication and integration over  $\Omega$  results in the inequality

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \{ \alpha \|\dot{v}\|_0^2 + \|\dot{\sigma}\|_0^2 \} + \alpha \|\dot{\sigma}\|_0^2 - \langle \alpha \varepsilon_{ij}(\dot{v}), \dot{\sigma}_{ij} + \dot{\sigma}_{ij,j}, \dot{v}_i \rangle - \\ & - \langle \varepsilon_{ij}(\dot{v}), \dot{\sigma}_{ij} - \dot{\xi}_{ij} \rangle - \langle \dot{f}, \dot{v} \rangle \leq \langle \alpha \dot{\sigma} + \dot{\sigma}, \dot{\xi} \rangle + \frac{1}{\delta} \left\langle \frac{d}{dt} p(\sigma), \dot{\xi} \right\rangle. \end{aligned}$$

Boundary conditions (20) can be taken into account in order to integrate by parts in the left-hand side of the latter inequality.

We integrate the last inequality with respect to  $t$  over the interval  $(0, t)$ . This implies

$$\begin{aligned} & \frac{1}{2} \{ \alpha \| \dot{v}(t) \|_0^2 + \| \dot{\sigma}(t) \|_0^2 \} + \alpha \sum_{i,j=1}^3 \int_0^t \| \varepsilon_{ij}(\dot{v}) \|_0^2 d\tau \leq \\ & \leq \frac{1}{\delta} \langle p(\sigma), \dot{\xi} \rangle_0^t - \frac{1}{\delta} \int_0^t \langle p(\sigma), \ddot{\xi} \rangle d\tau + \\ & + \langle \dot{\sigma}, \dot{\xi} \rangle_0^t - \int_0^t \langle \dot{\sigma}, \ddot{\xi} \rangle d\tau + \frac{\alpha}{2} \int_0^t \| \dot{\xi}(\tau) \|_0^2 d\tau . \end{aligned}$$

By estimates (29), (30) and the condition  $\dot{\xi}(0) = 0$  the  $(\alpha, \delta)$ -uniform estimate follows

$$\| \dot{\sigma} \|_{L^2(Q)}^2 + \alpha \| \dot{v} \|_{L^2(Q)}^2 \leq c. \quad (31)$$

Let  $v = v_N + \rho_N$  be the decomposition of  $v$  into the sum of two orthogonal elements,  $v_N \in L^2(0, T; H_N^1(\Omega))$ ,  $\rho_N \in L^2(0, T; R(\Omega))$ . We should note at this point that  $L^2(0, T; H^1(\Omega)) = L^2(0, T; R(\Omega)) \oplus L^2(0, T; H_N^1(\Omega))$  provided that the scalar product (4) is used in  $H^1(\Omega)$ . For almost all  $t \in (0, T)$ ,  $v_N(t) = (v_{N1}(t), v_{N2}(t), v_{N3}(t))$ ,

$$\int_{\Omega} v_N(t) = 0, \quad \int_{\Omega} (v_{Ni}(t)x_j - v_{Nj}(t)x_i) = 0, \quad i, j = 1, 2, 3.$$

Hence, by (29), (31) it follows from (19) that  $\varepsilon_{ij}(v_N)$ ,  $i, j = 1, 2, 3$ , are bounded in  $L^2(Q)$  uniformly in  $\alpha$  for any fixed  $\delta$ . This implies the estimate

$$\| v_N \|_{L^2(0, T; H_N^1(\Omega))} \leq c(\delta) \quad (32)$$

with the constant  $c(\delta)$  depending, in general, on  $\delta$ .

Now, observe that in view of the estimates (29), (31) we can use the Galerkin approximations of parabolic problems with monotone operators [14] in order to show that for any fixed  $\alpha, \delta$  the solution to (18)–(21) exists in the sense of (22)–(25).

The estimates obtained allow us to pass to the limit as  $\alpha \rightarrow 0$ . Indeed, denote by  $v^\alpha, \sigma^\alpha$  the solution to (22)–(25) and consider the decomposition  $v^\alpha = v_N^\alpha + \rho_N^\alpha$ ,  $v_N^\alpha \in L^2(0, T; H_N^1(\Omega))$ ,  $\rho_N^\alpha \in L^2(0, T; R(\Omega))$ . Note that the solution  $(v^\alpha, \sigma^\alpha)$  satisfies the estimates (29), (31), (32). Hence, from the sequence  $v^\alpha, \sigma^\alpha$  one can choose a subsequence (with the previous notation for the subsequence) such that for any fixed  $\delta > 0$  and  $\alpha \rightarrow 0$

$$\begin{aligned} \alpha v^\alpha & \rightarrow 0 \quad \text{weakly in } L^2(0, T; H^1(\Omega)), \\ v_N^\alpha & \rightarrow v^\delta \quad \text{weakly in } L^2(0, T; H_N^1(\Omega)), \\ \alpha \dot{v}^\alpha & \rightarrow 0 \quad \text{weakly in } L^2(Q), \\ \sigma^\alpha & \rightarrow \sigma^\delta \quad \text{weakly in } L^2(Q), \\ \dot{\sigma}^\alpha & \rightarrow \dot{\sigma}^\delta \quad \text{weakly in } L^2(Q). \end{aligned}$$

Passing to the limit in (18), (19) as  $\alpha \rightarrow 0$  it follows that

$$(\sigma_{ij}^\delta, \varepsilon_{ij}(\bar{v})) = (f, \bar{v}) \quad \forall \bar{v} \in L^2(0, T; H^1(\Omega)), \quad (33)$$

$$\dot{\sigma}_{ij}^\delta - \varepsilon_{ij}(v^\delta) + \frac{1}{\delta}p(\sigma^\delta)_{ij} = 0, \quad i, j = 1, 2, 3. \quad (34)$$

A justification of the convergence  $p(\sigma^\alpha) \rightarrow p(\sigma^\delta)$  can be done by the monotonicity arguments. We omit the details.

Now let us show that  $\frac{1}{\delta}p(\sigma^\delta)$  are bounded in  $L^2(0, T, L^1(\Omega))$  uniformly in  $\delta$ . It follows from (33) that for a.e.  $t \in (0, T)$ ,

$$-\langle \sigma_{ij}^\delta(t), \varepsilon_{ij}(\bar{v}) \rangle = \langle f(t), \bar{v} \rangle \quad \forall \bar{v} \in H^1(\Omega).$$

Hence, for almost all  $t \in (0, T)$

$$-\langle \sigma_{ij}^\delta(t), \varepsilon_{ij}(v^\delta(t)) \rangle = \langle f(t), v^\delta(t) \rangle. \quad (35)$$

Multiply (34) by  $\sigma_{ij}^\delta - \xi_{ij}$  and integrate over  $\Omega$ . This provides

$$\begin{aligned} \langle \varepsilon_{ij}(v^\delta(t)), \sigma_{ij}^\delta(t) - \xi_{ij}(t) \rangle + \frac{1}{\delta} \langle p(\sigma^\delta(t)), \xi(t) - \sigma^\delta(t) \rangle = \\ = \langle \dot{\sigma}^\delta(t), \sigma^\delta(t) - \xi(t) \rangle. \end{aligned} \quad (36)$$

By (29), (31) the right-hand side of (36) is bounded in  $L^2(0, T)$  uniformly in  $\delta$ . Summing up (35) and (36) we obtain that

$$\frac{1}{\delta} \langle p(\sigma^\delta(t)), \sigma^\delta(t) - \xi(t) \rangle \quad \text{are bounded in } L^2(0, T) \quad (37)$$

uniformly in  $\delta$ . Introduce the convex functional on the space  $L^2(\Omega)$ ,

$$F(\sigma) = \|\sigma - \pi\sigma\|_0^2, \quad \sigma = \{\sigma_{ij}\}, \quad i, j = 1, 2, 3.$$

The Gateaux derivative of  $F$  is given by  $F'(\sigma) = 2p(\sigma)$ . Let us take a function  $\tilde{\sigma} = \{\tilde{\sigma}_{ij}\} \in L^\infty(Q)$ . Then by the conditions imposed on  $\xi$  it follows that  $\xi(t) + \tilde{\sigma}(t)$  belongs to the set  $K, t \in (0, T)$ , provided that the norm  $\|\tilde{\sigma}\|_{L^\infty(Q)}$  is sufficiently small. By the convexity of  $F$  we have

$$\begin{aligned} \frac{1}{\delta} \langle p(\sigma^\delta(t)), \tilde{\sigma}(t) \rangle \leq \frac{1}{\delta} \langle p(\sigma^\delta(t)), \sigma^\delta(t) - \xi(t) \rangle + \\ + \frac{1}{2\delta} F(\xi(t) + \tilde{\sigma}(t)) - \frac{1}{2\delta} F(\sigma^\delta(t)). \end{aligned} \quad (38)$$

The second term of the right-hand side of (38) equals to zero by the inclusion  $\xi(t) + \tilde{\sigma}(t) \in K$  and consequently, in view of (37),

$$\frac{1}{\delta} \langle p(\sigma^\delta(t)), \tilde{\sigma}(t) \rangle \quad \text{are bounded in } L^2(0, T).$$

Since  $\tilde{\sigma}$  is an arbitrary element of the space  $L^\infty(Q)$  with a small norm we infer the desired estimate,

$$\frac{1}{\delta} p(\sigma^\delta(t)) \quad \text{are bounded in } L^2(0, T; L^1(\Omega)).$$

Hence, from (34) it follows that

$$\|\varepsilon_{ij}(v^\delta)\|_{L^2(0, T; L^1(\Omega))} \leq c, \quad i, j = 1, 2, 3. \quad (39)$$

The inequality (39) and the inclusion  $v^\delta \in L^2(0, T; H_N^1(\Omega))$  yield the estimate

$$\|v^\delta\|_{L^2(0, T; LD_N(\Omega))} \leq c,$$

uniformly in  $\delta$ . Moreover, the space  $L^1(\Omega)$  is continuously imbedded in  $\mathcal{M}^1(\Omega)$ , consequently

$$\|v^\delta\|_{L^2(0,T;BD_N(\Omega))} \leq c. \quad (40)$$

On the other hand the inequality

$$\|\sigma^\delta\|_{L^2(Q)} + \|\dot{\sigma}^\delta\|_{L^2(Q)} \leq c \quad (41)$$

is satisfied uniformly in  $\delta$ . Recall that  $BD(\Omega) \subset L^{3/2}(\Omega)$  in the three dimensional space. Moreover, we have the following estimate

$$\|\sigma^\delta(T)\|_0 \leq c\{\|\sigma^\delta\|_{L^2(Q)} + \|\dot{\sigma}^\delta\|_{L^2(Q)}\} \quad (42)$$

with the constant  $c$  independent of functions  $\sigma^\delta, \dot{\sigma}^\delta$ .

By (40), (41), (42) we can choose a subsequence (with the previous notation for the subsequence) such that as  $\delta \rightarrow 0$ ,

$$\begin{aligned} \sigma^\delta &\rightarrow \sigma \quad \text{weakly in } L^2(Q), \\ \dot{\sigma}^\delta &\rightarrow \dot{\sigma} \quad \text{weakly in } L^2(Q), \\ v^\delta &\rightarrow v \quad \text{weakly in } L^2(0,T;L^{3/2}(\Omega)), \\ \varepsilon_{ij}(v^\delta) &\rightarrow \varepsilon_{ij}(v) \quad (*)\text{-weakly in } L^2(0,T;\mathcal{M}^1(\Omega)), \quad i, j = 1, 2, 3, \\ \sigma^\delta(T) &\rightarrow \sigma(T) \quad \text{weakly in } L^2(\Omega). \end{aligned}$$

As a result, passing to the limit as  $\delta \rightarrow 0$  in (33), we obtain

$$(\sigma_{ij}, \varepsilon_{ij}(\bar{v})) = (f, \bar{v}) \quad \forall \bar{v} \in L^2(0,T;H^1(\Omega)). \quad (43)$$

The equations (34) imply

$$\begin{aligned} (\dot{\sigma}_{ij}^\delta, \bar{\sigma}_{ij} - \sigma_{ij}^\delta) - (\varepsilon_{ij}(v^\delta), \bar{\sigma}_{ij} - \sigma_{ij}^\delta) &\geq 0 \quad \forall \bar{\sigma} \in L^2(0,T;V_0(\Omega)), \\ \bar{\sigma}(t) &\in K, t \in (0,T). \end{aligned} \quad (44)$$

The identity (43) provides the fulfilment of the equations

$$-\sigma_{ij,j}^\delta = f_i, \quad i = 1, 2, 3, \quad (45)$$

in the sense of distributions.

Furthermore, by (33)

$$(\varepsilon_{ij}(v^\delta), \sigma_{ij}^\delta) = -(\sigma_{ij,j}^\delta, v_i^\delta)$$

and, moreover,

$$(\varepsilon_{ij}(v^\delta), \bar{\sigma}_{ij}) = -(v_i^\delta, \bar{\sigma}_{ij,j}) \quad \forall \bar{\sigma} \in L^2(0,T;V_0(\Omega)).$$

Hence, inequality (44) can be rewritten in the following form

$$\begin{aligned} (\dot{\sigma}_{ij}^\delta, \bar{\sigma}_{ij} - \sigma_{ij}^\delta) + (v_i^\delta, \bar{\sigma}_{ij,j} - \sigma_{ij,j}^\delta) &\geq 0 \quad \forall \bar{\sigma} \in L^2(0,T;V_0(\Omega)), \\ \bar{\sigma}(t) &\in K, t \in (0,T). \end{aligned} \quad (46)$$

By (45) the functions  $\sigma_{ij,j}^\delta$  can be replaced by  $-f_i$  and from (46) it follows that

$$(\dot{\sigma}_{ij}^\delta, \bar{\sigma}_{ij}) + (v_i^\delta, \bar{\sigma}_{ij,j} + f_i) \geq \frac{1}{2}\|\sigma^\delta(T)\|_0^2. \quad (47)$$

Passing to the lower limit in both sides of (47) and replacing  $f_i$  by  $-\sigma_{ij,j}$  we obtain (16). The inclusion  $\sigma(t) \in K, t \in (0, T)$ , can be verified by the standard arguments. Since  $v^\delta \in L^2(0, T; H_N^1(\Omega))$ , the convergence of  $v^\delta$  to  $v$  implies the inclusion  $v \in L^2(0, T; BD_N(\Omega))$ . The property  $\sigma(t) \in V_0(\Omega), t \in (0, T)$ , actually follows from (43) taking into account the equations

$$-\sigma_{ij,j} = f_i, \quad i = 1, 2, 3,$$

and the Green formula (6). Therefore, Theorem 1 is proved.

### 3. Domain with a nonsmooth boundary (body with a crack)

In this section we prove the existence theorem for elastoplastic problem in the case of the domain with nonsmooth boundary. In applications nonsmooth boundaries occur due to the presence of cracks in the body. In the framework of elasticity the properties of the solutions in domains with cracks are analysed in many papers, and we refer the reader to eg. [15–24] for the related results.

Again, let  $\Omega \subset \mathbb{R}^3$  be a bounded and connected domain with the smooth boundary  $\partial\Omega = \Gamma$  and  $\Sigma_c \subset \Omega$  be a smooth orientable two dimensional surface with a regular boundary. We assume that  $\Sigma_c$  can be extended in such a way that the domain  $\Omega$  is divided into two parts with Lipschitz boundaries. The surface  $\Sigma_c$  is described parametrically

$$x_i = x_i(y_1, y_2), \quad i = 1, 2, 3, \quad (48)$$

where  $(y_1, y_2) \in \bar{\omega}, \omega \subset \mathbb{R}^2$  is a bounded connected domain with smooth boundary  $\gamma, \bar{\omega} = \omega \cup \gamma$ . Assume that for any point  $(y_1, y_2) \in \bar{\omega}$  the rank of the Jakobi matrix  $\left[ \frac{\partial x_i}{\partial y_j} \right]$  equals 2 and the map (48) is one-to-one. In this case one can choose a unit normal vector to the surface  $\Sigma_c$ ,

$$\nu = \frac{\frac{\partial x}{\partial y_1} \times \frac{\partial x}{\partial y_2}}{\left| \frac{\partial x}{\partial y_1} \times \frac{\partial x}{\partial y_2} \right|}.$$

Denote  $\Omega_c = \Omega \setminus \Sigma_c, Q_c = \Omega_c \times (0, T), T > 0$ .

The equilibrium problem for elastoplastic body occupying the domain  $\Omega_c$  can be formulated as follows.

Find functions  $v = (v_1, v_2, v_3), \sigma = \{\sigma_{ij}\}, \eta_{ij}, i, j = 1, 2, 3$ , defined in  $Q_c$  and satisfying the following equations and inequalities

$$-\sigma_{ij,j} = f_i, \quad i = 1, 2, 3, \quad (49)$$

$$\varepsilon_{ij}(v) = c_{ijkl}\dot{\sigma}_{kl} + \eta_{ij}, \quad i, j = 1, 2, 3, \quad (50)$$

$$\Phi(\sigma) \leq 0, \quad \eta_{ij}(\bar{\sigma}_{ij} - \sigma_{ij}) \leq 0 \quad \forall \bar{\sigma}, \quad \Phi(\bar{\sigma}) \leq 0, \quad (51)$$

$$\sigma_{ij}n_j = 0, \quad i = 1, 2, 3, \quad \text{on } \Gamma \times (0, T), \quad (52)$$

$$\sigma_{ij}\nu_j = 0, \quad i = 1, 2, 3, \quad \text{on } \Sigma_c^\pm \times (0, T), \quad (53)$$

$$\sigma = 0, \quad t = 0. \quad (54)$$

The same notations are used as in sections 1,2. As we see, in this case the boundary of the domain  $\Omega_c$  consists of the parts  $\Gamma, \Sigma_c^+, \Sigma_c^-$ , where  $\Sigma_c^\pm$  correspond to the positive and negative directions of the normal  $\nu$ , respectively.

Introduce the space

$$H^1(\Omega_c) = \{v = (v_1, v_2, v_3) \mid v_i \in L^2(\Omega_c), i = 1, 2, 3; v_{i,j} \in L^2(\Omega_c), i, j = 1, 2, 3\}.$$

Note that functions from the space  $H^1(\Omega_c)$  have, in general, different boundary values on  $\Sigma_c^\pm$ . This holds for functions from functional spaces considered below provided that the boundary conditions are well defined even in a weak sense.

In this section we shall consider functions satisfying the following relations

$$\int_{\Omega_c} v = 0, \quad \int_{\Omega_c} (v_i x_j - v_j x_i) = 0, \quad i, j = 1, 2, 3, \quad v = (v_1, v_2, v_3). \quad (55)$$

Let

$$LD(\Omega_c) = \{v = (v_1, v_2, v_3) \mid v_i \in L^1(\Omega_c), i = 1, 2, 3, \varepsilon_{ij}(v) \in L^1(\Omega_c), i, j = 1, 2, 3\}.$$

The subspaces  $H_N^1(\Omega_c)$ ,  $LD_N(\Omega_c)$  consist of all function from  $H^1(\Omega_c)$  and  $LD(\Omega_c)$  respectively, satisfying (55). In the subspaces  $H_N^1(\Omega_c)$  and  $LD_N(\Omega_c)$  the following norms are introduced,

$$|v|_{H_N^1(\Omega_c)} = \sum_{i,j=1}^3 \|\varepsilon_{ij}(v)\|_{0,c}, \quad |v|_{LD_N(\Omega_c)} = \sum_{i,j=1}^3 \|\varepsilon_{ij}(v)\|_{L^1(\Omega_c)}$$

which are equivalent to the standard ones. Here  $\|\cdot\|_{0,c}$  stands for the norm in  $L^2(\Omega_c)$ . The proof of the norm equivalency uses the compactness of imbeddings  $H^1(\Omega_c) \subset L^2(\Omega_c)$ ,  $LD(\Omega_c) \subset L^1(\Omega_c)$  which are obtained under the conditions imposed on  $\Sigma_c$  and  $\Gamma$ . Indeed, let us prove that there exists a positive constant  $c$  such that

$$|v|_{LD(\Omega_c)} \geq c \|v\|_{LD(\Omega_c)} \quad \forall v \in LD(\Omega_c), \quad (56)$$

where

$$|v|_{LD(\Omega_c)} = \left| \int_{\Omega_c} v \right| + \sum_{i,j=1}^3 \left| \int_{\Omega_c} (v_i x_j - v_j x_i) \right| + \sum_{i,j=1}^3 \|\varepsilon_{ij}(v)\|_{L^1(\Omega_c)}.$$

Assume that the inequality (56) is false. In this case there exists a sequence  $v^k$ ,

$$\|v^k\|_{LD(\Omega_c)} = 1, \quad (57)$$

$$|v^k|_{LD(\Omega_c)} \leq \frac{1}{k}. \quad (58)$$

According to the assumptions on  $\Sigma_c$  we can divide the domain  $\Omega_c$  into two subdomains  $\Omega_1$  and  $\Omega_2$  with Lipschitz boundaries. Since the imbeddings  $LD(\Omega_i) \subset L^1(\Omega_i)$ ,  $i = 1, 2$ , are compact, it follows that the imbedding  $LD(\Omega_c) \subset L^1(\Omega_c)$  is also compact. Consequently, by (57) we can suppose that

$$v^k \rightarrow v \quad \text{strongly in } L^1(\Omega_c),$$

and hence, by (58)

$$v^k \rightarrow v \quad \text{strongly in } LD(\Omega_c). \quad (59)$$

In view of (58), we have  $\varepsilon_{ij}(v) = 0$  in  $\Omega_c$ , ie.  $v_i(x) = c_i + b_{ij}x_j$ , where  $c_i, b_{ij} \in \mathbb{R}$ ,  $b_{ij} = -b_{ji}$ ,  $i, j = 1, 2, 3$ . Moreover, the inequality (58) implies that  $v$  satisfies (55), whence  $v \equiv 0$  which contradicts (57), (59). Therefore, (56) is proved.

Consider the following spaces,

$$BD_N(\Omega_c) = \{v \in BD(\Omega_c) \mid v \text{ satisfies (55)}\},$$

$$U_0(\Omega_c) = \{\sigma = \{\sigma_{ij}\} \mid \sigma_{ij} \in H^1(\Omega_c), i, j = 1, 2, 3; \sigma_{ij,j} \in L^3(\Omega_c), i = 1, 2, 3; \\ \sigma_{ij}n_j = 0, i = 1, 2, 3, \text{ on } \Gamma; \sigma_{ij}\nu_j = 0, i = 1, 2, 3, \text{ on } \Sigma_c^\pm\}.$$



Again, to simplify the formulae we assume  $c_{ijkl} = \delta_k^i \delta_l^j$ . Recall that the set (12) contains zero as its interior point. The set  $K$  is introduced in the same way as in section 2,

$$K = \{\sigma = \{\sigma_{ij}\} | \sigma_{ij} \in L^2(\Omega_c), i, j = 1, 2, 3, \Phi(\sigma(x)) \leq 0 \text{ a.e. in } \Omega_c\}.$$

The scalar products in  $L^2(\Omega_c)$  and  $L^2(Q_c)$  are denoted by  $\langle \cdot, \cdot \rangle_c, (\cdot, \cdot)_c$ , respectively. The space of rigid displacements on  $\Omega_c$  is denoted by  $R(\Omega_c)$ .

We assume that there exists a function  $\xi = \{\xi_{ij}\}, i, j = 1, 2, 3, \xi \in C^2(\bar{Q}_c), \xi(0) = \dot{\xi}(0) = 0$ , satisfying the equation

$$(\xi_{ij}, \varepsilon_{ij}(\bar{v}))_c = (f, \bar{v})_c \quad \forall \bar{v} \in L^2(0, T; H^1(\Omega_c)) \quad (60)$$

and such that  $(1 + \kappa)\xi(t) \in K \cap U_0(\Omega_c), \kappa \in \mathbb{R}, \kappa > 0, t \in [0, T]$ . Now we are in a position to prove the theorem of existence of solutions to (49)-(54).

**Theorem 2.** Assume that  $f \in L^3(Q_c), \dot{f} \in L^2(Q_c), f(0) = 0, \langle f(t), \rho \rangle_c = 0 \forall \rho \in R(\Omega_c), t \in [0, T]$ , and the above assumption on  $\xi$  hold. Then there exist functions  $v = (v_1, v_2, v_3), \sigma = \{\sigma_{ij}\}$  such that

$$v \in L^2(0, T; BD_N(\Omega_c)), \quad \sigma, \dot{\sigma} \in L^2(Q_c), \sigma(t) \in K, t \in (0, T),$$

$$(\sigma_{ij}, \varepsilon_{ij}(\bar{v}))_c = (f, \bar{v})_c \quad \forall \bar{v} \in L^2(0, T; H^1(\Omega_c)), \quad (61)$$

$$(\dot{\sigma}_{ij}, \bar{\sigma}_{ij} - \sigma_{ij})_c + (v_i, \bar{\sigma}_{ij,j} - \sigma_{ij,j})_c \geq 0 \quad \forall \bar{\sigma} \in L^2(0, T; U_0(\Omega_c)), \quad (62)$$

$$\bar{\sigma}(t) \in K \quad \text{a.e. in } (0, T),$$

$$\sigma = 0, \quad t = 0. \quad (63)$$

**Proof.** The general scheme of the proof coincides with that used for Theorem 1 and our attention now focusses on details related to the nonsmoothness of the boundary.

Let  $p$  be the penalty operator related to the set  $K, p(\sigma) = \sigma - \pi\sigma, \pi$  is the orthogonal projection operator of the space  $[L^2(\Omega_c)]^9$  onto the set  $K$ .

Consider two positive parameters  $\alpha, \delta$  and auxiliary boundary value problem in  $Q_c$  of the following form. Find  $v = (v_1, v_2, v_3)$  and  $\sigma = \{\sigma_{ij}\}$ , such that

$$\alpha \dot{v}_i - \alpha \varepsilon_{ij}(v)_{,j} - \sigma_{ij,j} = f_i, \quad i = 1, 2, 3, \quad (64)$$

$$\dot{\sigma}_{ij} + \alpha \sigma_{ij} - \varepsilon_{ij}(v) + \frac{1}{\delta} p(\sigma)_{ij} = 0, \quad i, j = 1, 2, 3, \quad (65)$$

$$\sigma_{ij} n_j + \alpha \varepsilon_{ij}(v) n_j = 0, \quad i = 1, 2, 3, \quad \text{on } \Gamma \times (0, T), \quad (66)$$

$$\sigma_{ij} n_j + \alpha \varepsilon_{ij}(v) \nu_j = 0, \quad i = 1, 2, 3, \quad \text{on } \Sigma_c^\pm \times (0, T), \quad (67)$$

$$v = 0, \quad \sigma = 0, \quad t = 0. \quad (68)$$

We first obtain a priori estimates for solutions to (64)–(68). Multiply (64), (65) by  $v_i, \sigma_{ij} - \xi_{ij}$  and integrate over  $\Omega$ . In the same way as for (29) we derive

$$\sup_{0 \leq t \leq T} \|\sigma(t)\|_{0,c}^2 + \alpha \|v\|_{L^2(Q_c)}^2 + \alpha \sum_{i,j=1}^3 \int_0^T \|\varepsilon_{ij}(v)\|_{0,c}^2 d\tau \leq c \quad (69)$$

with a constant  $c$  uniform in  $\alpha, \delta, \alpha \leq \alpha_0$ .

>From (64), (65), (68) we have

$$\dot{v}_i(0) = 0, \quad i = 1, 2, 3, \quad \dot{\sigma}_{ij}(0) = 0, \quad i, j = 1, 2, 3.$$

Hence, the differentiation of (64), (65) with respect to  $t$  and multiplication by  $\dot{v}_i, \dot{\sigma}_{ij} - \dot{\xi}_{ij}$  result in the estimate

$$\|\dot{\sigma}\|_{L^2(Q_c)}^2 + \alpha \|\dot{v}\|_{L^2(Q_c)}^2 \leq c. \quad (70)$$

Moreover, if  $v = v_N + \rho_N$ ,  $v_N \in L^2(0, T; H_N^1(\Omega_c))$ ,  $\rho_N \in L^2(0, T; R(\Omega_c))$ , then (65) provides

$$\|v_N\|_{L^2(0, T; H_N^1(\Omega_c))} \leq c(\delta) \quad (71)$$

where the constant  $c(\delta)$  depends, in general, on  $\delta$ .

The estimates (69)–(70) allow us to prove the solvability of (64)–(68) for the fixed parameters  $\alpha, \delta$  in the following sense

$$\begin{aligned} v^\alpha &\in L^2(0, T; H^1(\Omega_c)), \quad \dot{v}^\alpha, \sigma^\alpha, \dot{\sigma}^\alpha \in L^2(Q_c), \\ \alpha(\dot{v}^\alpha, \bar{v})_c + \alpha(\varepsilon_{ij}(v^\alpha), \varepsilon_{ij}(\bar{v}))_c + (\sigma_{ij}^\alpha, \varepsilon_{ij}(\bar{v}))_c &= (f, \bar{v})_c \quad \forall \bar{v} \in L^2(0, T; H^1(\Omega_c)), \end{aligned} \quad (72)$$

$$\dot{\sigma}_{ij}^\alpha + \alpha \sigma_{ij}^\alpha - \varepsilon_{ij}(v^\alpha) + \frac{1}{\delta} p(\sigma^\alpha)_{ij} = 0, \quad i, j = 1, 2, 3, \quad (73)$$

$$v^\alpha = 0, \quad \sigma^\alpha = 0, \quad t = 0. \quad (74)$$

The solution of the above problem is denoted by  $v^\alpha, \sigma^\alpha$ , and the following step is a passage to the limit as  $\alpha \rightarrow 0$ . Note that boundary conditions (66), (67) are included in the identity (72).

In view of the estimates (69)–(71) for any fixed  $\delta > 0$  one can choose a subsequence  $v^\alpha, \sigma^\alpha$  such that as  $\alpha \rightarrow 0$

$$\begin{aligned} \alpha v^\alpha &\rightarrow 0 \quad \text{weakly in } L^2(0, T; H^1(\Omega_c)), \\ v_N^\alpha &\rightarrow v^\delta \quad \text{weakly in } L^2(0, T; H_N^1(\Omega_c)), \\ \alpha \dot{v}^\alpha &\rightarrow 0 \quad \text{weakly in } L^2(Q_c), \\ \sigma^\alpha, \dot{\sigma}^\alpha &\rightarrow \sigma^\delta, \dot{\sigma}^\delta \quad \text{weakly in } L^2(Q_c). \end{aligned}$$

After the passage to the limit as  $\alpha \rightarrow 0$  we obtain

$$(\sigma_{ij}^\delta, \varepsilon_{ij}(\bar{v}))_c = (f, \bar{v})_c \quad \forall \bar{v} \in L^2(0, T; H^1(\Omega_c)), \quad (75)$$

$$\dot{\sigma}_{ij}^\delta - \varepsilon_{ij}(v^\delta) + \frac{1}{\delta} p(\sigma^\delta)_{ij} = 0, \quad i, j = 1, 2, 3. \quad (76)$$

Analogously to (40) the following estimate holds

$$\|v^\delta\|_{L^2(0, T; BD_N(\Omega_c))} \leq c \quad (77)$$

being uniform in  $\delta$ . Consequently, without any loss of generality we can assume that there exists a subsequence still denoted by  $v^\delta, \sigma^\delta$  such that as  $\delta \rightarrow 0$

$$\begin{aligned} \sigma^\delta, \dot{\sigma}^\delta &\rightarrow \sigma, \dot{\sigma} \quad \text{weakly in } L^2(Q_c), \\ v^\delta &\rightarrow v \quad \text{weakly in } L^2(0, T; L^{3/2}(\Omega_c)), \\ \varepsilon_{ij}(v^\delta) &\rightarrow \varepsilon_{ij}(v) \quad (*)\text{-weakly in } L^2(0, T; \mathcal{M}^1(\Omega_c)), \quad i, j = 1, 2, 3, \\ \sigma^\delta(T) &\rightarrow \sigma(T) \quad \text{weakly in } L^2(\Omega_c). \end{aligned}$$

>From (75) it follows that

$$(\sigma_{ij}, \varepsilon_{ij}(\bar{v}))_c = (f, \bar{v})_c \quad \forall \bar{v} \in L^2(0, T; H^1(\Omega_c)), \quad (78)$$

hence, the equations

$$-\sigma_{ij,j} = f_i, \quad i = 1, 2, 3,$$

are satisfied in  $Q_c$  in the sense of distributions. Also, (75) implies that

$$-\sigma_{ij,j}^\delta = f_i, \quad i = 1, 2, 3. \quad (79)$$

Whence, by (75), (79) we have

$$(\varepsilon_{ij}(v^\delta), \sigma_{ij}^\delta)_c = -(\sigma_{ij,j}^\delta, v_i^\delta)_c = (f_i, v_i^\delta)_c.$$

Moreover,

$$(\varepsilon_{ij}(v^\delta), \bar{\sigma}_{ij})_c = -(v_i^\delta, \bar{\sigma}_{ij,j})_c$$

for all  $\bar{\sigma} \in L^2(0, T; U_0(\Omega_c))$ . As a result, it follows from (76) for any  $\bar{\sigma} \in L^2(0, T; U_0(\Omega_c))$ ,  $\bar{\sigma}(t) \in K$  a. e. in  $(0, T)$ , that

$$(\dot{\sigma}_{ij}^\delta, \bar{\sigma}_{ij} - \sigma_{ij}^\delta)_c + (v_i^\delta, \bar{\sigma}_{ij,j} - \sigma_{ij,j}^\delta)_c \geq 0. \quad (80)$$

The passage to the limit as  $\delta \rightarrow 0$  can be performed in (80) in the same way as in (46). Therefore, we arrive at (62). The property  $\sigma(t) \in K, t \in (0, T)$ , is obtained in the standard way. Boundary conditions (52) are fulfilled in the sense of the space  $H^{-1/2}(\Gamma)$ , and the conditions (53) are satisfied by identity (61) in the weak sense. This completes the proof of Theorem 2.

**Acknowledgement.** A.M. Khludnev was supported by Institut Elie Cartan, Université Henri Poincaré Nancy I during his stay in April 1997 and by the Russian Fund for Basic Research under the grant 97-01-00896.

## References

- [1] Temam, R., *Problèmes mathématiques en plasticité*, Gauthier-Villars, Paris, 1983.
- [2] Khludnev, A.M., Sokolowski, J., *Modelling and Control in Solid Mechanics*, Birkhäuser, Basel, Boston, Berlin, 1997.
- [3] Anzellotti, G., On the existence of the rates of stress and displacement for Prandtl-Reuss plasticity, *Quart. Appl. Math. J.*, **41**, 181 - 208 (1983).
- [4] Anzellotti, G., Giaquinta, M., On the existence of the fields of stresses and displacements for an elasto-perfectly plastic body in static equilibrium, *J. Math. Pure Appl.*, **61**, 219 - 244 (1982).
- [5] Demengel, F., Problèmes variationnels en plasticité parfaite des plaques, *Numer. Func. Anal. and Optim.*, **6**, 73 - 119 (1983).
- [6] Khludnev, A.M., On variational inequalities in contact plastic problems, *Differential Equations*, **24**, 1622 - 1628 (1988). (In Russian.)
- [7] Johnson, C., Existence theorems for plasticity problems, *J. Math. Pures et Appl.*, **55**, 431 - 444 (1976)
- [8] Suquet, P.M., Evolution problems for a class of dissipative materials, *Quart. Appl. Math.*, **38**, 391 - 414 (1981).
- [9] Temam, R. A generalized Norton-Hoff model and the Prandtl-Reuss law of plasticity, *Arch. Rat. Mech. Anal.*, **95**, 137-181 (1986).
- [10] Khludnev, A.M., Contact elastoplastic problem for a beam, *Siberian Math. J.*, **34**, 173 - 179 (1993). (In Russian.)
- [11] Khludnev, A.M., Contact viscoelastoplastic problem for a beam, in: *Free boundary problems in continuum mechanics*, eds. S.N. Antontsev, K.-H. Hoffmann, A.M. Khludnev, Birkhäuser, Basel, 1992, 159 - 166.
- [12] Khludnev, A.M., Contact elastoplastic problem for a rod under the creep condition, in: *Non-linear Math. Problems in Industry, Math. Sciences and Applications*, **1**, 311 - 319 (1993).
- [13] Khludnev, A.M., Hoffmann, K.-H., A variational inequality in a contact elastoplastic problem for a bar, *Advances in Mathematical Sciences and Applications*, **1**, 127 - 136 (1992).
- [14] Lions, J.-L., *Quelques méthodes de résolution des problèmes aux limites non linéaires*, Dunot and Gauthier-Villars, Paris, 1969.
- [15] Oleinik, O.A., Kondratiev, V.A., Kopaček, J., Asymptotic properties of solutions to a biharmonic equation, *Differential Equations*, **17**, 1886 - 1889 (1981). (In Russian.)
- [16] Kondratiev, V.A., Kopaček, J., Oleinik, O.A., On behaviour of solutions to the second order elliptic equations and elasticity equations in a neighbourhood of boundary points, in: *Proceedings of Petrovsky's seminar*, Moscow University Publishers, Moscow, **8**, 135 - 152 (1982). (In Russian.)
- [17] Grisvard, P., *Singularities in Boundary Value Problems*, Springer, 1992.

- [18] Duduchava, R., Wendland, W., The Wiener-Hopf method for system of pseudodifferential equations with applications to crack problems, *Integral Equations and Operator Theory*, **23**, 294 - 335 (1995).
- [19] Nazarov, S.A., Plamenevskii, B.A., *Elliptic problems in domains with piecewise smooth boundaries*, Nauka, Moscow, 1991. (In Russian.)
- [20] Morozov, N.F., *Mathematical foundation of the crack theory*, Nauka, Moscow, 1984. (In Russian.)
- [21] Khludnev, A.M., The contact problem for a shallow shell with a crack, *Prikladnaya Matematika i Mekhanika*, **59**, 318 - 326 (1995). Translated in: *J. Appl. Maths. Mechs.*, **59**, 299 - 306 (1995).
- [22] Khludnev, A.M., On a Signorini problem for inclusions in shells, *Europ. J. Appl. Math.*, **7**, 499-510 (1996).
- [23] Khludnev, A.M., Contact problem for a plate having a crack of minimal opening, *Control and Cybernetics*, **25**, 605 - 620 (1996).
- [24] Nicaise, S., About the Lamé system in a polygonal or polyhedral domain and coupled problem between the Lamé system and the plate equation. 1. Regularity of the solutions, *Annali Scuola Norm. Super. Pisa, serie IV*, **19**, 327 - 361 (1992).



---

Unit e de recherche INRIA Lorraine, Technop ole de Nancy-Brabois, Campus scientifique,  
615 rue du Jardin Botanique, BP 101, 54600 VILLERS L ES NANCY  
Unit e de recherche INRIA Rennes, Irista, Campus universitaire de Beaulieu, 35042 RENNES Cedex  
Unit e de recherche INRIA Rh one-Alpes, 655, avenue de l'Europe, 38330 MONTBONNOT ST MARTIN  
Unit e de recherche INRIA Rocquencourt, Domaine de Voluceau, Rocquencourt, BP 105, 78153 LE CHESNAY Cedex  
Unit e de recherche INRIA Sophia-Antipolis, 2004 route des Lucioles, BP 93, 06902 SOPHIA-ANTIPOLIS Cedex

---

 diteur  
INRIA, Domaine de Voluceau, Rocquencourt, BP 105, 78153 LE CHESNAY Cedex (France)  
ISSN 0249-6399