

The Boundary-Value Problem in Domains with Very Rapidly Oscillating Boundary

Gregory Chechkin, Avner Friedman, Andrey Piatnitski

► **To cite this version:**

Gregory Chechkin, Avner Friedman, Andrey Piatnitski. The Boundary-Value Problem in Domains with Very Rapidly Oscillating Boundary. RR-3062, INRIA. 1996. <inria-00073630>

HAL Id: inria-00073630

<https://hal.inria.fr/inria-00073630>

Submitted on 24 May 2006

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

***The boundary-value problem in domains with
very rapidly oscillating boundary***

Gregory Chechkin

Avner Friedman

Andrey Piatnitski

N° 3062

Décembre 1996

————— THÈME 4 —————

 ***rapport
de recherche***

The boundary-value problem in domains with very rapidly oscillating boundary

Gregory Chechkin^{*} Avner Friedman^{**} Andrey Piatnitski^{***}

Thème 4 — Simulation et optimisation
de systèmes complexes
Projet SYSDYS

Rapport de recherche n° 3062 — Décembre 1996 — 30 pages

Abstract: The asymptotic behavior of solutions to the boundary-value problems posed in domains with very rapidly oscillating locally periodic boundary, is studied. For various kinds of boundary conditions the limiting problems are found and the estimates of the residual are given. Also, we construct an additional term of the asymptotics to improve the estimates of the residual.

Key-words: Boundary homogenization, Rapidly oscillating boundary

(Résumé : tsvp)

The final version of this report was achieved during the stay of Andrey Piatnitski from July to December 1996 in the SYSDYS group. This stay was supported by INRIA (UR de Sophia–Antipolis), the SYSDYS project and the “Direction Générale du Développement Economique de la Ville de Marseille”.

^{*} Department of Differential Equations, Faculty of Mechanics and Mathematics, Moscow State University — Moscow 119899, RUSSIA. E-mail: chechkin@sci.lpi.ac.ru

^{**} Institute for Mathematics and its Applications — 513 Vincent Hall, 206 Church street S.E., Minneapolis, MN 55455, USA. E-mail: friedman@ima.umn.edu

^{***} Lebedev Physical Institute, Russian Academy of Sciences — Leninsky pr., 53, Moscow 117924, RUSSIA. E-mail: andrey@sci.lpi.msk.su

Le problème de conditions au bord dans un domaine à frontière rapidement oscillante

Résumé : Nous nous intéressons au comportement asymptotique des solutions d'un problème de conditions aux bords dans un domaine à frontière localement rapidement oscillante. Pour différents types de conditions aux bords, nous déterminons le problème limite et proposons des estimées des résidus. Nous construisons également un terme asymptotique supplémentaire afin d'améliorer les estimées des résidus.

Mots-clé : Homogénéisation des conditions au bords, frontières rapidement oscillantes

Contents

1	Introduction	5
2	Setting of the problem	6
3	The case $\beta > \alpha - 1$	7
4	The case $\beta = \alpha - 1$	15
5	The case $\beta < \alpha - 1$	18
6	The “smooth” corrector	24
	Bibliography	29

1 Introduction

The paper is devoted to the boundary-value problems in domains with rapidly oscillating boundary. Recent years the interest to this kind of problems appears in connection with the development of technologies of porous, composite and other micro-inhomogeneous materials, and also as a result of various physical experiments. For example, the morphology of contacting surfaces plays the important role in the frictional behavior of deformable bodies. The roughness of the contact surface and the material properties near this surface are micro characteristics which influence the large scale behavior. The mathematical analysis of these problems based on boundary homogenization, was presented in [SS], [BLS], [BLMS], [SPE] and others. Different boundary homogenization problems were considered in [SP], [B1], [B2], [NM], [O], [BMS].

In the present paper the authors study an elliptic problem with the inhomogeneous Fourier boundary condition in domains with very rapidly oscillating locally periodic boundary, depending on a small parameter. The peculiarity of the problems considered is the unlimited growth of $(n-1)$ -dimensional volume of the boundary as the small parameter tends to zero, while in the cases studied earlier [B2], [BLMS], [SPE], the $(n-1)$ -dimensional volume of the oscillating part of the boundary remained uniformly bounded. Thus, in the case under consideration the height of the "cogs" forming the boundary, is much greater than the size of their bases. This leads to the appearance of special normalizing factors both in the coefficients and in the right hand side of boundary operators. Depending on these normalizing factors different limiting boundary conditions can be obtained.

Also, an additional term of the asymptotic expansion which allows to precise the limiting behavior of solution and improve the estimate of the residual, is constructed. This term is defined with the help of a family of auxiliary problems posed in a fixed domain, with non-oscillating boundary conditions. In this connection we call it smooth corrector. Earlier, in different framework the corrector and the asymptotic expansion questions were considered in [B2], [NM].

For simplicity we suppose the oscillating part of the boundary to lie in a small neighborhood of a hyperplane. Without loss of generality one can consider the coordinate hyperplane $\{x \mid x_n = 0\}$.

2 Setting of the problem

Let Ω be a bounded domain in \mathbf{R}^n , $n \geq 2$, with a smooth boundary $\partial\Omega = \Gamma_1 \cup \Gamma_2$, where Ω lies in the half-space $x_n > 0$ and $\Gamma_1 \subset \{x : x_n = 0\}$.

Given smooth non-positive 1-periodic in $\hat{\xi}$ function $F(\hat{x}, \hat{\xi})$, $\hat{x} = (x_1, \dots, x_{n-1})$, $\hat{\xi} = (\xi_1, \dots, \xi_{n-1})$, define the domain Ω_ε as follows: $\partial\Omega_\varepsilon = \Gamma_1^\varepsilon \cup \Gamma_2$, where we set $\Gamma_1^\varepsilon = \{x : \hat{x} \in \Gamma_1, x_n = \varepsilon F(\hat{x}, \frac{\hat{x}}{\varepsilon^\alpha})\}$, $\alpha > 1$. Usually, we will assume $F(\hat{x}, \hat{\xi})$ to be compactly supported on Γ_1 uniformly in $\hat{\xi}$. Consider the following boundary-value problem

$$\begin{cases} -\Delta u_\varepsilon = f(x) & \text{in } \Omega_\varepsilon, \\ \frac{\partial u_\varepsilon}{\partial \nu_\varepsilon} + \varepsilon^\beta p(\hat{x}, \frac{\hat{x}}{\varepsilon^\alpha}) u_\varepsilon = \varepsilon^{\alpha-1} g(\hat{x}, \frac{\hat{x}}{\varepsilon^\alpha}) & \text{on } \Gamma_1^\varepsilon, \\ u_\varepsilon = 0 & \text{on } \Gamma_2, \end{cases} \quad (1)$$

where ν_ε is an outer normal to Γ_1^ε and $p(\hat{x}, \hat{\xi})$, $g(\hat{x}, \hat{\xi})$ are 1-periodic in $\hat{\xi}$.

Definition 1. *The function $u_\varepsilon \in H^1(\Omega_\varepsilon, \Gamma_2)$ is a solution of problem (1) if the following integral identity*

$$\begin{aligned} \int_{\Omega_\varepsilon} \nabla u_\varepsilon(x) \nabla v(x) dx + \varepsilon^\beta \int_{\Gamma_1^\varepsilon} p(\hat{x}, \frac{\hat{x}}{\varepsilon^\alpha}) u_\varepsilon(x) v(x) ds = \\ = \int_{\Omega_\varepsilon} f(x) v(x) dx + \varepsilon^{\alpha-1} \int_{\Gamma_1^\varepsilon} g(\hat{x}, \frac{\hat{x}}{\varepsilon^\alpha}) v(x) ds \end{aligned}$$

holds for all $v \in H^1(\Omega_\varepsilon, \Gamma_2)$. Here we use the standard notation $H^1(\Omega_\varepsilon, \Gamma_2)$ for the closure by the $H^1(\Omega_\varepsilon)$ norm of the set $C^\infty(\overline{\Omega_\varepsilon})$ -functions vanishing in a neighborhood of Γ_2 .

In what follows we study the limiting behavior of solution u_ε of (1) as $\varepsilon \rightarrow 0$. It will be shown that the boundary condition of the limit problem depends on the relation between α and β . In particular, the Dirichlet boundary condition as well as the Neumann and the Fourier boundary conditions can be obtained.

3 The case $\beta > \alpha - 1$

In this section we consider the case $\beta > \alpha - 1$. The main result here is the following theorem.

Theorem 1. *Let $\beta > \alpha - 1$, $f \in L_2(\mathbf{R}^n)$ and $F(\hat{x}, \hat{\xi})$, $g(\hat{x}, \hat{\xi})$, $p(\hat{x}, \hat{\xi})$ be periodic in $\hat{\xi}$ smooth functions. Suppose that $F(\hat{x}, \hat{\xi})$ is compactly supported in $x \in \Gamma_1$ uniformly in ξ . Then, for all sufficiently small ε problem (1) does have unique solution and the inequality holds*

$$\|u_0 - u_\varepsilon\|_{H^1(\Omega_\varepsilon)} \leq K_1 \left(\max(\varepsilon^{\beta-\alpha+1}, \varepsilon^{\alpha-1}, \sqrt{\varepsilon}) \right), \quad (2)$$

where K_1 does not depend on ε , u_0 satisfies the following limiting equation

$$\begin{cases} -\Delta u_0 = f & \text{in } \Omega, \\ \frac{\partial u_0}{\partial x_n} = G(\hat{x}) & \text{on } \Gamma_1, \\ u_0 = 0 & \text{on } \Gamma_2, \end{cases} \quad (3)$$

and

$$G(\hat{x}) = \int_0^1 \int_0^1 \cdots \int_0^1 \sqrt{|\nabla_{\hat{\xi}} F(\hat{x}, \hat{\xi})|^2} g(\hat{x}, \hat{\xi}) d\hat{\xi}.$$

Remark 1. *The definition of solution to problem (3) is analogous to that given for problem (1).*

Denote by V a n -dimensional open neighborhood of $(\text{supp } F)$ such that $\text{dist}((V \cap \Gamma_1), \partial\Gamma_1) > 0$, here ∂ means $(n - 2)$ -dimensional boundary of Γ_1 . It is clear that for sufficiently small ε the oscillating part of Γ_1^ε belongs to V . Due to the smoothness of the boundary $\partial\Omega$ the solution u_0 belongs to $H^2(\Omega \cap V)$ [L], and, hence, can be continued on V to belong to $H^2(V)$ [MV].

Preliminary lemma

Lemma 1. *There exist such constants C_1, C_2 that for any $v \in H^1(\Omega_\varepsilon, \Gamma_2)$ the inequalities*

$$\|v(\hat{x}, \varepsilon F(\hat{x}, \frac{\hat{x}}{\varepsilon^\alpha})) - v(\hat{x}, 0)\|_{L_2(\Gamma_1)} \leq C_1 \sqrt{\varepsilon} \|v\|_{H^1(\Omega_\varepsilon)}, \quad (4)$$

$$\|v\|_{L_2(\Omega_\varepsilon \setminus \Omega)} \leq C_2 \sqrt{\varepsilon} \|v\|_{H^1(\Omega_\varepsilon)} \quad (5)$$

take place.

Proof. Without loss of generality we assume $v \in C^\infty(\overline{\Omega_\varepsilon})$. Note that

$$v(\hat{x}, y) - v(\hat{x}, 0) = \int_0^y \frac{\partial v(\hat{x}, t)}{\partial x_n} dt$$

for any $(\hat{x}, y) \in \overline{\Omega_\varepsilon \setminus \Omega}$. Then, we have

$$|v(\hat{x}, y) - v(\hat{x}, 0)|^2 = \left| \int_0^y \frac{\partial v(\hat{x}, t)}{\partial x_n} dt \right|^2 \leq y \int_0^y \left| \frac{\partial v}{\partial x_n} \right|^2 dt.$$

Substituting $y = \varepsilon F(\hat{x}, \frac{\hat{x}}{\varepsilon^\alpha})$ and integrating over Γ_1 , we obtain (4). Further, for $(\hat{x}, y) \in \overline{\Omega_\varepsilon \setminus \Omega}$ the following inequality

$$\begin{aligned} v^2(\hat{x}, y) &\leq 2v^2(\hat{x}, 0) + 2 \left(\int_0^y \frac{\partial v(\hat{x}, t)}{\partial x_n} dt \right)^2 \leq \\ &\leq 2v^2(\hat{x}, 0) + 2\varepsilon F(\hat{x}, \frac{\hat{x}}{\varepsilon^\alpha}) \int_0^{\varepsilon F(\hat{x}, \frac{\hat{x}}{\varepsilon^\alpha})} \left| \frac{\partial v}{\partial x_n} \right|^2 dt \end{aligned}$$

holds. Consequently,

$$\begin{aligned} \int_{\Gamma_1} d\hat{x} \int_0^{\varepsilon F(\hat{x}, \frac{\hat{x}}{\varepsilon^\alpha})} v^2(\hat{x}, y) dy &\leq 2\varepsilon \max F \|v\|_{L_2(\Gamma_1)}^2 + \\ &+ \varepsilon^2 \max F^2 \int_{\Gamma_1} d\hat{x} \int_0^{\varepsilon F(\hat{x}, \frac{\hat{x}}{\varepsilon^\alpha})} \left| \frac{\partial v}{\partial x_n} \right|^2 dt. \end{aligned}$$

This implies (5). The lemma is proved.

Lemma 2. Let (ds) be an element of $(n-1)$ -dimensional volume of Γ_1^ε . Then

$$\begin{aligned} ds &= \left(\sqrt{1 + \varepsilon^{2-2\alpha} |\nabla_{\hat{\xi}} F(\hat{x}, \hat{\xi})|^2} \Big|_{\hat{\xi} = \frac{\hat{x}}{\varepsilon^\alpha}} \right) d\hat{x} (1 + O(\varepsilon)) = \\ &= \varepsilon^{1-\alpha} \left(\sqrt{|\nabla_{\hat{\xi}} F(\hat{x}, \hat{\xi})|^2} \Big|_{\hat{\xi} = \frac{\hat{x}}{\varepsilon^\alpha}} + O(\varepsilon^{\alpha-1}) \right) d\hat{x}. \end{aligned} \quad (6)$$

Proof. Due to our assumptions Γ_1^ε is defined by the equation

$$x_n - \varepsilon F\left(\hat{x}, \frac{\hat{x}}{\varepsilon^\alpha}\right) = 0.$$

Therefore,

$$ds = \sqrt{(\varepsilon F'_{x_1} + \varepsilon^{1-\alpha} F'_{\xi_1})^2 + \dots + (\varepsilon F'_{x_{n-1}} + \varepsilon^{1-\alpha} F'_{\xi_{n-1}})^2 + 1} d\hat{x},$$

where $\hat{\xi} = \frac{\hat{x}}{\varepsilon^\alpha}$. The direct calculations show that

$$\begin{aligned} &\sqrt{\varepsilon^2 |\nabla_{\hat{x}} F|^2 + 2\varepsilon^{2-\alpha} (\nabla_{\hat{x}} F, \nabla_{\hat{\xi}} F) + \varepsilon^{2-2\alpha} |\nabla_{\hat{\xi}} F|^2 + 1} - \sqrt{1 + \varepsilon^{2-2\alpha} |\nabla_{\hat{\xi}} F|^2} = \\ &= \frac{\varepsilon^2 |\nabla_{\hat{x}} F|^2 + 2\varepsilon^{2-\alpha} (\nabla_{\hat{x}} F, \nabla_{\hat{\xi}} F)}{\sqrt{\varepsilon^2 |\nabla_{\hat{x}} F|^2 + 2\varepsilon^{2-\alpha} (\nabla_{\hat{x}} F, \nabla_{\hat{\xi}} F) + \varepsilon^{2-2\alpha} |\nabla_{\hat{\xi}} F|^2 + 1} + \sqrt{1 + \varepsilon^{2-2\alpha} |\nabla_{\hat{\xi}} F|^2}} = O(\varepsilon) \end{aligned}$$

and

$$\begin{aligned} &\sqrt{\varepsilon^2 |\nabla_{\hat{x}} F|^2 + 2\varepsilon^{2-\alpha} (\nabla_{\hat{x}} F, \nabla_{\hat{\xi}} F) + \varepsilon^{2-2\alpha} |\nabla_{\hat{\xi}} F|^2 + 1} - \varepsilon^{1-\alpha} \sqrt{|\nabla_{\hat{\xi}} F(\hat{x}, \hat{\xi})|^2} = \\ &= \frac{\varepsilon^2 |\nabla_{\hat{x}} F|^2 + 2\varepsilon^{2-\alpha} (\nabla_{\hat{x}} F, \nabla_{\hat{\xi}} F) + 1}{\sqrt{\varepsilon^2 |\nabla_{\hat{x}} F|^2 + 2\varepsilon^{2-\alpha} (\nabla_{\hat{x}} F, \nabla_{\hat{\xi}} F) + \varepsilon^{2-2\alpha} |\nabla_{\hat{\xi}} F|^2 + 1} + \varepsilon^{1-\alpha} \sqrt{|\nabla_{\hat{\xi}} F(\hat{x}, \hat{\xi})|^2}} \leq \\ &\leq C(1 + O(\varepsilon)) = \varepsilon^{1-\alpha} O(\varepsilon^{\alpha-1}), \end{aligned}$$

where the constant C is independent of ε .

These inequalities imply (6). The lemma is proved.

Proposition 3. *Uniformly in $u, v \in H^{\frac{1}{2}}(\Gamma_1)$*

$$\left| \int_{\Gamma_1} uv \, d\hat{x} \right| \leq C_3 \|u\|_{H^{\frac{1}{2}}(\Gamma_1)} \|v\|_{H^{\frac{1}{2}}(\Gamma_1)}. \quad (7)$$

Proof. This inequality is a direct consequence of the Cauchy-Schwartz-Bunyakovskii inequality and the compactness of embedding $L_2(\Gamma_1)$ in $H^{\frac{1}{2}}(\Gamma_1)$.

Lemma 4. *There exists such a positive constant M , independent of ε , that*

$$\int_{\Omega_\varepsilon} |\nabla v|^2 dx + \varepsilon^\beta \int_{\Gamma_1^\varepsilon} p\left(\hat{x}, \frac{\hat{x}}{\varepsilon^\alpha}\right) v^2 ds \geq M \|v\|_{H^1(\Omega_\varepsilon)}^2$$

for any $v \in H^1(\Omega_\varepsilon)$.

Proof. In view of (4) and Lemma 2

$$\begin{aligned} \varepsilon^\beta \int_{\Gamma_1^\varepsilon} p\left(\hat{x}, \frac{\hat{x}}{\varepsilon^\alpha}\right) v^2 \left(x, \varepsilon F\left(\hat{x}, \frac{\hat{x}}{\varepsilon^\alpha}\right)\right) ds &\leq c \varepsilon^\beta \varepsilon^{\alpha-1} \int_{\Gamma_1^\varepsilon} v^2 \left(x, \varepsilon F\left(\hat{x}, \frac{\hat{x}}{\varepsilon^\alpha}\right)\right) d\hat{x} \leq \\ &\leq c' \left(\varepsilon^{\beta-\alpha+1} \|v\|_{L_2(\Gamma_1)}^2 + \varepsilon^{\beta-\alpha+3/2} \|v\|_{H^1(\Omega_\varepsilon)}^2 \right) \leq c'' \varepsilon^{\beta-\alpha+1} \|v\|_{H^1(\Omega_\varepsilon)}^2 \end{aligned}$$

Then, the assertion of the lemma is a consequence of the Friedrichs type inequality (see [M]).

Lemma 5. *Let $h(\hat{x}, \hat{\xi})$ be 1-periodic in ξ Lipschitz function such that*

$$\int_0^1 \int_0^1 \cdots \int_0^1 h(\hat{x}, \hat{\xi}) \, d\hat{\xi} \equiv 0. \quad (8)$$

Then, an inequality

$$\left| \int_{\Gamma_1} h\left(\hat{x}, \frac{\hat{x}}{\varepsilon^\alpha}\right) u(\hat{x}) v(\hat{x}) \, d\hat{x} \right| \leq C_4 \sqrt{\varepsilon^\alpha} \|u\|_{H^{\frac{1}{2}}(\Gamma_1)} \|v\|_{H^{\frac{1}{2}}(\Gamma_1)} \quad (9)$$

is satisfied, C_4 depending only on the Lipschitz constant of h .

Remark 2. *This statement for smooth $h(\hat{x}, \frac{\hat{x}}{\varepsilon^\alpha})$ and $n = 2$ was proved in [B2] and, independently, in [Br]. See also [BM]. Here we use the technique developed in [B2].*

Proof. In view of (8) an equation

$$\Delta_{\hat{\xi}} \Psi(\hat{x}, \hat{\xi}) = h(\hat{x}, \hat{\xi})$$

is solvable in the space of 1-periodic in $\hat{\xi}$ functions. The solution is unique up to an additive constant.

Suppose, initially, that $v(\hat{x}, 0)$ and $u(\hat{x}, 0)$ belong to $\overset{\circ}{H}^1(\Gamma_1)$ and $h(\hat{x}, \hat{\xi})$ is C^1 -function. Then, substituting $h(\hat{x}, \hat{\xi}) = \Delta_{\hat{\xi}} \Psi(\hat{x}, \hat{\xi})$, integrating by parts, and using the Cauchy-Schwartz-Bunyakovskii inequality, we have

$$\begin{aligned} \left| \int_{\Gamma_1} h(\hat{x}, \frac{\hat{x}}{\varepsilon^\alpha}) u(\hat{x}) v(\hat{x}) d\hat{x} \right| &= \left| \int_{\Gamma_1} (\Delta_{\hat{\xi}} \Psi(\hat{x}, \hat{\xi})) \Big|_{\hat{\xi}=\frac{\hat{x}}{\varepsilon^\alpha}} u(\hat{x}) v(\hat{x}) d\hat{x} \right| = \\ &= \left| \int_{\Gamma_1} \left(\varepsilon^\alpha \nabla_{\hat{x}} \left[\nabla_{\hat{\xi}} \Psi(\hat{x}, \hat{\xi}) \Big|_{\hat{\xi}=\frac{\hat{x}}{\varepsilon^\alpha}} \right] - \varepsilon^\alpha \left((\nabla_{\hat{x}}, \nabla_{\hat{\xi}}) \Psi(\hat{x}, \hat{\xi}) \Big|_{\hat{\xi}=\frac{\hat{x}}{\varepsilon^\alpha}} \right) \right| \leq \quad (10) \\ &\leq \varepsilon^\alpha \left| \int_{\Gamma_1} (\nabla_{\hat{\xi}} \Psi(\hat{x}, \hat{\xi}) \Big|_{\hat{\xi}=\frac{\hat{x}}{\varepsilon^\alpha}}, \nabla_{\hat{x}}(uv)) d\hat{x} \right| + \varepsilon^\alpha \left| \int_{\Gamma_1} \left((\nabla_{\hat{x}}, \nabla_{\hat{\xi}}) \Psi(\hat{x}, \hat{\xi}) \Big|_{\hat{\xi}=\frac{\hat{x}}{\varepsilon^\alpha}} uv d\hat{x} \right| \leq \\ &\leq C_6 \varepsilon^\alpha \|u\|_{H^1(\Gamma_1)} \|v\|_{H^1(\Gamma_1)}. \end{aligned}$$

Now, approximating arbitrary Lipschitz function $h(\hat{x}, \hat{\xi})$ by a sequence of C^1 -functions we extend the last inequality to the space of Lipschitz functions. Clearly, C_6 depends only on Lipschitz constant.

Let $p \in [0, 1]$. On the base of a bilinear form

$$\int_{\Gamma_1} h(\hat{x}, \frac{\hat{x}}{\varepsilon^\alpha}) u(\hat{x}) v(\hat{x}) d\hat{x}, \quad u, v \in H^p(\Gamma_1)$$

we define an operator $T_\varepsilon : H^p(\Gamma_1) \rightarrow H^{-p}(\Gamma_1)$ by the formula

$$\langle T_\varepsilon u, v \rangle_p = \int_{\Gamma_1} h(\hat{x}, \frac{\hat{x}}{\varepsilon^\alpha}) u(\hat{x}, 0) v(\hat{x}, 0) d\hat{x}.$$

From (10) it follows

$$\|T_\varepsilon\|_{L(H^1(\partial\Omega), H^{-1}(\partial\Omega))} \leq C_6 \varepsilon^\alpha.$$

Then, clearly

$$\|T_\varepsilon\|_{L(L_2(\partial\Omega), L_2(\partial\Omega))} \leq C_7$$

with the constant, independent of ε .

Using the space interpolation technique ([LM], [RS]) we derive from the above estimates

$$\|T_\varepsilon\|_{L(H^p(\partial\Omega), H^{-p}(\partial\Omega))} \leq C_7^{1-p} C_6^p \varepsilon^{p\alpha}. \quad (11)$$

Setting $p = \frac{1}{2}$ in (11), we obtain (9). The lemma is proved.

Lemma 6. *There exists $C_8 > 0$, such that for all $v \in H^1(\Omega_\varepsilon)$*

$$\left| \int_{\Gamma_1^\varepsilon} \varepsilon^{\alpha-1} g\left(\hat{x}, \frac{\hat{x}}{\varepsilon^\alpha}\right) v\left(\hat{x}, \varepsilon F\left(\hat{x}, \frac{\hat{x}}{\varepsilon^\alpha}\right)\right) ds - \int_{\Gamma_1} G(\hat{x}) v(\hat{x}, 0) d\hat{x} \right| \leq \quad (12)$$

$$C_8 (\varepsilon^{\alpha-1} + \sqrt{\varepsilon}) \|v\|_{H^1(\Omega_\varepsilon)}.$$

Proof. According to Lemma 2 we have

$$\begin{aligned} & \left| \int_{\Gamma_1^\varepsilon} \varepsilon^{\alpha-1} g\left(\hat{x}, \frac{\hat{x}}{\varepsilon^\alpha}\right) v\left(\hat{x}, \varepsilon F\left(\hat{x}, \frac{\hat{x}}{\varepsilon^\alpha}\right)\right) ds - \int_{\Gamma_1} G(\hat{x}) v(\hat{x}, 0) d\hat{x} \right| = \\ & = \left| \int_{\Gamma_1} \varepsilon^{\alpha-1} g\left(\hat{x}, \frac{\hat{x}}{\varepsilon^\alpha}\right) v\left(\hat{x}, \varepsilon F\left(\hat{x}, \frac{\hat{x}}{\varepsilon^\alpha}\right)\right) \varepsilon^{1-\alpha} \left(\sqrt{|\nabla_{\hat{\xi}} F\left(\hat{x}, \frac{\hat{x}}{\varepsilon^\alpha}\right)|^2} + O(\varepsilon^{\alpha-1}) \right) d\hat{x} - \right. \\ & \left. \int_{\Gamma_1} G(\hat{x}) v(\hat{x}, 0) d\hat{x} \right| \leq \left| \int_{\Gamma_1} \left(g\left(\hat{x}, \frac{\hat{x}}{\varepsilon^\alpha}\right) v\left(\hat{x}, \varepsilon F\left(\hat{x}, \frac{\hat{x}}{\varepsilon^\alpha}\right)\right) \sqrt{|\nabla_{\hat{\xi}} F\left(\hat{x}, \frac{\hat{x}}{\varepsilon^\alpha}\right)|^2} - \right. \quad (13) \\ & \quad \left. - g\left(\hat{x}, \frac{\hat{x}}{\varepsilon^\alpha}\right) v(\hat{x}, 0) \sqrt{|\nabla_{\hat{\xi}} F\left(\hat{x}, \frac{\hat{x}}{\varepsilon^\alpha}\right)|^2} \right) d\hat{x} \right| + \\ & + \left| \int_{\Gamma_1} \left(g\left(\hat{x}, \frac{\hat{x}}{\varepsilon^\alpha}\right) v(\hat{x}, 0) \sqrt{|\nabla_{\hat{\xi}} F\left(\hat{x}, \frac{\hat{x}}{\varepsilon^\alpha}\right)|^2} - G(\hat{x}) v(\hat{x}, 0) \right) d\hat{x} \right| + C_9 \varepsilon^{\alpha-1} \|v\|_{H^1(\Omega_\varepsilon)}. \end{aligned}$$

Set $h(\hat{x}, \hat{\xi}) = \sqrt{|\nabla_{\hat{\xi}} F(\hat{x}, \hat{\xi})|^2} g(\hat{x}, \hat{\xi}) - G(\hat{x})$. Then, by the definition of $G(\hat{x})$

$$\int_0^1 \int_0^1 \cdots \int_0^1 h(\hat{x}, \hat{\xi}) \, d\xi \equiv 0.$$

Hence, the inequality (12) follows if we estimate the first integral in the right-hand side by means of Lemma 1 and the second one by means of Lemma 5.

Proof of Theorem 1

Due to Lemma 4 the existence and the uniqueness of solution to problem (1) can be obtained on the base of the Lax-Milgram lemma [Y].

Then, after simple transformations we find

$$\begin{aligned} & \int_{\Omega_\varepsilon} \nabla(u_0 - u_\varepsilon) \nabla v \, dx + \varepsilon^\beta \int_{\Gamma_1^\varepsilon} p(u_0 - u_\varepsilon) v \, ds = \int_{\Omega_\varepsilon} \nabla u_0 \nabla v \, dx - \\ & - \int_{\Omega_\varepsilon} f v \, dx - \varepsilon^{\alpha-1} \int_{\Gamma_1^\varepsilon} g v \, ds + \varepsilon^\beta \int_{\Gamma_1^\varepsilon} p u_0 v \, ds = \int_{\Omega} \nabla u_0 \nabla v \, dx - \int_{\Omega_\varepsilon} f v \, dx - \\ & - \varepsilon^{\alpha-1} \int_{\Gamma_1^\varepsilon} g v \, ds + \int_{\Omega_\varepsilon \setminus \Omega} \nabla u_0 \nabla v \, dx + \varepsilon^\beta \int_{\Gamma_1^\varepsilon} p u_0 v \, ds = \int_{\Omega_\varepsilon \setminus \Omega} \nabla u_0 \nabla v \, dx - \\ & - \varepsilon^{\alpha-1} \int_{\Gamma_1^\varepsilon} g v \, ds + \int_{\Gamma_1} G v d\hat{x} - \int_{\Omega_\varepsilon \setminus \Omega} f v \, dx + \varepsilon^\beta \int_{\Gamma_1^\varepsilon} p u_0 v \, ds. \end{aligned} \quad (14)$$

According to Lemma 2 and Proposition 3 the last integral in the right-hand side of (14) is estimated as follows

$$\begin{aligned} & \varepsilon^\beta \left| \int_{\Gamma_1^\varepsilon} p u_0 v \, ds \right| = \varepsilon^\beta \left| \int_{\Gamma_1} p u_0 v \left[\varepsilon^{1-\alpha} \left(\sqrt{|\nabla_{\hat{\xi}} F(\hat{x}, \frac{\hat{x}}{\varepsilon^\alpha})|^2} + O(\varepsilon^{\alpha-1}) \right) \right] d\hat{x} \right| \leq \\ & \leq \varepsilon^{\beta-\alpha+1} C_{10} \left| \int_{\Gamma_1} p u_0 v \, d\hat{x} \right| \leq \varepsilon^{\beta-\alpha+1} C_{10} \|v\|_{H^{\frac{1}{2}}(\Gamma_1)} \leq \varepsilon^{\beta-\alpha+1} C_{11} \|v\|_{H^1(\Omega_\varepsilon)}. \end{aligned}$$

By (5) considering the uniform boundedness of $\|u_0\|_{H^2(\Omega_\varepsilon)}$, we have

$$\left| \int_{\Omega_\varepsilon \setminus \Omega} \nabla u_0 \nabla v \, dx \right| \leq \|\nabla u_0\|_{L_2(\Omega_\varepsilon \setminus \Omega)} \|v\|_{H^1(\Omega_\varepsilon)} \leq C_2 \sqrt{\varepsilon} \|u_0\|_{H^2(\Omega_\varepsilon)} \|v\|_{H^1(\Omega_\varepsilon)}$$

and

$$\left| \int_{\Omega_\varepsilon \setminus \Omega} f v \, dx \right| \leq \|f\|_{L_2(\Omega_\varepsilon \setminus \Omega)} \|v\|_{L_2(\Omega_\varepsilon \setminus \Omega)} \leq C_2 \sqrt{\varepsilon} \|f\|_{L_2(\Omega_\varepsilon)} \|v\|_{H^1(\Omega_\varepsilon)}.$$

Then, Lemma 6 implies

$$\left| \varepsilon^{\alpha-1} \int_{\Gamma_1^\varepsilon} g v \, ds - \int_{\Gamma_1} G v \, d\hat{x} \right| \leq C_8 (\varepsilon^{\alpha-1} + \sqrt{\varepsilon}) \|v\|_{H^1(\Omega_\varepsilon)}.$$

Combining these inequalities with (14) we deduce

$$\left| \int_{\Omega_\varepsilon} \nabla(u_0 - u_\varepsilon) \nabla v \, dx + \varepsilon^\beta \int_{\Gamma_1^\varepsilon} p(u_0 - u_\varepsilon) v \, ds \right| \leq K_1 \|v\|_{H^1(\Omega_\varepsilon)} (\sqrt{\varepsilon} + \varepsilon^{\alpha-1} + \varepsilon^{\beta-\alpha+1}).$$

It remains to substitute $v = u_0 - u_\varepsilon$. Then, (2) follows from Lemma 4 and the Friedrichs type inequality (see, for example, [M]). The theorem is proved.

4 The case $\beta = \alpha - 1$

In the section we study problem (1) in the case $\beta = \alpha - 1$. We show that the limit problem has an inhomogeneous Fourier boundary condition.

Theorem 2. *Let $\beta = \alpha - 1$, f belong to $L_2(\mathbf{R}^n)$ and $F(\hat{x}, \hat{\xi})$, $p(\hat{x}, \hat{\xi})$, $g(\hat{x}, \hat{\xi})$ be periodic in ξ smooth functions. Suppose that $F(\hat{x}, \hat{\xi})$ is compactly supported in $x \in \Gamma_1$ uniformly in ξ and that the function*

$$P(\hat{x}) = \int_0^1 \int_0^1 \cdots \int_0^1 \sqrt{|\nabla_{\hat{\xi}} F(\hat{x}, \hat{\xi})|^2} p(\hat{x}, \hat{\xi}) d\hat{\xi}$$

is nonnegative. Then, for all sufficiently small $\varepsilon > 0$ the existence and the uniqueness of solution to problem (1) follow and

$$\|u_0 - u_\varepsilon\|_{H^1(\Omega_\varepsilon)} \leq K_2 (\sqrt{\varepsilon} + \varepsilon^{\alpha-1}), \quad (15)$$

where K_2 does not depend on ε , $u_0(x)$ satisfies the following limiting equation

$$\begin{cases} -\Delta u_0 = f & \text{in } \Omega, \\ \frac{\partial u_0}{\partial x_n} + P(\hat{x})u_0 = G(\hat{x}) & \text{on } \Gamma_1, \\ u_0 = 0 & \text{on } \Gamma_2, \end{cases} \quad (16)$$

and $G(\hat{x})$ was defined in Theorem 1.

Remark 3. *The definition of solution to problem (16) is analogous to that given for problem (1). Problem (16) is well-posed because of the assumptions of the theorem.*

Due to the smoothness of the boundary $\partial\Omega$ the solution u_0 belongs to $H^2(\Omega \cap V)$ [L], and, hence, can be continued on V to belong to $H^2(V)$ [MV].

Lemma 7. *There exists $C_{12} > 0$, such that*

$$\begin{aligned} & \left| \int_{\Gamma_1^\varepsilon} \varepsilon^{\alpha-1} p\left(\hat{x}, \frac{\hat{x}}{\varepsilon^\alpha}\right) u_0\left(\hat{x}, \varepsilon F\left(\hat{x}, \frac{\hat{x}}{\varepsilon^\alpha}\right)\right) v\left(\hat{x}, \varepsilon F\left(\hat{x}, \frac{\hat{x}}{\varepsilon^\alpha}\right)\right) ds - \int_{\Gamma_1} P(\hat{x}) u_0(\hat{x}, 0) v(\hat{x}, 0) d\hat{x} \right| \leq \\ & \leq C_{12} (\varepsilon^{\alpha-1} + \sqrt{\varepsilon}) \|u_0\|_{H^1(\Omega_\varepsilon)} \|v\|_{H^1(\Omega_\varepsilon)}. \end{aligned} \quad (17)$$

for all $u, v \in H^1(\Omega_\varepsilon)$.

The proof is similar to that of Lemma 6.

Proof of Theorem 2. The existence and uniqueness of u_ε are due to the positiveness of $P(\hat{x})$, Lemma 7 and the Lax-Milgram lemma. Then, according to (1) and (16)

$$\begin{aligned}
& \int_{\Omega_\varepsilon} \nabla(u_0 - u_\varepsilon) \nabla v \, dx + \varepsilon^{\alpha-1} \int_{\Gamma_1^\varepsilon} p(u_0 - u_\varepsilon) v \, ds = \\
& = \int_{\Omega_\varepsilon} \nabla u_0 \nabla v \, dx - \int_{\Omega_\varepsilon} f v \, dx - \varepsilon^{\alpha-1} \int_{\Gamma_1^\varepsilon} g v \, ds + \varepsilon^{\alpha-1} \int_{\Gamma_1^\varepsilon} p u_0 v \, ds = \\
& = \int_{\Omega} \nabla u_0 \nabla v \, dx - \int_{\Omega_\varepsilon} f v \, dx - \varepsilon^{\alpha-1} \int_{\Gamma_1^\varepsilon} g v \, ds + \int_{\Omega_\varepsilon \setminus \Omega} \nabla u_0 \nabla v \, dx + \varepsilon^{\alpha-1} \int_{\Gamma_1^\varepsilon} p u_0 v \, ds = \\
& = \int_{\Omega_\varepsilon \setminus \Omega} \nabla u_0 \nabla v \, dx - \varepsilon^{\alpha-1} \int_{\Gamma_1^\varepsilon} g v \, ds + \int_{\Gamma_1} G(\hat{x}) v \, d\hat{x} - \int_{\Omega_\varepsilon \setminus \Omega} f v \, dx + \\
& \quad + \varepsilon^{\alpha-1} \int_{\Gamma_1^\varepsilon} p u_0 v \, ds - \int_{\Gamma_1} P(\hat{x}) u_0 v \, d\hat{x}.
\end{aligned}$$

Let us estimate all the terms in the right hand side of the last relation. By (5) considering the smoothness of u_0 , we have

$$\left| \int_{\Omega_\varepsilon \setminus \Omega} \nabla u_0 \nabla v \, dx \right| \leq \|\nabla u_0\|_{L_2(\Omega_\varepsilon \setminus \Omega)} \|v\|_{H^1(\Omega_\varepsilon)} \leq C_2 \sqrt{\varepsilon} \|u_0\|_{H^2(\Omega_\varepsilon)} \|v\|_{H^1(\Omega_\varepsilon)}$$

and

$$\left| \int_{\Omega_\varepsilon \setminus \Omega} f v \, dx \right| \leq \|f\|_{L_2(\Omega_\varepsilon \setminus \Omega)} \|v\|_{L_2(\Omega_\varepsilon \setminus \Omega)} \leq C_2 \sqrt{\varepsilon} \|f\|_{L_2(\Omega_\varepsilon)} \|v\|_{H^1(\Omega_\varepsilon)}.$$

Then, according to Lemma 6 and 7 the inequalities

$$\left| \varepsilon^{\alpha-1} \int_{\Gamma_1^\varepsilon} g v \, ds - \int_{\Gamma_1} G v \, d\hat{x} \right| \leq C_8 (\varepsilon^{\alpha-1} + \sqrt{\varepsilon}) \|v\|_{H^1(\Omega_\varepsilon)}$$

and

$$\left| \varepsilon^{\alpha-1} \int_{\Gamma_1^\varepsilon} p u_0 v \, ds - \int_{\Gamma_1} P u_0 v \, d\hat{x} \right| \leq C_{12} (\varepsilon^{\alpha-1} + \sqrt{\varepsilon}) \|u_0\|_{H^1(\Omega_\varepsilon)} \|v\|_{H^1(\Omega_\varepsilon)}$$

hold. With the help of these inequalities we obtain

$$\left| \int_{\Omega_\varepsilon} \nabla(u_0 - u_\varepsilon) \nabla v \, dx + \int_{\Gamma_1^\varepsilon} p(u_0 - u_\varepsilon) v \, ds \right| \leq K_2 \|v\|_{H^1(\Omega_\varepsilon)} (\sqrt{\varepsilon} + \varepsilon^{\alpha-1}).$$

Substituting $v = u_0 - u_\varepsilon$ and using Lemma 7 and the Friedrichs type inequality (see [M]), we obtain (15). The theorem is proved.

5 The case $\beta < \alpha - 1$

This section is devoted to the case of the Dirichlet limiting problem. Here we suppose the function $P(\hat{x})$ to be uniformly positive on Γ_1 . This means, in particular, that $F(\hat{x}, \hat{\xi})$ is not compactly supported in $x \in \Gamma_1$.

Theorem 3. *Let $\beta < \alpha - 1$, f belong to $L_2(\mathbf{R}^n)$ and $F(\hat{x}, \hat{\xi})$, $p(\hat{x}, \hat{\xi})$, $g(\hat{x}, \hat{\xi})$ be periodic in ξ smooth functions. Assume $P(\hat{x}) \geq C_0 > 0$ on Γ_1 and, also, at least one of the following conditions: $p(\hat{x}, \hat{\xi}) \geq 0$ or $\beta > \alpha - 2$ to be satisfied. Then, for each sufficiently small $\varepsilon > 0$ problem (1) does have unique solution, the family of solutions is uniformly bounded in $H^1(\Omega_\varepsilon)$ -norm and an estimate*

$$\|u_\varepsilon - u_0\|_{L^2(\Omega_\varepsilon)} \leq K_3(\varepsilon^{\alpha-\beta-1} + \sqrt{\varepsilon})^{1/2} \quad (18)$$

holds, where u_0 satisfies the following Dirichlet problem

$$\begin{cases} -\Delta u_0 = f & \text{in } \Omega, \\ u_0 = 0 & \text{on } \partial\Omega. \end{cases} \quad (19)$$

If $\beta > \alpha - 2$ then, in addition,

$$\|u_\varepsilon - u_0\|_{H^1(\Omega_\varepsilon)} \leq K_3 \left(\varepsilon^{\frac{\beta-\alpha+2}{2}} + \varepsilon^{\frac{\alpha-\beta-1}{2}} + \varepsilon^{\frac{1}{4}} \right)^{1/2}, \quad (20)$$

In both inequalities above the constant K_3 is independent of ε .

Remark 4. *In general one has only weak convergence of u_ε in $H^1(\Omega_\varepsilon)$.*

Due to the smoothness of the boundary $\partial\Omega$ the solution u_0 belongs to $H^2(\Omega)$ [L], and, hence, can be continued on \mathbf{R}^n to belong to $H^2(\mathbf{R}^n)$ [MV].

Proof of Theorem 3. In case of positive $p(\hat{x}, \hat{\xi})$ the existence, uniqueness and the uniform boundedness of u_ε follow directly from standard energy estimates and Lax-Milgram lemma (see [Mi]). Otherwise, they are based on the following statement. Denote

$$P_1(\varepsilon, \hat{x}) = \int_0^1 \int_0^1 \dots \int_0^1 \sqrt{\varepsilon^{2\alpha-2} + |\nabla_{\hat{\xi}} F(\hat{x}, \hat{\xi})|^2} p(\hat{x}, \hat{\xi}) d\hat{\xi}.$$

Lemma 8. For all $u, v \in H^1(\Omega_\varepsilon)$

$$\begin{aligned} & \left| \int_{\Gamma_1} P_1(\varepsilon, \hat{x}) u(x) v(x) d\hat{x} - \varepsilon^{\alpha-1} \int_{\Gamma_1^\varepsilon} p(\hat{x}, \frac{\hat{x}}{\varepsilon^\alpha}) u(x) v(x) ds \right| \leq \\ & \leq C_1 \sqrt{\varepsilon} (\|u\|_{H^1(\Omega_\varepsilon)} \|v\|_{L_2(\Gamma_1)} + \|u\|_{L_2(\Gamma_1)} \|v\|_{H^1(\Omega_\varepsilon)}) + \varepsilon \|u\|_{H^1(\Omega_\varepsilon)} \|v\|_{H^1(\Omega_\varepsilon)} \leq \\ & \leq C_1 \sqrt{\varepsilon} \|u\|_{H^1(\Omega_\varepsilon)} \|v\|_{H^1(\Omega_\varepsilon)} \end{aligned}$$

where C_1 does not depend on ε .

Proof. The lemma can be proved in the same way as Lemma 6.

Now, let us transform the integral identity for the solution of problem (1) as follows

$$\begin{aligned} & \int_{\Omega_\varepsilon} \nabla u_\varepsilon(x) \nabla v(x) dx + \varepsilon^{\beta-\alpha+1} \int_{\Gamma_1} P_1(\varepsilon, \hat{x}) u_\varepsilon(x) v(x) d\hat{x} = \\ & = \int_{\Omega_\varepsilon} f(x) v(x) dx + \left(\varepsilon^{\alpha-1} \int_{\Gamma_1^\varepsilon} g(\hat{x}, \frac{\hat{x}}{\varepsilon^\alpha}) v(x) ds - \int_{\Gamma_1} G(\hat{x}) v(x) d\hat{x} \right) + \quad (21) \\ & + \int_{\Gamma_1} G(\hat{x}) v(x) d\hat{x} + \left(\varepsilon^{\beta-\alpha+1} \int_{\Gamma_1} P_1(\varepsilon, \hat{x}) u_\varepsilon(x) v(x) d\hat{x} - \varepsilon^\beta \int_{\Gamma_1^\varepsilon} p(\hat{x}, \frac{\hat{x}}{\varepsilon^\alpha}) u_\varepsilon(x) v(x) ds \right) \end{aligned}$$

Substituting here $v = u_\varepsilon$ and estimating all the terms in the right hand side with the help of Lemma 6, Lemma 8 and Cauchy-Schwartz-Bunyakovskii inequality we find

$$\begin{aligned} & \int_{\Omega_\varepsilon} |\nabla u_\varepsilon(x)|^2 dx + \varepsilon^{\beta-\alpha+1} \int_{\Gamma_1} P_1(\varepsilon, \hat{x}) u_\varepsilon^2(x) d\hat{x} \leq \\ & \leq \|f\|_{L_2(\Omega_\varepsilon)} \|u_\varepsilon\|_{L_2(\Omega_\varepsilon)} + C \left((\varepsilon^{\alpha-1} + \sqrt{\varepsilon}) \|u_\varepsilon\|_{H^1(\Omega_\varepsilon)} + \|u_\varepsilon\|_{H^1(\Omega_\varepsilon)} + \right. \quad (22) \\ & \left. \varepsilon^{\beta-\alpha+3/2} \|u_\varepsilon\|_{H^1(\Omega_\varepsilon)} \|u_\varepsilon\|_{L_2(\Gamma_1)} + \varepsilon^{\beta-\alpha+2} \|u_\varepsilon\|_{H^1(\Omega_\varepsilon)}^2 \right). \end{aligned}$$

Next, considering an evident estimate $|P_1(\varepsilon, \hat{x}) - P(\hat{x})| \leq C\varepsilon^{\alpha-1}$ it is easy to see that under conditions of the theorem

$$P_1(\varepsilon, \hat{x}) \geq C'_0 > 0. \quad (23)$$

Thus, under the assumption $\beta > \alpha - 2$ we have for all sufficiently small ε

$$\begin{aligned} C(\varepsilon^{\beta-\alpha+3/2} \|u_\varepsilon\|_{H^1(\Omega_\varepsilon)} \|u_\varepsilon\|_{L_2(\Gamma_1)} + \varepsilon^{\beta-\alpha+2} \|u_\varepsilon\|_{H^1(\Omega_\varepsilon)}^2) &\leq \\ &\leq \frac{1}{2} \int_{\Omega_\varepsilon} |\nabla u_\varepsilon(x)|^2 dx + \varepsilon^{\beta-\alpha+1} \int_{\Gamma_1} P_1(\varepsilon, \hat{x}) u_\varepsilon^2(x) d\hat{x}; \end{aligned}$$

the constant C here is taken from (22). Then, the existence and the uniqueness of solution to problem (1) as well as its uniform boundedness in $H^1(\Omega_\varepsilon)$ -norm follow from (21), (22) and (23) by standard arguments [Mi].

In order to prove (18) let us divide (22) by $\varepsilon^{\beta-\alpha+1}$:

$$\begin{aligned} \varepsilon^{\alpha-\beta-1} \int_{\Omega_\varepsilon} |\nabla u_\varepsilon(x)|^2 dx + \int_{\Gamma_1} P_1(\hat{x}) u_\varepsilon^2(x) d\hat{x} &\leq \\ &\leq \varepsilon^{\alpha-\beta-1} \|f\|_{L_2(\Omega_\varepsilon)} \|u_\varepsilon\|_{L_2(\Omega_\varepsilon)} + C \left((\varepsilon^{2\alpha-\beta-2} + \varepsilon^{\alpha-\beta-1/2}) \|u_\varepsilon\|_{H^1(\Omega_\varepsilon)} + \right. \\ &\quad \left. + \varepsilon^{\alpha-\beta-1} \|u_\varepsilon\|_{H^1(\Omega_\varepsilon)} + \sqrt{\varepsilon} \|u_\varepsilon\|_{H^1(\Omega_\varepsilon)}^2 \right). \end{aligned} \quad (24)$$

From this relation taking into account (23) and the uniform boundedness of $\|u_\varepsilon\|_{H^1(\Omega_\varepsilon)}$, we derive

$$\|u_\varepsilon\|_{L_2(\Gamma_1)} \leq (\varepsilon^{\alpha-\beta-1} + \sqrt{\varepsilon})^{1/2}. \quad (25)$$

Since the family $\{u_\varepsilon\}$ is uniformly bounded in $H^1(\Omega_\varepsilon)$ it is compact in $H^{\frac{1}{2}}(\Omega)$. Consider arbitrary convergent subsequence u_{ε_k} , $\varepsilon_k \rightarrow 0$. It is evident that the limiting function $u'(x)$ satisfies the equation $\Delta u' = f$ and, in view of (25), the boundary condition $u'|_{\Gamma_1} = 0$. Hence, $u'(x) = u_0(x)$. Then, the whole family $\{u_\varepsilon\}$ does converge and the estimate (18) holds.

The estimate (20) is based on the following transformations

$$\int_{\Omega_\varepsilon} |\nabla(u_\varepsilon - u_0)|^2 dx = \int_{\Omega_\varepsilon} \nabla u_\varepsilon \nabla(u_\varepsilon - u_0) dx -$$

$$\begin{aligned}
 & - \int_{\Omega} \nabla u_0 \nabla (u_\varepsilon - u_0) dx - \int_{\Omega_\varepsilon \setminus \Omega} \nabla u_0 \nabla (u_\varepsilon - u_0) dx = \\
 & = \int_{\Omega_\varepsilon} f(u_\varepsilon - u_0) dx - \varepsilon^\beta \int_{\Gamma_1^\varepsilon} p(x, \frac{\hat{x}}{\varepsilon^\alpha}) u_\varepsilon (u_\varepsilon - u_0) ds - \int_{\Omega} f(u_\varepsilon - u_0) dx + \quad (26) \\
 & + \int_{\Gamma_1} \frac{\partial}{\partial x_n} u_0 (u_\varepsilon - u_0) d\hat{x} - \int_{\Omega_\varepsilon \setminus \Omega} \nabla u_0 \nabla (u_\varepsilon - u_0) dx + \varepsilon^{\alpha-1} \int_{\Gamma_1^\varepsilon} g(x, \frac{\hat{x}}{\varepsilon^\alpha}) (u_\varepsilon - u_0) ds = \\
 & = \int_{\Omega_\varepsilon \setminus \Omega} f(u_\varepsilon - u_0) dx - \int_{\Omega_\varepsilon \setminus \Omega} \nabla u_0 \nabla (u_\varepsilon - u_0) dx - \varepsilon^\beta \int_{\Gamma_1^\varepsilon} p(x, \frac{\hat{x}}{\varepsilon^\alpha}) u_\varepsilon^2 ds + \\
 & + \varepsilon^{\alpha-1} \int_{\Gamma_1^\varepsilon} g(x, \frac{\hat{x}}{\varepsilon^\alpha}) (u_\varepsilon - u_0) ds + \varepsilon^\beta \int_{\Gamma_1^\varepsilon} p(x, \frac{\hat{x}}{\varepsilon^\alpha}) u_\varepsilon u_0 ds + \int_{\Gamma_1} u_\varepsilon \frac{\partial}{\partial x_n} u_0 d\hat{x}.
 \end{aligned}$$

In the further considerations we use the following statement that can be proved in the same way as Lemma 1.

Proposition 9. *Let $u(x)$ belong to $H^2(\mathbf{R}^n)$. Then,*

$$\|u(\hat{x}, \varepsilon F(\hat{x}, \frac{\hat{x}}{\varepsilon^\alpha})) - u(\hat{x}, 0)\|_{L_2(\Gamma_1)} \leq C_5 \varepsilon \|u\|_{H^2(\Omega_\varepsilon)}.$$

Now, in case of nonnegative $p(\hat{x}, \hat{\xi})$ we deduce from (26) by means of (18), Lemma 1, Lemma 2 and Proposition 9

$$\begin{aligned}
 & \int_{\Omega_\varepsilon} |\nabla (u_\varepsilon - u_0)|^2 dx + \varepsilon^\beta \int_{\Gamma_1^\varepsilon} p(x, \frac{\hat{x}}{\varepsilon^\alpha}) u_\varepsilon^2 ds \leq \\
 & \leq C \left(\sqrt{\varepsilon} \|f\|_{L_2(\Omega_\varepsilon)} \|u_\varepsilon - u_0\|_{H^1(\Omega_\varepsilon)} + \sqrt{\varepsilon} \|u_0\|_{H^2(\mathbf{R}^n)} \|u_\varepsilon - u_0\|_{H^1(\Omega_\varepsilon)} + \right. \\
 & \quad \left. + \varepsilon^{\beta-\alpha+1} \varepsilon \|u_0\|_{H^2(\mathbf{R}^n)} (\varepsilon^{\frac{\alpha-\beta-1}{2}} + \varepsilon^{\frac{1}{4}}) \|u_\varepsilon\|_{H^1(\Omega_\varepsilon)} + \right. \\
 & \quad \left. + (\varepsilon^{\frac{\alpha-\beta-1}{2}} + \varepsilon^{\frac{1}{4}}) \|u_\varepsilon\|_{H^1(\Omega_\varepsilon)} + \varepsilon \|u_0\|_{H^2(\mathbf{R}^n)} \right).
 \end{aligned}$$

Since $\|u_\varepsilon\|_{H^1(\Omega_\varepsilon)}$ is uniformly bounded, this implies

$$\int_{\Omega_\varepsilon} |\nabla(u_\varepsilon - u_0)|^2 dx \leq C(\varepsilon^{\frac{1}{4}} + \varepsilon^{\frac{\alpha-\beta-1}{2}} + \varepsilon^{\frac{\beta-\alpha+3}{2}} + \varepsilon^{\beta-\alpha+9/4})$$

and (20) follows.

In the case $\beta > \alpha - 2$ we start with the following identity

$$\begin{aligned} & \int_{\Omega_\varepsilon} |\nabla(u_0 - u_\varepsilon)|^2 dx + \varepsilon^{\beta-\alpha+1} \int_{\Gamma_1} P_1(\varepsilon, \hat{x})(u_0 - u_\varepsilon)^2 d\hat{x} = \\ & = \int_{\Omega_\varepsilon} \nabla u_0 \nabla(u_0 - u_\varepsilon) dx - \int_{\Omega_\varepsilon} f(u_0 - u_\varepsilon) dx - \\ & - \varepsilon^{\alpha-1} \int_{\Gamma_1^\varepsilon} g(u_0 - u_\varepsilon) ds + \varepsilon^\beta \int_{\Gamma_1^\varepsilon} p u_0 (u_0 - u_\varepsilon) ds + \\ & + \varepsilon^{\beta-\alpha+1} \int_{\Gamma_1} P_1(\varepsilon, \hat{x})(u_0 - u_\varepsilon)^2 d\hat{x} - \varepsilon^\beta \int_{\Gamma_1^\varepsilon} p(u_0 - u_\varepsilon)^2 ds. \end{aligned} \quad (27)$$

Then, taking into account the assumption $\beta > \alpha - 2$ we obtain by (23) and Lemma 8

$$\begin{aligned} & \left| \varepsilon^{\beta-\alpha+1} \int_{\Gamma_1} P_1(\varepsilon, \hat{x})(u_0 - u_\varepsilon)^2 d\hat{x} - \varepsilon^\beta \int_{\Gamma_1^\varepsilon} p(u_0 - u_\varepsilon)^2 ds \right| \leq \\ & \leq \varepsilon^{\beta-\alpha+1} C \left(\sqrt{\varepsilon} \|u_\varepsilon - u_0\|_{L_2(\Gamma_1)} \|u_\varepsilon - u_0\|_{H^1(\Omega_\varepsilon)} + \varepsilon \|u_\varepsilon - u_0\|_{H^1(\Omega_\varepsilon)}^2 \right) \leq \\ & \leq \frac{1}{2} \left(\int_{\Omega_\varepsilon} |\nabla(u_0 - u_\varepsilon)|^2 dx + \varepsilon^{\beta-\alpha+1} \int_{\Gamma_1} P_1(\varepsilon, \hat{x})(u_0 - u_\varepsilon)^2 d\hat{x} \right) \end{aligned}$$

for all sufficiently small $\varepsilon > 0$. Combining this with (27) we find

$$\int_{\Omega_\varepsilon} |\nabla(u_0 - u_\varepsilon)|^2 dx + \varepsilon^{\beta-\alpha+1} \int_{\Gamma_1} P_1(\varepsilon, \hat{x})(u_0 - u_\varepsilon)^2 d\hat{x} \leq$$

$$\begin{aligned} &\leq 2 \left| \int_{\Omega_\varepsilon} \nabla u_0 \nabla (u_0 - u_\varepsilon) dx - \int_{\Omega_\varepsilon} f(u_0 - u_\varepsilon) dx - \right. \\ &\quad \left. - \varepsilon^{\alpha-1} \int_{\Gamma_1^\varepsilon} g(u_0 - u_\varepsilon) ds + \varepsilon^\beta \int_{\Gamma_1^\varepsilon} p u_0 (u_0 - u_\varepsilon) ds \right| \end{aligned}$$

Now, (20) can be obtained in the same way as above. The theorem is proved.

6 The “smooth” corrector

In this section we find more precise asymptotics of u^ε all over the domain Ω_ε in the case $\beta \geq \alpha - 1$. For this aim we introduce a family of auxiliary problems posed in the fixed domain Ω , with non-oscillating boundary conditions depending on ε . With the help of these auxiliary problems we define the corrector term which is regular in ε and allows to improve the estimate of the residual. For simplicity we suppose $p(\hat{x}, \hat{\xi}) \equiv 0$. One can easily generalize the results to the case of arbitrary smooth $p(\hat{x}, \hat{\xi})$ and $\beta \geq \alpha - 1$.

Theorem 4. *Let all the conditions of Theorem 1 be satisfied and $p(\hat{x}, \hat{\xi}) \equiv 0$. Then,*

$$\|U_0 - u_\varepsilon\|_{H^1(\Omega_\varepsilon)} \leq K_4 \sqrt{\varepsilon} \quad (28)$$

where K_4 does not depend on ε , $U_0(\varepsilon, x)$ is the solution of a problem

$$\begin{cases} -\Delta U_0 = f & \text{in } \Omega, \\ \frac{\partial U_0}{\partial x_n} = G_1(\varepsilon, \hat{x}) & \text{on } \Gamma_1, \\ U_0 = 0 & \text{on } \Gamma_2, \end{cases} \quad (29)$$

and

$$G_1(\varepsilon, \hat{x}) = \int_0^1 \int_0^1 \cdots \int_0^1 \sqrt{\varepsilon^{2\alpha-2} + |\nabla_{\hat{\xi}} F(\hat{x}, \hat{\xi})|^2} g(\hat{x}, \hat{\xi}) d\hat{\xi}.$$

The relation

$$\|U_0(\varepsilon, x) - u_0(x)\|_{H^1(\Omega)} = O(\varepsilon^{\alpha-1}) \quad (30)$$

holds.

Proof of Theorem 4. We start with the following statement that can be proved in the same way as Lemma 6.

Proposition 10. *For any $v \in H^1(\Omega_\varepsilon)$*

$$\left| \int_{\Gamma_1^\varepsilon} \varepsilon^{\alpha-1} g\left(\hat{x}, \frac{\hat{x}}{\varepsilon^\alpha}\right) v\left(\hat{x}, \varepsilon F\left(\hat{x}, \frac{\hat{x}}{\varepsilon^\alpha}\right)\right) ds - \int_{\Gamma_1} G_1(\varepsilon, \hat{x}) v(\hat{x}, 0) d\hat{x} \right| \leq C \sqrt{\varepsilon} \|v\|_{H^1(\Omega_\varepsilon)},$$

Due to the smoothness of the boundary $\partial\Omega$ the solution U_0 belongs to $H^2(\Omega \cap V)$ [L], and, hence, can be continued on V to belong to $H^2(V)$ [MV].

Let us write the integral identity for problem (29)

$$\begin{aligned} \int_{\Omega_\varepsilon} \nabla(U_0 - u_\varepsilon) \nabla v \, dx &= \int_{\Omega_\varepsilon} \nabla U_0 \nabla v \, dx - \int_{\Omega_\varepsilon} f v \, dx - \varepsilon^{\alpha-1} \int_{\Gamma_1^\varepsilon} g v \, ds = \\ &= \int_{\Omega} \nabla U_0 \nabla v \, dx - \int_{\Omega_\varepsilon} f v \, dx - \varepsilon^{\alpha-1} \int_{\Gamma_1^\varepsilon} g v \, ds + \int_{\Omega_\varepsilon \setminus \Omega} \nabla U_0 \nabla v \, dx = \\ &= \int_{\Omega_\varepsilon \setminus \Omega} \nabla U_0 \nabla v \, dx - \varepsilon^{\alpha-1} \int_{\Gamma_1^\varepsilon} g v \, ds + \int_{\Gamma_1} G_1 v \, d\hat{x} - \int_{\Omega_\varepsilon \setminus \Omega} f v \, dx. \end{aligned}$$

By Lemma 1

$$\left| \int_{\Omega_\varepsilon \setminus \Omega} \nabla U_0 \nabla v \, dx \right| \leq \|\nabla U_0\|_{L_2(\Omega_\varepsilon \setminus \Omega)} \|v\|_{H^1(\Omega_\varepsilon)} \leq C_2 \sqrt{\varepsilon} \|U_0\|_{H^2(\Omega_\varepsilon)} \|v\|_{H^1(\Omega_\varepsilon)}$$

and

$$\left| \int_{\Omega_\varepsilon \setminus \Omega} f v \, dx \right| \leq \|f\|_{L_2(\Omega_\varepsilon \setminus \Omega)} \|v\|_{L_2(\Omega_\varepsilon \setminus \Omega)} \leq C_2 \sqrt{\varepsilon} \|f\|_{L_2(\Omega_\varepsilon)} \|v\|_{H^1(\Omega_\varepsilon)}.$$

Then, due to Proposition 10

$$\left| \varepsilon^{\alpha-1} \int_{\Gamma_1^\varepsilon} g v \, ds - \int_{\Gamma_1} G_1 v \, d\hat{x} \right| \leq C \sqrt{\varepsilon} \|v\|_{H^1(\Omega_\varepsilon)}.$$

From these inequalities we deduce

$$\left| \int_{\Omega_\varepsilon} \nabla(U_0 - u_\varepsilon) \nabla v \, dx \right| \leq K_4 \sqrt{\varepsilon} \|v\|_{H^1(\Omega_\varepsilon)}.$$

Substituting $v = U_0 - u_\varepsilon$ and applying the Friedrichs type inequality, we obtain (28). The theorem is proved.

In the remaining part of this section we analyze in more detail the asymptotics of corrector term $(u_0 - U_0)$ depending on the structure of function $F(\hat{x}, \hat{\xi})$. It is clear that the principal term of this asymptotics depends only on the behavior of F in a small neighborhood of the singular points of F (zero-points of ∇F). For simplicity we consider the case $n = 2$. Multidimensional case can be studied in analogous way.

The following assertion takes place.

Proposition 11. *Suppose that $n = 2$, $(\alpha - 1)$ is sufficiently small and*

$$\text{dist} \left(\text{supp}_x g(\hat{x}, \hat{\xi}), \Gamma_1 \setminus \text{supp}_x F(\hat{x}, \hat{\xi}) \right) > 0.$$

Then,

- (i) *If F is a Morse function of ξ uniformly in $x \in \text{supp}_x g$ and all its singular points are quadratically nondegenerate, then*

$$\|u_0 - U_0\|_{H^1(\Omega)} \leq M_1 \varepsilon^{2(\alpha-1)} |\ln \varepsilon|.$$

- (ii) *If F has a finite number of singular points and the degeneracy of $|\nabla F|^2$ at these points is of order $2k$, then*

$$\|u_0 - U_0\|_{H^1(\Omega)} \leq M_2 \varepsilon^{(1+\frac{1}{k})(\alpha-1)}.$$

Remark 6. *In general estimate (30) cannot be improved.*

Proof. Let us introduce a new small parameter $\omega = \varepsilon^{\alpha-1}$ and estimate the derivative in ω of the function $G_1(\omega, \hat{x})$ for small ω . We have

$$\frac{\partial G_1(\omega, \hat{x})}{\partial \omega} = \int_0^1 \frac{g(\hat{x}, \hat{\xi}) \omega d\xi}{\sqrt{\omega^2 + |\nabla_{\xi} F|^2}}.$$

Now,

- (i) If $F(\hat{x}, \hat{\xi})$ is a Morse function one can easily verify that the main contribution to the derivative $\left. \frac{\partial G_1(\omega, \hat{x})}{\partial \omega} \right|_{\omega=0}$ is given by the small neighborhood of the

singular points. Let us estimate the contribution of one of them. We denote it by ξ_0 . By our assumption $|\nabla_\xi F|^2 = C|\xi - \xi_0|^2(1 + o(1))$ as $\xi - \xi_0 \rightarrow 0$; therefore,

$$\begin{aligned} \left| \int_{\xi_0-\delta}^{\xi_0+\delta} \frac{g(\omega, \hat{x}) \omega d\xi}{\sqrt{\omega^2 + |\nabla_\xi F|^2}} \right| &\leq C_1 \left| \int_{\xi_0-\delta}^{\xi_0+\delta} \frac{g(\omega, \hat{x}) \omega d\xi}{\sqrt{\omega^2 + C|\xi - \xi_0|^2}} \right| + O(\omega) \leq \\ &\leq C_2 \int_0^{\frac{\delta}{\omega}} \frac{\omega ds}{\sqrt{1 + Cs^2}} \leq C_3 \omega |\ln \omega|. \end{aligned}$$

Hence,

$$|G_1(\omega, \hat{x}) - G(\hat{x})| \leq C_4 \omega^2 |\ln \omega|,$$

and by standard energy estimate

$$\|u_0 - U_0\|_{H^1(\Omega)} \leq M_1 \varepsilon^{2(\alpha-1)} |\ln \varepsilon|.$$

(ii) By the same reason as in 1 we will estimate the contribution of one of the singular points. If $|\nabla_\xi F|^2 = C|\xi - \xi_0|^{2k}(1 + o(1))$ as $\xi - \xi_0 \rightarrow 0$, then

$$\begin{aligned} \left| \int_{\xi_0-\delta}^{\xi_0+\delta} \frac{g(\omega, \hat{x}) \omega d\xi}{\sqrt{\omega^2 + |\nabla_\xi F|^2}} \right| &\leq C_1 \int_{\xi_0-\delta}^{\xi_0+\delta} \frac{\omega d\xi}{\sqrt{\omega^2 + C|\xi - \xi_0|^{2k}}} \leq \\ &\leq C_2 \int_0^{\delta/\omega^{\frac{1}{k}}} \frac{\omega^{\frac{1}{k}} ds}{\sqrt{1 + Cs^{2k}}} \leq C_3 \omega^{\frac{1}{k}}. \end{aligned}$$

Thus, in this case

$$|G_1(\omega, \hat{x}) - G(\hat{x})| \leq C_4 \omega^{1+\frac{1}{k}}.$$

The proposition is proved.

Acknowledgements. The research was initiated while G.A.Checkkin and A.L.Piatnitski were enjoying their stay at the Institute for Mathematics and its Applications of the University of Minnesota, whose support is gratefully

acknowledged. It is a pleasure to thank professor N.V.Krylov for stimulating our interest to the problem. We also wish to express our thanks to A.G.Belyaev for the useful remarks. The first author was partially supported by RFFI grant No 95-01-00165.

Bibliography

- [B1] Belyaev, A.G. 1988. Average of the Third Boundary-Value Problem for the Poisson Equation in a Domain with Rapidly Oscillating Boundary (Russian). *Vestnik Moskov. Univ. (Ser. I Mat. Mekh)* **6**: 63 - 66.
- [B2] Belyaev, A.G. 1990. *On Singular Perturbations of Boundary-Value Problems* (Russian). PhD Thesis. Moscow State University.
- [Br] Briane, M. 1990. *Homogénéisation de Matériaux Fibres et Multi-Couches* (French). PhD Thesis. University Paris 6.
- [BLMS] Bouchitte, G., Lidouh, A., Michel, J.C. and Suquet, P. 1993. Might Boundary Homogenization Help to Understand Friction? Trieste: *International Centre for Theoretical Physics*, SMR. 719/9.
- [BLS] Bouchitte, G., Lidouh, A., and Suquet, P. 1991. Homogénéisation de Frontière pour la Modélisation du Contact entre un Corps déformable non Linéaire et un Corps rigide. *C. R. Acad. Sc. Paris* **313** (Ser. I): 967 - 972.
- [BM] Ball, J.M., and Murat, F. 1990. Remarks on Rank-One Convexity and Quasiconvexity. Paris: *Université Pierre et Marie Curie, Centre National de la Recherche Scientifique*, R 90043.
- [BMS] Belyaev, A.G., Mikheev, A.G., and Shamaev, A.S. 1992. Plane Wave Diffraction by a Rapidly Oscillating Surface. *Zh. Vychisl. Mat. i Mat. Fiz.* **32**(8): 1258 - 1272. (translation *Comput. Math. Math. Phys.* **32**(8): 1121 - 1133.)
- [L] Ladyzhenskaya, O.A. 1973. *Boundary-Value Problems of Mathematical Physics*. Moscow: Nauka.
- [LM] Lions, J.L., and Magenes, E. 1973. *Non-homogeneous Boundary-Value Problems and Applications*. Berlin - New York: Springer-Verlag.
- [M] Maz'ya, V.G. 1984. *The S.L.Sobolev's Spaces*. Leningrad: Leningrad State University Press.

- [Mi] Mikhailov, V.P. 1984. *Partial Differential Equations*. Moscow: Nauka.
- [MV] Maz'ya, V.G. 1987. Classes of Spaces, Measures, and Capacities in the theory of spaces of differentiable functions. In: *Modern Problems of Mathematics. Fundamental investigations. (Itogi Nauki i Tekhniki VINITI AN SSSR)* **26**: 159 - 228.
- [NM] Movchan, A.B., and Nazarov, S.A. 1991. The Influence of Small Surface Irregularities on the Stress State of a Body and the Energy Balance for a Growing Crack. *Prikl. Mat. Mech.* **55**(5): 819 - 828. (translation *J. Appl. Maths Mechs.* **55**(5): 691 - 701.)
- [O] Oleinik, O.A., Shamaev, A.S., and Yosifian, G.A. 1992. *Mathematical Problems in Elasticity and Homogenization*. Amsterdam: North-Holland.
- [RS] Reed, M., and Simon, B. 1975. *Methods of Modern Mathematical Physics. II Fourier analysis. Self-Adjointness*. New York - San Francisco - London: Academic Press.
- [SP] Sanchez-Palencia, E. 1987. *Homogenization Techniques for Composite Media*. Berlin, New York: Springer-Verlag.
- [SPE] Sanchez-Palencia, E. 1980. *Non Homogeneous Media and Vibration Theory*. Berlin, New York: Springer-Verlag.
- [SS] Sanchez-Palencia, E., and Suquet, P. 1983. Friction and homogenization of a boundary. In: *Free Boundary Problems: Theory and Applications*, ed. A.Fasano, M.Primicerio, 561 - 571. London: Pitman.
- [Y] Yosida, K. 1995. *Functional Analysis*. Berlin - New York: Springer-Verlag.



Unité de recherche INRIA Lorraine, Technopôle de Nancy-Brabois, Campus scientifique,
615 rue du Jardin Botanique, BP 101, 54600 VILLERS LÈS NANCY
Unité de recherche INRIA Rennes, Irisa, Campus universitaire de Beaulieu, 35042 RENNES Cedex
Unité de recherche INRIA Rhône-Alpes, 655, avenue de l'Europe, 38330 MONTBONNOT ST MARTIN
Unité de recherche INRIA Rocquencourt, Domaine de Voluceau, Rocquencourt, BP 105, 78153 LE CHESNAY Cedex
Unité de recherche INRIA Sophia Antipolis, 2004 route des Lucioles, BP 93, 06902 SOPHIA ANTIPOLIS Cedex

Éditeur
INRIA, Domaine de Voluceau, Rocquencourt, BP 105, 78153 LE CHESNAY Cedex (France)
ISSN 0249-6399