

Optimization Problems with perturbations: A Guided Tour

J. Frederic Bonnans, Alexander Shapiro

► **To cite this version:**

J. Frederic Bonnans, Alexander Shapiro. Optimization Problems with perturbations: A Guided Tour. [Research Report] RR-2872, INRIA. 1996. inria-00073819

HAL Id: inria-00073819

<https://hal.inria.fr/inria-00073819>

Submitted on 24 May 2006

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

*Optimization Problems with perturbations,
A guided tour*

J. Frédéric Bonnans, Alexander Shapiro

N° 2872

Avril 1996

_____ THÈME 4 _____



*Rapport
de recherche*





Optimization Problems with perturbations, A guided tour

J. Frédéric Bonnans*, Alexander Shapiro**

Thème 4 — Simulation
et optimisation
de systèmes complexes
Projet Programmation mathématique

Rapport de recherche n° 2872 — Avril 1996 — 43 pages

Abstract: This paper presents an overview of some recent and significant progress in the theory of optimization with perturbations. We put the emphasis on methods based on upper and lower estimates of the value of the perturbed problems. These methods allow to compute expansions of the value function and approximate solutions in situations where the set of Lagrange multipliers may be unbounded, or even empty. We give rather complete results for nonlinear programming problems, and describe some partial extensions of the method to more general problems. We illustrate the results by computing the equilibrium position of a chain that is almost vertical or horizontal.

Key-words: Constrained optimization, Stability, sensitivity, duality, expansion of solutions, semi-infinite programming, semi-definite programming, second order analysis.

(Résumé : tsvp)

*INRIA-Rocquencourt, Domaine de Voluceau, B.P. 105, 78153 Rocquencourt, France, and Ecole Polytechnique, France

**School of Industrial and Systems Engineering, Georgia Institute of Technology, Atlanta, Georgia 30332-0205, USA

Unité de recherche INRIA Rocquencourt
Domaine de Voluceau, Rocquencourt, BP 105, 78153 LE CHESNAY Cedex (France)
Téléphone : (33 1) 39 63 55 11 – Télécopie : (33 1) 39 63 53 30

Problèmes d'optimisation avec perturbation : Une visite guidée

Résumé : Cet article présente une vue d'ensemble de progrès récents et significatifs en théorie de l'optimisation avec perturbation. Nous mettons l'accent sur les méthodes basées sur les estimations supérieures et inférieures de la valeur des problèmes perturbés. Ces méthodes permettent de calculer des développements de la fonction valeur et des solutions approchées dans des situations où l'ensemble des multiplicateurs de Lagrange peut être non borné, ou même vide. Nous donnons des résultats assez complets pour la programmation non linéaire, et décrivons des extensions partielles des méthodes à des problèmes plus généraux. Nous illustrons les résultats en calculant la position d'équilibre d'une chaîne presque verticale ou horizontale.

Mots-clé : Optimisation avec contraintes, stabilité, sensibilité, dualité, développement des solutions, optimisation semi-infinie, optimisation semi définie, analyse du deuxième ordre.

AMS 49K40, 90C31, 26B25, 41A58, 65K05, 49M45.

1 Introduction

In this paper we present a survey of some results on stability and sensitivity analysis of optimization problems which are subject to perturbations. We consider problems of the form

$$\underset{x \in X}{\text{Min}} f(x, u) \quad \text{subject to } x \in \Phi(u). \quad (P_u) \quad (1.1)$$

In most examples, considered in this paper, X is a Banach space, while the perturbation parameter u can be a scalar, a finite dimensional vector, or even an element of an appropriate normed or metric space U . We investigate continuity and differentiability properties of the optimal value $v(u)$ and the set $S(u)$ of optimal solutions of (P_u) considered as functions of the parameter u .

In order to proceed in our analysis we need a constructive method for describing the feasible set $\Phi(u)$. We say that the feasible set is defined by *abstract constraints* if it is given in the following form

$$\Phi(u) = \{x : G(x, u) \in K\}, \quad (1.2)$$

where K is a closed convex subset of a Banach space Y and $G : X \times U \rightarrow Y$. This includes the situations where K is a closed convex cone, in which case we speak of *cone constraints*.

An important particular case is when the feasible set is *finitely constrained*, i.e. is of the form

$$\Phi(u) = \{x : g_i(x, u) = 0, i = 1, \dots, q; g_i(x, u) \leq 0, i = q + 1, \dots, p\}. \quad (1.3)$$

Then $G(x, u) = (g_1(x, u), \dots, g_p(x, u)) : X \times U \rightarrow \mathbb{R}^p$ and $K = \{0\} \times \mathbb{R}_-^{p-q}$. In that case, if in addition X is finite dimensional, (P_u) becomes a *nonlinear programming problem*. Another important case is when Y is the linear space of $n \times n$ symmetric matrices and $K \subset Y$ is the cone of positive semidefinite matrices. This example corresponds to the so-called semidefinite programming. Another example where the ‘‘cone constraints’’ formulation appears in a natural way is the example of semi-infinite programming. That is, let T be a compact metric space, let for every u , $G(x, u)$ be a mapping from X into the space $Y = C(T)$ of continuous real valued functions on T , and let $K = \{g \in C(T) : g(t) \leq 0, \forall t \in T\}$ be the set of nonpositive valued functions in $C(T)$. For such defined $G(x, u)$ and K , equation (1.2) is equivalent to

$$\Phi(u) = \{x : g_t(x, u) \leq 0, t \in T\}, \quad (1.4)$$

where $g_t(x, u) = G(x, u)(t)$. In case the feasible set is defined by an infinite number of constraints, as in (1.4), and the space X is finite dimensional, the corresponding program (P_u) is called a semi-infinite program.

Investigation of max-min problems goes back to the classical work of Chebyshev on uniform approximations by algebraic polynomials. It was also one of the driving forces behind systematic development of nonsmooth analysis in the sixties and seventies. There are numerous studies where various aspects of max-min optimization are discussed (see, e.g.,

[20, 21]). Quite surprisingly it was discovered recently by Gauvin and Janin [27] that in some cases first-order differentiability properties of the optimal value function are closely related to a second-order analysis of the corresponding optimization program.

Differentiability properties of the optimal solutions were first derived by applying the classical Implicit Function Theorem to the first-order optimality conditions written in a form of (nonlinear) equations. This approach is discussed at length and relevant references can be found in Fiacco [24]. The equations based approach was extended and generalized in works of Robinson [54, 55] where sensitivity analysis of variational inequalities (generalized equations in the terminology of Robinson) was developed. In recent years substantial advances in our understanding of asymptotic behavior of optimal solutions were made by using a somewhat different approach. The main idea of that approach is based on second-order upper and lower estimates of the (optimal) value function. When these estimates are sufficiently close, they allow to state an expansion of the optimal value function that, in turn, give a formula for the expansion of optimal (suboptimal) solutions, that is related to the optimal solutions of an auxiliary optimization problem [6, 10, 12, 15, 31, 59, 60, 68]. In this paper we mainly follow this approach.

1.1 Basic notation and terminology

\mathbb{R}^n n -dimensional Euclidean space

$\mathbb{R}_+^n = \{x \in \mathbb{R}^n : x_i \geq 0, i = 1, \dots, n\}$ nonnegative orthant

$\mathbb{R}_-^n = -\mathbb{R}_+^n = \{x \in \mathbb{R}^n : x_i \leq 0, i = 1, \dots, n\}$ nonpositive orthant

$x \cdot y = \sum_{i=1}^n x_i y_i$ scalar product of $x, y \in \mathbb{R}^n$

$\nabla g(x)$ gradient of the function $g : \mathbb{R}^n \rightarrow \mathbb{R}$ at the point $x \in \mathbb{R}^n$

X^* dual of the normed space X

$\langle \alpha, x \rangle$ value of the linear functional $\alpha \in X^*$ on $x \in X$

$B_X(x, r) = \{x' \in X : \|x' - x\| < r\}$ open ball of radius $r > 0$ centered at x

$B_X = B_X(0, 1)$ is the unit ball in X

$[x] = \{tx : t \in \mathbb{R}\}$ linear space generated by vector x

2^X the set of subsets of X

$\Psi : X \rightarrow 2^Y$ is a multifunction (point-to-set mapping), mapping X into the set of subsets of Y

$\text{dom}\Psi = \{x \in X : \Psi(x) \neq \emptyset\}$ is the domain of the multifunction Ψ

$\text{range}\Psi = \{y \in Y : y \in \Psi(x), x \in X\}$ is the range of the multifunction Ψ

$\text{gph}\Psi = \{(x, y) \in X \times Y : y \in \Psi(x), x \in X\}$ is the graph of the multifunction Ψ

$\Psi^{-1}(y) = \{x \in X : y \in \Psi(x)\}$ graph inverse of the multifunction Ψ

$I(x, u) = \{i : g_i(x, u) = 0, i = q + 1, \dots, p\}$ set of active at (x, u) inequality constraints

$L(x, \lambda, u) = f(x, u) + \langle \lambda, G(x, u) \rangle$ the Lagrangian function

$\Lambda(x, u)$ the set of Lagrange multipliers of (P_u) at the point $x \in S(u)$

$L^g(x, \lambda_0, \lambda, u) = \lambda_0 f(x, u) + \langle \lambda, G(x, u) \rangle$ generalized Lagrangian function

$\text{val}(P)$ optimal value of the program (P)

$\mathcal{S}(P)$ set of optimal solutions of the program (P)

$v(u) = \text{val}(P_u) = \inf_{x \in \Phi(u)} f(x, u)$ optimal value function
 $S(u) = \mathcal{S}(P_u) = \arg \min_{x \in \Phi(u)} f(x, u)$ set of optimal solutions
 $\bar{x}(u) \in S(u)$ an optimal solution
 $\text{dist}(x, S) = \inf_{z \in S} \|x - z\|$ distance from the point $x \in X$ to set $S \subset X$
 $Dg(x) \in \mathcal{L}(X, Y)$ derivative (Gâteaux, Hadamard or Fréchet, depending on the context) of the mapping $g : X \rightarrow Y$ at the point $x \in X$
 $D^2g(x) : X \rightarrow \mathcal{L}(X, Y)$ second-order derivative of the mapping g at the point x
 $D^2g(x)(h, h) = (D^2g(x)h)h$ quadratic form corresponding to D^2g
 $D_xg(x, u) \in \mathcal{L}(X, Y)$ partial derivative of $g : X \times U \rightarrow Y$
 $Dg(x, u)(h, d) = D_xg(x, u)h + D_u g(x, u)d$
 $g'(x, d) = \lim_{t \rightarrow 0^+} [g(x + td) - g(x)]/t$ directional derivative of g at x in direction d
 $f'_+(x, d) = \limsup_{t \rightarrow 0^+} [f(x + td) - f(x)]/t$ upper directional derivative of $f : X \rightarrow \mathbb{R}$
 $\partial f(x) = \{\alpha \in X^* : f(y) - f(x) \geq \langle \alpha, y - x \rangle, \forall y \in X\}$ subdifferential of the convex function $f : X \rightarrow \mathbb{R} \cup \{-\infty\} \cup \{+\infty\}$
 $C^- = \{\alpha \in X^* : \langle \alpha, x \rangle \leq 0, \forall x \in C\}$ polar (negative dual) of the convex cone $C \subset X$
 $\Sigma^- = \{x \in X : \langle \alpha, x \rangle \leq 0, \forall \alpha \in \Sigma\}$ polar (negative dual) of the convex cone $\Sigma \subset X^*$
 $a \preceq_C b$ order relation imposed by the cone C , i.e. $b - a \in C$
 $\mathbb{R}_+(S) = \{tx : x \in S, t \geq 0\}$ cone generated by the set S
 $S^\perp = \{\alpha \in X^* : \langle \alpha, x \rangle = 0, \forall x \in S\}$ orthogonal set to $S \subset X$
 $\text{Sp}(S) = \mathbb{R}_+(S - S)$ linear space generated by $S \subset X$
 $T_S(x) = \{h \in X : \text{dist}(x + th, S) = o(t), t \geq 0\}$ tangent cone to the convex set S at the point $x \in S$
 $T_S^2(x, h) = \{w \in X : \text{dist}(x + th + \frac{1}{2}t^2w, S) = o(t^2), t \geq 0\}$ second-order tangent set
 $N_S(x) = \{\alpha \in X^* : \langle \alpha, z - x \rangle \leq 0, \forall z \in S\}$ normal cone to the set $S \subset X$ at $x \in S$
 $N_\Omega(\alpha) = \{x \in X : \langle \omega - \alpha, x \rangle \leq 0, \forall \omega \in \Omega\}$ normal cone to the set $\Omega \subset X^*$ at $\alpha \in \Omega$
 $\text{int}(S)$ interior of the set S
 $\text{core}(S) = \{x \in S : \forall x' \in X, \exists \varepsilon > 0, \forall t \in [-\varepsilon, \varepsilon], x + tx' \in S\}$
 $\sigma(\lambda, S) = \sup_{x \in S} \langle \lambda, x \rangle$ is the support function of the set S
 $C(T)$ Banach space of continuous functions $\psi : T \rightarrow \mathbb{R}$ defined on the compact metric space T and equipped with the sup-norm $\|\psi\| = \sup_{\tau \in T} |\psi(\tau)|$
 For $\varepsilon \geq 0$ we say that \bar{x} is an ε -optimal solution of (P_u) if $\bar{x} \in \Phi(u)$ and $f(\bar{x}, u) \leq v(u) + \varepsilon$. Consider a mapping $g : X \rightarrow Y$. We say that g is directionally differentiable, at x , in the Hadamard sense if the directional derivative $g'(x, d)$ exists for all d and, moreover,

$$g'(x, d) = \lim_{\substack{t \rightarrow 0^+ \\ d' \rightarrow d}} \frac{g(x + td') - g(x)}{t}. \quad (1.5)$$

Note that if g is directionally differentiable, at x , in the Hadamard sense, then $g'(x, d)$ is continuous in d and if, in addition, X is finite dimensional, then $g(x + h) = g(x) + g'(x, h) + o(\|h\|)$. For discussion and comparison of various concepts of directional differentiability see [7, 47, 62].

Consider a multifunction $\Psi : X \rightarrow 2^Y$. It is said that Ψ is upper Lipschitz at a point $\bar{x} \in X$, if $\Psi(x) \subset \Psi(\bar{x}) + c\|x - \bar{x}\|B_Y$ for some $c > 0$ and all x in a neighborhood of \bar{x} . The

multifunction Ψ is said to be closed at a point $x \in X$, if $x_n \rightarrow x$, $y_n \in \Psi(x_n)$ and $y_n \rightarrow y$, imply that $y \in \Psi(x)$. It is said that Ψ is closed if it is closed at every point of X . Note that Ψ is closed iff its graph $\text{gph}\Psi$ is a closed subset of $X \times Y$.

2 The chain problem

Let us now introduce a simple problem in static mechanics, known as the chain or catenary problem, on which some of the theoretical results presented here will be illustrated. We will deal with situations where the classical regularity results do not hold. For instance, in one of them, the set of Lagrange multipliers is empty.

Consider a chain with m rigid links ($m > 1$), each of length one. Suppose that the two endpoints of the chain are fixed. Assuming that the mass of each link is concentrated in its middle, we consider the problem of computing the equilibrium position of the chain. A formal description of the problem follows. Denote by (y_k, z_k) , the increment, horizontally and vertically, along the link. The links are numbered from 1 to m . We assume that one endpoint is fixed at $(0,0)$. The position of the other endpoint is $\sum_{i=1}^m (y_i, z_i)$, and is constrained to be $(y^e(u), z^e(u))$, where $u \in \mathbb{R}$ is viewed as the perturbation parameter.

The potential energy of the chain, is given by

$$E(y, z) = \frac{1}{2} \sum_{k=1}^m \left(\sum_{i=1}^k z_i + \sum_{i=1}^{k-1} z_i \right) = \sum_{k=1}^m \alpha_k z_k, \quad (2.1)$$

where $\alpha_k = m - k$, for $k = 1, \dots, m - 1$, and $\alpha_m = 1/2$. The equilibrium position of the chain can be calculated by minimizing the potential energy subject to involved physical constraints. This leads to the following optimization problem:

$$\begin{aligned} & \text{Min}_{y,z} E(y, z) \\ & \text{s.t. } y_k^2 + z_k^2 - 1 = 0, \quad k = 1, \dots, m, \quad \sum_{k=1}^m (y_k, z_k) = (y^e(u), z^e(u)). \end{aligned} \quad (2.2)$$

This example has been discussed by various authors, e.g. in the context of numerical algorithms [42] or as an illustration of optimality conditions [71] in finite-dimensional optimization. The above two references cover the situation when the solution is stable with respect to a perturbation of the position of the endpoints. Suppose now that for $u = 0$ the distance between the two endpoints is exactly m . Then the unperturbed problem has a unique feasible point (the chain is a straight line) which is therefore the unique optimal solution. However, the corresponding minimum is not always stable under a small perturbation of u (the set of feasible points vanishes when the distance between the two endpoints increases). These limit cases are perhaps those for which the perturbation analysis is the most interesting. For instance, let us consider the two situations in which the chain is either vertical or horizontal in the null position (i.e. when $u = 0$). That is, assume that either $y^e(u) = 0$, $z^e(u) = m - u$, or $y^e(u) = m - u$, $z^e(u) = 0$, and let one of the endpoints make a small step toward the other, i.e. let u increase from zero a small positive number. Is it possible then to compute an expansion of the position of the chain ?

As an application of the results reviewed here, we will show that the answer is positive. Note that, although these two cases may seem very similar, they are of fundamentally different nature. The chain is a model of a mechanical system, in which the force due to the weight of each link should be compensated by the forces applied at the endpoints of each link by the neighbouring links. The latter are colinear to the axes of the neighbouring links. The relation of equilibrium of forces coincides with the first-order optimality system. In the case of the vertical chain, the set of Lagrange multipliers is nonempty but unbounded (and so is the set of mechanical forces for which the equilibrium is satisfied). In the case of the horizontal chain, however, the forces between links are horizontal, while the weight forces are vertical. Therefore, no mechanical equilibrium is possible. Expressed in terms of optimization theory, this means that the set of Lagrange multipliers is empty. The ability of computing an expansion of the optimal solution with an empty set of Lagrange multipliers is a striking example of the strength of the tools presented here.

Related to the lack of regularity of these problems is the fact that in both cases, the variation of the optimal solution is of the order of the square root of the perturbation parameter. This can be seen easily in the case of a two links chain. Take for instance the case of a vertical chain. Let us denote by $\bar{y}_k(u), \bar{z}_k(u)$, $k = 1, \dots, m$, an optimal solution of (2.2), and consider the case of two links, i.e. $m = 2$. Then for $u > 0$ we have that the feasible set of (2.2) consists of two symmetric points, each one giving an optimal solution, and $\bar{y}_1(u) = \sqrt{u - u^2/4}$. We see here that (2.2) has an optimal solution $(\bar{y}(u), \bar{z}(u))$ which is not Lipschitz continuous at $u = 0$ and that this solution is of order $O(\sqrt{u})$. As we shall see later such behavior of $(\bar{y}(u), \bar{z}(u))$ persists in cases the chain has more than two links, and that $(\bar{y}(u), \bar{z}(u))$ can be expanded in terms of \sqrt{u} .

3 Lagrange multipliers and constraint qualifications

3.1 First order optimality systems

In this section we assume that the feasible set $\Phi(u)$ is defined by constraints and that $f(x, u)$ and $G(x, u)$ are continuously differentiable. If x is a local solution of (P_u) , then certain relations between the differential properties of the objective function f and the boundary of the feasible set must hold. In order to motivate the following development, let us give a mechanical interpretation of problem (P_u) , assuming for the moment that $X = \mathbb{R}^n$ and $\Phi(u)$ is of the form (1.3). Let us view the objective function as a potential energy to be minimized. The mechanical force deriving from it is $-\nabla_x f(x, u)$. Each inequality constraint is viewed as the non penetration of a rigid body of equation $g_i(x, u) \geq 0$. With this constraint is associated a repulsion force which is normal to the surface of the body. If $\nabla_x g_i(x, u)$ is nonzero, then the outward normal to the body is $-\nabla_x g_i(x, u)$. Taking into account the fact that the force should be null if the point is not in contact with the body, we may write the

equilibrium of forces as

$$\begin{cases} \nabla_x f(x, u) + \sum_{i=1}^p \lambda_i \nabla_x g_i(x, u) = 0, \\ g_i(x, u) = 0, \quad i = 1, \dots, q, \\ g_i(x, u) \leq 0, \quad \lambda_i \geq 0, \quad \lambda_i g_i(x, u) = 0, \quad i = q + 1, \dots, p. \end{cases} \quad (3.1)$$

This is the celebrated *first-order optimality system*. Note that this way of deriving the optimality system is opposite to the classical argument, where one analyses the outward normal directions to the feasible set rather than to the complementary set (both points of view are equivalent only under regularity assumptions).

If (x, λ) is a solution of the first-order optimality system, then λ is called a *Lagrange multiplier*. The set of Lagrange multipliers is a (possibly empty) polyhedron. We say that a local solution of (P_u) is *qualified*, whenever the set of associated Lagrange multiplier is not empty. A simple example of a convex problem with a non qualified solution is $\text{Min}_x \{x : x^2 \leq 0\}$. F. John [32] proved that with any local solution of a finitely constrained problem is associated at least one *generalized Lagrange multiplier*, defined as a nonzero vector $(\lambda_0, \lambda_1, \dots, \lambda_p)$ satisfying the *generalized optimality system*

$$\begin{cases} \lambda_0 \nabla_x f(x, u) + \sum_{i=1}^p \lambda_i \nabla_x g_i(x, u) = 0, \\ g_i(x, u) = 0, \quad i = 1, \dots, q, \\ g_i(x, u) \leq 0, \quad \lambda_i \geq 0, \quad \lambda_i g_i(x, u) = 0, \quad i = q + 1, \dots, p; \quad \lambda_0 \geq 0. \end{cases} \quad (3.2)$$

A generalized multiplier with $\lambda_0 = 0$ is said to be *singular*. With a nonsingular generalized multiplier is associated the Lagrange multiplier $(\lambda_0)^{-1}(\lambda_1, \dots, \lambda_p)$. Therefore, John's result may be restated as follows: with any local solution of a finitely constrained problem is associated either a Lagrange multiplier or a singular multiplier. Equivalently, a sufficient condition for qualification is that the set of singular multipliers is empty. The latter is a condition that depends only on the parametrization of the feasible set, and is equivalent to the Mangasarian-Fromovitz constraint qualification [44]:

$$\begin{cases} \text{the gradients } \nabla_x g_i(x, u), \quad i = 1, \dots, q, \text{ are linearly independent,} \\ \exists h \in \mathbb{R}^n : \quad h \cdot \nabla_x g_i(x, u) = 0, \quad i = 1, \dots, q; \\ \quad \quad \quad h \cdot \nabla_x g_i(x, u) < 0, \quad \forall i \in I(x, u), \end{cases} \quad (3.3)$$

where $I(x, u)$ denotes the set of active inequality constraints at (x, u) . If all constraints are of equality (resp. inequality) type, then (3.3) reduces to the more familiar condition that $D_x G(x, u)$ is onto (resp. $\exists h \in \mathbb{R}^n : G(x, u) + D_x G(x, u)h < 0$).

We now turn to the problem with abstract constraints. The first-order generalized optimality system

$$D_x L^g(x, \lambda_0, \lambda, u) = 0, \quad G(x, u) \in K, \quad \lambda \in N_K(G(x, u)), \quad \lambda_0 \geq 0, \quad (3.4)$$

where $L^g(x, \lambda_0, \lambda, u) = \lambda_0 f(x, u) + \langle \lambda, G(x, u) \rangle$, is a natural generalization of (3.2) to problems with abstract constraints. If $\lambda_0 \neq 0$, then one can take $\lambda_0 = 1$ and hence (3.4) becomes the following first-order optimality system

$$D_x L(x, \lambda, u) = 0, \quad G(x, u) \in K, \quad \lambda \in N_K(G(x, u)). \quad (3.5)$$

Note that if K is a convex closed cone and $y \in K$, then $N_K(y) = K^- \cap \{y\}^\perp$. Therefore in that case the optimality condition $\lambda \in N_K(G(x, u))$ is equivalent to $\lambda \in K^-$ and $\langle \lambda, G(x, u) \rangle = 0$.

We now discuss some qualification conditions that apply to problems with abstract constraints.

3.2 A general qualification condition

The following *regularity condition* is due to Robinson [52]:

$$0 \in \text{int} \{G(x, u) + D_x G(x, u)X - K\}. \quad (3.6)$$

If the feasible set is defined by a finite number of constraints in the form (1.3), then regularity conditions (3.3) and (3.6) are equivalent. The above condition recovers many special cases of interest. When K is a singleton (i.e., in the case of equality constraints), (3.6) reduces to the condition that $D_x G(x, u)$ is onto. When K has a non empty interior, (3.6) reduces (e.g. [48, Prop. 3.9(f)]) to the condition

$$\exists h \in X : G(x, u) + D_x G(x, u)h \in \text{int}(K).$$

The latter applies in particular to semi-infinite programming problems, with $\Phi(u)$ defined in the form (1.4) (see [64]). These two characterizations are very similar to those given before for (3.3) when all constraints are of equality (resp. inequality) type.

An interesting case is when $\Phi(u)$ is in the form (1.2) with Y being the Cartesian product of two Banach spaces Y_1 and Y_2 , and $K = \{0\} \times K_2 \subset Y_1 \times Y_2$, where K_2 is a closed convex subset of Y_2 with non empty interior. Then $G(x, u) = (G_1(x, u), G_2(x, u))$, with $G_i(x, u) \in Y_i$, $i = 1, 2$. In this case (3.6) is equivalent to the condition

$$\begin{aligned} D_x G_1(x, u) \text{ is onto;} \\ \exists h \in X : D_x G_1(x, u)h = 0; G_2(x, u) + D_x G_2(x, u)h \in \text{int}(K_2). \end{aligned} \quad (3.7)$$

Yet another particular form of the constraints is useful when dealing with optimal control problems. Consider the *abstract optimal control problem*

$$\text{Min}_{x \in X} f(x, u) \quad \text{subject to } x \in \mathcal{C} \text{ and } \mathcal{G}(x, u) \in \mathcal{K}, \quad (3.8)$$

where X is a space of controls, \mathcal{C} is a closed convex subset of X , representing control constraints, and \mathcal{K} is a closed convex subset of Banach space \mathcal{Y} ; the relation $\mathcal{G}(x, u) \in \mathcal{K}$ might represent state constraints. This is a particular form of (1.1) with the feasible set given in the form (1.2), as is seen by setting $Y := X \times \mathcal{Y}$, $G(x, u) := (x, \mathcal{G}(x, u))$, and $K := \mathcal{C} \times \mathcal{K}$. The regularity condition in this case turns out to be equivalent to

$$0 \in \text{int} \{G(x, u) + D_x G(x, u)(\mathcal{C} - x) - \mathcal{K}\}.$$

If in addition \mathcal{K} has a nonempty interior, then regularity is equivalent to a condition often used in optimal control (e.g. [11])

$$\exists x' \in \mathcal{C} : \mathcal{G}(x, u) + D_x \mathcal{G}(x, u)(x' - x) \in \text{int}(\mathcal{K}).$$

Let us mention some more abstract conditions equivalent to (3.6), that may be found in the literature. Because the core of a convex set includes its interior, (3.6) implies

$$0 \in \text{core} \{G(x, u) + D_x G(x, u)X - K\}. \quad (3.9)$$

It is not difficult to see that (3.9) is equivalent to the relation $\mathcal{A}(x, u) = Y$, where $\mathcal{A}(x, u)$ is defined by

$$\mathcal{A}(x, u) := \mathbb{R}_+[G(x, u) + D_x G(x, u)X - K]. \quad (3.10)$$

It happens that condition $\mathcal{A}(x, u) = Y$ implies (3.6). Therefore conditions (3.6), (3.9) and $\mathcal{A}(x, u) = Y$, in fact, are equivalent. The above implication may be proved by considering the multifunction $\mathcal{M} : \mathbb{R} \times X \rightarrow 2^Y$ defined by $\mathcal{M}(a, h) = D_x G(x, u)h + a(G(x, u) - K)$, if $a \geq 0$, and $\mathcal{M}(a, h) = \emptyset$, if $a < 0$. Condition $\mathcal{A}(x, u) = Y$ means that \mathcal{M} is onto, i.e. $\text{range}(\mathcal{M}) = Y$. It follows then from the generalized open mapping theorem (see Theorem 3.1 below) that $0 \in Y$ is an interior point of the set $\mathcal{M}([0, 1] \times X)$ and hence (3.6) follows.

Theorem 3.1 (Generalized Open Mapping Theorem [50],[53, Theorem 1]) *Let X and Y be Banach spaces, and let $\Psi : X \rightarrow 2^Y$ be a closed convex multifunction (i.e. $\text{gph}\Psi$ is a closed convex subset of $X \times Y$). Let $y \in \text{int}(\text{range}\Psi)$. Then for every $x \in \Psi^{-1}(y)$ and all $r > 0$, it follows that $y \in \text{int}\Psi(B_X(x, r))$.*

The following combination of results by [12, Part I, Prop. 4.1], [45, 51], [72, Th. 4.1]), states that *regularity is a qualification condition*.

Proposition 3.1 *Let x be a locally optimal solution of (P_u) , and $\mathcal{A}(x, u)$ be given in (3.10). Then: (a) If $\mathcal{A}(x, u) = Y$ (i.e. (3.6) holds), then the set of Lagrange multipliers is nonempty and bounded, and there exists no singular multiplier. (b) If $\mathcal{A}(x, u)$ is not dense in Y , then there exists a singular multiplier.*

Note that we do not have an alternative as before: it may happen that there exist no generalized Lagrange multipliers. This can occur only if $\mathcal{A}(x, u)$ is dense in Y but is not equal to Y .

Regularity is also related to stability with respect to perturbations of the feasible set, as it follows from the next result which is a consequence of the Robinson [53] - Ursescu [70] stability theorem (cf. [52, Corollary 1]).

Proposition 3.2 *Suppose that the constraint qualification (3.6) holds at (x^0, u^0) . Then for all (x, u) in a neighborhood of (x^0, u^0) , one has*

$$\text{dist}(x, \Phi(u)) = O(\text{dist}(G(x, u), K)). \quad (3.11)$$

The stability property (3.11) (called metric regularity) is particularly useful in deriving upper estimates for the optimal value function. Indeed, for a direction $d \in U$ let us consider a path $u(t) := u^0 + td + o(t)$ in the parameter space U . Suppose that we want to obtain upper estimates of the optimal value function $v(u(t))$ by constructing a path $x(t) \in X$ that is *feasible* with respect to the considered perturbations, i.e. $x(t) \in \Phi(u(t))$. We say that $h \in X$ is a (first-order) *feasible direction*, relative to the direction $d \in U$, if there exists $r(t) = o(t)$ such that $x^0 + th + r(t) \in \Phi(u(t))$.

It is easily checked that a necessary condition for h to be a feasible direction is

$$DG(x^0, u^0)(h, d) \in T_K(G(x^0, u^0)). \quad (3.12)$$

This condition is obviously sufficient if (3.11) holds. Therefore, by Proposition 3.2, (3.12) characterizes feasible directions whenever the constraints are regular.

3.3 Directional regularity

We continue the analysis of the set of directions that are feasible with respect to perturbations of the form $u(t) = u^0 + td + o(t)$, with $d \in U$. Whereas regularity is a condition that does not depend on the perturbation, we may search conditions depending on d , under which (3.12) characterizes the set of feasible directions.

The following condition, due to Gollan [28], applies to finitely constrained problems,

$$\left\{ \begin{array}{l} \text{the gradients } \nabla g_i(x^0, u^0), i = 1, \dots, q, \text{ are linearly independent,} \\ \exists h \in \mathbb{R}^n : \quad (h, d) \cdot \nabla g_i(x^0, u^0) = 0, i = 1, \dots, q; \\ \quad \quad \quad (h, d) \cdot \nabla g_i(x^0, u^0) < 0, \forall i \in I(x^0, u^0). \end{array} \right. \quad (3.13)$$

A natural extension that applies to general constraints is [12, Part I]:

$$0 \in \text{int} \{G(x^0, u^0) + DG(x^0, u^0)(X \times \mathbb{R}_+(d)) - K\}. \quad (3.14)$$

These conditions may be interpreted as regularity conditions with respect to the set $\{(x, t) : G(x, u^0 + td) \in K, t \geq 0\}$. It is clear that they explicitly depend on the chosen direction d , while the regularity conditions (3.3) and (3.6) (applied at (x^0, u^0)) do not. We refer to (3.13) and (3.14) as *directional regularity* condition because of the statement (c) in the following proposition.

Proposition 3.3 [12, Part I] *(a) Regularity implies directional regularity (i.e., (3.6) implies (3.14)). (b) The latter is equivalent to (3.13) in the case of finitely many constraints. (c) If directional regularity holds, then h is a first-order feasible direction iff it satisfies (3.12).*

Note that directional regularity is not a qualification condition (it does not imply existence of a Lagrange multiplier). It has useful characterizations similar to those of (3.6) when the constraints have particular forms. In the case of a product form discussed just before (3.7), such a characterization is

$$D_x G_1(x^0, u^0) \text{ is onto; } \exists h \in X, \exists \varepsilon > 0 : DG_1(x^0, u^0)(h, d) = 0, \\ G_2(x^0, u^0) + DG_2(x^0, u^0)(h, \varepsilon d) \in \text{int}(K_2). \quad (3.15)$$

In the case of the abstract optimal control problem when \mathcal{K} has a nonempty interior, we get the characterization

$$\exists x' \in \mathcal{C}, \exists \varepsilon > 0 : \mathcal{G}(x^0, u^0) + D\mathcal{G}(x^0, u^0)(x' - x^0, \varepsilon d) \in \text{int}(\mathcal{K}).$$

4 Analysis of the optimal value function

4.1 Problems with unperturbed feasible sets

In this section we discuss first-order differentiability properties of the optimal value function $v(u)$. We start our analysis by considering the case when the feasible set is unperturbed.

Example 4.1 *Let $f(x, u) = xu$, with $x, u \in \mathbb{R}$, and $\Phi(u) = [-1, 1]$ for all u . It is easily seen that $v(u) = -|u|$. Although the feasible set here is unperturbed and compact, the optimal value function is not differentiable at $u = 0$.*

This example shows how easily, starting with a smooth data, we ended up with the nondifferentiable optimal value function. Although $v(u)$ is nondifferentiable at zero, it nevertheless possesses directional derivatives in both, positive and negative, directions. Such *directionally differentiable* behavior is typical for the optimal value function. Quite a complete characterization of first-order differentiability properties of the optimal value function can be given when the feasible set is unperturbed, i.e. $\Phi(u) = \Phi$ for all $u \in U$, as shows the following result, essentially due to Danskin [20].

Theorem 4.1 *Let X be a metric space and U be a normed space. Suppose that for all $x \in X$ the function $f(x, \cdot)$ is differentiable, that $f(x, u)$ and $D_u f(x, u)$ are continuous on $X \times U$ and let Φ be a compact subset of X . Then the optimal value function $v(u) := \inf_{x \in \Phi} f(x, u)$ is Hadamard directionally differentiable and*

$$v'(u, d) = \text{Min}_{x \in S(u)} D_u f(x, u)d. \quad (4.1)$$

It follows from (4.1) that $v'(u^0, d)$ is linear in $d \in U$ if $f(\cdot, u^0)$ has a unique minimizer x^0 over Φ . In that case the optimal value function is Hadamard differentiable at u^0 and $Dv(u^0) = D_u f(x^0, u^0)$.

Application to semi-infinite programming. As an example let us briefly outline how the above result can be used to derive first-order optimality conditions in semi-infinite optimization (cf. [21, 49]). Consider the problem

$$\text{Min}_{x \in \mathbb{R}^n} f(x) \quad \text{subject to} \quad g_t(x) \leq 0, \quad t \in T,$$

where $f(\cdot)$ is a continuously differentiable function, T is a compact metric space, the function $g_t(\cdot)$ is differentiable for every $t \in T$, and that $g_t(x)$ and $\nabla g_t(x)$ are continuous jointly in x and t .

Let x^0 be a locally optimal solution of the above semi-infinite problem. Let $T(x^0) = \{t \in T : g_t(x^0) = 0\}$, be the set of active at x_0 constraints. We assume that $T(x^0) \neq \emptyset$, otherwise the problem, at least locally, is unconstrained. Note that then $T(x^0) = \arg \max_{t \in T} g_t(x^0)$. Let us observe that if x^0 is a locally optimal solution of the above problem, then the max-function

$$h(x) := \max \left\{ f(x) - f(x^0), \max_{t \in T} g_t(x) \right\} \quad (4.2)$$

attains its local minimum (equal 0) at x^0 . Setting $g_0(x) := f(x) - f(x^0)$, we may write $h(x) = \max_{t \in T'} g_t(x)$, where $T' = T \cup \{0\}$. This max-function $h(x)$ can be considered as an optimal value function with t being the optimization variable. By Danskin's theorem, $h(x)$ is directionally differentiable and

$$h'(x^0, d) = \max \left\{ d \cdot \nabla f(x^0), \max_{t \in T(x^0)} d \cdot \nabla g_t(x^0) \right\}. \quad (4.3)$$

Optimality of x^0 implies that the directional derivative $\eta(d) := h'(x^0, d)$ is nonnegative for all $d \in \mathbb{R}^n$. By (4.3) the directional derivative $\eta(d)$ is a maximum of linear functions and hence is convex. Therefore $\eta(d) \geq 0$ for all d iff $0 \in \partial\eta(0)$. It also follows from (4.3) that $\eta(\cdot)$ is the support function of the set formed by vectors $\nabla f(x^0)$ and $\nabla g_t(x^0)$, $t \in T(x^0)$, and hence the subdifferential $\partial\eta(0)$ is given by the convex hull of these vectors. (Note that since $T(x^0)$ is compact, this convex hull is also compact and hence is closed.) Together with the optimality condition $0 \in \partial\eta(0)$ this implies that there exist points $t_i \in T(x^0)$, $i = 1, \dots, n$, and (generalized Lagrange) multipliers $\lambda_i \geq 0$, $i = 0, 1, \dots, n$, not all of them zero, such that

$$\lambda_0 \nabla f(x^0) + \sum_{i=1}^n \lambda_i \nabla g_{t_i}(x^0) = 0. \quad (4.4)$$

Moreover, under a constraint qualification, the multiplier λ_0 is not zero and can be taken $\lambda_0 = 1$. Note that Robinson's constraint qualification (3.6) is equivalent here (e.g. [64]) to existence of a vector w such that $w \cdot \nabla g_t(x^0) < 0$ for all $t \in T(x^0)$, which in turn can be considered as an extended Mangasarian-Fromovitz constraint qualification.

It is not true in general that if the max-function $h(x)$ attains its (local) minimum at x_0 , then x_0 is a (locally) optimal solution of the corresponding semi-infinite program. Think, for example, about a case where $g_t(x) = 0$ for all $x \in \mathbb{R}^n$ and all $t \in T$. However, it is not difficult to see that if x_0 is a strict local minimizer of $h(x)$, i.e. $h(x) > 0$ for all $x \neq x_0$ in a neighborhood of x_0 , then x_0 is a strict local minimizer of the corresponding semi-infinite program. Since the space $X = \mathbb{R}^n$ is finite dimensional, it follows from Hadamard directional differentiability of $h(x)$ that a sufficient condition for x_0 to be a strict local minimizer of $h(x)$ is that $h'(x_0, d) > 0$ for all $d \neq 0$. This, in turn, is equivalent to the condition $0 \in \text{int}\{\partial\eta(0)\}$. Therefore we obtain the following first-order sufficient conditions for strict local optimality of x_0 ,

$$0 \in \text{int} \left\{ \text{conv} \left(\nabla f(x^0), \cup_{t \in T(x^0)} \nabla g_t(x^0) \right) \right\}. \quad (4.5)$$

It is possible to conduct a general perturbation analysis using this idea of reduction to composite unconstrained optimization [31].

4.2 Optimal value functions of convex problems

Suppose now that X and U are Banach spaces and that $\Phi(u)$ is defined by cone constraints in the form (1.2) with K being a closed convex cone in the Banach space Y . We say that a mapping $g : X \rightarrow Y$ is convex with respect to a convex closed cone $C \subset Y$ if the inequality

$$g(tx + (1-t)y) \preceq_C tg(x) + (1-t)g(y), \quad (4.6)$$

holds for any $x, y \in X$ and any $t \in [0, 1]$. For example, if $Y = \mathbb{R}^n$, then the mapping $g(x)$ can be written componentwise $g(x) = (g_1(x), \dots, g_n(x))$ and it is convex with respect to the cone $C = \mathbb{R}_+^n$ iff $g_i(x)$, $i = 1, \dots, n$, are convex (real valued) functions. Let us observe that (4.6) means that

$$tg(x) + (1-t)g(y) - g(tx + (1-t)y) \in C,$$

which, by duality, is equivalent to the condition that

$$\langle \lambda, tg(x) + (1-t)g(y) - g(tx + (1-t)y) \rangle \leq 0,$$

for any $\lambda \in C^\circ$. This in turn is equivalent to convexity of $\langle -\lambda, g(\cdot) \rangle$ for any $\lambda \in C^\circ$.

Consider now the unperturbed program

$$\text{Min}_{x \in X} f(x) \quad \text{subject to } G(x) \in K, \quad (P_0) \quad (4.7)$$

with $f(x) = f(x, u^0)$ and $G(x) = G(x, u^0)$. Suppose that the unperturbed program (P_0) is *convex*. That is, the function $f(x)$ is convex and the mapping $G(x)$ is convex with respect to the cone $C = -K$. For example, if $K = \mathbb{R}_-^p$, and hence $C = \mathbb{R}_+^p$, program (P_0) becomes a standard nonlinear convex programming problem. Consider the Lagrangian $L(x, \lambda) = f(x) + \langle \lambda, G(x) \rangle$ of the program (P_0) . We have then that $\text{Max}_{\lambda \in K^\circ} L(x, \lambda)$ is $f(x)$ if $G(x) \in K$, and is $+\infty$ otherwise. Therefore program (P_0) can be written as the min-max problem

$$\text{Min}_{x \in X} \text{Max}_{\lambda \in K^\circ} L(x, \lambda). \quad (4.8)$$

The (Lagrangian) dual of (P_0) is obtained by reversing the order in which min and max are applied, that is

$$\text{Max} \left\{ \psi(\lambda) := \inf_{x \in X} L(x, \lambda) \right\} \quad \text{subject to } \lambda \in K^\circ. \quad (D_0) \quad (4.9)$$

Note that it is always true that $\text{val}(P_0) \geq \text{val}(D_0)$. It is said that there is no duality gap between (P_0) and (D_0) if $\text{val}(P_0) = \text{val}(D_0)$.

Suppose now that $f(x, u) = f(x)$, i.e. the objective function is independent of u , that $U = Y$ and that the mapping $G(x, u)$ is of the form $G(x, u) = G(x) + u$. In that case the program (P_u) becomes

$$\text{Min}_{x \in X} f(x) \quad \text{subject to } G(x) + u \in K. \quad (4.10)$$

Properties of the optimal value function $v(u)$ of the program (4.10) were thoroughly investigated in Rockafellar [57, 58]. Since (P_0) is convex, the optimal value function $v(u)$ is also convex. Let us denote by Λ_0 the set of optimal solutions of the dual program (D_0) . We have the following result due to Rockafellar [58, p.41].

Theorem 4.2 *Suppose that: (i) the program (P_0) is convex, (ii) its optimal value $v(0)$ is finite, and (iii) the feasible set of the program (4.10) is nonempty for all u in a neighborhood of $0 \in Y$. Then: (a) there is no duality gap between (P_0) and its dual (D_0) , (b) the set Λ_0 of optimal solutions of the dual problem is nonempty and bounded, (c) $v(u)$ is continuous at $u = 0$ and $\partial v(0) = \Lambda_0$, (d) $v(u)$ is Hadamard directionally differentiable at $u = 0$ and*

$$v'(0, d) = \text{Max}_{\lambda \in \Lambda_0} \langle \lambda, d \rangle. \quad (4.11)$$

We note that if $G(x)$ is continuously differentiable, then assumption (iii) in the above theorem is equivalent to Robinson's constraint qualification (3.6). Whenever K has a nonempty interior, assumption (iii) is equivalent to the Slater condition, that is, existence of a point $\bar{x} \in X$ such that $G(\bar{x}) \in \text{int}(K)$.

It is important to understand a relation between the set Λ_0 of optimal solutions of (D_0) and the set of Lagrange multipliers of (P_0) . Let $x^0 \in X$ and $\lambda^0 \in Y^*$ be such that

$$x^0 \in \arg \min_{x \in X} L(x, \lambda^0), \quad G(x^0) \in K, \quad \langle \lambda^0, G(x^0) \rangle = 0, \quad \lambda^0 \in K^-. \quad (4.12)$$

Then (x^0, λ^0) is a saddle point of the min-max problem (4.8) and hence $\text{val}(P_0) = \text{val}(D_0)$, and x^0 and λ^0 solve (P_0) and (D_0) , respectively. Moreover, in that case $\lambda \in \Lambda_0$ iff it satisfies the first, third and fourth conditions of (4.12). Note also that if $f(x)$ and $G(x)$ are continuously differentiable then, by convexity, the first condition in (4.12) is equivalent to $D_x L(x^0, \lambda^0) = 0$. We obtain an important result that, in the convex case, if the set of Lagrange multipliers is nonempty for an optimal solution of (P_0) then it is the same for any other optimal solution of (P_0) and coincides with the set of optimal solutions of the dual problem (D_0) .

Example 4.2 *Consider the following optimization problem*

$$\text{Min}_{x \in \mathbb{R}^2} x_1 \quad \text{subject to} \quad -x_1 + e^{x_2} + u \leq 0, \quad (4.13)$$

parameterized by $u \in \mathbb{R}$. This is a convex problem and its dual, for $u = 0$, has only one solution $\lambda = 1$. The Slater condition holds and there is no duality gap between the primal and dual problems here. It can be easily seen that $v(u) = u$ and hence formula (4.11) holds, although the primal problem (4.13) does not possess an optimal solution.

4.3 A general first-order estimate

Let us come back now to the general problem under abstract constraints. Suppose that $f(x, u)$ and $G(x, u)$ are continuously differentiable and let $x^0 \in S(u^0)$. For a given direction

$d \in U$ consider the following linearization of (P_u) ,

$$(PL_d) \quad \underset{h \in X}{\text{Min}} Df(x^0, u^0)(h, d) \quad \text{s.t.} \quad DG(x^0, u^0)(h, d) \in T_K(G(x^0, u^0)).$$

Since $[T_K(G(x^0, u^0))]^- = N_K(G(x^0, u^0))$ it is not difficult to see that the (Lagrangian) dual program of (PL_d) can be written in the form

$$(DL_d) \quad \underset{\lambda \in \Lambda(x^0, u^0)}{\text{Max}} D_u L(x^0, \lambda, u^0)d.$$

It is possible to show by convex analysis that under the directional constraint qualification (3.14) there is no duality gap between (PL_d) and (DL_d) [12, Part I, Prop. 3.1]. The following upper bound (4.14) is then a direct consequence of proposition 3.3 (cf. [12, Part I],[38]).

Proposition 4.1 *Suppose that the directional constraint qualification (3.14) holds in the direction d . Then $\text{val}(PL_d) = \text{val}(DL_d)$ and*

$$v'_+(u^0, d) \leq \text{val}(PL_d). \quad (4.14)$$

Moreover, $\text{val}(PL_d) > -\infty$ if and only if $\Lambda(x^0, u^0) \neq \emptyset$, in which case the set of optimal solutions of (DL_d) is a nonempty weak* compact subset of $\Lambda(x^0, u^0)$.

A natural question arises, whether the upper bound given in (4.14) is tight in the sense that $v(u)$ is directionally differentiable at u^0 in the direction d and the directional derivative is equal to $\text{val}(DL_d)$? As the following example (due to Gauvin and Tolle [25, pp. 308-309]) shows, however, this may be not the case and the above question has a negative answer.

Example 4.3 *Let $X = \mathbb{R}^2$, $U = \mathbb{R}^2$ and consider the problem of minimization of $f(x) = -x_2$ subject to $x_2 + x_1^2 \leq u_1$ and $x_2 - x_1^2 \leq u_2$. For $u^0 = (0, 0)$ this program has the unique optimal solution $x^0 = (0, 0)$. The Mangasarian-Fromovitz constraint qualification holds here and the (bounded) Lagrange multipliers set is given by*

$$\Lambda(x^0, u^0) = \{(\lambda_1, \lambda_2) : \lambda_1 + \lambda_2 = 1, \lambda_1 \geq 0, \lambda_2 \geq 0\}.$$

Consider the direction $d = (1, 0)$. Then for $u = td$, $t \geq 0$, this program has two optimal solutions $((t/2)^{1/2}, t/2)$ and $(-(t/2)^{1/2}, t/2)$ and the optimal value function $v(td) = -t/2$, and hence $v'(u^0, d) = -1/2$. On the other hand,

$$\text{val}(PL_d) = \text{val}(DL_d) = \max_{\lambda \in \Lambda(x^0, u^0)} D_u L(x^0, \lambda, u^0)d = 0.$$

As we shall see later lower bounds for the directional derivatives of $v(u)$ are based on a second-order analysis. It is still possible, however, to obtain directional derivatives of $v(u)$ by a first-order analysis in the convex case. Suppose now that for $u = u^0$ the corresponding program (P_0) is convex. That is, the function $f(\cdot, u^0)$ is convex and the mapping $G(\cdot, u^0)$

is convex with respect to the cone $C = -K$. Recall that this implies that the Lagrangian $L(\cdot, \lambda, u^0)$ is convex for any $\lambda \in K^-$. As we have seen, in the convex case the set $\Lambda(x^0, u^0)$ is the set of optimal solutions of the dual program of (P_0) and hence is independent of a particular $x^0 \in S(u^0)$. Therefore we can omit x^0 and denote this set by $\Lambda(u^0)$. We now give an extension of Gol'shtein's Theorem [29] to the Banach space setting. This result is easily proved combining the techniques of [12, Prop. 3.2, Part I] and [65].

Theorem 4.3 *Suppose that: (i) the problem (P_0) is convex, (ii) the optimal set $S(u^0)$ is nonempty and compact, (iii) the directional regularity condition holds for all $x^0 \in S(u^0)$, and (iv) for sufficiently small $t > 0$ the program (P_{u^0+td}) possesses an $o(t)$ -optimal solution $x(t)$ such that $\text{dist}(x(t), S(u^0)) \rightarrow 0$ as $t \rightarrow 0^+$. Then the optimal value function is directionally differentiable at u^0 in the direction d and*

$$v'(u^0, d) = \inf_{x \in S(u^0)} \sup_{\lambda \in \Lambda(u^0)} D_u L(x, \lambda, u^0)d. \quad (4.15)$$

Recall that if the Slater condition, for the program (P_0) , is satisfied, then the constraint qualification (3.6) (and hence the directional regularity condition (3.14)) holds for all $x^0 \in S(u^0)$ and the set $\Lambda(u^0)$ of Lagrange multipliers is nonempty and bounded. In that case $v(u)$ is directionally differentiable at u^0 (in all directions) in the *Hadamard* sense.

Consider, for example, a parameterized semi-infinite program with $X = \mathbb{R}^n$ and the feasible set $\Phi(u)$ given in the form (1.4). In this case the problem (P_0) is convex if the functions $f(\cdot, u^0)$ and $g_t(\cdot, u^0)$, $t \in T$, are convex. Slater condition is equivalent here to existence of a point $\bar{x} \in \mathbb{R}^n$ such that $g_t(\bar{x}, u^0) < 0$ for all $t \in T$ and assumptions (ii) and (iv) of Theorem 4.3 can be easily ensured by compactness arguments (cf. [65]).

Consider the linearization (PL_d) of the parameterized (not necessarily convex) problem (P_u) . Suppose that the optimal value of (PL_d) is finite and let h be an ε -optimal solution of (PL_d) . (Note that it can happen that the linearized program (PL_d) does not possess an optimal solution even if its optimal value is finite.) Then it follows from Proposition 3.3 that there exists a path $x(t) = x^0 + th + o(t) \in \Phi(u^0 + td)$ and hence

$$v(u^0 + td) - v(u^0) \leq f(x(t), u^0 + td) - f(x^0, u^0) \leq t \text{val}(PL_d) + t\varepsilon + o(t).$$

Therefore equality $v'(u^0, d) = \text{val}(DL_d)$ will imply existence, for any $\varepsilon > 0$, of $(t\varepsilon)$ -optimal solution $x(t)$ of (P_{u^0+td}) such that $\|x(t) - x^0\| = O(t)$. It is possible to show that in a sense the converse of that is also true, i.e. existence of Lipschitz stable at x^0 , nearly optimal, solutions of (P_{u^0+td}) implies $v'(u^0, d) = \text{val}(DL_d)$, [38, 39, 40].

Let us finish this section by giving the following (first-order) result which holds for general (not necessarily convex) programs. In finite dimensional setting it was given in [25, 26] and extended to Banach spaces in [38].

Theorem 4.4 *Suppose that: (i) $S(u^0) \neq \emptyset$, (ii) Robinson's constraint qualification holds at every $x \in S(u^0)$, and (iii) for every $\varepsilon > 0$ there exists a $t\varepsilon$ -optimal solution of (P_{u^0+td})*

converging to a point of the optimal set $S(u^0)$. Then for every $d \in U$,

$$\begin{aligned} \inf_{x \in S(u^0)} \inf_{\lambda \in \Lambda(x, u^0)} D_u L(x, \lambda, u^0) d &\leq v'_-(u^0, d) \\ &\leq v'_+(u^0, d) \leq \inf_{x \in S(u^0)} \sup_{\lambda \in \Lambda(x, u^0)} D_u L(x, \lambda, u^0) d. \end{aligned} \quad (4.16)$$

In particular, if in addition $\Lambda(x, u^0) = \{\bar{\lambda}(x)\}$ is a singleton for every $x \in S(u^0)$, then $v(u)$ is directionally differentiable at u^0 and

$$v'(u^0, d) = \inf_{x \in S(u^0)} D_u L(x, \bar{\lambda}(x), u^0) d. \quad (4.17)$$

Note that if the optimal set $S(u^0)$ is compact, then assumption (iii) in the above theorem can be replaced by the assumption of existence of an $o(t)$ -optimal solution $x(t)$ of (P_{u^0+td}) such that $\text{dist}(x(t), S(u^0))$ tends to zero as $t \rightarrow 0^+$. A discussion of conditions ensuring uniqueness of Lagrange multipliers under cone constraints is given in [66].

5 Strong stability

5.1 Strong stability for generalized equations

Consider the parameterized problem (P_u) with the corresponding feasible set defined by abstract constraints in the form (1.2). We assume in this section that X , Y and U are Banach spaces, that K is a closed convex subset of Y , and that $f(x, u)$ and $G(x, u)$ are twice continuously differentiable (jointly in x and u). We may assume, without loss of generality, that the unperturbed problem is formulated for $u = 0$. The approach followed in this section is to conduct a perturbation analysis by expanding the associated optimality system representing the corresponding first-order necessary conditions

$$D_x L(x, \lambda, u) = 0, \quad \lambda \in N_K(G(x, u)). \quad (5.1)$$

It is advantageous at this point to imbed the above optimality system into the following abstract problem

$$F(z, u) \in N(z), \quad (5.2)$$

where $F : Z \times \mathbb{R} \rightarrow W$, Z and W are Banach spaces, and N is a closed multifunction $N : Z \rightarrow 2^W$. Relation (5.2) is called a *generalized equation*. Using N_K^{-1} , the graph inverse of N_K , we may write (5.1) as

$$(D_x L(x, \lambda, u), G(x, u)) \in (0, N_K^{-1}(\lambda)), \quad (5.3)$$

which is a particular case of (5.2) with $z = (x, \lambda) \in X \times Y^*$. Note that if K is a convex closed cone, then $N_K(y)$ is given by $\{\alpha \in K^- : \langle \alpha, y \rangle = 0\}$, if $y \in K$, and is the empty set if $y \notin K$, and that

$$N_K^{-1}(\lambda) = N_{K^-}(\lambda) = \begin{cases} y \in K : \langle \lambda, y \rangle = 0, & \text{if } \lambda \in K^-, \\ \emptyset, & \text{otherwise.} \end{cases} \quad (5.4)$$

Suppose that for $u = 0$ the generalized equation (5.2) has a solution z^0 . Since we know how to linearize the mapping F , and not the multifunction N , we may try to approximate the solution $\bar{z} = \bar{z}(u)$ of (5.2) by $z^0 + z^1(u)$, where $z^1(u)$ is a solution of the following generalized equation (with unknown ζ), representing partial linearization of (5.2),

$$F(z^0, 0) + DF(z^0, 0)(\zeta, u) \in N(z^0 + \zeta). \quad (5.5)$$

The “first-order term” $z^1(u)$, if it exists, is in general a nonlinear function of u . This is a price to pay for the full generality of the framework (5.2). Therefore we call $z^0 + z^1(u)$ a *pseudo expansion* of the solution $\bar{z}(u)$.

Definition 5.1 *Let, for $u = 0$, z^0 be a solution of the generalized equation (5.2). We say that the strong stability condition holds (that z^0 is a strongly stable solution), if there exist $\varepsilon > 0$ and $M > 0$ such that for all $w \in B_W(0, \varepsilon)$ the linearized generalized equation*

$$F(z^0, 0) + D_z F(z^0, 0)z + w \in N(z^0 + z) \quad (5.6)$$

has a unique solution $\hat{z}(w) \in B_Z(0, M)$, and $\hat{z}(\cdot) : B_W(0, \varepsilon) \rightarrow B_Z(0, M)$ is Lipschitz continuous.

The above strong stability condition, introduced by Robinson [54], essentially states that the linearized generalized equation is well posed, for arbitrary but small perturbations. The following result is due to Robinson [54] (the result stated there is for the case when the multifunction $N(\cdot)$ is a certain normal cone, but the proof goes without modification for any closed multifunction).

Theorem 5.1 *Suppose that, for $u = 0$, z^0 is a solution of the generalized equation (5.2), and that the strong stability condition holds. Then, for all u in a neighborhood of 0, the mappings $\bar{z}(u)$ and $z^1(u)$ are well defined in the vicinity of z^0 and in $B_Z(0, M)$, respectively. In addition, $\bar{z}(u)$ is Lipschitz continuous, $z^1(u) = O(\|u\|)$, and $\bar{z}(u) = z^0 + z^1(u) + o(\|u\|)$.*

The above theorem gives effective means to check well-posedness and Lipschitz stability of $\bar{z}(u)$. It does not give, however, any formula for the directional derivative of $\bar{z}(u)$, as it may happen that $t^{-1}z^1(tu)$ has no limit when $t \downarrow 0$.

If the data have enough regularity, then we may write a higher-order pseudo expansion along the same lines [16]. Suppose, for instance, that the mapping F is twice continuously differentiable, and z^0 is a strongly stable solution of (5.2). Then $\bar{z}(u) = z^0 + z^1(u) + \frac{1}{2}z^2(u) + o(\|u\|^2)$, where $z^1(u)$ is as above and $z^2(u)$ is a solution of the following linear generalized equation (with unknown ζ) obtained by using a second-order expansion of $F(z(u), u)$:

$$F(z^0, 0) + DF(z^0, 0)(z^1(u), u) + \frac{1}{2} [D_z F(z^0, 0)\zeta + D^2 F(z^0, 0)(z^1(u), u)(z^1(u), u)] \in N(z^0 + z^1(u) + \frac{1}{2}\zeta).$$

5.2 Strongly stable solutions of the optimality system

When applying Theorem 5.1 to the optimality system (5.1), two questions arise. The first is to establish a relation between the locally optimal solutions of (P_u) and solutions of the optimality system, under the strong stability condition. The second is to give necessary and/or sufficient conditions for strong regularity.

The result below deals with the first question. The only delicate part of the proof consists in checking that the perturbed optimization problem has a solution. The argument uses Ekeland's variational principle [5]; it deals with the scalar perturbation case, but the argument is valid in the case when U is a Banach space.

Proposition 5.1 [16, Thm 4.6] *If (x^0, λ^0) is a strongly stable solution of (5.1), then there exists a neighborhood N of x^0 such that, for all u sufficiently close to $0 \in U$, the perturbed problem (P_u) has a unique local minimizer $\bar{x}(u) \in N$ and an associated unique Lagrange multiplier $\bar{\lambda}(u)$. It follows that $(\bar{x}(u), \bar{\lambda}(u))$ is the locally unique solution of (5.1).*

We now discuss a relation between strong stability, constraint qualifications and second-order conditions. A necessary condition for strong stability is that the regularity condition (3.6) holds ([16, Lemma 4.3]). Below is given a sufficient condition for strong regularity ([16, Theorem 4.9]), which reduces in a finite dimensional setting to Theorem 4.1 in [54].

Proposition 5.2 *Sufficient conditions for strong stability are: (i) $D_x G(x^0, 0) : X \rightarrow Y$ is onto, and (ii) there exists $\alpha > 0$ such that*

$$D_{xx}^2 L(x^0, \lambda^0, 0)(h, h) \geq \alpha \|h\|^2, \quad \forall h \in X \text{ satisfying } D_x G(x^0, 0)h \in \text{Sp}(K). \quad (5.7)$$

The expansion of the solution and multiplier has a nice interpretation. Expanding $\bar{x}(u)$ and $\bar{\lambda}(u)$ as $\bar{x}(u) = x^0 + x^1(u) + o(\|u\|)$ and $\bar{\lambda}(u) = \lambda^0 + \lambda^1(u) + o(\|u\|)$, respectively, and specializing (5.5) to the linearized optimality system of (P_u) , we obtain that $x^1(u)$ and $\lambda^1(u)$ are solutions of the generalized equation with unknown $(\chi, \lambda) \in X \times Y^*$:

$$\begin{aligned} D_x f(x^0, 0) + D_{(x,u)} [D_x L(x^0, \lambda^0, 0)](\chi, u) + [D_x G(x^0, 0)]^*(\lambda^0 + \lambda) &= 0, \\ G(x^0, 0) + DG(x^0, 0)(\chi, u) &\in N_K^{-1}(\lambda^0 + \lambda). \end{aligned}$$

Observe that the above (generalized) equations represent the first-order optimality system for the problem

$$\begin{aligned} \text{Min}_{h \in X} \quad & Df(x^0, 0)(h, u) + \frac{1}{2} D_{(x,u)^2}^2 L(x^0, \lambda^0, 0)((h, u), (h, u)), \\ \text{s.t.} \quad & G(x^0, 0) + DG(x^0, 0)(h, u) \in K. \end{aligned} \quad (Q_u)$$

It can be easily checked that conditions for strong stability of (P_u) and (Q_u) are identical. Consequently, if (x^0, λ^0) is a strongly stable solution of (P_0) , then (locally) the optimality system of (Q_u) has the unique solution $(x^1(u), \lambda^0 + \lambda^1(u))$, and $x^1(u)$ is a local minimizer of (Q_u) . A practical consequence of this result is that, whenever u is sufficiently close to 0, we may compute $x^1(u)$ by applying any locally convergent algorithm for solving (Q_u) , starting from the point x^0 (or (x^0, λ^0) for a primal-dual algorithm).

5.3 Polyhedral case

In the case of scalar perturbations, i.e. $u(t) = td$, with $d \in U$ and $t \geq 0$, and if $\Phi(u)$ is of the form (1.3), then the pseudo-expansion appears to be a Taylor expansion. That is, the coefficients of the pseudo expansion of $(\bar{x}(td), \bar{\lambda}(td))$ of order k are, when t is small enough, of the form $t^k(\hat{x}^k, \hat{\lambda}^k)$ for some $(\hat{x}^k, \hat{\lambda}^k) \in X \times Y^*$. Let us verify this for the first-order term. Recall that $I(x, u)$ denotes the set of active at (x, u) inequality constraints. For u small enough, only the inequality constraints in $I(x^0, 0)$ may be active for local solutions of (Q_u) . Therefore, by setting $\hat{h} := h/t$, we can write an equivalent of (Q_u) in the form

$$\begin{aligned} \text{Min}_{\hat{h} \in X} \quad & Df(x^0, 0)(\hat{h}, d) + \frac{1}{2}D^2L(x^0, \lambda^0, 0)((\hat{h}, d), (\hat{h}, d)), \\ \text{s.t.} \quad & Dg_i(x^0, 0)(\hat{h}, d) = 0, \quad i = 1, \dots, q, \\ & Dg_i(x^0, 0)(\hat{h}, d) \leq 0, \quad i \in I(x^0, 0). \end{aligned} \quad (\hat{Q})$$

As the data of (\hat{Q}) do not depend on t , the unique associated solution $x^1(td)/t$ is also independent of u . Therefore, under the strong stability condition, we obtain that $\bar{x}(td)$ is directionally differentiable at $u = 0$ and the corresponding directional derivatives $\bar{x}'(0, d)$ and $\bar{\lambda}'(0, d)$ are given by the optimal solution of the program (\hat{Q}) (cf. [56, 59]).

If in addition the data of (P_u) are analytic (i.e. locally the involved function can be expanded in power series), then for $t > 0$ small enough $\bar{x}(u)$ and $\bar{\lambda}(u)$ are themselves analytic functions ([16, Theorem 4.13]).

In the case of finitely many inequality constraints another interesting result is that we know several characterizations for strong stability [22]. One of them reduces to the sufficient condition of Proposition 5.2, if we observe that a) only active constraints at x^0 should be taken into account and b) those inequality constraints associated with a positive multiplier will remain active for the perturbed problem, and we do not change the solution nor the multiplier by setting these as equalities. Define

$$J := \{1, \dots, q\} \cup I(x, u); \quad J^0 := \{1, \dots, q\} \cup \{i = q + 1, \dots, p; \lambda_i^0 > 0\}.$$

By G_J we denote the restriction of G to components in J . We have

Proposition 5.3 *Let $\Phi(u)$ be given by a finite number of constraints in the form (1.3). Then (x^0, λ^0) is a strongly stable solution if and only if $D_x G_J(x^0, 0)$ is onto, and there exists $\alpha > 0$ such that*

$$D_{xx}^2 L(x^0, \lambda^0)(h, h) \geq \alpha \|h\|^2, \quad \forall h \in X, \quad D_x g_i(x^0, 0)h = 0, \quad i \in J^0. \quad (5.8)$$

The above Theorem is a consequence of [33, 36, 54]; see [16] for a simple proof, and [35, 37] for related results. The strong regularity approach is applied to the sensitivity analysis of optimal control problems in [43]. In a nonlinear programming setting, this approach is extended to a case when the linearized optimality system has a finite number of solutions in [9].

6 Quantitative stability of optimal solutions

We assume in this section that X and U are Banach spaces and discuss continuity properties of optimal solutions of the problem (P_u) from a quantitative point of view. In particular, we will be interested in Hölder and Lipschitzian behavior of the optimal-solution multifunction $S(\cdot) : U \rightarrow 2^X$. We say that $S(u)$ is *Hölder stable* of degree α ($\alpha > 0$), at a point $u^0 \in U$, if there exists a constant $\kappa > 0$ such that

$$\sup_{x \in S(u)} \text{dist}(x, S(u^0)) \leq \kappa \|u - u^0\|^\alpha \quad (6.1)$$

for all u in a neighborhood of u^0 . If $S(u)$ is Hölder stable of degree $\alpha = 1$, i.e. $S(u)$ is upper Lipschitzian at u^0 , we say that $S(u)$ is *Lipschitz stable*. We also discuss Hölder and Lipschitz stability of ε -optimal solutions of (P_u) .

Example 6.1 Consider $f(x, u) = x^{2m}/(2m) - ux$, $u \in \mathbb{R}$, m is a positive integer, and let $\Phi(u) = [-1, 1]$ for all $u \in \mathbb{R}$. Then $S(u) = \{u^{1/(2m-1)}\}$ is a singleton for any $u \in (-1, 1)$. Therefore, $S(u)$ is not Lipschitz stable at $u = 0$ if $m > 1$.

The above example demonstrates that in order to establish Lipschitzian behavior of optimal solutions we need some additional assumptions beside continuity and first-order differentiability properties of the involved functions. It turns out that quantitative stability of $S(u)$ is closely related to a second-order analysis of (P_u) . Let us start by introducing the following second-order abstract condition. Consider the unperturbed program (P_0) , and suppose that its optimal solution set $S_0 = S(u^0)$ is not empty. Denote $\Phi_0 = \Phi(u^0)$ and $f_0 = v(u^0) = \inf_{x \in \Phi_0} f(x)$, where $f(x) = f(x, u^0)$.

Definition 6.1 We say that the growth condition of order $\alpha > 0$ (for the program (P_0) , in a neighborhood N of the optimal set S_0) holds, if there exists a constant $c > 0$ such that

$$f(x) \geq f_0 + c [\text{dist}(x, S_0)]^\alpha, \quad \forall x \in \Phi_0 \cap N. \quad (6.2)$$

When $\alpha = 1$ (resp. 2) we speak of the first-order (resp. second-order) growth condition.

If the set S_0 is finite, then the second-order growth condition is strongly connected to certain second-order sufficient conditions discussed below. In case S_0 has non isolated connected parts, verification of (6.2) may be tricky. If the space X is finite dimensional and Φ_0 is defined by a finite number of constraints, then we know a characterization of the second-order growth condition in terms of the Hessian of Lagrangian and qualification conditions for convex problems [13], whereas for nonconvex problems, we know only some necessary or sufficient conditions [14].

Under (6.2), it is relatively easy to give, quite general, sufficient conditions for Lipschitzian stability of $S(u)$ when the feasible set $\Phi(u) \equiv \Phi_0$ is unperturbed, i.e. independent of u .

Proposition 6.1 [63] *Suppose that $\Phi(u) \equiv \Phi_0$, that the second-order growth condition (6.2) holds in a neighborhood N of the optimal set S_0 , that the difference function $f(\cdot, u) - f(\cdot, u^0)$ is Lipschitz continuous on N modulus $\ell(u)$, and let $\bar{x} \in N$ be an ε -optimal solution of (P_u) . Then*

$$\text{dist}(\bar{x}, S(u^0)) \leq c^{-1}\ell(u) + c^{-1/2}\varepsilon^{1/2}. \quad (6.3)$$

In particular if, in addition, $f(x, u)$ is continuously differentiable in x with $D_x f(x, \cdot)$ being Lipschitz continuous on N , with the corresponding Lipschitz constant independent of $x \in N$, then $S(u)$ is Lipschitz stable at u^0 .

We now discuss some necessary and/or sufficient conditions for quadratic growth, assuming $\Phi(u)$ to be defined by constraints. A central object is the *critical cone*, defined as follows:

$$C_0 = \{h \in X : D_x G(x^0)h \in T_K(G(x^0)), D_x f(x^0)h \leq 0\}. \quad (6.4)$$

The critical cone C_0 represents those directions for which the linearization of cost function and constraints does not provide information about optimality of x^0 . We may interpretate the “feasibility relation” $d \in C_0$ as the optimization problem $\text{Min}_{h \in X} 0$ s.t. $d \in C_0$. Computing the dual of this problem, we find the problem of maximizing 0 over the set of generalized Lagrange multipliers. In that sense, the critical cone is dual to the set of generalized Lagrange multipliers. In particular, if d is a critical direction and (λ_0, λ) is a generalized Lagrange multiplier, then (this is also a consequence of (3.4)) we have $\lambda_0 D_x f(x^0, u^0)h = 0$ and $\langle \lambda, D_x G(x^0)h \rangle = 0$. Therefore, if the set of Lagrange multipliers is non empty, then every critical direction h satisfies $D_x f(x^0, u^0)h = 0$.

The relation between the critical cone and the set of Lagrange multipliers is closely related to the following linearization of the problem (P_0) :

$$\text{Min}_{h \in X} Df(x^0)h \text{ s.t. } DG(x^0)h \in T_K(G(x^0)). \quad (6.5)$$

As for any problem of minimization of a linear function over a cone, the optimal value of this problem is either 0 or $-\infty$. The dual (in the sense of convex analysis) of (6.5) is $\text{Max } 0$ subject to $\lambda \in \Lambda(x^0)$, i.e. it consists in maximizing 0 over the (possibly empty) set of Lagrange multipliers. Therefore the optimal value of the dual problem is also either 0 or $-\infty$, and is equal to 0 iff the set of Lagrange multipliers is non empty. Consequently there is no duality gap between these problems if there exist Lagrange multipliers.

The case when $C_0 = \{0\}$ is a very specific situation, strongly related to the first-order growth condition as the following statement shows.

Proposition 6.2 *Suppose that $S(u^0) = \{x^0\}$. Then: (i) if Robinson’s constraint qualification and the first-order growth condition hold at x^0 , then $C_0 = \{0\}$, (ii) if the feasible set is finitely constrained and $C_0 = \{0\}$, then the first-order growth condition holds at x^0 .*

For a constant $\eta > 0$, consider the cone

$$C_\eta = \{h \in X : \text{dist}(DG(x^0)h, T_K(G(x^0))) \leq \eta\|h\|, Df(x^0)h \leq \eta\|h\|\}, \quad (6.6)$$

which we call the *approximate critical cone*. The notation is consistent, in the sense that if $\eta = 0$, then C_η reduces to C_0 . It can be easily verified that if $\{x^k\}$ is a sequence of feasible points converging to x^0 such that $f(x^k) \leq f(x^0) + o(\|x^k - x^0\|)$, then for any $\eta > 0$ the associated normalized directions $d^k := (x^k - x^0)/\|x^k - x^0\|$ belong to C_η for k large enough. This explains the relevance of the approximate critical cones in second-order analysis.

Definition 6.2 *We say that second-order sufficient conditions hold at x^0 if there exist $\eta > 0$, $\beta > 0$ and a bounded set $\Omega \subset \Lambda^g(x^0)$ (of generalized Lagrange multipliers) such that*

$$\sup_{(\lambda_0, \lambda) \in \Omega} D_{xx}L^g(x^0, \lambda_0, \lambda, u^0)(h, h) \geq \beta\|h\|^2, \quad \forall h \in C_\eta. \quad (6.7)$$

Theorem 6.1 ([45], [12, Prop. 6.1, Part I and Prop. 3.6, Part II]) *The second-order sufficient conditions (6.7) imply the second-order growth condition, for the problem (P_0) at the point x^0 . Both conditions are equivalent if the feasible set is finitely constrained and the Mangasarian-Fromovitz constraint qualification holds.*

There are situations where it is possible to state some second-order sufficient optimality condition that are more specific. Since $\cap_{\eta>0} C_\eta = C_0$, it is possible to show by compactness arguments that if X is a finite-dimensional space, then the approximate critical cone C_η can be replaced by C_0 . The second case is when Robinson's constraint qualification holds. Then it can be shown that every $d \in C_\eta$ is at distance $O(\eta)$ of the smaller cone

$$\hat{C}_\eta = \{h \in X : D_x G(x^0)h \in T_K(G(x^0)), D_x f(x^0)h \leq \eta\|h\|\}. \quad (6.8)$$

We may then replace C_η by \hat{C}_η and choose $\Omega = \{1\} \times \Lambda(x^0)$, i.e. to use the Lagrangian instead of the generalized Lagrangian, since in that case $\Lambda(x^0)$ is bounded.

Note also that $T_K(G(x^0))$ is a subset of the topological closure of the linear space $\text{Sp}(K)$, generated by K , and hence (6.7) is weaker than the second-order conditions (5.7), even when $\Lambda(x^0) = \{\lambda^0\}$ is a singleton.

In the case of an unconstrained optimization problem, the second-order conditions may be motivated as follows. If x^0 is solution of (P_0) , then $D_x f(x^0) = 0$ and $D_{xx}f(x^0)(h, h) \geq 0$ for all $h \in X$. Moreover, if the second order sufficient condition $D_{xx}f(x^0)(h, h) \geq \beta\|h\|^2$ (for some $\beta > 0$ and all $h \in X$) holds, then $\|h\|_1 := (D_{xx}f(x^0)(h, h))^{1/2}$ defines a norm on X which is equivalent to the original norm of X . Therefore in that case X is Hilbertisable, i.e. there exists a norm that is equivalent to the one of X and for which X is a Hilbert space. We end up with a surprising consequence. If X is a non Hilbertisable Banach space, then the second-order growth condition can never hold at a unique point x_0 for an unconstrained optimization of a smooth function! Similarly, if the space X is non Hilbertisable, then $D_{xx}^2L(x^0, \lambda)(h, h)$ cannot be positive definite for any Lagrange multiplier λ . This of course does not prevent the second-order growth condition to hold for a constrained problem.

When the feasible set is defined by the abstract constraints, the second-order sufficient conditions of definition 6.2 are rather crude since they do not take into account the curvature of the boundary of K . Indeed, by Theorem 6.1, they imply that quadratic growth holds

at x^0 for the problem in which we replace constraints $G(x) \in K$ by constraints $G(x) \in G(x^0) + T_K(G(x^0))$, where the set K is “linearized” at the point $G(x^0)$. Second-order necessary conditions, that take into account the curvature of K , have been obtained in [19] based on an approach in [34]. In a finite dimensional setting, under the assumption of second order regularity introduced in [12], it is possible to formulate second order sufficient conditions that close the gap with the corresponding second order necessary conditions [17].

6.1 Hölder and Lipschitz estimates

Example 4.3 shows that under the hypotheses of regularity of constraints and some second-order sufficient conditions, one may hope to establish Hölder stability of optimal solutions of degree $\alpha = 1/2$ at most. Such Hölder stability, in various frameworks of generality, was established in a number of publications [1, 2, 12, 15, 27, 63].

Proposition 6.3 [63] *Suppose that: (i) the second-order growth condition holds in a neighborhood N of x^0 , (ii) $f(\cdot, u)$ is Lipschitz continuous on N with the Lipschitz constant independent of u for all u in a neighborhood of u^0 , (iii) the difference function $f(\cdot, u) - f(\cdot, u^0)$ is Lipschitz continuous on N modulus $\ell(u) = O(\|u - u^0\|)$, (iv) the multifunction $u \rightarrow \Phi(u) \cap N$ is upper Lipschitzian at u^0 and $\text{dist}(x^0, \Phi(u)) = O(\|u - u^0\|)$. Then for any $\varepsilon(u)$ -optimal solution $\bar{x}(u) \in N$, with $\varepsilon(u) = O(\|u - u^0\|)$, it follows that*

$$\|\bar{x}(u) - x^0\| = O(\|u - u^0\|^{1/2}). \quad (6.9)$$

In case $f(x, u)$ is sufficiently smooth, assumptions (ii) and (iii) in the above proposition can be easily ensured by appropriate bounds on the corresponding derivatives of f . If $\Phi(u)$ is defined by abstract constraints, assumption (iv) holds under Robinson’s constraint qualification as it follows from the Robinson-Ursescu stability theorem.

Now for a given $d \in U$ consider the set $\Lambda_1(x^0, d)$ of optimal solutions of the problem (DL_d) , defined in section 4.3. Let us consider the following strengthened form of second-order sufficient conditions: there exist constants $\beta > 0$ and $\eta > 0$ such that

$$\text{Max}_{\lambda \in \Lambda_1(x^0, d)} D_{xx}^2 L(x^0, \lambda)(h, h) \geq \beta \|h\|^2, \quad \forall h \in C_\eta. \quad (6.10)$$

Since $\Lambda_1(x^0, d) \subset \Lambda(x^0)$, we have that condition (6.10) is stronger than (6.7). Also condition (6.10) depends on the direction $d \in U$. In a finite dimensional setting condition (6.10) was introduced in [60].

Proposition 6.4 [6, 15, 60] *Suppose that the feasible set $\Phi(u)$ is defined by a finite number of constraints in the form (1.3) and that, for a given $d \in U$, Gollan’s regularity condition (3.13) is satisfied. Let $\bar{x}(t)$, $t \geq 0$, be an $\varepsilon(t)$ -optimal solution of (P_{u^0+td}) with $\varepsilon(t) = O(t^2)$ and such that $\bar{x}(t) \rightarrow x^0$ as $t \rightarrow 0^+$. Then: (a) under the second-order sufficient conditions (6.7), Hölder stability $\|\bar{x}(t) - x^0\| = O(t^{1/2})$, $t \geq 0$, holds (b) under the second-order sufficient conditions (6.10), Lipschitz stability $\|\bar{x}(t) - x^0\| = O(t)$, $t \geq 0$, holds.*

It turns out that in the general case, where the set K is not polyhedral, regularity condition (3.6) and second-order sufficient conditions alone do not guarantee Lipschitz stability of optimal solutions.

Example 6.2 [67] *Consider the linear space $Y = \mathcal{S}_2$ of 2×2 symmetric matrices and the cone $K \subset \mathcal{S}_2$ of 2×2 positive semidefinite matrices. Consider the linear mapping $G : \mathbb{R}^2 \times \mathcal{S}_2 \rightarrow \mathcal{S}_2$ given by $G(x_1, x_2, A) := \text{diag}(x_1, x_2) + A$, where $\text{diag}(x_1, x_2)$ denotes the diagonal matrix with diagonal elements x_1 and x_2 , and the parameterized problem*

$$\min_{x \in \mathbb{R}^2} x_1 + x_1^2 + x_2^2 \quad \text{subject to } G(x_1, x_2, A) \in K. \quad (P_A) \quad (6.11)$$

For $A = 0$ the feasible set of the corresponding (unperturbed) problem is \mathbb{R}_+^2 and its optimal solution is $x_0 = (0, 0)$. Also we have that the Slater condition holds here and the unperturbed problem has unique Lagrange multiplier $\lambda_0 = \text{diag}(-1, 0)$. Moreover, the problem (6.11) is convex and the Hessian of Lagrangian is positive definite, on the whole space \mathbb{R}^2 , i.e. the strongest form of second-order sufficient conditions holds. However, it is not difficult to verify here that for any nondiagonal matrix $A \in \mathcal{S}_2$, the problem (P_{tA}) , $t \geq 0$, has a unique optimal solution $\bar{x}(t) = (\bar{x}_1(t), \bar{x}_2(t))$ with $\bar{x}_2(t)$ is of order of $t^{2/3}$ as $t \rightarrow 0^+$. Therefore the optimal solution $\bar{x}(t)$ is not Lipschitz stable along any nondiagonal matrix-direction A .

Various additional conditions, ensuring Lipschitz stability of optimal solutions, were suggested in the literature (see [67] for a survey). The following condition was suggested in [64].

Assumption A The linearized problem (PL_d) , has an optimal solution $\hat{h} = \hat{h}(d)$ such that for $t > 0$,

$$\text{dist} \left(G(x^0, u^0) + tDG(x^0, u^0)(\hat{h}, d), K \right) = O(t^2). \quad (6.12)$$

Note that the constraint $DG(x^0, u^0)(h, d) \in T_K(G(x^0, u^0))$, of the linearized problem (PL_d) , means that the left hand side of equation (6.12) is of order $o(t)$. Assumption A is stronger and requires it to be of order $O(t^2)$.

Under the directional regularity condition (3.14) and assumption A, the following upper bound for the optimal value function holds [12, 64],

$$v(u^0 + td) \leq v(u^0) + t \text{val}(PL_d) + O(t^2). \quad (6.13)$$

Note that the above inequality is stronger than the upper bound (4.14) of Proposition 4.1. Together with the second-order conditions (6.10), the upper bound (6.13) imply (directional) Lipschitz stability of the optimal solutions.

Theorem 6.2 [12, 64, 68] *Suppose that the directional regularity condition (3.14), assumption A and second-order conditions (6.10) hold, and let $\bar{x}(t)$, $t \geq 0$, be an $\varepsilon(t)$ -optimal solution of (P_{u^0+td}) , with $\varepsilon(t) = O(t^2)$, converging to x^0 as $t \rightarrow 0^+$. Then $\|\bar{x}(t) - x^0\| = O(t)$.*

It is possible to show that if the space X is reflexive, then existence of an optimal solution of the linearized problem (PL_d) is a necessary condition for directional Lipschitz stability of optimal solutions of (P_u) (see [12, Proposition 3.3], [67]). It is not difficult to see that this condition does not hold in the example 6.2 for any nondiagonal matrix-direction A .

The above result does not say anything about continuity properties of the associated Lagrange multipliers. In fact it may happen that under the assumptions of Theorem 6.2 the corresponding Lagrange multipliers are not continuous at u^0 even if $\Lambda(x^0, u^0)$ is a singleton (this may happen, for example, in semi-infinite optimization). Therefore, in order to establish Lipschitz stability of $(\bar{x}(u), \bar{\lambda}(u))$, a regularity condition, stronger than assumption A, is required. Let K be a closed convex cone and for $\lambda \in K^\circ$ consider the cone $K(\lambda) := \{y \in K : \langle \lambda, y \rangle = 0\}$.

Assumption B Let K be a closed convex cone and $\lambda^0 \in \Lambda(x^0, u^0)$. Suppose that the point x^0 is regular with respect to the cone $K_0 := K(\lambda^0)$, i.e.

$$0 \in \text{int}\{G(x^0) + DG(x^0)X - K_0\}. \quad (6.14)$$

Clearly assumption B implies the regularity condition (3.6). Moreover, assumption B implies uniqueness of the Lagrange multiplier λ^0 , and if the cone K is polyhedral, then λ^0 is unique iff assumption B holds, [63, Lemma 4.3]. It is possible to show that assumption B is stronger than assumption A. Another property that follows from assumption B is that the Lagrange multipliers have the same order of perturbation as the optimal solutions [63, Lemma 4.4]. Therefore, under assumption B, Lipschitz stability of Lagrange multipliers follows from Lipschitz stability of optimal solutions.

7 Second-order analysis of the optimal value function

In this section we investigate directional differentiability of $v(u)$ by employing second-order expansions. We assume that x^0 is the unique solution of (P_{u^0}) , and that $d \in U$ satisfies the directional regularity condition. Then we have to deal with three basic situations. The first is when the set of solutions is upper-Lipschitz, which is the case if some strong second-order condition holds. Then the upper bound (4.14) is exact. A second-order expansion allows to state a second-order upper estimate of the value function that, for nonlinear programming problems, happens to be exact and allows to compute the expansion of $o(t^2)$ -optimal paths.

The second situation is when estimate (4.14) is not tight, as is the case in example 4.3, but the set of solutions is Hölder stable of degree $1/2$. Then an expansion of paths that vary like $O(\sqrt{t})$ allows to obtain a sharper upper bound of the value function that, for nonlinear programming problems, happens to be exact and allows to compute the expansion of $o(t)$ -optimal paths.

The third situation is when the set of Lagrange multiplier is empty, and the set of solutions is Hölder stable of degree $1/2$ again, which is as before the consequence of some second-order conditions. Then the solution varies in general as \sqrt{t} . An expansion of paths that vary like $O(\sqrt{t})$ allows to estimate the upper limit of $t^{-1/2}(\text{val}(u^0 + td) - \text{val}(u^0))$. For

nonlinear programming problems, this estimate is exact and allows to compute the expansion of $o(\sqrt{t})$ -optimal paths.

7.1 Second order expansion of the optimal value function and first-order expansion of optimal solutions

Let x^0 be an optimal solution of the problem (P_{u^0}) and d be a direction of perturbations for which directional regularity condition holds. We have established in section 4 that if $h \in \mathcal{S}(PL_d)$ (i.e. h is an optimal solution of the linearized problem (PL_d)) and $x(t)$ is a feasible path of the form $x(t) = x^0 + th + o(t)$, then $x(t)$ is an $o(t)$ -optimal solution of (P_{u^0+td}) . However, the set $\mathcal{S}(PL_d)$ is usually far too large to give a useful information about the variation of solutions. Take for instance the case of unconstrained minimization problems: then $\mathcal{S}(PL_d)$ is the whole space X . We obtain more precise upper estimates of the optimal value function, that give some insight into the variation of solutions, by considering feasible paths of the form $x(t) = x^0 + th + \frac{1}{2}t^2w + o(t^2)$.

Define the second order tangent set, to K at a point y in a direction z , as

$$T_K^2(y, z) := \left\{ w \in Y : \text{dist}(y + tz + \frac{1}{2}t^2w, K) = o(t^2) \right\}. \quad (7.15)$$

Note that $T_K^2(y, z)$ is empty if $y \notin K$ or $z \notin T_K(y)$. For the sake of brevity denote

$$T_K^2(h, d) := T_K^2(G(x^0, u^0), DG(x^0, u^0)(h, d)). \quad (7.16)$$

Feasibility of the path $x(t)$ implies that

$$DG(x^0, u^0)(h, d) \in T_K G((x^0, u^0)), \quad (7.17)$$

$$D_x G(x^0, u^0)w + D^2 G(x^0, u^0)((h, d), (h, d)) \in T_K^2(h, d). \quad (7.18)$$

We assume that $h \in \mathcal{S}(PL_d)$. The expansion (7.18) suggests that in order to derive a tight upper estimate for the directional variation of the optimal value function, the best choice for w is to minimize the second order term of the Taylor expansion of the objective function, while satisfying the constraint (7.18). That is, to take w as an optimal solution of

$$(\mathcal{PQ}_{d,h}) \quad \begin{array}{ll} \text{Min}_w & D_x f(x^0, u^0)w + D^2 f(x^0, u^0)((h, d), (h, d)) \\ \text{s.t.} & D_x G(x^0, u^0)w + D^2 G(x^0, u^0)((h, d), (h, d)) \in T_K^2(h, d). \end{array}$$

Then the best choice for h is to take it as an optimal solution of

$$(\mathcal{PQ}_d) \quad \text{Min}_{h \in \mathcal{S}(PL_d)} \text{val}(\mathcal{PQ}_{d,h}).$$

Theorem 7.1 *Let x^0 be an optimal solution of the problem (P_{u^0}) , with which is associated at least one Lagrange multiplier. Let d be a direction of perturbations for which the directional regularity condition holds. Then*

$$\limsup_{t \downarrow 0} \frac{v(u^0 + td) - v(u^0) - t \text{val}(PL_d)}{\frac{1}{2}t^2} \leq \text{val}(\mathcal{PQ}_d). \quad (7.19)$$

Note that if $\text{val}(\mathcal{PQ}_d) > -\infty$, then (7.19) is equivalent to

$$v(u^0 + td) \leq v(u^0) + t \text{val}(PL_d) + \frac{1}{2}t^2 \text{val}(\mathcal{PQ}_d) + o(t^2). \quad (7.20)$$

Note that $\mathcal{S}(PL_d)$ may be empty even if (PL_d) has a finite value. Also, it may happen that $T_K^2(h, d)$ is empty for all $h \in \mathcal{S}(PL_d)$. In that case the above inequality reduces to the trivial estimate $v(u^0 + td) \leq +\infty$.

Under the assumptions of Theorem 7.1, it can be proved that, for any $h \in \mathcal{S}(PL_d)$, the optimal value of $(\mathcal{PQ}_{d,h})$ is equal to the optimal value of the problem

$$(\mathcal{DQ}_{d,h}) \quad \text{Max}_{\lambda \in \Lambda_1(x^0, d)} \{D^2L(x^0, \lambda, u^0)((h, d), (h, d)) - \sigma(\lambda, T_K^2(h, d))\},$$

where $\Lambda_1(x^0, d) := \mathcal{S}(DL_d)$ and $\sigma(\cdot, T_K^2(h, d))$ denotes the support function of the set $T_K^2(h, d)$.

Theorem 7.2 *Let (P_u) be a (finite dimensional, finitely constrained) nonlinear programming problem, x^0 be the unique solution of (P_{u^0}) , and assume that $S(u^0 + td)$ is non empty and uniformly bounded for small enough $t > 0$. If d is a direction of perturbations for which both, the directional regularity condition and the directional second order sufficient conditions (6.10), hold, then*

$$v(u^0 + td) = v(u^0) + t \text{val}(PL_d) + \frac{1}{2}t^2 \text{val}(\mathcal{PQ}_d) + o(t^2). \quad (7.21)$$

In addition, the set of optimal solutions of (\mathcal{PQ}_d) coincides with the set

$$\{h \in X : \exists x(t) = x^0 + th + o(t); x(t) \text{ is an } o(t^2)\text{-optimal solution of } (P_{u^0+td})\}.$$

Note that in the case of finitely many constraints the “sigma term” vanishes and the optimal value $\text{val}(\mathcal{PQ}_d)$ in (7.21) is given by the optimal value of the problem

$$\text{Min}_{h \in \mathcal{S}(PL_d)} \left\{ \psi_d(h) := \text{Max}_{\lambda \in \Lambda_1(x^0, d)} D^2L(x^0, \lambda, u^0)((h, d), (h, d)) \right\}. \quad (7.22)$$

It follows that if $\bar{x}(t)$ is an $o(t^2)$ -optimal solution of (P_{u^0+td}) converging to x_0 , then under the assumptions of Theorem 7.2 every accumulation point of $t^{-1}(\bar{x}(t) - x_0)$ is an optimal solution of the program (7.22). In particular, if the optimization problem (7.22) possesses a unique optimal solution $\bar{h} = \bar{h}(d)$, then the directional derivative $\bar{x}'(u_0, d)$ exists and is equal to $\bar{h}(d)$.

These results were obtained first for nonlinear programming problems, under the assumption of linear independence of the gradients of the constraint functions and the strict complementarity condition, by applying the implicit function theorem to the first order optimality conditions considered as a system of nonlinear equations [23],[24]. The above approach of investigating directional differentiability of optimal solutions by using a second order expansion of the corresponding optimal value function, was employed in [59] under

the assumption of uniqueness of Lagrange multipliers and in [60] under the Mangasarian-Fromovitz constraint qualification. The duality method was suggested in [6] were the results were extended further under the directional regularity condition. The last statement of Theorem 7.2 is due to [15]. The results were extended to the Banach space setting in [12, Part I], taking advantage of the second order analysis of [19].

Example 7.1 [60] *Consider the following problem*

$$\text{Min}_{(x_1, x_2) \in \mathbb{R}^2} \frac{1}{2}(x_1 - 1)^2 + \frac{1}{2}x_2^2 \quad \text{s.t. } x_1 \leq 0, x_1 + u_1x_2 + u_2 \leq 0, \quad (7.23)$$

depending on the parameter vector $(u_1, u_2) \in \mathbb{R}^2$. For $u_0 = (0, 0)$ this problem has the unique optimal solution $x_0 = (0, 0)$. The Mangasarian-Fromovitz constraint qualification holds at x_0 and the corresponding set Λ_0 of Lagrange multipliers is

$$\Lambda_0 = \{(\lambda_1, \lambda_2) : \lambda_1 + \lambda_2 = 1, \lambda_1 \geq 0, \lambda_2 \geq 0\}.$$

Also the strong second order sufficient conditions hold and hence the results of Theorem 7.2 apply. We have here

$$D^2L(x_0, \lambda, u_0)((h, d)(h, d)) = \frac{1}{2}h_1^2 + \frac{1}{2}h_2^2 + \lambda_2 d_1 h_2,$$

and $\mathcal{S}(PL_d)$ is the optimal solutions set of the linearized problem

$$\text{Min}(-h_1) \quad \text{s.t. } h_1 \leq 0, h_1 + d_2 \leq 0. \quad (7.24)$$

Moreover, $D_u L(x_0, \lambda, u_0) = (0, \lambda_2)$ and hence $\Lambda_1(x_0, d)$ is formed by the maximizers of $d_2 \lambda_2$ over Λ_0 .

Let $d = (1, 0)$. Then $\Lambda_1(x_0, d) = \Lambda_0$,

$$\psi_d(h) = \max\{\frac{1}{2}h_1^2 + \frac{1}{2}h_2^2, \frac{1}{2}h_1^2 + \frac{1}{2}h_2^2 + h_2\}$$

and $\mathcal{S}(PL_d) = \{(h_1, h_2) : h_1 = 0\}$. It follows that in that case $\bar{h} = (0, 0)$ and hence $\bar{x}'(0, d) = (0, 0)$. Now let $d = (1, \gamma)$ for some $\gamma > 0$. Then $\Lambda_1(x_0, d) = \{(0, 1)\}$, $\psi_d(h)(h) = \frac{1}{2}h_1^2 + \frac{1}{2}h_2^2 + h_2$ and $\mathcal{S}(PL_d) = \{(h_1, h_2) : h_1 = -\gamma\}$. It follows that in that case $\bar{h} = (-\gamma, -1)$ and hence $\bar{x}'(0, d) = (-\gamma, -1)$. We see that in this example the optimal solution $\bar{x}(u)$ is directionally differentiable at $u = 0$, but the directional derivative $\bar{x}'(0, d)$ is a discontinuous function of the direction d . It follows that $\bar{x}(u)$ is not directionally differentiable in the Hadamard sense and is not Lipschitz continuous in any neighborhood of $u = 0$. A reason for such discontinuous behavior of $\bar{x}'(0, \cdot)$ is that the optimal set $\Lambda_1(x_0, d)$ depends on d in a discontinuous way.

Let us make the following observations. If the Lagrange multiplier is unique, then the function $\psi_d(\cdot)$ is quadratic and (7.22) becomes a quadratic programming problem. If the hypotheses of Proposition 5.3 hold, then the results of this section are a consequence of those proved for strongly regular solutions. If the strict complementarity condition holds, then we recover the result of Fiacco and McCormick [23],[24].

7.2 First-order differentiability of the optimal value function and Hölder expansion of optimal solutions by the second order analysis

Let x^0 be a solution of problem (P_{u^0}) , and d be a direction of perturbation for which directional regularity holds. As was observed in example 4.3, it may happen that the upper bound (4.14), for the first-order directional behavior of the optimal value function $v(u)$, is not tight. In this section we derive a sharper estimate by employing feasible paths of the form $x(t) = x^0 + t^{1/2}h + w + o(t)$. By expanding $G(x(t), u + td)$ we obtain that a necessary condition for this path to be feasible is that

$$\begin{aligned} D_x G(x^0, u^0)h &\in T_K(G(x^0, u^0)), \\ 2DG(x^0, u^0)(w, d) + D_{xx}^2 G(x^0, u^0)(h, h) &\in \mathcal{T}_K^2(h), \end{aligned} \quad (7.25)$$

where we have set

$$\mathcal{T}_K^2(h) := T_K^2(G(x^0, u^0), D_x G(x^0, u^0)h).$$

(Note that $\mathcal{T}_K^2(h)$ should not be confused with the set $T_K^2(h, d)$ introduced earlier.) In order to deal with this type of paths, we need the following condition. A direction of perturbations $d \in U$ is said to satisfy the *strong directional regularity* condition [12] if it satisfies the directional regularity condition and, in addition, whenever (h, w) is such that (7.25) holds, then for $\gamma < 1$, and arbitrary close to 1, one can find $w_\gamma \in X$ with $w_\gamma \rightarrow w$, and feasible paths of the form $x^\gamma(t) = x_0 + \gamma t^{1/2}d + tw_\gamma + o(t)$. This condition is equivalent to directional regularity whenever $K = \{0\} \times K_2$, with K_2 being a closed convex cone with nonempty interior. In particular, for nonlinear programming problems, this is equivalent to Gollan's condition (3.13). It is not known whether both conditions are equivalent in general.

The above expansion suggests that in order to derive an upper estimate for the directional behavior of the optimal value function the best choice is to take h in the critical cone C_0 , and then for a given h to take w that minimizes the second order term of the expansion of the objective function, while satisfying the above constraint. That is, to take w as an optimal solution of

$$\begin{aligned} (\mathcal{P}\mathcal{L}_{d,h}) \quad & \text{Min}_w \quad 2Df(x^0, u^0)(w, d) + D_{xx}^2 f(x^0, u^0)(h, h) \\ & \text{s.t.} \quad 2DG(x^0, u^0)(w, d) + D_{xx}^2 G(x^0, u^0)(h, h) \in \mathcal{T}_K^2(h), \end{aligned}$$

and then the best choice for h is to take it as an optimal solution of

$$(\mathcal{P}\mathcal{L}_d) \quad \text{Min}_{h \in C_0} \text{val}(\mathcal{P}\mathcal{L}_{d,h}).$$

Theorem 7.3 *Let x^0 be an optimal solution of the problem (P_{u^0}) , with which is associated at least one Lagrange multiplier. Let d be a direction of perturbations for which strong directional regularity holds. Then*

$$v(u^0 + td) \leq v(u^0) + \frac{1}{2}t \text{val}(\mathcal{P}\mathcal{L}_d) + o(t).$$

It may happen that $\mathcal{T}_K^2(h)$ is empty for certain critical directions h . However, for $h = 0$ we note that problem $(\mathcal{P}\mathcal{L}_{d,0})$ coincides with the linearized problem (PL_d) . Therefore, we have that $\text{val}(\mathcal{P}\mathcal{L}_d) \leq \text{val}(\mathcal{P}\mathcal{L}_{d,0}) \leq \text{val}(PL_d)$, the latter being less than $+\infty$ under the directional qualification condition.

The dual of problem $(\mathcal{P}\mathcal{L}_{d,h})$ is

$$(\mathcal{D}\mathcal{L}_{d,h}) \quad \text{Max}_{\lambda \in \Lambda(x^0, u^0)} \{2D_u L(x^0, \lambda, u^0) + D_{xx}^2 L(x^0, \lambda, u^0)(h, h) - \sigma(\lambda, \mathcal{T}_K^2(h))\}.$$

It is not known whether $\text{val}(\mathcal{P}\mathcal{L}_{d,h})$ is equal to $\text{val}(\mathcal{D}\mathcal{L}_{d,h})$ for any $h \in C_0$. It may be proved, however, that under the hypotheses of Theorem 7.3, we have

$$\text{val}(\mathcal{P}\mathcal{L}_d) = \inf_{h \in C_0} \text{val}(\mathcal{D}\mathcal{L}_{d,h}).$$

Theorem 7.4 *Let (P_u) be a (finite dimensional, finitely constrained) nonlinear programming problem, x^0 be the unique solution of (P_{u^0}) , with which is associated at least one Lagrange multiplier, and suppose that $S(u^0 + td)$ is non empty and uniformly bounded for small enough $t > 0$. If d is a direction of perturbation for which directional regularity and the second order condition (definition 6.2) both hold, then*

$$v(u^0 + td) = v(u^0) + \frac{1}{2}t \text{val}(\mathcal{P}\mathcal{L}_d) + o(t).$$

In addition, the set of solutions of $(\mathcal{P}\mathcal{L}_d)$ coincides with the set

$$\{h \in X : \exists x(t) = x^0 + t^{1/2}h + o(t^{1/2}); x(t) \text{ is an } o(t) \text{ optimal solution of } (P_{u^0+td})\}.$$

The above results were obtained first in the setting of nonlinear programming problems. The expansion of the value function (under a different form) is due to [27], while [15] obtained the expansion of approximate solutions. Part II of [12] extends the result to the Banach space setting.

7.3 Square root expansion of the optimal value function and Hölder expansion of optimal solutions

We now consider the case when directional regularity condition holds, but the set of Lagrange multipliers is *empty*. By Proposition 4.1, we already know that in that case the directional derivative $v'(u_0, d) = -\infty$. We show in this section that a more precise result holds. Namely, variations of the optimal value of (P_{u^0+td}) are of the order $t^{1/2}$, and $(v(u^0 + td) - v(u^0))/t^{1/2}$ has a limit in the case of nonlinear programming problems, assuming the second-order conditions (definition 6.2) to hold.

Again our analysis is based on considering feasible paths of the form $x(t) = x^0 + t^{1/2}h + tw + o(t)$. Then, as before, (7.25) holds. We wish to minimize the first term of the expansion of the objective function, i.e. $D_x f(x^0, u^0)h$. Since we want this amount to be negative, we assume that $h \in C_0$. We may write the corresponding problem as

$$(\mathcal{P}\mathcal{S}_{d,h}) \quad \text{Min}_w D_x f(x^0, 0)h \quad \text{s.t.} \quad 2DG(x^0, u^0)(w, d) + D_{xx}^2 G(x^0, u^0)(h, h) \in \mathcal{T}_K^2(h).$$

For a given h this is in fact a feasibility problem (since the objective function does not depend on w). Minimizing over h we obtain the problem

$$(\mathcal{PS}_d) \quad \underset{h \in C_0}{\text{Min val}}(\mathcal{PS}_{d,h}).$$

Theorem 7.5 *Let x^0 be an optimal solution of the problem (P_{u^0}) , with which is associated at least one singular multiplier, but no Lagrange multipliers. Let d be a direction of perturbations for which strong directional regularity condition holds. Then $\text{val}(\mathcal{PS}_d) < 0$ and*

$$v(u^0 + td) \leq v(u^0) + t^{1/2}\text{val}(\mathcal{PS}_d) + o(t^{1/2}).$$

Under the assumptions of Theorem 7.5, it can be shown that the optimal value of (\mathcal{PS}_d) is equal to the optimal value of the following problem

$$\underset{h \in C(x^0)}{\text{Min}} D_x f(x^0, 0)h + \sup_{(\lambda_0, \lambda) \in \Lambda^g(x^0)} \left\{ \begin{array}{l} 2D_u L^g(x^0, \lambda_0, \lambda, u^0)d + \\ D_{xx} L^g(x^0, \lambda_0, \lambda, u^0)(h, h) - \sigma(\lambda, \mathcal{T}_K^2(h)) \end{array} \right\}.$$

Theorem 7.6 *Let (P_u) be a (finite dimensional, finitely constrained) nonlinear programming problem, x^0 be the unique optimal solution of (P_{u^0}) , and suppose that $S(u^0 + td)$ is non empty and uniformly bounded for small enough $t > 0$. If d is a direction of perturbations for which directional regularity condition holds, no Lagrange multiplier is associated with x^0 and the second order condition (definition 6.2) is satisfied, then*

$$v(u^0 + td) = v(u^0) + t^{1/2}\text{val}(\mathcal{PS}_d) + o(t^{1/2}).$$

In addition, the set of solutions of (\mathcal{PS}_d) coincides with the set

$$\left\{ h \in X : \begin{array}{l} \exists x(t) = x^0 + t^{1/2}h + o(t^{1/2}), \\ x(t) \text{ is an } o(t^{1/2})\text{-optimal solution of } (P_{u^0+td}) \end{array} \right\}.$$

These results were obtained in [10], in the nonlinear programming setting, and generalized in [12, Part II]. A major difference with the previously studied situations is that the variation of the cost is of order $t^{1/2}$.

8 Some illustrations

We apply some of the previous results to the computation of the expansion of the chain when the unperturbed position is either vertical or horizontal.

8.1 Application to the perturbation of the vertical chain

Equivalence to a directionally qualified problem It is easy to show that the position of the chain is the same as if we change the last equality into an inequality, obtaining the

following optimization problem:

$$\begin{aligned} & \text{Min}_{y,z} E(y, z) \\ & \text{s.t. } y_k^2 + z_k^2 - 1 = 0, \quad k = 1, \dots, m, \quad \sum_{k=1}^m y_k = 0, \\ & m - u - \sum_{k=1}^m z_k \leq 0. \end{aligned} \quad (8.26)$$

It appears that the new problem (8.26) satisfies the directional qualification condition, as we now check, although problem (2.2) does not. Indeed, the linearized equality constraints are onto, as solving

$$d_k^z = \gamma_k, \quad k = 1, \dots, m; \quad \sum_{k=1}^m d_k^y = \beta; \quad \gamma \in \mathbb{R}^m, \quad \beta \in \mathbb{R} \quad (8.27)$$

obviously has a solution for any (γ, β) . If $(\gamma, \beta) = 0$, then $d_k^z = 0$. It follows that a displacement in the kernel of linearized equality constraints cannot satisfy strictly the linearized active inequality constraint $-\sum_k d_k^z \leq 0$. Therefore, qualification does not hold. On the other hand, directional qualification condition holds as (3.3) is satisfied with $d = (d^y, d^z) = 0$.

Lagrange multipliers, critical cone and second order conditions The expression for the Lagrangian function $L(y, z, \lambda, u)$ is

$$\sum_{k=1}^m \alpha_k z_k + \sum_{k=1}^m \lambda_k \left(\frac{1}{2} y_k^2 + \frac{1}{2} z_k^2 - 1 \right) + \lambda_{m+1} \sum_{k=1}^m y_k + \lambda_{m+2} \left(m - u - \sum_{k=1}^m z_k \right).$$

Its derivatives, with respect to (y, z) , are null whenever

$$\lambda_{m+1} = 0 \quad \text{and} \quad \alpha_k + \lambda_k - \lambda_{m+2} = 0, \quad k = 1, \dots, m.$$

Therefore, the set of Lagrange multipliers is

$$\Lambda = \{ \lambda \in \mathbb{R}^{m+2} : \lambda_{m+1} = 0; \lambda_{m+2} \geq 0; \lambda_k = \lambda_{m+2} - \alpha_k, \quad k = 1, \dots, m \}.$$

The equations of the critical cone are

$$\sum_{k=1}^{m-1} \alpha_k d_k^z \leq 0; \quad d_k^z = 0, \quad k = 1, \dots, m; \quad \sum_{k=1}^m d_k^y = 0; \quad \sum_{k=1}^m d_k^z \leq 0,$$

i.e.

$$C(x^0) = \left\{ (d^y, d^z) : d^z = 0 \text{ and } \sum_{k=1}^m d_k^y = 0 \right\}.$$

Let us now check the standard second order conditions. Given a critical direction d , using $d^z = 0$, we compute

$$D_{xx}^2 L(x^0, \lambda, 0)(d, d) = \sum_{k=1}^m \lambda_k (d_k^y)^2.$$

There exists a Lagrange multiplier such that $\lambda_{m+2} = p$. Then $\lambda_k \geq 1$, $k = 1, \dots, p$. In that case the second-order variation of the Lagrangian is non zero when d is a non zero critical direction, i.e. the weak second-order sufficient condition holds.

It may be checked that the strong directional second-order sufficient condition does not hold.

Study of the auxiliary problem The expression of the auxiliary problem is

$$\text{Min}_{d \in C(x^0)} \sup_{\lambda \in \Lambda(x^0)} \Delta(d, \lambda),$$

where

$$\Delta(d, \lambda) = -2\lambda_{m+2} + \sum_{k=1}^m \lambda_k (d_k^y)^2 = \lambda_{m+2} \left(\sum_{k=1}^m (d_k^y)^2 - 2 \right) - \sum_{k=1}^m \alpha_k (d_k^y)^2.$$

If $\sum_k (d_k^y)^2 > 2$, then $\sup\{\Delta(d, \lambda) ; \lambda \in \Lambda(x^0)\} = +\infty$. Otherwise, the sup is attained for $\lambda_{m+2} = 0$. Therefore the subproblem has the same solutions as

$$\text{Min}_{d \in C(x^0)} -\frac{1}{2} \sum_{k=1}^m \alpha_k (d_k^y)^2 ; \sum_{k=1}^m (d_k^y)^2 \leq 2.$$

Using the expression of $C(x^0)$, we formulate this problem as

$$\text{Min}_{d^y} -\frac{1}{2} \sum_{k=1}^m \alpha_k (d_k^y)^2 ; \frac{1}{2} \sum_{k=1}^m (d_k^y)^2 - 1 \leq 0 ; \sum_{k=1}^m d_k^y = 0.$$

The problem has solutions (compact nonempty feasible set), has a concave objective function and convex constraints, and every feasible point is qualified. With a solution d^y are associated multipliers (η, ν) with $\eta \geq 0$ such that

$$-\alpha_k d_k^y + \eta d_k^y + \nu = 0, \quad k = 1, \dots, m. \quad (8.28)$$

Let us prove that $\nu \neq 0$. If $\nu = 0$, then $(\eta - \alpha_k)d_k^y = 0$, $k = 1, \dots, m$. All components of α_k having different values, we deduce that $d^y = 0$ except for at most one component. From the equality constraint it follows that $d^y = 0$. Then $\eta = 0$ (as the inequality constraint is not active). However, the second-order necessary condition does not hold at $d^y = 0$. It follows that ν cannot be null, as had to be proved. As $\nu \neq 0$, we deduce from (8.28) that

$$d_k^y = \nu / (\alpha_k - \eta); \quad k = 1, \dots, m. \quad (8.29)$$

Summing over k , we get

$$0 = \sum_{k=1}^m d_k^y = \nu \sum_{k=1}^m \frac{1}{\alpha_k - \eta}. \quad (8.30)$$

As $\nu \neq 0$, this gives a scalar equation with the (one) scalar unknown η

$$\sum_{k=1}^m \frac{1}{\alpha_k - \eta} = 0.$$

Let us study this equation, remembering that α_k is a strictly decreasing sequence. If $\eta < \alpha_p$ or $\eta > \alpha_p$ all terms of the sum are of the same sign. Here a solution must be in (α_p, α_1) ; and cannot be equal to some α_k . Let us fix k in $\{1, \dots, m-1\}$. We study the solution for $\eta \in (\alpha_{k+1}, \alpha_k)$. The left hand side of the equation is strictly increasing from $-\infty$ to $+\infty$. It follows that (8.30) has a unique solution in each interval (α_{k+1}, α_k) . This solution will be said to be the k -th solution.

We determine the value of ν by plugging the expression of d^y given by (8.29) in the active inequality constraint. It follows that

$$\nu = \pm \sqrt{2 / \sum_{k=1}^m (\alpha_k - \eta)^{-2}}$$

By (8.29) the two values of ν are associated with opposite values of d^y of equal cost. This is natural due to the symmetry in the problem. The components of d^y corresponding to the k -th solution have the same sign from 1 to k , and then the opposite sign.

It is intuitively clear that the best solution is the first one. This is easy to prove if we observe, from the data of the subproblem, that $k \rightarrow |d_k^y|$ must be decreasing. Otherwise, applying a permutation to the components of d^y that makes them decreasing with k , we get another feasible point associated with a strictly inferior value of the cost. In summary, d_1^y is, say, positive, all other components of d^y are negative and strictly decreasing.

As the subproblem has two solutions, application of theorem 7.4 gives that any $o(u)$ optimal path is of the form $x^0 + s(t)\sqrt{u}d + o(\sqrt{u})$, where $s(t)$ has values in $\{-1, 1\}$.

8.2 Application to the perturbation of the horizontal chain

Equivalence to a directionally qualified problem It is easily shown that an equivalent problem is obtained by changing the one before the last equality into an inequality, i.e. optimization problem:

$$\begin{aligned} & \text{Min}_{y,z} E(y, z) \\ & \text{s.t. } y_k^2 + z_k^2 - 1 = 0, \quad k = 1, \dots, m, \quad \sum_{k=1}^m z_k = 0, \\ & m - u - \sum_{k=1}^m y_k \leq 0. \end{aligned} \tag{8.31}$$

By arguments similar to those used in the case of the vertical chain, we may check that the new problem (8.26) satisfies the directional qualification hypothesis.

Lagrange multipliers, critical cone and second order conditions The expression of the generalized Lagrangian function $L(y, z, \lambda, u)$ is

$$\lambda_0 \sum_{k=1}^m \alpha_k z_k + \sum_{k=1}^m \lambda_k \left(\frac{1}{2} y_k^2 + \frac{1}{2} z_k^2 - 1 \right) + \lambda_{m+1} \sum_{k=1}^m z_k + \lambda_{m+2} \left(m - u - \sum_{k=1}^m y_k \right).$$

Its derivatives, with respect to (y, z) , are null whenever

$$\lambda_k - \lambda_{m+2} = 0, \quad \lambda_0 \alpha_k + \lambda_{m+1} = 0 \quad k = 1, \dots, m,$$

Therefore, the set of Lagrange multipliers is empty, and

$$\Lambda^g = \{(\lambda_0, \lambda) \in \mathbb{R}^{m+2} : \lambda_0 = \lambda_{m+1} = 0; \lambda_{m+2} \geq 0; \lambda_k = \lambda_{m+2}, \quad k = 1, \dots, m\}.$$

The critical cone is

$$C(x^0) = \left\{ (d^y, d^z) ; d^y = 0; \sum_{k=1}^m \alpha_k d_k^z \leq 0; \sum_{k=1}^m d_k^z = 0 \right\}.$$

Let us now check the standard second order conditions. Given a critical direction d , using $d^y = 0$, we have

$$D_{xx}^2 L(x^0, \lambda, 0)(d, d) = \sum_{k=1}^m \lambda_k (d_k^z)^2.$$

Whenever $\lambda_{m+2} > 0$, as $\lambda_k = \lambda_{m+2}, k = 1, \dots, m$, this is non zero when d is a non zero critical direction, i.e. the (singular) weak second-order sufficient condition holds.

Study of the auxiliary problem The expression of the auxiliary problem is

$$\text{Min}_{d \in C(x^0)} \sup_{\lambda \in \Lambda^g(x^0)} \Delta(d, \lambda),$$

where

$$\Delta(d, \lambda) = \sum_{k=1}^m \alpha_k d_k^z + \lambda_{m+2} \left(\sum_{k=1}^m (d_k^z)^2 - 2 \right).$$

If $\sum_k (d_k^z)^2 > 2$, then $\sup\{\Delta(d, \lambda) ; \lambda \in \Lambda(x^0)\} = +\infty$. Otherwise, the sup is attained for $\lambda_{m+2} = 0$. Therefore the subproblem has the same solutions as

$$\text{Min}_{d \in C(x^0)} \sum_{k=1}^m \alpha_k d_k^z; \sum_{k=1}^m (d_k^z)^2 \leq 2.$$

Using the expression of $C(x^0)$, we formulate this problem as

$$\text{Min}_{d^z} - \sum_{k=1}^m \alpha_k d_k^z; \frac{1}{2} \sum_{k=1}^m (d_k^z)^2 - 1 \leq 0; \sum_{k=1}^m d_k^z = 0.$$

This is a qualified convex problem. The solution is better expressed in terms of the variation of altitude of the end point of the links. The altitude of the end of the k th link is $\sum_{i=1}^k z_k$. The associated displacement $\bar{d}_k := \sum_{i=1}^k d_k^z$ is solution of the problem

$$\text{Min}_{\bar{d}} - \sum_{k=1}^m \bar{d}_k; \quad \frac{1}{2} \sum_{k=1}^m (\bar{d}_k - \bar{d}_{k-1})^2 - 1 \leq 0.$$

Let $\alpha \neq 0$ be the Lagrange multiplier associated with the constraint. Then \bar{d} is solution of

$$\text{Min} \sum_{k=1}^p \bar{d}_k + \frac{\alpha}{2} \sum_{k=1}^p (\bar{d}_k - \bar{d}_{k-1})^2.$$

Therefore we may compute \bar{d} as follows: let \hat{d} be the solution of the linear system

$$\hat{d}_{k+1} - 2\hat{d}_k + \hat{d}_{k-1} = 1; \quad \hat{d}_0 = \hat{d}_p = 0.$$

Compute the associated energy $\eta := \frac{1}{2} \sum_{k=1}^p (\hat{d}_k - \hat{d}_{k-1})^2$, and set $\bar{d} := \eta^{-1} \hat{d}$. Then by theorem 7.6, we have $x(u) = x(0) + \sqrt{u} \bar{d} + o(\sqrt{u})$.

9 Comments and conclusion

Sensitivity analysis for optimization problems is a very active field, and when writing this article we had to be selective. Among the important contributions that were not quoted yet, let us mention the work based on epigraphical analysis [3, 4, 69] and the many applications of polyhedricity theory [46, 30] and its generalizations [8, 12].

We have carefully reviewed the approach of sensitivity analysis that uses lower and upper estimates of the value function. For nonlinear programming problems, this gives very strong results and the theory seems to be now more or less complete. It is anticipated that some of the strongest results might be extended to semi definite programming in the near future. At the same time, example 6.2 suggests that Hölder expansion of order different from 1/2 should be considered. The fact that many examples of semidefinite programs involve linear cost functions and constraints [18] is another challenge, as the set of solutions of the unperturbed program might typically be nonisolated.

References

- [1] W. ALT, *Stability of solutions for a class of nonlinear cone constrained optimization problems, Part 1: basic theory*, Numerical Functional Analysis and Optimization, 10 (1989), pp. 1053-1064.
- [2] W. ALT, *Local stability of solutions to differentiable optimization problems in Banach spaces*, Journal Optimization Theory and Applications, 70 (1991), pp. 443-466.

-
- [3] H. ATTOUCH, *Variational Convergence for Functions and Operators*, Pitman, Boston, MA, 1984.
- [4] H. ATTOUCH, *Viscosity solution of minimization problems*. SIAM J. Optimization, to appear.
- [5] J. P. AUBIN AND I. EKELAND, *Applied Nonlinear Analysis*, Wiley, New York, 1984.
- [6] A. AUSLENDER AND R. COMINETTI, *First and second order sensitivity analysis of nonlinear programs under directional constraint qualification conditions*, Optimization, 21 (1990), pp. 351-363.
- [7] V. I. AVERBUKH AND O. G. SMOLYANOV, *The various definitions of the derivative in linear topological spaces*, Russian Mathematical Surveys, 23 (1968), pp. 67-113.
- [8] L. BARBET AND R. JANIN, *Analyse de sensibilité différentielle pour un problème d'optimisation paramétré en dimension infinie*. Comptes Rendus de l'Académie des Sciences de Paris, 318 (1993), Série I, pp. 221-226.
- [9] J. F. BONNANS, *A semi-strong sufficiency condition for optimality in non convex programming and its connection to the perturbation problem*, Journal Optimization Theory and Applications, 60 (1989), pp. 7-18.
- [10] J. F. BONNANS, *Directional derivatives of optimal solutions in smooth nonlinear programming*, Journal Optimization Theory and Applications, 73 (1992), pp. 27-45.
- [11] J. F. BONNANS AND E. CASAS, *Contrôle de systèmes elliptiques semilinéaires comportant des contraintes sur l'état*, in "Nonlinear partial differential equations and their applications, Collège de France seminar vol. VIII", H. Brézis & J.L. Lions eds., Pitman Research Notes in Mathematics Series 166, Longman Scientific & Technical, New York, 69-86, 1988.
- [12] J. F. BONNANS, AND R. COMINETTI, *Perturbed optimization in Banach spaces*, SIAM J. Control Optimization, 34 (1996), to appear.
- [13] J. F. BONNANS, A. D. IOFFE, *Quadratic growth and stability in convex programming problems with multiple solutions*. J. Convex Analysis 2 (Special issue dedicated to R.T. Rockafellar), pp. 41-57.
- [14] J. F. BONNANS, A. D. IOFFE, *Second-order sufficiency and quadratic growth for non isolated minima*. Mathematics of Operations Research, to appear.
- [15] J. F. BONNANS, A. D. IOFFE AND A. SHAPIRO, *Expansion of exact and approximate solutions in nonlinear programming*, Proc. French-German Conference on Optimization, D. Pallaschke ed., Lecture Notes in Economics and Mathematical Systems, Springer, Berlin, 1992, pp. 103-117.

-
- [16] J. F. BONNANS, AND A. SULEM, *Pseudopower expansion of solutions of generalized equations and constrained optimization problems*, Mathematical Programming, 70 (1995), pp. 123-148.
- [17] J. F. BONNANS, AND A. SHAPIRO, *Perturbation Analysis of Optimization Problems*, in preparation.
- [18] S. BOYD, L. EL GHAOU, E. FERON AND V. BALAKRISHNAN, *Linear Matrix inequalities*. Studies in Applied Math. Vol. 15, SIAM, Philadelphia, PA, 1994.
- [19] R. COMINETTI, *Metric regularity, tangent sets and second order optimality conditions*, Applied Mathematics and Optimization, 21 (1990), pp. 265-287.
- [20] J. M. DANSKIN, *The Theory of Max-Min and Its Applications to Weapons Allocation Problems*, Springer, New York, 1967.
- [21] V. F. DEM'YANOV AND V. N. MALOZEMOV, *Introduction to Minimax*, Wiley, New York, 1974.
- [22] A. L. DONTCHEV AND R. T. ROCKAFELLAR, *Characterizations of strong regularity for variational inequalities over polyhedral convex sets*. SIAM J. Optimization, to appear.
- [23] A. V. FIACCO AND G. P. MCCORMICK, *Nonlinear Programming: Sequential Unconstrained Minimization Techniques*, Wiley, New York, 1968.
- [24] A. V. FIACCO, *Introduction to Sensitivity and Stability Analysis in Nonlinear Programming*, Academic Press, New York, 1983.
- [25] J. GAUVIN AND J. W. TOLLE, *Differential stability in nonlinear programming*, SIAM J. Control and Optimization, 15 (1977), pp. 294-311.
- [26] J. GAUVIN AND F. DUBEAU, *Differential properties of the marginal function in mathematical programming*, Mathematical Programming Study, 19 (1982), pp. 101-119.
- [27] J. GAUVIN AND R. JANIN, *Directional behaviour of optimal solutions in nonlinear mathematical programming*, Mathematics of Operations Research, 13 (1988), pp. 629-649.
- [28] B. GOLLAN, *On the marginal function in nonlinear programming*, Mathematics of Operations Research, 9 (1984), pp. 208-221.
- [29] E. G. GOL'SHTEIN, *Theory of Convex Programming*, Translations of Mathematical Monographs 36, American Mathematical Society, Providence, RI, 1972.
- [30] A. HARAUX, *How to differentiate the projection on a convex set in Hilbert space. Some applications to variational inequalities*, J. Mathematical Society of Japan (29) (1977), pp. 615-631.

-
- [31] A. D. IOFFE, *On sensitivity analysis of nonlinear programs in Banach spaces: the approach via composite unconstrained optimization*, SIAM J. Optimization, 4 (1994), pp. 1-43.
- [32] F. JOHN, *Extremum problems with inequalities as subsidiary conditions*, in Studies and Essays, R. Courant anniversary volume, Interscience, New York, 1948, pp. 187-204.
- [33] H. TH. JONGEN, D. KLATTE, K. TAMMER, *Implicit functions and sensitivity of stationary points*, Mathematical Programming, 19 (1990), pp. 123-138.
- [34] H. KAWASAKI, *An envelope-like effect of infinitely many inequality constraints on second order necessary conditions for minimization problems*, Mathematical Programming, 41 (1988), pp. 73-96.
- [35] D. KLATTE, K. TAMMER, *Strong stability of stationary solutions and Karush-Kuhn-Tucker points in nonlinear optimization*, Annals of Operations Research, 27 (1990), pp. 285-308.
- [36] M. KOJIMA, *Strongly stable stationary solutions in nonlinear programming*, in S. M. Robinson, ed., *Analysis and computation of fixed points*, Academic Press, New York, 1988, pp. 93-138.
- [37] B. KUMMER, *Lipschitzian inverse functions directional derivatives and application in $C^{1,1}$ optimization*, Journal of Optimization Theory and Applications, 70 (1991), pp. 561-582.
- [38] F. LEMPIO AND H. MAURER, *Differential stability in infinite-dimensional nonlinear programming*, Applied Mathematics and Optimization, 6 (1980), pp. 139-152.
- [39] E. S. LEVITIN, *On differential properties of the optimal value of parametric problems of mathematical programming*, Dokl. Akad. Nauk SSSR, 224 (1975), pp. 1354-1358.
- [40] E. S. LEVITIN, *Differentiability with respect to a parameter of the optimal value in parametric problems of mathematical programming*, Kibernetika, (1976), pp. 44-59.
- [41] E. S. LEVITIN, *Perturbation Theory in Mathematical Programming and Its Applications*, Wiley, New York, 1994.
- [42] D. G. LUENBERGER, *Introduction to linear and nonlinear programming*, Addison-Wesley, Reading, Massachusetts, 1973.
- [43] K. MALANOWSKI, *Second order conditions and constraint qualifications in stability and sensitivity analysis of solutions to optimization problems in Hilbert spaces*, Applied Mathematics and Optimization, 25 (1992), pp. 51-79.
- [44] O. MANGASARIAN AND S. FROMOVITZ, *The Fritz-John necessary optimality conditions in the presence of equality and inequality constraints*, J. Math. Anal. Appl., 7 (1967), pp. 37-47.

-
- [45] H. MAURER AND J. ZOWE, *First and second-order necessary and sufficient optimality conditions for infinite-dimensional programming problems*, Mathematical Programming, 16 (1979), pp. 98-110.
- [46] F. MIGNOT, *Contrôle dans les inéquations variationnelles elliptiques*, J. Functional Analysis 22 (1976), 25-39.
- [47] M. Z. NASHED, *Differentiability and Related Properties of Nonlinear Operators: Some Aspects of the Role of Differentials in Nonlinear Functional Analysis*, Nonlinear Functional Analysis and Applications, Edited by L.B. Rall, Academic Press, New York, pp. 103-309, 1971.
- [48] J. P. PENOT, *On regularity conditions in mathematical programming*, Mathematical Programming Study, 19 (1982), pp. 167-199.
- [49] B. N. PSHENICHNYI, *Necessary Conditions for an Extremum*, Marcel Dekker, New York, 1971.
- [50] S. M. ROBINSON, *Normed convex processes*, Transactions of the American Mathematical Society, 174 (1972), pp. 127-140.
- [51] S. M. ROBINSON, *First order conditions for general nonlinear optimization*, SIAM Journal on Applied Mathematics, 30 (1976), pp. 597-607.
- [52] S. M. ROBINSON, *Stability theorems for systems of inequalities, Part II: differentiable nonlinear systems*, SIAM J. Numerical Analysis, 13 (1976), pp. 497-513.
- [53] S. M. ROBINSON, *Regularity and stability for convex multivalued functions*, Mathematics of Operations Research, 1 (1976), pp. 130-143.
- [54] S. M. ROBINSON, *Strongly regular generalized equations*, Mathematics of Operations Research, 5 (1980), pp. 43-62.
- [55] S. M. ROBINSON, *Generalized equations and their solutions, Part II: applications to nonlinear programming*, Mathematical Programming Study, 19 (1982), pp. 200-221.
- [56] S. M. ROBINSON, *Implicit B-differentiability in generalized equations*, Technical Report 2854, Mathematics Research Center, University of Wisconsin-Madison, 1985.
- [57] R. T. ROCKAFELLAR, *Convex Analysis*, Princeton University Press, Princeton, NJ, 1970.
- [58] R. T. ROCKAFELLAR, *Conjugate Duality and Optimization*, Regional Conference Series in Applied Mathematics, SIAM, Philadelphia, 1974.
- [59] A. SHAPIRO, *Second-order sensitivity analysis and asymptotic theory of parametrized, nonlinear programs*, Math. Programming, 33 (1985), pp. 280-299.

- [60] A. SHAPIRO, *Sensitivity analysis of nonlinear programs and differentiability properties of metric projections*, SIAM J. Control and Optimization, 26 (1988), pp. 628-645.
- [61] A. SHAPIRO, *Perturbation theory of nonlinear programs when the set of optimal solutions is not a singleton*, Applied Mathematics & Optimization. An International Journal, 18 (1988), pp. 215-229.
- [62] A. SHAPIRO, *On concepts of directional differentiability*, Journal of Optimization Theory and Applications, 66 (1990), pp. 477-487.
- [63] A. SHAPIRO, *Perturbation analysis of optimization problems in Banach spaces*, Numerical Functional Analysis and Optimization, 13 (1992), pp. 97-116.
- [64] A. SHAPIRO, *On Lipschitzian stability of optimal solutions of parametrized semi-infinite programs*, Mathematics of Operations Research, 19 (1994), pp. 743-752.
- [65] A. SHAPIRO, *Directional differentiability of the optimal value function in convex semi-infinite programming*, Mathematical Programming, Series A, 70 (1995), pp. 149-157.
- [66] A. SHAPIRO, *On uniqueness of Lagrange multipliers in optimization problems subject to cone constraints*, SIAM Journal on Optimization, to appear.
- [67] A. SHAPIRO, *A variational principle and its applications*, to appear in: Proceedings of the Conference on "Parametric Optimization and Related Topics IV", Series "Approximation and Optimization", J. Guddat, H.Th. Jongen, F. Nozicka, G. Still and F. Twilt (Eds.), Verlag Peter Lang, Frankfurt.
- [68] A. SHAPIRO AND J. F. BONNANS, *Sensitivity analysis of parametrized programs under cone constraints*, SIAM J. Control and Optimization, 30 (1992), pp. 1409-1422.
- [69] D. TORRALBA, *Développements asymptotiques pour les méthodes d'approximation par viscosité*. Comptes Rendus de l'Académie des Sciences de Paris, 322 (1996), Série I, pp. 123-128.
- [70] C. URSESCU, *Multifunctions with convex closed graph*, Czechoslovak Mathematical Journal, 25 (1975), pp. 438-441.
- [71] K. VESELIĆ, *Finite catenary and the method of Lagrange*, SIAM Review, 30 (1995), pp. 224-229.
- [72] J. ZOWE AND S. KURCYUSZ, *Regularity and stability for the mathematical programming problem in Banach spaces*, Applied Mathematics and Optimization, 5 (1979), pp. 49-62.



Unité de recherche INRIA Lorraine, Technopôle de Nancy-Brabois, Campus scientifique,
615 rue du Jardin Botanique, BP 101, 54600 VILLERS LÈS NANCY
Unité de recherche INRIA Rennes, Irista, Campus universitaire de Beaulieu, 35042 RENNES Cedex
Unité de recherche INRIA Rhône-Alpes, 46 avenue Félix Viallet, 38031 GRENOBLE Cedex 1
Unité de recherche INRIA Rocquencourt, Domaine de Voluceau, Rocquencourt, BP 105, 78153 LE CHESNAY Cedex
Unité de recherche INRIA Sophia-Antipolis, 2004 route des Lucioles, BP 93, 06902 SOPHIA-ANTIPOLIS Cedex

Éditeur
INRIA, Domaine de Voluceau, Rocquencourt, BP 105, 78153 LE CHESNAY Cedex (France)
ISSN 0249-6399