

# Expansions for Steady-State Characteristics in (max, +)-Linear Systems

François Baccelli, Sven Hasenfuss, Volker Schmidt

► **To cite this version:**

| François Baccelli, Sven Hasenfuss, Volker Schmidt. Expansions for Steady-State Characteristics in  
| (max, +)-Linear Systems. RR-2785, INRIA. 1996. <inria-00073906>

**HAL Id: inria-00073906**

**<https://hal.inria.fr/inria-00073906>**

Submitted on 24 May 2006

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

INSTITUT NATIONAL DE RECHERCHE EN INFORMATIQUE ET EN AUTOMATIQUE

***Expansions for Steady-State Characteristics in  
(max, +)-Linear Systems***

François Baccelli – Sven Hasenfuss – Volker Schmidt

**N° 2785**

Janvier 1996

PROGRAMME 1



***Rapport  
de recherche***





## Expansions for Steady-State Characteristics in $(\max, +)$ -Linear Systems

François Baccelli \* – Sven Hasenfuss \*\* – Volker Schmidt \*\*\*

Programme 1 — Architectures parallèles, bases de données, réseaux et systèmes distribués  
Projet Mistral

Rapport de recherche n° 2785 — Janvier 1996 — 43 pages

**Abstract:** This paper gives finite and infinite expansion formulas for the expected value of functions of the steady state variables in open, stochastic  $(\max, +)$ -linear systems with Poisson input. Expansions for Laplace transforms, moments and tail functions of these steady state variables are considered as specific instances of our main formulas. Such  $(\max, +)$ -linear systems are known to allow to represent a class of discrete event networks called stochastic event graphs. A few examples of such event graphs pertaining to queueing theory are given in the paper in order to illustrate the proposed expansion method.

**Key-words:** Queueing network, stochastic Petri net, Poisson process, stochastic recurrence equation, stationary state variable, marked point process, factorial moment measure, admissibility, perturbation analysis.

*AMS classification:* 60K25, 60G55, 90B22.

(Résumé : *tsvp*)

\*INRIA-Sophia Antipolis (France), {Francois.Baccelli}@sophia.inria.fr

\*\* University of Ulm (Germany), {Sven.Hasenfuss}@mathematik.uni-ulm.de

\*\*\* University of Ulm (Germany), {Volker.Schmidt}@mathematik.uni-ulm.de

# Développements en série pour les caractéristiques stationnaires des systèmes $(\max, +)$ -linéaires

**Résumé :** Nous donnons des développements en série pour les valeurs moyennes de fonctions des variables d'état d'un système stochastique  $(\max, +)$ -linéaire ouvert avec des entrées formant un processus ponctuel de Poisson, en fonction de l'intensité de ce processus ponctuel.

Les développements en série des transformées de Laplace, des moments et des fonctions de répartition de ces variables sont présentés comme corollaires directs de ce résultat.

Les systèmes  $(\max, +)$ -linéaires correspondent à la classe des graphes d'événements stochastiques. Cette classe contient plusieurs exemples de réseaux de files d'attente dont nous étudions les développements en série à titre d'illustration.

**Mots-clé :** Réseau de files d'attente, réseau de Petri stochastique, processus de Poisson, équation de récurrence stochastique, régime stationnaire, processus ponctuel marqué, mesure factorielle, admissibilité, analyse de perturbation.

*Classification AMS:* 60K25, 60G55, 90B22.

## 1 Introduction

In the present paper, we focus on a certain class of functions  $f$  of the unique finite random vector solution  $W$  of a vectorial  $(\max, +)$ -linear functional equation of the form

$$W \circ \theta = A \otimes C \otimes W \oplus B \circ \theta, \quad (1)$$

which arises in the theory of open multidimensional discrete event systems. In this equation,

- $\theta$  is a shift operator defined on some probability space  $(\Omega, \mathcal{F}, \mathbb{P})$ ;
- $A, B$  and  $C$  are random matrices which characterize the topological and the stochastic structure of the system:  $A$  and  $B$  are associated with the system itself, and  $C$  with its input point process;
- The reference algebra is the so-called  $(\max, +)$  algebra: when operating on scalars,  $\oplus$  stands for  $\max$  and  $\otimes$  for  $+$ , whereas the  $\otimes$ -product of two matrices, say  $a$  of size  $p \times q$  and  $b$  of size  $q \times r$  is the  $p \times r$  matrix  $a \otimes b$  with entries

$$(a \otimes b)_{i,j} = \bigoplus_{k=1}^q a_{i,k} \otimes b_{k,j}$$

and the  $\oplus$ -sum of two matrices, say  $a$  and  $b$  both of size  $p \times q$  is the  $p \times q$  matrix  $a \oplus b$  with entries

$$(a \oplus b)_{i,j} = a_{i,j} \oplus b_{i,j}.$$

The aim of the present paper is to show that for all  $f : \mathbb{R} \rightarrow \mathbb{R}$  in this class of functions, under certain assumptions, whenever the input process is Poisson, the expectation  $\mathbb{E}[f(W^i)]$ , where  $W^i$  is the  $i$ -th component of the stationary state vector  $W$ , is finite and can be expanded into a finite or infinite power series with respect to the arrival intensity  $\lambda$ , where an algorithm is given to determine all coefficients of the expansion. This result will then be specialized to derive series expansions for

1. Laplace transforms  $\mathbb{E}[e^{-sW^i}]$ ,  $s \geq 0$ ;
2. Higher-order moments  $\mathbb{E}[(W^i)^\nu]$ ,  $\nu \in \mathbb{N} = \{1, 2, \dots\}$ ;
3. Tail functions  $\mathbb{P}(W^i > x)$ ,  $x \geq 0$ .

In all cases we derive explicit expressions for the coefficients of the expansion.

The present paper is a companion paper to [3], where the practical motivations for studying equations of type (1) are analyzed in great detail, and where similar expansions were first derived for the first moments of the  $W^i$  variables.

For sake of brevity, we have chosen not to recall these practical motivations in the present paper. So, for a more complete understanding of the general setting and of our examples, or for more examples, the reader should refer to [3].

## 2 General Setting

Let  $\alpha \in \mathbb{N}$  be an arbitrary natural number characterizing the dimension of the system. As it was shown in [3], the assumptions of the forthcoming three subsections are satisfied by  $\alpha$ -dimensional vectorial recurrence equations of the form

$$X_{n+1} = A_n \otimes X_n \oplus B_{n+1} \otimes T_{n+1} \quad (2)$$

which characterize the evolution of the state variables of any *open, input-connected stochastic event graph* with independent and i.i.d. firing times. Various incarnations of such equations will be considered in the examples.

### 2.1 Support and Monotonicity Assumptions

We assume that each entry of the  $\alpha \times \alpha$ -dimensional random matrix  $A_n$  is either a.s. non-negative or a.s. equal to  $\epsilon = -\infty$ , i.e.

$$(A_n)_{i,j} \geq 0 \quad \text{or} \quad (A_n)_{i,j} = \epsilon, \quad \text{a.s.} \quad (3)$$

and that all entries on the diagonal of  $A_n$  are non-negative, i.e.  $(A_n)_{i,i} \geq 0$ . We also assume that there exists an integer  $0 < \alpha' \leq \alpha$  such that the first  $\alpha'$  coordinates of the  $\alpha$ -dimensional random vectors  $B_n$  are non-negative, i.e.  $B_n^i \geq 0$  for all  $1 \leq i \leq \alpha'$ . Let  $D_0, D_1, \dots$  be the following  $\alpha$ -dimensional vectors:  $D_0 = B_0$ , and

$$D_k = \left( \bigotimes_{n=1}^k A_{-n} \right) \otimes B_{-k}, \quad \text{for } k \geq 1. \quad (4)$$

The  $\alpha'$  first coordinates of  $D_k$  are assumed to be non-decreasing in  $k$ :

$$0 \leq D_0^i \leq D_1^i \leq \dots, \quad \text{for all } i = 1, \dots, \alpha'. \quad (5)$$

### 2.2 Stochastic Assumptions

Throughout the paper, we assume that  $\{T_n\}$  is a stationary, homogeneous Poisson process with intensity  $\lambda$ , and  $\{Z_n\} = \{A_n, B_n\}$  is a stationary sequence of random matrices which is independent of  $\{T_n\}$ . We further assume that the underlying probability space  $(\Omega, \mathcal{F}, \mathbb{P})$  is equipped with a measurable shift  $\theta : \Omega \rightarrow \Omega$  which leaves the probability measure  $\mathbb{P}$  invariant, is  $\mathbb{P}$ -ergodic, and is such that  $\{Z_n\}$  is consistent with  $\theta$ , i.e.  $A_n = A \circ \theta^n$  and  $B_n = B \circ \theta^n$  for all integers  $n$ . Since the sequences  $\{T_n\}$  and  $\{Z_n\}$  are independent, we can also assume that  $T_{n+1} - T_n = T_1 \circ \theta^n$  for  $n \geq 1$  and  $T_{n+1} - T_n = (-T_0) \circ \theta^n$ , for  $n < 0$ , where  $T_1$  and  $T_0$  are the first positive and negative points of  $\{T_n\}$ , respectively. Put  $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$ . For  $n \in \mathbb{N}_0$  and  $q \in \mathbb{N}$ , let

$$H_{n,q} = \bigoplus_{i=1}^{\alpha'} \left\{ \left( A_{-(nq+1)} \otimes A_{-(nq+2)} \otimes \dots \otimes A_{-(n+1)q} \otimes (B_{-(n+1)q} \oplus O) \right)^i \right\}, \quad (6)$$

and

$$F_n = \bigoplus_{i=1}^{\alpha} \left\{ \left( A_{-1} \otimes A_{-2} \otimes \dots \otimes A_{-n} \otimes (B_{-n} \oplus O) \right)^i \right\}, \quad (7)$$

where  $O$  is the  $\alpha$ -dimensional column vector with all its components equal to zero. We assume that with probability one

$$\lim_{q \rightarrow \infty} H_{0,q} = \infty. \quad (8)$$

Besides this we will assume that for some  $r' \in \mathbb{N}$  large enough,

$$\lambda < r' \left[ \mathbb{E}[H_{0,r'}] \right]^{-1}, \quad (9)$$

and that  $\{H_{n,r'}\}_{n \geq 0}$  is a sequence of *1-dependent* random variables. An immediate consequence is the following result:

**Lemma 1** *There exists an  $r \in \mathbb{N}$ ,  $r \geq r'$  such that*

$$\lambda < (r-1) \left[ \mathbb{E}[H_{0,r}] \right]^{-1} \quad (10)$$

and  $\{H_{n,r}\}_{n \geq 0}$  is a sequence of *1-dependent* random variables.

In what follows, we will most often drop the index  $r$  and will only write  $H_n$  instead of  $H_{n,r}$ , but will always implicitly assume that  $r$  is chosen according to Lemma 1.

### 2.3 Steady-State Solution

Let  $\tau_n = T_{n+1} - T_n$ ,  $n \geq 0$ . By subtracting  $T_{n+1}$  on both sides of (2), it is easily checked that the state vector  $W_n$ , the components of which are given by  $W_n^i = X_n^i - T_n$ , satisfies the equation

$$W_{n+1} = A_n \otimes C(\tau_n) \otimes W_n \oplus B_{n+1}, \quad (11)$$

where, for all  $x \in \mathbb{R}$ ,  $C(x)$  is the  $\alpha \times \alpha$  matrix with all diagonal entries equal to  $-x$  and all non-diagonal entries equal to  $\epsilon$ . As it was shown in [2], under the above assumptions,  $\{W_n\}$  couples with a unique stationary sequence  $\{W \circ \theta^n\}$ , where  $W$  is the unique finite random variable solution of the functional equation

$$W \circ \theta = A \otimes C \otimes W \oplus B \circ \theta, \quad (12)$$

which is given by the series

$$W = B \oplus \bigoplus_{n \geq 1} A \circ \theta^{-1} \otimes C \circ \theta^{-1} \otimes \dots \otimes A \circ \theta^{-n} \otimes C \circ \theta^{-n} \otimes B \circ \theta^{-n}, \quad (13)$$



where  $C = C(-\tau_0)$ . Since  $C(x)$  commutes with any matrix,  $W$  admits the following equivalent representation:

$$W = B_0 \oplus \bigoplus_{n \geq 1} C(-T_{-n}) \otimes D_n. \quad (14)$$

## 2.4 Additional Notation

We now recall a class of polynomials which were obtained in the expansion for the expectation vector of  $W$  in [3]. These polynomials are given by the following formula

$$p_k(x_0, x_1, \dots, x_{k-1}) = \sum_{(i_0, i_1, \dots, i_{k-1}) \in S_k} (-1)^{\gamma_k(i_0, i_1, \dots, i_{k-1})} \frac{x_0^{i_0} x_1^{i_1}}{i_0! i_1!} \dots \frac{x_{k-1}^{i_{k-1}}}{i_{k-1}!}, \quad (15)$$

where

$$S_k = \left\{ (i_0, i_1, \dots, i_{k-1}) \in \{0, 1, \dots, k\}^k : i_0 + i_1 + \dots + i_{k-1} = k \text{ and if } \right. \\ \left. i_s = l > 1, \text{ then } i_{s-1 \bmod k} = i_{s-2 \bmod k} = \dots = i_{s-l+1 \bmod k} = 0 \right\}, \quad (16)$$

and

$$\gamma_k(i_0, i_1, \dots, i_{k-1}) = 1 + \sum_{s=0}^{k-1} \mathbf{1}(i_s > 0), \quad (17)$$

for all  $k \geq 1$ , with  $\mathbf{1}(i_s > 0) = 1$  whenever  $i_s > 0$  and  $\mathbf{1}(i_s > 0) = 0$  otherwise.

These polynomials satisfy the following properties, which can be found in [3]:

- *Translation Invariance:*

$$p_k(x_0 + u, x_1 + u, \dots, x_{k-1} + u) = p_k(x_0, x_1, \dots, x_{k-1}) \quad (18)$$

for all non-decreasing sequences of real numbers  $0 \leq x_0 \leq x_1 \leq \dots \leq x_{k-1}$ , for all  $u \geq 0$  and  $k = 2, 3, \dots$

- *Integral Representation:*

$$p_k(x_0, \dots, x_{k-1}) = \sum_{n=0}^{k-2} \int_{x_n}^{x_{n+1}} \left\{ p_{k-1}(\underbrace{v, \dots, v}_n, x_{n+1}, \dots, x_{k-1}) \right. \\ \left. - p_{k-1}(\underbrace{v, \dots, v}_{n+1}, x_{n+1}, \dots, x_{k-2}) \right\} dv, \quad (19)$$

for all non-decreasing sequences of real numbers  $0 \leq x_0 \leq x_1 \leq \dots \leq x_{k-1}$ ,  $k \geq 2$ , with  $p_1(x_0) = x_0$ .

### 3 Main Results

Throughout the paper, it will always be supposed that all the assumptions made on  $\{A_n, B_n\}$  in Section 2 hold. In this section we first give an explicit finite series expansion for the expectation of a class of functions of the stationary state variables  $W^i$ . This result will then be applied to get expansions for Laplace transforms, higher-order moments and tail functions of the variables  $W^i$ .

#### 3.1 General Theorem

Put  $\mathbb{R}^+ = [0, \infty)$  and let  $f : \mathbb{R}^+ \rightarrow \mathbb{R}^+$  be some Riemann-integrable function (e.g.  $f(x)$  monotone or continuous). Associated with  $f$ , we define the following class of functions:  $q_1(x_0) = f(x_0)$ , and

$$\begin{aligned}
 & q_{k+1}(x_0, x_1, \dots, x_k) \\
 = & \sum_{n=0}^k \binom{k}{n} (-1)^{k-n} F^{[k]}(x_n) - \sum_{n=0}^{k-1} \sum_{j=n}^{k-1} \binom{j}{n} (-1)^{j-n} F^{[j]}(x_n) \\
 & \left\{ p_{k-j}(x_{n+1}, \dots, x_{k-j+n}) - p_{k-j}(x_n, \dots, x_{k-j+n-1}) \right\}, \quad (20)
 \end{aligned}$$

for  $1 \leq k \leq m$ , with  $F^{[0]}(x) = f(x)$  and  $F^{[n]}(x)$ ,  $n \geq 1$ , is recursively defined by the a suitably chosen version of the indefinite Riemann-integral  $\int F^{[n-1]}(x) dx$ ,  $x \in \mathbb{R}^+$ . The functions  $p_k$  in (20) are the polynomials defined in (15).

**Theorem 1** *Let  $f : \mathbb{R}^+ \rightarrow \mathbb{R}^+$  be a Riemann-integrable, non-negative function which satisfies  $f(x) \leq c x^\nu$  for all  $x \geq 0$ , where  $\nu \in \mathbb{N}_0$  is a non-negative integer and  $c \in \mathbb{R}^+$  is a finite constant.*

1. *If the  $(\nu + m + 2)$ -th moment of  $H_n$  is finite, i.e.  $\mathbb{E}[(H_n)^{\nu+m+2}] < \infty$ , then, for  $1 \leq i \leq \alpha'$ ,*

$$\mathbb{E}[f(W^i)] = \sum_{k=0}^m \lambda^k \mathbb{E}[q_{k+1}(D_0^i, D_1^i, \dots, D_k^i)] + \mathcal{O}(\lambda^{m+1}). \quad (21)$$

2. *If there exists a  $\theta^* > 0$  such that for all  $\theta \in [0, \theta^*)$ ,*

$$\mathbb{E}[e^{\theta F_n}] \leq L_\theta (\phi(\theta))^n, \quad (22)$$

*for some finite functions  $\phi(\theta) > 1$  and  $L_\theta$ , then, for  $1 \leq i \leq \alpha'$ , the expectation  $\mathbb{E}[f(W^i)]$  is infinitely differentiable in  $\lambda$  for all  $\lambda$  in a right neighborhood of 0,*

$$\lim_{\lambda \downarrow 0} \frac{d^n}{d\lambda^n} \mathbb{E}[f(W^i)](\lambda) = \mathbb{E}[q_{n+1}(D_0^i, D_1^i, \dots, D_n^i)] \quad (23)$$

and

$$\mathbb{E}[f(W^i)] = \sum_{k=0}^{\infty} \lambda^k \mathbb{E}[q_{k+1}(D_0^i, D_1^i, \dots, D_k^i)]. \quad (24)$$

**Remark** In principle, it is possible to consider any version of the indefinite integral  $\int F^{[n-1]}(x) dx$  for  $F^{[n]}(x)$  in Theorem 1. However, depending on the particular function  $f(x)$ , a suitably chosen version of the indefinite integral may help to reduce computational efforts significantly.

For instance, taking  $f(x) = x$ , gives a function which satisfies the boundedness requirement on  $f(x)$  with  $c = 1$  and  $\nu = 1$ . For this function, it is advantageous to take  $F^{[n]}(x) = \int_0^x F^{[n-1]}(u) du$ , i.e.  $F^{[n]}(x) = \frac{x^{n+1}}{(n+1)!}$ , for  $n \in \mathbb{N}_0$ . Thus,

$$q_{k+1}(x_0, x_1, \dots, x_k) = \sum_{n=0}^k \binom{k}{n} (-1)^{k-n} \frac{(x_n)^{k+1}}{(k+1)!} - \sum_{n=0}^{k-1} \sum_{j=n}^{k-1} \binom{j}{n} (-1)^{j-n} \frac{(x_n)^{j+1}}{(j+1)!} \\ \left\{ p_{k-j}(x_{n+1}, \dots, x_{k-j+n}) - p_{k-j}(x_n, \dots, x_{k-j+n-1}) \right\}. \quad (25)$$

Elementary but tedious calculations show that the polynomials obtained in (25) are equal to those given by (15), see Section 6.2 of [3], i.e. Theorem 1 is actually a generalization of the result obtained in [3] and (25) gives another recurrence equation for the polynomials  $p_k$ .

The proof of Theorem 1 will be the object of Sections 4–6. In the following subsections, Theorem 1 will be applied to obtain expansion results for various functions of the stationary state vector  $W$ .

### 3.2 Laplace Transform

In order to derive an explicit finite series expansion of the Laplace transform of the components  $W^i$  of the stationary state vector  $W$ , we choose  $f(x) = e^{-sx}$ , for all  $x \in \mathbb{R}^+$  and  $s \geq 0$ . Since  $f(x)$  is bounded by 1 for all  $x \in \mathbb{R}^+$ , the following corollary is an immediate consequence of Theorem 1. In this case, choosing  $F^{[n]}(x) = \int_x^\infty F^{[n-1]}(u) du$  saves a lot of computational steps later on.

**Corollary 1** *In addition to the assumptions made on  $\{A_n, B_n\}$  above, assume the  $(m+2)$ -th moment of  $H_n$  to be finite for some  $m \in \mathbb{N}$ , i.e.  $\mathbb{E}[(H_n)^{m+2}] < \infty$ . Then, for all  $s > 0$ , and for  $1 \leq i \leq \alpha'$ ,*

$$\mathbb{E}\left[e^{-sW^i}\right] = \sum_{k=0}^m \lambda^k \mathbb{E}\left[q_{k+1}(s; D_0^i, D_1^i, \dots, D_k^i)\right] + \mathcal{O}(\lambda^{m+1}), \quad (26)$$

where  $q_1(s; x_0) = e^{-sx_0}$ , and

$$q_{k+1}(s; x_0, x_1, \dots, x_k) \\ = \frac{1}{s^k} \sum_{n=0}^k \binom{k}{n} (-1)^n e^{-sx_n} - \sum_{j=0}^{k-1} \frac{1}{s^j} \sum_{n=0}^j \binom{j}{n} (-1)^n e^{-sx_n} \\ \left\{ p_{k-j}(x_{n+1}, \dots, x_{k-j+n}) - p_{k-j}(x_n, \dots, x_{k-j+n-1}) \right\} \quad (27)$$

$$\begin{aligned}
 &= \frac{q_k(s; x_0, x_1, \dots, x_{k-1}) - q_k(s; x_1, x_2, \dots, x_k)}{s} \\
 &\quad - e^{-sx_0} [p_k(x_1, x_2, \dots, x_k) - p_k(x_0, x_1, \dots, x_{k-1})], \quad (28)
 \end{aligned}$$

for  $k = 1, \dots, m$ , and the functions  $p_k$  are the polynomials defined in (15).

Equation (28) gives a recurrence equation for the exponential-polynomial functions  $q_{k+1}$ , where the polynomials  $p_j$  of order  $1 \leq j \leq k$  have to be known in order to calculate  $q_{k+1}$ . To see that (28) is actually the same as (27), we split up the summations in (27) as follows:

$$\begin{aligned}
 &q_{k+1}(s; x_0, x_1, \dots, x_k) \\
 &= \frac{1}{s} \frac{1}{s^{k-1}} \left\{ e^{-sx_0} + \sum_{n=1}^{k-1} \left\{ \binom{k-1}{n} + \binom{k-1}{n-1} \right\} (-1)^n e^{-sx_n} + (-1)^k e^{-sx_k} \right\} \\
 &\quad - \frac{1}{s} \sum_{j=1}^{k-1} \frac{1}{s^{j-1}} \left\{ \sum_{n=0}^{j-1} \binom{j-1}{n} (-1)^n e^{-sx_n} \left\{ p_{k-j}(x_{n+1}, \dots, x_{k-j+n}) \right. \right. \\
 &\quad \quad \quad \left. \left. - p_{k-j}(x_n, \dots, x_{k-j+n-1}) \right\} \right. \\
 &\quad \quad \quad \left. - \sum_{n=0}^{j-1} \binom{j-1}{n} (-1)^n e^{-sx_{n+1}} \left\{ p_{k-j}(x_{n+2}, \dots, x_{k-j+n+1}) \right. \right. \\
 &\quad \quad \quad \left. \left. - p_{k-j}(x_{n+1}, \dots, x_{k-j+n}) \right\} \right\} \\
 &\quad - e^{-sx_0} \left\{ p_k(x_1, \dots, x_k) - p_{k-j}(x_0, \dots, x_{k-1}) \right\},
 \end{aligned}$$

which gives (28) after substituting  $j \rightarrow j + 1$  and grouping corresponding parts together.

### 3.2.1 Example: Pollaczek-Khinchine Formula for the $M/GI/1$ Queue

Corollary 1 can be specialized to the Pollaczek-Khinchine formula for the  $M/GI/1$  queue, which is the simplest instance of a one-dimensional open event graph. As it is known from [3], in this case we have  $\alpha = \alpha' = 1$ ,  $D_0 = 0$  and  $D_n = \sum_{i=1}^n \sigma_{-i}$ , for  $n \geq 1$ , where  $\{\sigma_i\}$  is the sequence of service times. From [3] we also know that the polynomials  $p_k(x_0, \dots, x_{k-1})$  are translation-invariant for  $k \geq 2$ , which implies that  $p_k(x_1, x_2, \dots, x_k) = p_k(0, x_2 - x_1, \dots, x_k - x_1)$ , for all  $k \geq 2$ .

By further noticing that the sequence  $\{\sigma_i\}$  forms a sequence of i.i.d. random variables, and therefore  $D_n - D_1 = \sum_{i=2}^n \sigma_{-i} \stackrel{d}{=} D_{n-1}$ , for all  $n \geq 1$ , we conclude that

$$\mathbb{E}[p_k(D_1, \dots, D_k)] = \mathbb{E}[p_k(D_0, \dots, D_{k-1})], \quad (29)$$

for all  $k \geq 2$ .

Now let  $S^*(s) := \mathbb{E}[e^{-s\sigma_0}]$ . The first coefficient of the expansion provided in Corollary 1 is given by  $\mathbb{E}[q_1(s; D_0)] = 1$ . Using (28) of Corollary 1, we get

$$\begin{aligned} \mathbb{E}[q_2(s; D_0, D_1)] &= \mathbb{E}\left[\frac{q_1(s; D_0) - q_1(s; D_1)}{s} - e^{-sD_0}\{p_1(D_1) - p_1(D_0)\}\right] \\ &= \mathbb{E}\left[\frac{1 - e^{-s\sigma_0}}{s} - \sigma_0\right] = \frac{1 - S^*(s)}{s} - \frac{1}{\mu}, \end{aligned} \quad (30)$$

where  $\mu^{-1} = \mathbb{E}[\sigma_0]$ . Making use of (29), we obtain from (28) that for  $k \geq 2$

$$\begin{aligned} \mathbb{E}[q_{k+1}(s; D_0, D_1, \dots, D_k)] &= \frac{\mathbb{E}[q_k(s; D_0, \dots, D_{k-1}) - q_k(s; D_1, \dots, D_k)]}{s} \\ &= \mathbb{E}[q_k(s; D_0, \dots, D_{k-1})] \frac{1 - S^*(s)}{s}, \end{aligned} \quad (31)$$

because from (27) and the translation-invariance property (18) of the polynomials  $p_k$ , it is easily seen that  $q_k(s; x_0 - u, \dots, x_{k-1} - u) = e^{su} q_k(s; x_0, \dots, x_{k-1})$  for  $k \geq 1$ , and therefore

$$\begin{aligned} \mathbb{E}[q_k(s; D_1, \dots, D_k)] &= \mathbb{E}\left[q_k\left(s; \sigma_{-1}, \dots, \sum_{i=1}^k \sigma_{-i}\right)\right] \\ &= \mathbb{E}\left[e^{-s\sigma_{-1}}\right] \mathbb{E}\left[q_k\left(s; 0, \sigma_{-2}, \dots, \sum_{i=2}^k \sigma_{-i}\right)\right] \\ &= \mathbb{E}\left[e^{-s\sigma_0}\right] \mathbb{E}[q_k(s; D_0, D_1, \dots, D_{k-1})]. \end{aligned}$$

Iterating (31)  $(k-1)$ -times and using (30) finally yields

$$\mathbb{E}[q_{k+1}(s; D_0, D_1, \dots, D_k)] = \left(\frac{1 - S^*(s)}{s} - \frac{1}{\mu}\right) \left(\frac{1 - S^*(s)}{s}\right)^{k-1}, \quad (32)$$

for all  $k \geq 1$ .

Since for the  $M/GI/1$  queue we have an infinite-order expansion of  $\mathbb{E}[e^{-sW}]$  in  $\lambda$ , we obtain from Corollary 1 that

$$\begin{aligned} \mathbb{E}[e^{-sW}] &= 1 + \sum_{k=1}^{\infty} \lambda^k \mathbb{E}[q_{k+1}(s; D_0, D_1, \dots, D_k)] \\ &= 1 + \lambda \left(\frac{1 - S^*(s)}{s} - \frac{1}{\mu}\right) \sum_{k=0}^{\infty} \lambda^k \left(\frac{1 - S^*(s)}{s}\right)^k \end{aligned}$$

$$\begin{aligned}
 &= 1 + \left( \frac{\lambda(1 - S^*(s))}{s} - \rho \right) \left( \frac{s}{s - \lambda(1 - S^*(s))} \right) \\
 &= \frac{(1 - \rho)s}{s - \lambda(1 - S^*(s))},
 \end{aligned}$$

i.e.

$$\mathbb{E}\left[e^{-sW}\right] = \frac{(1 - \rho)s}{s - \lambda(1 - S^*(s))},$$

a formula in which we recognize the Pollaczek-Khinchine formula for the Laplace transform of stationary waiting times in the  $M/GI/1$  queue.

### 3.2.2 Example: Queues in Tandem

Consider a network of  $\alpha$  single-server FIFO queues with infinite capacity in tandem, with all queues initially empty,  $\alpha = \alpha' \geq 2$ . Let  $\sigma_n^i$  denote the service time of customer  $n$  in station  $i$ . The  $i$ -th component  $W^i$  of the random vector  $W$  describes the stationary waiting time of a randomly chosen customer until the beginning of his service on station  $i$ . It was shown in [3] that in this case, the vectors  $D_n$  have the following form:

$$D_n^i = \max_{1 \leq l_n \leq \dots \leq l_1 \leq i} \left\{ \sum_{k=l_1}^{i-1} \sigma_0^k + \sum_{k=l_2}^{l_1} \sigma_{-1}^k + \dots + \sum_{k=l_n}^{l_{n-1}} \sigma_{-n+1}^k + \sum_{k=1}^{l_n} \sigma_{-n}^k \right\}. \quad (33)$$

**Expansions — Deterministic Service Times Case** Let  $\sigma^i$  denote the service time in queue  $i \in \{1, \dots, \alpha\}$ . Without loss of generality we can and will assume that  $\sigma^1 \leq \sigma^2 \leq \dots \leq \sigma^\alpha$ . In the other case, say  $\sigma^i > \sigma^{i+1}$  for some  $i < \alpha$ , we can consider the  $i$ -th queue and the  $(i+1)$ -th queue as *one* single-server queue with service time  $\sigma^i + \sigma^{i+1}$  because in front of the  $(i+1)$ -th server the waiting room is always empty. In this case

$$D_n = \begin{pmatrix} n\sigma^1 \\ \sigma^1 + n\sigma^2 \\ \sigma^1 + \sigma^2 + n\sigma^3 \\ \sigma^1 + \sigma^2 + \sigma^3 + n\sigma^4 \\ \vdots \\ \sum_{k=1}^{\alpha-1} \sigma^k + n\sigma^\alpha \end{pmatrix}. \quad (34)$$

Due to the translation-invariance (18) of the polynomials  $p_k$ , for all  $j = 0, \dots, k-2$ , and for all  $n = 0, \dots, j$ ,

$$p_{k-j}(D_{n+1}^i, \dots, D_{k-j+n}^i) = p_{k-j}(D_n^i, \dots, D_{k-j+n-1}^i).$$

So, from Corollary 1, for all  $m \geq 1$ ,  $\mathbb{E}[e^{-sW^i}]$  admits an expansion of order  $m$  in  $\lambda$ , with the  $k$ -th coefficient  $c_k = q_{k+1}(D_0^i, D_1^i, \dots, D_k^i)$ ,  $k = 0, 1, \dots, m$ , given by the formula

$$c_k = \frac{1}{s^k} e^{-s \sum_{l=1}^{i-1} \sigma^l} \sum_{n=0}^k \binom{k}{n} (-1)^n e^{-sn\sigma^i} - \frac{1}{s^{k-1}} \sigma^i e^{-s \sum_{l=1}^{i-1} \sigma^l} \sum_{n=0}^{k-1} \binom{k-1}{n} (-1)^n e^{-sn\sigma^i}. \quad (35)$$

**Expansions — Random Identical Service Times Case** Assume now that  $\sigma_{-n}^i = \sigma(n)$  for all  $i$ , a model which was for instance considered in [3]. Assume that

$$\mathbb{E}[(\sigma(n))^{m+2}] < \infty. \quad (36)$$

The integrability condition of Corollary 1 is then satisfied. From (33) it follows that in this case the random variables  $D_n^i$  are given by the following expressions:

$$\begin{aligned} D_0^i &= (i-1)\sigma(0) \\ D_n^i &= \sigma(1) + \dots + \sigma(n) + (i-1) \max\{\sigma(0), \dots, \sigma(n)\}, \quad n \geq 1. \end{aligned}$$

So, for instance, under moment condition (36) for  $m = 2$ , the coefficients  $c_k$  of the second order expansion

$$\mathbb{E}[e^{-sW^i}] = c_0 + \lambda c_1 + \lambda^2 c_2 + \mathcal{O}(\lambda^3)$$

are given by the following formulas:

$$\begin{aligned} c_0 &= \mathbb{E}[e^{-s(i-1)\sigma(0)}], \\ c_1 &= \frac{\mathbb{E}[e^{-s(i-1)\sigma(0)}] - \mathbb{E}[e^{-s\{\sigma(1)+(i-1)(\sigma(0) \vee \sigma(1))\}}]}{s} \\ &\quad - \mathbb{E}[e^{-s(i-1)\sigma(0)} \{\sigma(1) + (i-1)(\sigma(1) - \sigma(0))^+\}], \\ c_2 &= \frac{\mathbb{E}[e^{-s(i-1)\sigma(0)}] - 2\mathbb{E}[e^{-s\{\sigma(1)+(i-1)(\sigma(0) \vee \sigma(1))\}}]}{s^2} + \mathbb{E}[e^{-s\{\sigma(1)+\sigma(2)+(i-1)(\sigma(0) \vee \sigma(1) \vee \sigma(2))\}}] \\ &\quad - \frac{\mathbb{E}[e^{-s(i-1)\sigma(0)} \{\sigma(1) + (i-1)(\sigma(1) - \sigma(0))^+\}]}{s} \\ &\quad + \frac{\mathbb{E}[e^{-s\{\sigma(1)+(i-1)(\sigma(0) \vee \sigma(1))\}} \{\sigma(2) + (i-1)(\sigma(2) - (\sigma(0) \vee \sigma(1)))^+\}]}{s} \\ &\quad - \frac{\mathbb{E}[e^{-s(i-1)\sigma(0)} \{(\sigma(2) + (i-1)(\sigma(2) - (\sigma(0) \vee \sigma(1)))^+)^2 - (\sigma(1) + (i-1)(\sigma(1) - \sigma(0))^+)^2\}]}{2}. \end{aligned}$$

### 3.3 Higher-Order Moments

We can derive a similar expansion for higher-order moments of  $W^i$ ,  $1 \leq i \leq \alpha'$ . Choosing  $f(x) = x^\nu$  and setting  $c = 1$  allows us to apply Theorem 1 to get the following result:

**Corollary 2** *In addition to the foregoing assumptions on  $\{A_n, B_n\}$  above, assume that the  $(\nu + m + 2)$ -th moment of  $H_n$  is finite for some  $m, \nu \in \mathbb{N}$ , i.e.  $\mathbb{E}[(H_n)^{\nu+m+2}] < \infty$ . Then*

$$\mathbb{E}[(W^i)^\nu] = \sum_{k=0}^m \lambda^k \mathbb{E}\left[p_{k+1}^{(\nu)}(D_0^i, D_1^i, \dots, D_k^i)\right] + \mathcal{O}(\lambda^{m+1}), \quad (37)$$

for  $1 \leq i \leq \alpha'$ , where  $p_1^{(\nu)}(x_0) = x_0^\nu$ , and

$$\begin{aligned} & p_{k+1}^{(\nu)}(x_0, x_1, \dots, x_k) \\ = & \frac{\nu!}{(\nu+k)!} \sum_{n=0}^k \binom{k}{n} (-1)^{k-n} x_n^{\nu+k} - \sum_{j=0}^{k-1} \frac{\nu!}{(\nu+j)!} \sum_{n=0}^j \binom{j}{n} (-1)^{j-n} x_n^{\nu+j} \\ & \left\{ p_{k-j}(x_{n+1}, \dots, x_{k-j+n}) - p_{k-j}(x_n, \dots, x_{k-j+n-1}) \right\}, \quad (38) \end{aligned}$$

for  $1 \leq k \leq m$ , and the functions  $p_k$  are the polynomials given by (15).

**Remark** Having in mind that

$$\mathbb{E}[(W^i)^\nu] = \lim_{s \downarrow 0} \left\{ (-1)^\nu \frac{d^\nu}{ds^\nu} \mathbb{E}\left[e^{-s W^i}\right] \right\},$$

whenever the  $\nu$ -th moment of  $W^i$  is finite, one is tempted to think that the polynomials  $p_k^{(\nu)}(x_0, \dots, x_{k-1})$  can be obtained from the exponential-polynomial functions  $q_k(s; x_0, \dots, x_{k-1})$  by differentiating and taking the limit  $s \downarrow 0$ , i.e.

$$p_k^{(\nu)}(x_0, x_1, \dots, x_{k-1}) = \lim_{s \downarrow 0} \left\{ (-1)^\nu \frac{d^\nu}{ds^\nu} q_k(s; x_0, x_1, \dots, x_{k-1}) \right\}, \quad (39)$$

for all  $k \geq 1$ . Elementary calculations show that equality (39) holds indeed.

**Example: Blocking Queues** The following finite capacity queueing system with blocking was considered in [3]: the system consists of four single-server FIFO stations in tandem. The first station, which is fed by the arrival point process, has an infinite capacity buffer, whereas all other stations have no buffering capacity. The blocking mechanism is that of “blocking after service”, i.e. in each station, a customer can always start his service but once his service is completed, the customer can only proceed to



the downstream station whenever this one is empty (this is also called *manufacturing blocking*). Taking as state variables the variables  $W_n^i$ , where  $W_n^i$  gives the time which elapses between the  $n$ -th external arrival and the time when customer  $n$  leaves station  $i$ , it was shown in [3] that the corresponding  $D_n$  vectors are given by the following formulas: whenever the system starts empty and whenever the service times are deterministic and equal to  $\sigma^i$  in station  $i$  with  $\sigma^1 \leq \sigma^2 \leq \sigma^3 \leq \sigma^4$ :

$$D_0 = \begin{pmatrix} \sigma^1 \\ \sigma^1 + \sigma^2 \\ \sigma^1 + \sigma^2 + \sigma^3 \\ \sigma^1 + \sigma^2 + \sigma^3 + \sigma^4 \end{pmatrix}, \quad D_1 = \begin{pmatrix} \sigma^1 + \sigma^2 \\ \sigma^1 + \sigma^2 + \sigma^3 \\ \sigma^1 + \sigma^2 + \sigma^3 + \sigma^4 \\ \sigma^1 + \sigma^2 + \sigma^3 + 2\sigma^4 \end{pmatrix}, \quad (40)$$

$$D_2 = \begin{pmatrix} \sigma^1 + \sigma^2 + \sigma^3 \\ \sigma^1 + \sigma^2 + \sigma^3 + \sigma^4 \\ \sigma^1 + \sigma^2 + \sigma^3 + 2\sigma^4 \\ \sigma^1 + \sigma^2 + \sigma^3 + 3\sigma^4 \end{pmatrix}, \quad D_n = \begin{pmatrix} \sigma^1 + \sigma^2 + \sigma^3 + (n-2)\sigma^4 \\ \sigma^1 + \sigma^2 + \sigma^3 + (n-1)\sigma^4 \\ \sigma^1 + \sigma^2 + \sigma^3 + (n)\sigma^4 \\ \sigma^1 + \sigma^2 + \sigma^3 + (n+1)\sigma^4 \end{pmatrix}, \quad \forall n \geq 3. \quad (41)$$

So, as a direct application of Corollary 2, the coefficients  $c_k$  of the expansion of the second moment of the stationary “waiting time” in station 1

$$\mathbb{E}[(W^1)^2] = c_0 + c_1\lambda + c_2\lambda^2 + \mathcal{O}(\lambda^3)$$

are given by the following formulas:

$$\begin{aligned} c_0 &= (\sigma^1)^2 \\ c_1 &= (\sigma^1)(\sigma^2)^2 + \frac{1}{3}(\sigma^2)^3 \\ c_2 &= \frac{(\sigma^3)^4 - (\sigma^1)^4}{12} - \frac{(\sigma^1)(\sigma^2)^3 - (\sigma^1)(\sigma^3)^3 - (\sigma^2)(\sigma^3)^3}{3} + \frac{1}{2}(\sigma^2)^2(\sigma^3)^2 + (\sigma^1)(\sigma^2)(\sigma^3)^2. \end{aligned}$$

### 3.4 The Tail Function of $W^i$

Another interesting function of  $W^i$  is its distribution function, or equivalently its tail function  $\mathbb{P}(W^i > \xi)$ ,  $\xi \geq 0$ . Before we can apply Theorem 1 to derive a finite series expansion of  $\mathbb{P}(W^i > \xi)$ , we first notice that  $\mathbb{P}(W^i > \xi) = \mathbb{E}[\mathbf{1}(W^i > \xi)]$  and thus choose  $f(x) = \mathbf{1}(x > \xi)$  for all  $x \geq 0$ . Clearly,  $0 \leq f(x) \leq 1$ , for all  $x \in \mathbb{R}^+$ . Applying Theorem 1, we use the version  $F^{[n]}(x) = \frac{1}{n!}(x - \xi)^n \mathbf{1}(x > \xi)$ , for all  $n \in \mathbb{N}_0$ , of the indefinite integral  $\int F^{[n-1]}(x) dx$ . So we obtain

**Corollary 3** *In addition to the foregoing assumptions on  $\{A_n, B_n\}$ , assume the  $(m+2)$ -th moment of  $H_n$  to be finite for some  $m \in \mathbb{N}$ , i.e.  $\mathbb{E}[(H_n)^{m+2}] < \infty$ . Then, for all  $\xi \geq 0$  and for  $1 \leq i \leq \alpha'$ ,*

$$\mathbb{P}(W^i > \xi) = \sum_{k=0}^m \lambda^k \mathbb{E}[\tilde{q}_{k+1}(D_0^i, D_1^i, \dots, D_k^i)] + \mathcal{O}(\lambda^{m+1}), \quad (42)$$

where  $\tilde{q}_1(x_0) = \mathbf{1}(x_0 > \xi)$ ,

$$\begin{aligned} & \tilde{q}_{k+1}(x_0, x_1, \dots, x_k) \\ &= \sum_{n=0}^k \mathbf{1}(x_n > \xi) \frac{(-1)^{k-n}}{(k-n)! n!} (x_n - \xi)^k - \sum_{n=0}^{k-1} \mathbf{1}(x_n > \xi) \sum_{j=n}^{k-1} \frac{(-1)^{j-n}}{(j-n)! n!} (x_n - \xi)^j \\ & \quad \left\{ p_{k-j}(x_{n+1}, \dots, x_{k-j+n}) - p_{k-j}(x_n, \dots, x_{k-j+n-1}) \right\}, \end{aligned} \quad (43)$$

for  $1 \leq k \leq m$ , and the functions  $p_k$  are the polynomials defined in (15).

**Example: Queues in Tandem — Continued** We continue the example of tandem queues with random and identical service times in all stations. Under the assumption that the service times have moments of order  $m + 2$ , the first three coefficients  $c_k$ ,  $k = 0, 1, 2$ , of the expansion

$$\mathbb{P}(W^i > \xi) = \sum_{k=0}^m c_k \lambda^k + \mathcal{O}(\lambda^{m+1})$$

are given by the formulas:

$$\begin{aligned} c_0 &= \mathbb{P}\left((i-1)\sigma(0) > \xi\right), \\ c_1 &= \mathbb{E}\left[\mathbf{1}\left((i-1)\sigma(0) \leq \xi\right) \cdot \left(\sigma(1) + (i-1)(\sigma(0) \vee \sigma(1)) - \xi\right)^+\right], \\ c_2 &= \mathbb{E}\left[\mathbf{1}\left((i-1)\sigma(0) \leq \xi\right) \cdot \left\{ \frac{1}{2} \left( (\sigma(1) + \sigma(2) + (i-1)(\sigma(0) \vee \sigma(1) \vee \sigma(2)) - \xi)^+ \right)^2 \right. \right. \\ & \quad \left. \left. - \left( \sigma(1) + (i-1)(\sigma(0) \vee \sigma(1)) - \xi \right)^+ \left( \sigma(1) + \sigma(2) + (i-1)(\sigma(0) \vee \sigma(1) \vee \sigma(2)) - \xi \right)^+ \right\} \right]. \end{aligned}$$

## 4 Factorial Moment Expansion

### 4.1 Preliminary Results

In this subsection we will recall the general method for proving part 1 of our main theorem and also give some simple inequalities which turn out to be useful later on.

#### 4.1.1 General Expansion Theorem

In order to prove the series expansion stated in Theorem 1 we will use a general idea which consists in expanding the expectation of vector-valued functionals of marked point processes. This method, which was introduced in [4], was shown to be quite useful for approximating the characteristics of several stochastic models (see e.g. [3], [5], [6], [7] or [8]).

**Expansion Kernels** For any given natural number  $\alpha$ , let  $\psi$  be an  $\mathbb{R}^\alpha$ -valued functional of a marked point process, i.e. a measurable mapping  $\psi : \mathcal{M} \times \mathcal{K}^\infty \rightarrow \mathbb{R}^\alpha$ , where  $\mathcal{M}$  is the space of all realizations of the point process  $\{T_n\}$  and  $\mathcal{K}^\infty$  is the space of all sequences  $Z = \{Z_n\}$  of potential marks. We assume that the *mark space*  $\mathcal{K}$  is a complete separable metric space. Note that the sequence  $\{T_n\}$  of points may be infinite, finite or empty, whereas the sequence  $Z = \{Z_n\}$  of potential marks is always two-sided infinite. Let  $Z_n$  denote the mark of point  $T_n$ . We represent a realization  $\{t_n\}$  of the point process  $\{T_n\}$  by the counting measure  $\mu = \sum_n \delta_{t_n}$ . By  $o$  we denote the null measure, representing an input with no arrivals (i.e.  $o(\mathbb{R}) = 0$ ).

For every  $x \in \mathbb{R}$ , let the restriction  $\mu|_x$  of  $\mu \in \mathcal{M}$  be defined by

$$\mu|_x(D) = \mu(D \cap (x, \infty)).$$

Furthermore, for any  $x \in \mathbb{R}$  and  $z \in \mathcal{K}^\infty$ , let

$$\psi_x(\mu, z) = \psi(\mu|_x + \delta_x, z) - \psi(\mu|_x, z). \quad (44)$$

Let  $k \geq 1$  be an arbitrary, but fixed integer. For any  $x_1, \dots, x_k \in \mathbb{R}$ , let  $\psi_{x_1, \dots, x_k}$  be defined by iteration of the mapping  $\psi \rightarrow \psi_x$ , i.e.

$$\psi_{x_1, \dots, x_k}(\mu, z) = \left( \dots (\psi_{x_1})_{x_2} \dots \right)_{x_k}(\mu, z).$$

Note that the functional  $\psi_{x_1, \dots, x_k}$  can be written in the form

$$\psi_{x_1, \dots, x_k}(\mu, z) = \begin{cases} \sum_{j=0}^k (-1)^{k-j} \sum_{\pi \in K_{k,j}} \psi(\mu|_{x_\pi} + \sum_{i \in \pi} \delta_{x_i}, z) & \text{for } x_1 < \dots < x_k, \\ 0 & \text{otherwise,} \end{cases} \quad (45)$$

where  $K_{k,j}$  denotes the collection of all subsets of  $\{1, \dots, k\}$  containing  $j$  elements. Following [4], we call the functional  $\psi$  *continuous at infinity* if

$$\lim_{x \rightarrow \infty} \psi(\mu|_x + \nu, z) = \psi(\nu, z), \quad \lim_{x \rightarrow -\infty} \psi(\mu|_x, z) = \psi(\mu, z) \quad (46)$$

for all  $\mu, \nu \in \mathcal{M}, z \in \mathcal{K}^\infty$  with  $\nu(\mathbb{R}) < \infty$ .

**Representation Formula** For the stationary Poisson process  $\{T_n\}$  with intensity  $\lambda$ , and for the stationary sequence  $Z = \{Z_n\}$  of  $\mathcal{K}$ -valued random variables which is independent of  $\{T_n\}$ , let  $P_\lambda$  denote the distribution of  $\{T_n\}$ , and  $Q$  the distribution of  $Z$ .

A slight variant of the following result is given in [8], where the general expansion derived in [4] is specified to the Poisson case. With  $\psi^i$  we denote the  $i$ -th component of the  $\mathbb{R}^\alpha$ -valued functional  $\psi$ .

**Theorem 2** Let  $m \geq 1$  be a fixed integer. If the functional  $\psi$  is continuous at infinity, if

$$\int_{\mathbb{R}^k} \int_{\mathcal{K}^\infty} \int_{\mathcal{M}} \left| \psi_{x_1, \dots, x_k}^i(\mu, z) \right| P_\lambda(d\mu) Q(dz) d(x_1, \dots, x_k) < \infty, \quad (47)$$

for all  $k = 1, \dots, m$ , and if

$$\limsup_{\lambda \rightarrow 0} \int_{\mathbb{R}^{m+1}} \int_{\mathcal{K}^\infty} \int_{\mathcal{M}} \left| \psi_{x_1, \dots, x_{m+1}}^i(\mu, z) \right| P_\lambda(d\mu) Q(dz) d(x_1, \dots, x_{m+1}) < \infty, \quad (48)$$

then

$$\begin{aligned} & \mathbb{E} \left[ \psi^i(\{(T_n, Z_n)\}) \right] \\ &= \mathbb{E} \left[ \psi^i(o, \{Z_n\}) \right] + \sum_{k=1}^m \lambda^k \int_{\mathbb{R}^k} \mathbb{E} \left[ \psi_{x_1, \dots, x_k}^i(o, \{Z_n\}) \right] d(x_1, \dots, x_k) + \mathcal{O}(\lambda^{m+1}). \end{aligned} \quad (49)$$

#### 4.1.2 Simple Inequalities

A useful bound is given by

**Lemma 2** For  $k, r \in \mathbb{N}$  and  $1 \leq i \leq \alpha$

$$D_{kr}^i \leq H_0 + H_1 + \dots + H_{k-1}. \quad (50)$$

*Proof* It suffices to note that

$$\begin{aligned} D_{kr}^i &= \left( \left( \bigotimes_{n=1}^{kr} A_{-n} \right) \otimes B_{-kr} \right)^i = \left( \left( \bigotimes_{p=0}^{k-1} \bigotimes_{q=0}^{r-1} A_{-(pr+q+1)} \right) \otimes B_{-kr} \right)^i \\ &\leq \bigotimes_{p=0}^{k-1} \left( \bigotimes_{q=0}^{r-1} A_{-(pr+q+1)} \otimes (B_{-(p+1)r} \otimes O) \right)^i \leq \bigotimes_{p=0}^{k-1} H_p. \end{aligned}$$

□

We will also use the next simple result later on.

**Lemma 3** For  $x_1, \dots, x_k \in \mathbb{R}_0^+$ ,  $i_1, \dots, i_k, k \in \mathbb{N}$  and  $n \in \mathbb{N}_0$  the following (conventional algebra) inequalities hold:

$$(a) \quad x_1^{i_1} \cdots x_k^{i_k} \leq x_1^{i_1 + \dots + i_k} + \dots + x_k^{i_1 + \dots + i_k}, \quad (51)$$

$$(b) \quad (x_1 + \dots + x_k)^n \leq k^n (x_1^n + \dots + x_k^n). \quad (52)$$

*Proof* In order to prove (a) note that

$$\prod_{j=1}^k x_j^{i_j} \leq \prod_{j=1}^k \left( \max_{1 \leq j \leq k} \{x_j\} \right)^{i_j} = \left( \max_{1 \leq j \leq k} \{x_j\} \right)^{\sum_{j=1}^k i_j} \leq \sum_{j=1}^k x_j^{\sum_{j=1}^k i_j}.$$

On the other hand, using (a) gives (b) for  $n \geq 1$  by the following reasoning:

$$\left( \sum_{j=1}^k x_j \right)^n = \sum_{\substack{(i_1, \dots, i_k): \\ i_1 + \dots + i_k = n}} \binom{k}{i_1 \dots i_k} \prod_{j=1}^k x_j^{i_j} \leq \sum_{\substack{(i_1, \dots, i_k): \\ i_1 + \dots + i_k = n}} \binom{k}{i_1 \dots i_k} \sum_{j=1}^k x_j^n = k^n \sum_{j=1}^k x_j^n.$$

For  $n = 0$ , (b) is immediate. □

## 4.2 Proof of the Finite Expansion Formula

In this subsection we will use Theorem 2 in order to show that the finite expansion of  $\mathbb{E}[f(W^i)]$  given in (21) holds. To prove that the conditions of Theorem 2 are fulfilled under the assumptions made above, we can proceed in a way which is similar to that used in [3] for expanding the first moment  $\mathbb{E}[W^i]$ . Consider the following functional  $\psi$  given by

$$\begin{aligned} \psi(\mu, z) &= (\psi^i(\mu, z))_{i=1, \dots, \alpha} \\ &= \left( f \left( d_0^i \oplus \bigoplus_{n=1}^{\mu((-\infty, 0))} (d_n^i \otimes t_{-n}) \right) \right)_{i=1, \dots, \alpha}, \end{aligned} \quad (53)$$

where  $\mu = \sum_n \delta_{t_n}$  denotes the realization of the arrival process  $\{T_n\}$ . Furthermore, let  $z = \{a_n, b_n\}$ , where  $a_n$  and  $b_n$  denote the realizations of the random matrices  $A_n$  and  $B_n$ , respectively. In the same vein, let  $d_0^i = b_0^i$  and  $d_k^i = \left( \left( \bigotimes_{n=1}^k a_{-n} \right) \otimes b_{-k} \right)^i$ , for  $k \geq 1$ , denote the realizations of the  $i$ -th component of the random vectors  $D_k$  defined in (4). In what follows, we are going to look only at those components  $\psi^i$  of the functional  $\psi$ , where  $1 \leq i \leq \alpha'$ , in accordance with the statement of Theorem 1.

The fact that  $\psi$  is a.s. continuous at infinity whenever  $\rho < 1$  follows directly from the backward monotone construction used in Chapter 7 of [2], where  $\rho = \lambda a$  and  $a$  is the maximal  $(\max, +)$ -Lyapunov exponent of the sequence  $\{A_n\}$ .

### 4.2.1 Integrability

We first show that the expectation  $\mathbb{E}[f(W^i)]$  is finite for each  $i \in \{1, \dots, \alpha'\}$  whenever the conditions of Theorem 1 are satisfied. Let  $r \in \mathbb{N}$  be chosen according to Lemma 1. Since  $f(\cdot)$  is assumed to be

bounded by  $f(x) \leq cx^\nu$ , for all  $x \in \mathbb{R}^+$ , we get

$$\begin{aligned}
 \psi^i(\mu, z) &= f\left(d_0^i \oplus \bigoplus_{n=1}^{\mu((-\infty, 0))} (d_n^i + t_{-n})\right) \leq c \left(d_0^i \oplus \bigoplus_{n=1}^{\mu((-\infty, 0))} (d_n^i + t_{-n})\right)^\nu \\
 &\leq c \left(b_0^i + \sup_{p \geq 0} \max_{1 \leq q \leq r} \left\{d_{pr+q} + t_{-(pr+q)}\right\}^+\right)^\nu \leq c \left(b_0^i + \sup_{p \geq 0} \left\{d_{(p+1)r} + t_{-(p+1)r}\right\}^+\right)^\nu \\
 &\leq c \left(b_0^i + \sup_{p \geq 0} \left\{h_0 + \dots + h_p + t_{-pr}\right\}^+\right)^\nu \\
 &= c \left(b_0^i + \sup_{p \geq 0} \left\{h_0 + \sum_{\{1 \leq 2k \leq p\}} (h_{2k} + (t_{-2kr} - t_{-(2k-1)r})) \right. \right. \\
 &\quad \left. \left. + \sum_{\{1 \leq 2k-1 \leq p\}} (h_{2k-1} + (t_{-(2k-1)r} - t_{-(2k-2)r}))\right\}^+\right)^\nu \\
 &\leq c \left(\max_j \{b_0^j\} + h_0 + \varphi^{(0)}(\mu, z) + \varphi^{(1)}(\mu, z)\right)^\nu, \tag{54}
 \end{aligned}$$

where  $\varphi^{(j)}(\mu, z)$ ,  $j = 0, 1$ , denotes the realization of the random variable

$$\varphi^{(j)}(\{T_n\}, \{Z_n\}) = \sup_{p \geq 1} \left\{ \sum_{k=1}^p (H_{2k-j} + (T_{-(2k-j)r} - T_{-(2k-j-1)r})) \right\}^+, \quad j = 0, 1, \tag{55}$$

for a given  $(\mu, z)$  and  $\{x\}^+ = \max\{0, x\}$  for all  $x \in \mathbb{R}$ . Applying Lemma 3 on (54) yields

$$\left| \psi^i(\mu, z) \right| \leq c 4^\nu \left\{ \left(\max_j \{b_0^j\}\right)^\nu + (h_0)^\nu + \left(\varphi^{(0)}(\mu, z)\right)^\nu + \left(\varphi^{(1)}(\mu, z)\right)^\nu \right\}.$$

Because of the general stochastic assumptions made in Section 2, we see that  $\varphi^{(j)}(\{T_n\}, \{Z_n\})$ ,  $j = 0, 1$ , is actually the maximum of a random walk with negative drift, since under (10), for  $j = 0, 1$ , we have

$$\mathbb{E} \left[ H_{2k-j} + (T_{-(2k-j)r} - T_{-(2k-j-1)r}) \right] = \mathbb{E} [H_0] - \frac{r}{\lambda} < \mathbb{E} [H_0] - \frac{r-1}{\lambda} < 0.$$

The finiteness of  $\mathbb{E} [(W^i)^\nu]$  now follows from the well known fact that the  $\nu$ -th moment of the maximum of a random walk with negative drift is finite whenever the  $(\nu + 1)$ -th moment of its increments is finite (see e.g. Theorem VIII.2.1 in [1]). But this is true since

$$\begin{aligned}
 \mathbb{E} \left[ \left( H_{2k-j} + (T_{-(2k-j)r} - T_{-(2k-j-1)r}) \right)^{\nu+1} \right] &\leq 2^{\nu+1} \left\{ \mathbb{E} [(H_0)^{\nu+1}] + (\nu+1)! \left(\frac{r}{\lambda}\right)^{\nu+1} \right\} \\
 &< \infty,
 \end{aligned}$$

for  $j = 0, 1$ , whenever  $\mathbb{E} [(H_0)^{\nu+1}] < \infty$ .

### 4.2.2 Proof of Condition (47): $k = 1$

We first prove that condition (47) holds for  $k = 1$ . Let  $x \in [t_{-l}, t_{-(l-1)})$ , for some  $l \in \{1, 2, \dots\}$ . The first expansion kernel of the functional  $\psi$  can then be written as

$$\begin{aligned} \psi_x^i(\mu, z) &= \psi^i(\mu|^{x+\delta_x}, z) - \psi^i(\mu|^{x}, z) \\ &= f\left(d_0^i \oplus \bigoplus_{n=1}^{l-1} (d_n^i + t_{-n}) \oplus (d_l^i + x)\right) - f\left(d_0^i \oplus \bigoplus_{n=1}^{l-1} (d_n^i + t_{-n})\right). \end{aligned}$$

It is easily seen that  $\psi_x^i(\mu, z)$  is equal to zero whenever the last term  $(d_l^i + x)$  of the sum in the argument of  $f(\cdot)$  corresponding to  $\psi^i(\mu|^{x+\delta_x}, z)$  is smaller than or equal to all other terms in this sum; so

$$\begin{aligned} \left| \psi_x^i(\mu, z) \right| &= \left| f(d_l^i + x) - f\left(d_0^i \oplus \bigoplus_{n=1}^{l-1} (d_n^i + t_{-n})\right) \right| \mathbf{1}\left(d_l^i + x > d_0^i \oplus \bigoplus_{n=1}^{l-1} (d_n^i + t_{-n})\right) \\ &\leq c \left(d_l^i + x\right)^\nu \mathbf{1}\left(d_l^i + x > 0\right), \end{aligned} \tag{56}$$

since  $d_l^i + x > d_0^i \geq 0$ .

We assumed that  $l$  was chosen such that  $x \in [t_{-l}, t_{-(l-1)})$ . Let us first consider the case when  $l \leq (r-1)^2$ . Then

$$\left(d_l^i + x\right)^\nu \mathbf{1}\left(d_l^i + x > 0\right) \leq \left(d_{(r-1)^2}^i\right)^\nu \mathbf{1}\left(d_{(r-1)^2}^i + x > 0\right) \leq \left(d_{r^2}^i\right)^\nu \mathbf{1}\left(-x < d_{r^2}^i\right),$$

where these inequalities hold for a.e.  $(\mu, z)$  due to (5). Hence

$$\begin{aligned} &\int_{\mathbb{R}} \int_{\mathcal{K}} \int_{\mathcal{M}} \left| \psi_x^i(\mu, z) \right| \mathbf{1}\left(x : x \geq t_{-(r-1)^2}\right) P_\lambda(d\mu) Q(dz) dx \\ &\leq c \int_{\mathcal{K}} \int_{\mathcal{M}} \int_{-\infty}^0 \left(d_{r^2}^i(\mu, z)\right)^\nu \mathbf{1}\left((\mu, z) : -x < d_{r^2}^i(\mu, z)\right) dx P_\lambda(d\mu) Q(dz) \\ &= c \mathbb{E}\left[\left(D_{r^2}^i\right)^{\nu+1}\right] \leq c \mathbb{E}\left[(H_0 + H_1 + \dots + H_{r-1})^{\nu+1}\right] \\ &\leq c r^{\nu+2} \mathbb{E}\left[(H_0)^{\nu+1}\right] < \infty, \end{aligned}$$

where the second and third inequalities follow from Lemma 2 and 3, respectively.

For  $l > (r-1)^2$ , we choose  $p \in \mathbb{N}$  to be such that  $p(r-1) < l \leq (p+1)(r-1)$ . Therefore both  $l$  and  $p$  are actually functions of  $\mu, z$  and  $x$ . In particular, for a given realization  $(\mu, z)$  of the arrival process, the values of  $l = l(x)$  and  $p = p(x)$  will possibly change when  $x$  is changed.

We first derive an upper bound on  $(d_l^i + x)$  using the fact that for a.e. realization  $(\mu, z)$ , the sequence  $\{d_n^i\}_{n \geq 0}$  is non-decreasing, thanks to (5). Note that in this case,  $l \leq pr$ . This is true, because we have  $l > (r-1)(r-1)$  and also  $l \leq (p+1)(r-1)$  which implies  $r-1 \leq p$  and therefore  $pr + (r-1) - p \leq pr$ , or, equivalently,  $(p+1)(r-1) \leq pr$ . Making use of these considerations and applying Lemma 2 once more, we obtain

$$\begin{aligned}
 d_l^i + x &\leq d_{pr}^i + t_{-(l-1)} \leq d_{pr}^i + t_{-p(r-1)} \leq h_0 + h_1 + \dots + h_{p-1} + t_{-p(r-1)} \\
 &\leq \sum_{k=0}^{p-1} \left( h_k + (t_{-(k+1)(r-1)} - t_{-k(r-1)}) \right) \\
 &= \sum_{\{k: 0 \leq 2k \leq p-1\}} \left( h_{2k} + (t_{-(2k+1)(r-1)} - t_{-2k(r-1)}) \right) \\
 &\quad + \sum_{\{k: 0 \leq 2k-1 \leq p-1\}} \left( h_{2k-1} + (t_{-2k(r-1)} - t_{-(2k-1)(r-1)}) \right) \\
 &= \xi_p^{(0)} + \xi_p^{(1)},
 \end{aligned}$$

where  $\xi_p^{(j)} = \xi_p^{(j)}(\mu, z)$ ,  $j = 0, 1$ , denote the realizations of the random variables

$$\xi_p^{(j)}(\{T_n\}, \{Z_n\}) = \sum_{\{k: 0 \leq 2k-j \leq p-1\}} \left( H_{2k-j} + (T_{-(2k+1-j)(r-1)} - T_{-(2k-j)(r-1)}) \right), \quad (57)$$

for  $j = 0, 1$  and  $n \in \mathbb{N}$ , where  $\{T_n\}$  is the stationary Poisson process with intensity  $\lambda$ , and  $\{H_n\}$  is defined in (6). So in this second case,

$$\left( d_l^i + x \right)^\nu \mathbf{1}(d_l^i + x > 0) \leq \left( \xi_p^{(0)} + \xi_p^{(1)} \right)^\nu \mathbf{1}(\xi_p^{(0)} + \xi_p^{(1)} > 0). \quad (58)$$

We now derive upper bounds on each term of the last product.

**Lemma 4** *Let  $\eta^{(j)} = \eta^{(j)}(\mu, z)$ ,  $j = 0, 1$  denote the realizations of the random variables*

$$\eta^{(j)}(\{T_n\}, \{Z_n\}) = \sup_{p \geq 1} \left\{ -T_{-(p+1)(r-1)} : \xi_p^{(j)}(\{T_n\}, \{Z_n\}) > 0 \right\}, \quad j = 0, 1. \quad (59)$$

and let

$$\phi^{(j)} = \sup_{p \geq 1} \left\{ \xi_p^{(j)} \right\}, \quad j = 0, 1. \quad (60)$$

Then

$$\begin{aligned}
 \left( d_l^i + x \right)^\nu \mathbf{1}(d_l^i + x > 0) &\leq \left( \xi_p^{(0)} + \xi_p^{(1)} \right)^\nu \mathbf{1}(\xi_p^{(0)} + \xi_p^{(1)} > 0) \\
 &\leq \left( \phi^{(0)} + \phi^{(1)} \right)^\nu \mathbf{1}(-x < \eta^{(0)} + \eta^{(1)}).
 \end{aligned}$$



*Proof* The fact that  $\xi_p^{(0)} + \xi_p^{(1)} \leq \phi^{(0)} + \phi^{(1)}$  is immediate, and we now concentrate on the second term. Because  $\{H_n\}$  is a sequence of 1-dependent random variables, the random variables  $H_0, H_2, \dots$  are i.i.d. and independent of the i.i.d. random variables  $-(T_{-(r-1)} - T_0), -(T_{-3(r-1)} - T_{-2(r-1)}), \dots$ , which are Erlang distributed with expectation  $(r-1) \cdot \lambda^{-1}$ . Under (10), we see that

$$\mathbb{E} \left[ H_{2k} + (T_{-(2k+1)(r-1)} - T_{-2k(r-1)}) \right] = \mathbb{E} [H_0] - \frac{r-1}{\lambda} < 0, \quad (61)$$

so that  $\xi_p^{(0)}(\{T_n\}, \{Z_n\})$  is a random walk with negative drift. Therefore  $\xi_p^{(0)} \rightarrow -\infty$  a.s. for  $p \rightarrow \infty$ . Since the sequences  $H_1, H_3, \dots$  and  $-(T_{-2(r-1)} - T_{-(r-1)}), -(T_{-4(r-1)} - T_{-3(r-1)}), \dots$  have the same properties as the sequences considered above, also  $\xi_p^{(1)}(\{T_n\}, \{Z_n\})$  exhibits the same limiting behavior. Because the arrival process is Poisson, we know that for almost every realisation  $(\mu, z)$ ,  $p = p(x) \rightarrow \infty$  whenever  $x \rightarrow -\infty$  and as a consequence both  $\xi_p^{(0)} \rightarrow -\infty$  and  $\xi_p^{(1)} \rightarrow -\infty$  a.s. for  $x \rightarrow -\infty$ . Therefore, the inequality  $\xi_{p(x)}^{(0)} + \xi_{p(x)}^{(1)} > 0$  cannot hold for  $x$  arbitrarily small. More precisely, since the relation between  $x$  and  $p(x)$  is that  $x \in [t_{-(p(x)+1)(r-1)}, t_{-p(x)(r-1)})$ ,  $\{\xi_{p(x)}^{(0)} + \xi_{p(x)}^{(1)} > 0\}$  implies that

$$x > \inf_{\zeta \leq 0} \left\{ t_{-(p(\zeta)+1)(r-1)} : \xi_{p(\zeta)}^{(0)} + \xi_{p(\zeta)}^{(1)} > 0 \right\}, \quad (62)$$

or, equivalently,

$$-x < \sup_{\zeta \leq 0} \left\{ -t_{-(p(\zeta)+1)(r-1)} : \xi_{p(\zeta)}^{(0)} + \xi_{p(\zeta)}^{(1)} > 0 \right\}. \quad (63)$$

An obvious bound for the righthand side of (63) is given by

$$\begin{aligned} & \sup_{\zeta \leq 0} \left\{ -t_{-(p(\zeta)+1)(r-1)} : \xi_{p(\zeta)}^{(0)} + \xi_{p(\zeta)}^{(1)} > 0 \right\} \\ & \leq \sup_{\zeta \leq 0} \left\{ -t_{-(p(\zeta)+1)(r-1)} : \xi_{p(\zeta)}^{(0)} > 0 \text{ or } \xi_{p(\zeta)}^{(1)} > 0 \right\} \\ & \leq \sup_{\zeta \leq 0} \left\{ -t_{-(p(\zeta)+1)(r-1)} : \xi_{p(\zeta)}^{(0)} > 0 \right\} + \sup_{\zeta \leq 0} \left\{ -t_{-(p(\zeta)+1)(r-1)} : \xi_{p(\zeta)}^{(1)} > 0 \right\} \\ & = \sup_{p \geq 1} \left\{ -t_{-(p+1)(r-1)} : \xi_p^{(0)} > 0 \right\} + \sup_{p \geq 1} \left\{ -t_{-(p+1)(r-1)} : \xi_p^{(1)} > 0 \right\}. \\ & = \eta^{(0)} + \eta^{(1)}. \end{aligned} \quad (64)$$

Using (63) and (64) we see that

$$\left\{ (\mu, z) : \xi_p^{(0)} + \xi_p^{(1)} > 0 \right\} \subseteq \left\{ (\mu, z) : -x < \eta^{(0)} + \eta^{(1)} \right\}, \quad (65)$$

which concludes the proof. □

Therefore, up to this point we have shown that

$$\begin{aligned}
 & \int_{\mathbb{R}} \int_{\mathcal{K}} \int_{\mathcal{M}} \left| \psi_x^i(\mu, z) \right| \mathbf{1}(x : x < t_{-(r-1)^2}) P_\lambda(d\mu) Q(dz) dx \\
 & \leq c \int_{\mathcal{K}} \int_{\mathcal{M}} \int_{-\infty}^0 \left( \phi^{(0)} + \phi^{(1)} \right)^\nu \mathbf{1}(-x < \eta^{(0)} + \eta^{(1)}) dx P_\lambda(d\mu) Q(dz) \\
 & = c \mathbb{E} \left[ \left( \phi^{(0)} + \phi^{(1)} \right)^\nu \left( \eta^{(0)} + \eta^{(1)} \right) \right] \leq c 2^\nu \mathbb{E} \left[ \left( \left( \phi^{(0)} \right)^\nu + \left( \phi^{(1)} \right)^\nu \right) \left( \eta^{(0)} + \eta^{(1)} \right) \right] \\
 & \leq c 2^{\nu+1} \left\{ \mathbb{E} \left[ \left( \phi^{(0)} \right)^{\nu+1} \right] + \mathbb{E} \left[ \left( \phi^{(1)} \right)^{\nu+1} \right] + \mathbb{E} \left[ \left( \eta^{(0)} \right)^{\nu+1} \right] + \mathbb{E} \left[ \left( \eta^{(1)} \right)^{\nu+1} \right] \right\}, \quad (66)
 \end{aligned}$$

where the arguments have been dropped if no misunderstanding is possible. We used Lemma 3 to obtain the last two inequalities and the fact that  $P(\phi^{(j)} \leq y) = P(\varphi^{(j)} \leq y)$ , for all  $y \in \mathbb{R}$ , and  $j = 0, 1$ , to get the final line.

Our last step consists in proving the integrability of  $(\eta^{(j)}(\{T_n\}, \{Z_n\}))^{\nu+1}$ ,  $j = 0, 1$ . Before doing so, let us first introduce some additional notation. For  $i, j = 0, 1$  let

$$\zeta^{(i,j)}(\{T_n\}, \{Z_n\}) = \sup_{p \geq 1} \left\{ - \sum_{\{k: 0 \leq 2k-i \leq p\}} (T_{-(2k-i)(r-1)} - T_{-(2k-i)(r-1)}) : \xi_p^{(j)} > 0 \right\}, \quad (67)$$

where we write  $\xi_p^{(j)}$  instead of  $\xi_p^{(j)}(\{T_n\}, \{Z_n\})$  for sake of simplicity; as usual, let  $\zeta^{(i,j)} = \zeta^{(i,j)}(\mu, z)$ , for  $i, j = 0, 1$ , denote the realizations of the corresponding random variables. For  $\zeta^{(i,j)}(\{T_n\}, \{Z_n\})$  the following result holds:

**Lemma 5** *The  $q$ -th moment of  $\zeta^{(i,j)}(\{T_n\}, \{Z_n\})$ ,  $i, j = 0, 1$  is finite provided that the  $(q+1)$ -th moment of  $H_n$  is finite, i.e.  $\mathbb{E}[(H_n)^{q+1}] < \infty$ .*

A proof for this statement in case  $i = j$  is given in Lemma 6 in [3]. For  $i \neq j$  see Lemma 7 in the same paper.

**Lemma 6** *The  $q$ -th moment of  $\eta^{(j)}(\{T_n\}, \{Z_n\})$ ,  $j = 0, 1$  is finite whenever the  $(q+1)$ -th moment of  $H_n$  is finite, i.e.  $\mathbb{E}[(H_n)^{q+1}] < \infty$ .*

*Proof* For  $j = 0, 1$

$$\eta^{(j)} = \sup_{p \geq 1} \left\{ -t_{-(p+1)(r-1)} : \xi_p^{(j)} > 0 \right\}$$

$$\begin{aligned}
&= \sup_{p \geq 1} \left\{ - \sum_{\{k: 0 \leq 2k \leq p\}} (t_{-(2k+1)(r-1)} - t_{-2k(r-1)}) \right. \\
&\quad \left. - \sum_{\{k: 0 \leq 2k-1 \leq p\}} (t_{-2k(r-1)} - t_{-(2k-1)(r-1)}) : \xi_p^{(j)} > 0 \right\} \\
&\leq \zeta^{(0,j)} + \zeta^{(1,j)}.
\end{aligned} \tag{68}$$

So we conclude that

$$\begin{aligned}
\mathbb{E} \left[ \left( \eta^{(j)}(\{T_n\}, \{Z_n\}) \right)^q \right] &\leq \mathbb{E} \left[ \left( \zeta^{(0,j)}(\{T_n\}, \{Z_n\}) + \zeta^{(1,j)}(\{T_n\}, \{Z_n\}) \right)^q \right] \\
&\leq 2^q \left\{ \mathbb{E} \left[ \left( \zeta^{(0,j)}(\{T_n\}, \{Z_n\}) \right)^q \right] + \mathbb{E} \left[ \left( \zeta^{(1,j)}(\{T_n\}, \{Z_n\}) \right)^q \right] \right\},
\end{aligned} \tag{69}$$

where we used (52) to get the last inequality. The desired statement follows immediately from Lemma 5.  $\square$

Applying Lemma 6 to (66) yields

$$\int_{\mathcal{R}} \int_{\mathcal{K}} \int_{\mathcal{M}} \left| \psi_x^i(\mu, z) \right| P_\lambda(d\mu) Q(dz) dx < \infty,$$

provided that  $\mathbb{E} \left[ (H_n)^{\nu+2} \right] < \infty$ , which completes the proof of the validity of (47) for  $k = 1$ .

#### 4.2.3 Proof of Condition (47): $k \geq 1$

Now we want to look at condition (47) for  $k \geq 1$ . Therefore, let  $x_1 < x_2 < \dots < x_k$  for  $k \in \{1, \dots, m\}$  and  $x_k \in [t_{-l}, t_{-(l-1)})$  for some  $l \in \{1, 2, \dots\}$ . By applying (45) we obtain

$$\begin{aligned}
\psi_{x_1, \dots, x_k}^i(\mu, z) &= \sum_{j=0}^k (-1)^{k-j} \sum_{\pi \in K_{k,j}} \psi(\mu|^{x_k} + \sum_{i \in \pi} \delta_{x_i}, z) \mathbf{1}(x_1 < \dots < x_k) \\
&= \sum_{j=0}^k (-1)^{k-j} \sum_{\pi \in K_{k,j}} f \left( d_0^i \oplus \bigoplus_{n=1}^{l-1} (d_n^i + t_{-n}) \oplus \bigoplus_{n=1}^j (d_{l-1+n}^i + x_{\pi(j+1-n)}) \right),
\end{aligned} \tag{70}$$

where  $\pi_{(n)}$  denotes the  $n$ -th smallest element of the  $j$ -tuple  $\pi$ . The summation can now be organized in the following way (with the three dots being placeholders for the function  $f(\cdot)$  in (70)):

$$\psi_{x_1, \dots, x_k}^i(\mu, z) = \sum_{j=0}^{k-1} (-1)^{k-j} \left\{ \sum_{\pi \in K_{k,j}, \pi \not\ni 1} \dots - \sum_{\pi \in K_{k,j+1}, \pi \ni 1} \dots \right\}$$

$$\begin{aligned}
 = & \sum_{j=0}^{k-1} (-1)^{k-j} \sum_{\pi \in K_{k,j}, \pi \not\geq 1} \left\{ f \left( d_0^i \oplus \bigoplus_{n=1}^{l-1} (d_n^i + t_{-n}) \oplus \bigoplus_{n=1}^j (d_{l-1+n}^i + x_{\pi(j+1-n)}) \right) \right. \\
 & \left. - f \left( d_0^i \oplus \bigoplus_{n=1}^{l-1} (d_n^i + t_{-n}) \oplus \bigoplus_{n=1}^j (d_{l-1+n}^i + x_{\pi(j+1-n)}) \oplus (d_{l+j} + x_1) \right) \right\}. \quad (71)
 \end{aligned}$$

If we assume that  $d_{l-1+k}^i + x_1 \leq 0$ , then we also have  $d_{l+j}^i + x_1 \leq 0$ , for all  $1 \leq j \leq k-1$ , since the  $d_n^i$ 's are non-decreasing in  $n$ . Using this information together with  $d_0^i \geq 0$ , the above representation of the functional  $\psi_{x_1, \dots, x_k}^i$  immediately yields

$$\psi_{x_1, \dots, x_k}^i(\mu, z) = 0 \quad \text{whenever} \quad d_{l-1+k}^i + x_1 \leq 0,$$

and therefore (with the three dots representing the corresponding functions in (71) )

$$\begin{aligned}
 \left| \psi_{x_1, \dots, x_k}^i(\mu, z) \right| & \leq \sum_{j=0}^{k-1} \sum_{\pi \in K_{k,j}, \pi \not\geq k} \left| \dots - \dots \right| \\
 & \leq \sum_{j=0}^{k-1} \binom{k-1}{j} c \left( d_{l-1+k}^i + x_1 \right)^\nu \mathbf{1} \left( d_{l-1+k}^i + x_1 > 0 \right) \\
 & = 2^{k-1} c \left( d_{l-1+k}^i + x_1 \right)^\nu \mathbf{1} \left( d_{l-1+k}^i + x_1 > 0 \right), \quad (72)
 \end{aligned}$$

We first consider the case  $l \leq (r-1)^2 + (r-1)(k-1)$ . Then

$$\begin{aligned}
 \left( d_{l-1+k}^i + x_1 \right)^\nu \mathbf{1} \left( d_{l-1+k}^i + x_1 > 0 \right) & \leq \left( d_{(r-1)^2 + r(k-1)}^i \right)^\nu \mathbf{1} \left( d_{(r-1)^2 + r(k-1)}^i + x_1 > 0 \right) \\
 & \leq \left( d_{kr^2}^i \right)^\nu \mathbf{1} \left( -x_1 < d_{kr^2}^i \right),
 \end{aligned}$$

and, as a consequence, for  $n = (r-1)^2 + (r-1)(k-1)$ ,

$$\begin{aligned}
 & \int_{\mathbb{R}^k} \int_{\mathcal{K}} \int_{\mathcal{M}} \left| \psi_{x_1, \dots, x_k}^i(\mu, z) \right| \mathbf{1} \left( x_k : x_k \geq t_{-n} \right) P_\lambda(d\mu) Q(dz) d(x_1 \dots x_k) \\
 & \leq \int_{\mathcal{K}} \int_{\mathcal{M}} \int_{-\infty}^0 \int_{x_1}^0 \dots \int_{x_{k-1}}^0 2^{k-1} c \left( d_{kr^2}^i \right)^\nu \mathbf{1} \left( -x_1 < d_{kr^2}^i \right) dx_k \dots dx_2 dx_1 P_\lambda(d\mu) Q(dz) \\
 & = c 2^{k-1} \int_{\mathcal{K}} \int_{\mathcal{M}} \left( d_{kr^2}^i \right)^\nu \int_{-d_{kr^2}^i}^0 \int_{x_1}^0 \dots \int_{x_{k-2}}^0 (-x_{k-1}) dx_{k-1} \dots dx_2 dx_1 P_\lambda(d\mu) Q(dz) \\
 & \vdots
 \end{aligned}$$

$$\begin{aligned}
&= c 2^{k-1} \int_{\mathcal{K}} \int_{\mathcal{M}} \left(d_{kr^2}^i\right)^\nu (-1)^k \frac{(-d_{kr^2}^i)^k}{k!} P_\lambda(d\mu) Q(dz) = c 2^{k-1} \frac{1}{k!} \mathbb{E} \left[ \left(D_{kr^2}^i\right)^{\nu+k} \right] \\
&\leq c 2^{k-1} \frac{1}{k!} \mathbb{E} \left[ (H_0 + H_1 + \dots + H_{kr-1})^{\nu+k} \right] \leq c 2^{k-1} \frac{(kr)^{\nu+k+1}}{k!} \mathbb{E} \left[ (H_0)^{\nu+k} \right] \\
&< \infty,
\end{aligned}$$

provided that  $\mathbb{E} \left[ (H_0)^{\nu+k} \right] < \infty$ , where we used Lemma 2 and Lemma 3 b) for the last inequalities.

Next we will look at the case  $l > (r-1)^2 + (r-1)(k-1)$ . As we did for  $k=1$ , we choose  $p \in \mathbb{N}$  to be such that  $p(r-1) < l \leq (p+1)(r-1)$ . From  $l > (r-1)^2 + (r-1)(k-1)$  and  $l \leq (p+1)(r-1)$  we conclude that  $(r-1) + (k-1) \leq p$  and thus  $pr + (r-1) - p \leq pr - (k-1)$  or, equivalently,  $(p+1)(r-1) \leq pr + 1 - k$ . So we note that under the given assumptions  $l-1+k \leq pr$ . By applying this information we obtain the following inequalities for a.e.  $(\mu, z)$ :

$$\begin{aligned}
d_{l-1+k}^i + x_1 &\leq d_{pr}^i + x_k - (x_k - x_1) \leq d_{pr}^i + t_{-(l-1)} - (x_k - x_1) \\
&\leq d_{pr}^i + t_{-p(r-1)} - (x_k - x_1) \leq \xi_p^{(0)} + \xi_p^{(1)} - (x_k - x_1) \\
&\leq \phi^{(0)} + \phi^{(1)},
\end{aligned}$$

where we use the same notation as in the  $k=1$  part above. Only this time we find  $x_k \in [t_{-(p+1)(r-1)}, t_{-p(r-1)})$  and by the same arguments as for  $k=1$  we conclude that

$$\begin{aligned}
&\left\{ (\mu, z) : \xi_p^{(0)} + \xi_p^{(1)} > x_k - x_1 \right\} \\
&\subseteq \left\{ (\mu, z) : -x_k < \sup_{p \geq 1} \left\{ -t_{-(p+1)(r-1)} : \xi_p^{(0)} + \xi_p^{(1)} > x_k - x_1 \right\} \right\} \\
&= \left\{ (\mu, z) : -x_1 < \sup_{p \geq 1} \left\{ -t_{-(p+1)(r-1)} + (x_k - x_1) : \xi_p^{(0)} + \xi_p^{(1)} > x_k - x_1 \right\} \right\} \\
&\subseteq \left\{ (\mu, z) : -x_1 < \sup_{p \geq 1} \left\{ -t_{-(p+1)(r-1)} : \xi_p^{(0)} + \xi_p^{(1)} > 0 \right\} \right\} \\
&\subseteq \left\{ (\mu, z) : -x_1 < \eta^{(0)} + \eta^{(1)} \right\}.
\end{aligned}$$

So we have learned that, for  $n = (r-1)^2 + (r-1)(k-1)$ ,

$$\begin{aligned}
&\int_{\mathbb{R}^k} \int_{\mathcal{K}} \int_{\mathcal{M}} \left| \psi_{x_1, \dots, x_k}^i(\mu, z) \right| \mathbf{1}(x_k : x_k < t_{-n}) P_\lambda(d\mu) Q(dz) d(x_1 \dots x_k) \\
&\leq c 2^{k-1} \int_{\mathcal{K}} \int_{\mathcal{M}} \int_{-\infty}^0 \int_{x_1}^0 \dots \int_{x_{k-1}}^0 \left( \phi^{(0)} + \phi^{(1)} \right)^\nu \mathbf{1}(-x_1 < \eta^{(0)} + \eta^{(1)}) \\
&\hspace{25em} dx_k \dots dx_2 dx_1 P_\lambda(d\mu) Q(dz)
\end{aligned}$$



can be obtained by thinning from a stationary Poisson process with a larger intensity. Thus, (73) can be bounded by fixing  $\lambda$  at any permissible value  $\lambda'$ , i.e.  $0 < \lambda' < a$ , where  $a$  denotes the maximal  $(\max, +)$ -Lyapunov exponent of  $\{A_n\}$ . The finiteness of (73), provided that  $\mathbb{E}[(H_n)^{\nu+m+2}] < \infty$ , then follows from Lemma 6. Therefore, (48) also holds under the assumptions made.

## 5 Differentiability, Admissibility

In general, the conditions of Theorem 2 are not sufficient to ensure any differentiability property. However, it was shown in [3] that under Cramer type assumptions on the random variables  $D_n^i$ , the mean values of the random variables  $W^i$  are such that (23) holds and are infinitely differentiable in  $\lambda$  in a right neighborhood of 0.

We show below that this infinite differentiability result can also be extended to the more general functions of  $W$  considered here. In this sense, at least in this particular Cramer case, the expansion given in Theorem 1 is a Taylor type expansion.<sup>1</sup>

The proof of this property is based on the notion of *admissibility* defined in [9], which is sufficient to grant (23) and (24).

In order to prove the admissibility of  $\psi$ , where  $\psi$  is the function defined in (53), we have to show that there exist constants  $K, N < \infty$  and  $1 < a < \infty, \theta > 0$ , such that for all  $s < t < 0$ ,

$$\int \int |\psi(\mu|^t, z) - \psi(\mu|^s, z)| P_\lambda(d\mu | \mathcal{C}(l, j))Q(dz) \leq K(j+l)^N a^{j+l} e^{-\theta t}, \quad (74)$$

where  $\mathcal{C}(l, j) = \{\mu' : \mu'([s, t]) = l, \mu'([t, 0]) = j\}$ . Let  $\mathcal{B}^i(l, j)$  be the event

$$\mathcal{B}^i(l, j) = \left\{ \bigcap_{n=j+1}^{j+l} \{d_n^i - t_{-n} < 0\} \right\}. \quad (75)$$

On  $\mathcal{B}^i(l, j) \cap (\mathcal{C}(l, j) \times \mathcal{K}^\infty)$ , we have  $\psi(\mu|^t, z) = \psi(\mu|^s, z)$ , so that

$$\begin{aligned} & \int \int |\psi(\mu|^t, z) - \psi(\mu|^s, z)| P_\lambda(d\mu | \mathcal{C}(l, j))Q(dz) \\ &= \int \int |\psi(\mu|^t, z) - \psi(\mu|^s, z)| \mathbf{1}_{\overline{\mathcal{B}^i(l, j)}} P_\lambda(d\mu | \mathcal{C}(l, j))Q(dz) \leq h^{\frac{1}{2}} g^{\frac{1}{2}}, \end{aligned}$$

where

$$h = \int \int (\psi(\mu|^t, z) - \psi(\mu|^s, z))^2 P_\lambda(d\mu | \mathcal{C}(l, j))Q(dz) \quad (76)$$

<sup>1</sup>We actually conjecture that this is always the case.

and

$$g = \mathbb{P} \left( \overline{\mathcal{B}^i(l, j)} \mid \mathcal{C}(l, j) \times \mathcal{K}^\infty \right). \quad (77)$$

Using now the special form of  $\psi$  and the independence assumptions, we obtain that

$$h \leq 2c \mathbb{E} \left[ (D_{l+j}^i)^{2\nu} \mid \mathcal{C}(l, j) \times \mathcal{K}^\infty \right] = 2c \mathbb{E} \left[ (D_{l+j}^i)^{2\nu} \right]. \quad (78)$$

Let  $\chi_n = \max_{i,j} \left\{ (A_{-n})_{i,j} + (B_{-n})_j \oplus 0 \right\}$ . We have, for all  $n \geq 1$ ,

$$(D_n^i)^{2\nu} \leq \left( \sum_{p=1}^n \chi_p \right)^{2\nu} \leq n^{2\nu} \sum_{p=1}^n (\chi_p)^{2\nu}$$

and so  $h \leq n^{2\nu+1} \kappa$ , with  $\kappa = 2c \mathbb{E}(\chi_1)^{2\nu} = 2c \mathbb{E}[(F_1)^{2\nu}] < \infty$ , where  $F_n$  is defined in (7).

As for  $h$ , we have

$$\begin{aligned} g &= \mathbb{P} \left( \bigcup_{n=j+1}^{j+l} \{D_n^i - T_{-n} > 0\} \mid \mathcal{C}(l, j) \times \mathcal{K}^\infty \right) \\ &\leq \mathbb{P} (D_{j+l}^i > t \mid \mathcal{C}(l, j) \times \mathcal{K}^\infty) = \mathbb{P} (D_{j+l}^i > t) \\ &= \mathbb{P} (e^{uD_{j+l}^i} > e^{ut}) \leq \mathbb{E} [e^{uD_{j+l}^i}] e^{-ut} \\ &\leq L_u \phi(u)^{j+l} e^{-ut}, \end{aligned}$$

where  $u$  is any real number in the interval  $(0, \theta^*)$ , and where we used Chebyshev's inequality and Assumption (22) in order to derive the last two inequalities. Finally admissibility is proved with  $\theta = \frac{u}{2}$ ,  $a = (\phi(u))^{\frac{1}{2}}$ ,  $N = \nu + 1$  and  $K = \sqrt{\kappa L_u}$ .

## 6 Calculation of the Coefficients

Now that we have seen that Theorem 2 can be applied to derive the expansion of  $\mathbb{E} [f(W^i)]$ , we will calculate the coefficients of  $\lambda^k$ . We will first give a recursive representation of these coefficients and will later on use this result to reveal their explicit form.

### 6.1 Recursion Formula

From Theorem 2 or from admissibility, we know that the coefficients of  $\lambda^k$  in the expansions are given by

$$\mathbb{E} \left[ \psi^i(o, \{Z_n\}) \right], \quad \int_{\mathbb{R}^k} \mathbb{E} \left[ \psi_{x_1, \dots, x_k}^i(o, \{Z_n\}) \right] d(x_1, \dots, x_k), \quad (79)$$



for  $k = 0$ ,  $k \geq 1$ , respectively, where  $\psi$  is the functional defined in (53). Introducing the notation  $q_1(d_0^i) = \psi^i(o, z)$  and

$$q_{k+1}(d_0^i, d_1^i, \dots, d_k^i) = \int_{\mathbb{R}^k} \psi_{x_1, \dots, x_k}^i(o, z) d(x_1, \dots, x_k), \quad (80)$$

for  $k \geq 1$ , we obtain the following result.

**Theorem 3** *Under the monotonicity assumption (5) it holds for each  $i \in \{1, \dots, \alpha'\}$  that  $q_1(d_0^i) = f(d_0^i)$  and*

$$q_{k+1}(d_0^i, d_1^i, \dots, d_k^i) = \sum_{p=0}^{k-1} \int_{d_p^i - d_0^i}^{d_{p+1}^i - d_0^i} \left\{ q_k(\underbrace{d_0^i, \dots, d_0^i}_p, d_{p+1}^i - u, \dots, d_k^i - u) \right. \\ \left. - q_k(\underbrace{d_0^i, \dots, d_0^i}_{p+1}, d_{p+1}^i - u, \dots, d_{k-1}^i - u) \right\} du, \quad (81)$$

for all  $k \geq 1$ .

*Proof* From (53) it is immediate to see that

$$q_1(d_0^i) = \psi^i(o, \{Z_n\}) = f(d_0^i).$$

Using (45), (53) and (80) for  $k \geq 1$  we obtain

$$q_{k+1}(d_0^i, d_1^i, \dots, d_k^i) \\ = \int_{-\infty}^0 \int_{x_1}^0 \dots \int_{x_{k-1}}^0 \sum_{j=0}^k (-1)^{k-j} \sum_{\pi \in K_{k,j}} f\left(d_0^i \oplus \bigoplus_{n=1}^j (d_n^i + x_{\pi(j+1-n)})\right) dx_k \dots dx_2 dx_1, \\ = \int_0^\infty \int_{s_1}^\infty \dots \int_{s_{k-1}}^\infty \sum_{j=0}^k (-1)^{k-j} \sum_{\pi \in K_{k,j}} f\left(d_0^i \oplus \bigoplus_{n=1}^j (d_n^i - s_{\pi(n)})\right) ds_k \dots ds_2 ds_1, \quad (82)$$

where  $\pi_{(n)}$  denotes the  $n$ -th order statistic of  $\pi$ , i.e., the  $n$ -th smallest element of the  $j$ -tuple  $\pi \in K_{k,j}$ . The second equality is due to the substitution  $x_j \rightarrow -x_j$ ,  $j \in \{1, \dots, k\}$ , an interchange of the order of integration and finally by another substitution  $x_j \rightarrow s_{k+1-j}$ , for all  $1 \leq j \leq k$ .

As a next step, we decompose the outer integral in the following way:

$$\int_0^\infty \dots = \sum_{p=0}^{k-1} \int_{d_p^i - d_0^i}^{d_{p+1}^i - d_0^i} \dots + \int_{d_k^i - d_0^i}^\infty \dots = \sum_{p=0}^{k-1} \int_{d_p^i - d_0^i}^{d_{p+1}^i - d_0^i} \dots \quad (83)$$

because from  $s_1 \leq \dots \leq s_k$ ,  $d_0^i \leq \dots \leq d_k^i$  and  $s_1 \geq d_k^i - d_0^i$  it follows that  $s_m \geq s_1 \geq d_k^i - d_0^i \geq d_n^i - d_0^i$  and therefore  $d_n^i - s_m \leq d_0^i$  for all  $1 \leq m, n \leq k$ . But this means that the integrand of the integral ranging from  $d_k^i - d_0^i$  to  $\infty$  is equal to

$$\sum_{j=0}^k (-1)^{k-j} \sum_{\pi \in K_{k,j}} f(d_0^i) = f(d_0^i) \sum_{j=0}^k \binom{k}{j} (-1)^{k-j} = 0.$$

For each of the remaining summands in (83), we decompose the summation according to

$$\sum_{j=0}^k (-1)^{k-j} \sum_{\pi \in K_{k,j}} \dots = \sum_{j=1}^k (-1)^{k-j} \sum_{\pi \in K_{k,j}, \pi \ni 1} \dots + \sum_{j=0}^{k-1} (-1)^{k-j} \sum_{\pi \in K_{k,j}, \pi \not\ni 1} \dots, \quad (84)$$

where the dots are placeholders for the function  $f(\cdot)$  given in (82).

We first consider that part of the integration involving only the first of these two sums, i.e., for  $0 \leq p \leq k-1$ ,

$$\begin{aligned} & \int_{d_p^i - d_0^i}^{d_{p+1}^i - d_0^i} \int_{s_1}^{\infty} \dots \int_{s_{k-1}}^{\infty} \sum_{j=1}^k (-1)^{k-j} \sum_{\pi \in K_{k,j}, \pi \ni 1} f\left(d_0^i \oplus \bigoplus_{n=1}^j (d_n^i - s_{\pi(n)})\right) ds_k \dots ds_2 ds_1 \\ &= \int_{d_p^i - d_0^i}^{d_{p+1}^i - d_0^i} \int_{s_1}^{\infty} \dots \int_{s_{k-1}}^{\infty} \sum_{j=1}^k (-1)^{k-j} \sum_{\pi \in K_{k,j}, \pi \ni 1} f\left(d_0^i \oplus \bigoplus_{n=\min\{j,p\}+1}^j (d_n^i - s_{\pi(n)})\right) \\ & \hspace{15em} ds_k \dots ds_2 ds_1, \quad (85) \end{aligned}$$

because from  $s_1 \geq d_p^i - d_0^i$ , it follows that  $d_n^i - s_{\pi(n)} \leq d_n^i - s_1 \leq d_p^i - s_1 \leq d_0^i$ , for all  $1 \leq n \leq p$ , and therefore, in particular, for all  $1 \leq n \leq \min\{j, p\}$ .

We will first consider the case  $p = 0$ . Because  $\pi \ni 1$ , we see that  $s_{\pi(1)} = s_1$ , and since in this case  $d_1^i - s_1 \geq d_0^i$  we get that (85) is equal to

$$\begin{aligned} & \int_0^{d_1^i - d_0^i} \int_{s_1}^{\infty} \dots \int_{s_{k-1}}^{\infty} \sum_{j=1}^k (-1)^{k-j} \sum_{\pi \in K_{k,j}, \pi \ni 1} f\left((d_1^i - s_1) \oplus \bigoplus_{n=2}^j (d_n^i - s_{\pi(n)})\right) ds_k \dots ds_2 ds_1 \\ &= \int_0^{d_1^i - d_0^i} \int_0^{\infty} \int_{s_2}^{\infty} \dots \int_{s_{k-1}}^{\infty} \sum_{j=0}^{k-1} (-1)^{k-1-j} \sum_{\pi \in K_{k,j+1}, \pi \ni 1} f\left((d_1^i - u) \oplus \bigoplus_{n=1}^j (d_{n+1}^i - u - s_{\pi(n+1)})\right) \\ & \hspace{15em} ds_k \dots ds_3 ds_2 du \end{aligned}$$

$$\begin{aligned}
&= \int_0^{d_1^i - d_0^i} \int_0^\infty \int_{s_1}^\infty \dots \int_{s_{k-2}}^\infty \sum_{j=0}^{k-1} (-1)^{k-1-j} \sum_{\pi \in K_{k-1,j}} f\left((d_1^i - u) \oplus \bigoplus_{n=1}^j ((d_{n+1}^i - u) - s_{\pi(n)})\right) \\
&\hspace{20em} ds_{k-1} \dots ds_2 ds_1 du \\
&= \int_0^{d_1^i - d_0^i} q_k(d_1^i - u, d_2^i - u, \dots, d_k^i - u) du, \tag{86}
\end{aligned}$$

where the first of the above equalities follows from the substitutions  $s_1 \rightarrow u$ ,  $s_n \rightarrow s_n - u$  for all  $2 \leq n \leq k$ ,  $j \rightarrow j + 1$  and  $n \rightarrow n + 1$ . The last but one equality is a consequence of the substitution  $s_n \rightarrow s_{n-1}$  for all  $n = 2, \dots, k$ .

Now we come back to the remaining cases  $1 \leq p \leq k - 1$ . Applying the transformations we just used for  $p = 0$  yields

$$\begin{aligned}
&\int_{d_p^i - d_0^i}^{d_{p+1}^i - d_0^i} \int_{s_1}^\infty \dots \int_{s_{k-1}}^\infty \sum_{j=1}^k (-1)^{k-j} \sum_{\pi \in K_{k,j}, \pi \ni 1} f\left(d_0^i \oplus \bigoplus_{n=\min\{j,p\}+1}^j (d_n^i - s_{\pi(n)})\right) ds_k \dots ds_2 ds_1 \\
&= \int_{d_p^i - d_0^i}^{d_{p+1}^i - d_0^i} \int_0^\infty \int_{s_1}^\infty \dots \int_{s_{k-2}}^\infty \sum_{j=0}^{k-1} (-1)^{k-1-j} \sum_{\pi \in K_{k-1,j}} \\
&\hspace{15em} f\left(d_0^i \oplus \bigoplus_{n=1}^{\min\{j,p-1\}} (d_0^i - s_{\pi(n)}) \oplus \bigoplus_{n=\min\{j,p-1\}+1}^j (d_{n+1}^i - u - s_{\pi(n)})\right) \\
&\hspace{20em} ds_{k-1} \dots ds_2 ds_1 du \\
&= \int_{d_p^i - d_0^i}^{d_{p+1}^i - d_0^i} q_k(\underbrace{d_0^i, \dots, d_0^i}_p, d_{p+1}^i - u, \dots, d_k^i - u) du, \tag{87}
\end{aligned}$$

for  $p = 1, \dots, k - 1$ . Aggregating (86) and (87), we obtain for (85)

$$\begin{aligned}
&\int_{d_p^i - d_0^i}^{d_{p+1}^i - d_0^i} \int_{s_1}^\infty \dots \int_{s_{k-1}}^\infty \sum_{j=1}^k (-1)^{k-j} \sum_{\pi \in K_{k,j}, \pi \ni 1} f\left(d_0^i \oplus \bigoplus_{n=1}^j (d_n^i - s_{\pi(n)})\right) ds_k \dots ds_2 ds_1 \\
&= \int_{d_p^i - d_0^i}^{d_{p+1}^i - d_0^i} q_k(\underbrace{d_0^i, \dots, d_0^i}_p, d_{p+1}^i - u, \dots, d_k^i - u) du, \tag{88}
\end{aligned}$$

for all  $p = 0, \dots, k - 1$ .

Using analogous reasoning for the remaining second part of the integration, we get

$$\begin{aligned}
 & \int_{d_p^i - d_0^i}^{d_{p+1}^i - d_0^i} \int_{s_1}^{\infty} \dots \int_{s_{k-1}}^{\infty} \sum_{j=0}^{k-1} (-1)^{k-j} \sum_{\pi \in K_{k,j}, \pi \not\equiv 1} f\left(d_0^i \oplus \bigoplus_{n=1}^j (d_n^i - s_{\pi(n)})\right) ds_k \dots ds_2 ds_1 \\
 &= - \int_{d_p^i - d_0^i}^{d_{p+1}^i - d_0^i} q_k(\underbrace{d_0^i, \dots, d_0^i}_{p+1}, d_{p+1}^i - u, \dots, d_{k-1}^i - u) du,
 \end{aligned} \tag{89}$$

for all  $0 \leq p \leq k-1$ , and thus

$$\begin{aligned}
 q_{k+1}(d_0^i, d_1^i, \dots, d_k^i) &= \sum_{p=0}^{k-1} \int_{d_p^i - d_0^i}^{d_{p+1}^i - d_0^i} \left\{ q_k(\underbrace{d_0^i, \dots, d_0^i}_p, d_{p+1}^i - u, \dots, d_k^i - u) \right. \\
 &\quad \left. - q_k(\underbrace{d_0^i, \dots, d_0^i}_{p+1}, d_{p+1}^i - u, \dots, d_{k-1}^i - u) \right\} du,
 \end{aligned}$$

which is the statement of Theorem 3. □

## 6.2 Explicit Solution

In this section we want to derive the explicit representation (20) of the functions  $q_k(x_0, \dots, x_{k-1})$ ,  $k \geq 1$ , that show up in the calculation of the coefficients  $\mathbb{IE}[q_{k+1}(D_0^i, \dots, D_k^i)]$  of  $\lambda^k$  in the series expansion of  $\mathbb{IE}[f(W^i)]$  considered in Theorem 1.

**Theorem 4** For  $k \geq 1$  and  $0 \leq x_0 \leq x_1 \leq \dots \leq x_k$  a solution to the integral recursion equation (81) stated in Theorem 3 is given by the functions  $q_k$  outlined in Theorem 1, particularly in (20).

*Proof* We show (20) by induction with respect to  $k \geq 1$ . Using (81) for  $k = 1$  yields

$$\begin{aligned}
 q_2(x_0, x_1) &= \int_0^{x_1 - x_0} \{q_1(x_1 - u) - q_1(x_0)\} du = \int_0^{x_1 - x_0} \{f(x_1 - u) - f(x_0)\} du \\
 &= \left[ -F(x_1 - u) - f(x_0)u \right]_0^{x_1 - x_0} \\
 &= (F(x_1) - F(x_0)) - f(x_0)[p_1(x_1) - p_1(x_0)],
 \end{aligned}$$

where we wrote  $F(x)$  for  $F^{[1]}(x)$  and thus, (20) holds for  $k = 1$ .

In the following  $k$  will always be assumed to be greater than one, i.e.  $k \geq 2$ . Equation (81) can be rewritten in the form:

$$q_{k+1}(x_0, x_1, \dots, x_k) = \sum_{p=0}^{k-1} \int_{x_p - x_0}^{x_{p+1} - x_0} q_k(\underbrace{x_0, \dots, x_0}_p, x_{p+1} - u, \dots, x_k - u) du \quad (90)$$

$$- \sum_{p=0}^{k-1} \int_{x_p - x_0}^{x_{p+1} - x_0} q_k(\underbrace{x_0, \dots, x_0}_{p+1}, x_{p+1} - u, \dots, x_{k-1} - u) du. \quad (91)$$

In what follows, (90) and (91) will be evaluated separately.

Now let (20) be true for some natural  $k \geq 2$ , i.e.

$$\begin{aligned} & q_k(x_1, x_2, \dots, x_k) \\ &= \sum_{n=0}^{k-1} \binom{k-1}{n} (-1)^{k-1-n} F^{[k-1]}(x_{n+1}) - \sum_{n=0}^{k-2} \sum_{j=n}^{k-2} \binom{j}{n} (-1)^{j-n} F^{[j]}(x_{n+1}) \\ & \quad \left\{ p_{k-1-j}(x_{n+2}, \dots, x_{k-j+n}) - p_{k-1-j}(x_{n+1}, \dots, x_{k-j+n-1}) \right\}, \quad (92) \end{aligned}$$

for some  $k \geq 2$ . Since the polynomials  $p_n(x_0, \dots, x_{n-1})$  are translation-invariant with respect to a constant translation of all arguments, for  $n \geq 2$ , and since  $p_1(x) = x$  implies  $p_1(x_1 - u) - p_1(x_0 - u) = p_1(x_1) - p_1(x_0)$ , it follows that for  $p = 0$

$$\begin{aligned} & \int_0^{x_1 - x_0} q_k(x_1 - u, \dots, x_k - u) du \\ &= \sum_{n=0}^{k-1} \binom{k-1}{n} (-1)^{k-1-n} \int_0^{x_1 - x_0} F^{[k-1]}(x_{n+1} - u) du \\ & \quad - \sum_{n=0}^{k-2} \sum_{j=n}^{k-2} \binom{j}{n} (-1)^{j-n} \left\{ p_{k-1-j}(x_{n+2}, \dots, x_{k-j+n}) \right. \\ & \quad \left. - p_{k-1-j}(x_{n+1}, \dots, x_{k-j+n-1}) \right\} \int_0^{x_1 - x_0} F^{[j]}(x_{n+1} - u) du \\ &= \sum_{n=0}^{k-1} \binom{k-1}{n} (-1)^{k-1-n} \left( F^{[k]}(x_{n+1}) - F^{[k]}(x_{n+1} + x_0 - x_1) \right) \\ & \quad - \sum_{n=0}^{k-2} \sum_{j=n}^{k-2} \binom{j}{n} (-1)^{j-n} \left\{ p_{k-1-j}(x_{n+2}, \dots, x_{k-j+n}) - p_{k-1-j}(x_{n+1}, \dots, x_{k-j+n-1}) \right\} \quad (93) \end{aligned}$$

$$\left( F^{[j+1]}(x_{n+1}) - F^{[j+1]}(x_{n+1} + x_0 - x_1) \right). \quad (94)$$

For  $1 \leq p \leq k-1$  the integrand is given by:

$$\begin{aligned} & q_k(\underbrace{x_0, \dots, x_0}_p, x_{p+1} - u, \dots, x_k - u) \\ = & \sum_{n=0}^{p-1} \binom{k-1}{n} (-1)^{k-1-n} F^{[k-1]}(x_0) + \sum_{n=p}^{k-1} \binom{k-1}{n} (-1)^{k-1-n} F^{[k-1]}(x_{n+1} - u) \\ & - \sum_{n=0}^{p-1} \sum_{j=n}^{k-2} \binom{j}{n} (-1)^{j-n} F^{[j]}(x_0) \left\{ p_{k-1-j}(\underbrace{x_0, \dots, x_0}_{p-n-1}, x_{p+1} - u, \dots, x_{k-j+n} - u) \right. \\ & \quad \left. - p_{k-1-j}(\underbrace{x_0, \dots, x_0}_{p-n}, x_{p+1} - u, \dots, x_{k-j+n-1} - u) \right\} \\ & - \sum_{n=p}^{k-2} \sum_{j=n}^{k-2} \binom{j}{n} (-1)^{j-n} F^{[j]}(x_{n+1} - u) \left\{ p_{k-1-j}(x_{n+2} - u, \dots, x_{k-j+n} - u) \right. \\ & \quad \left. - p_{k-1-j}(x_{n+1} - u, \dots, x_{k-j+n-1} - u) \right\}, \end{aligned}$$

and therefore, if we use the translation-invariance property (18) of the polynomials  $p_n$  to shift all arguments by  $u$  and then substitute  $x_0 + u \rightarrow v$ , we see that

$$\begin{aligned} & \int_{x_p - x_0}^{x_{p+1} - x_0} q_k(\underbrace{x_0, \dots, x_0}_p, x_{p+1} - u, \dots, x_k - u) du \\ = & \sum_{n=0}^{p-1} \binom{k-1}{n} (-1)^{k-1-n} F^{[k-1]}(x_0) \left\{ p_1(x_{p+1}) - p_1(x_p) \right\} \quad (95) \end{aligned}$$

$$+ \sum_{n=p}^{k-1} \binom{k-1}{n} (-1)^{k-1-n} \left\{ F^{[k]}(x_{n+1} + x_0 - x_p) - F^{[k]}(x_{n+1} + x_0 - x_{p+1}) \right\} \quad (96)$$

$$\begin{aligned} & - \sum_{n=0}^{p-1} \sum_{j=n}^{k-2} \binom{j}{n} (-1)^{j-n} F^{[j]}(x_0) \int_{x_p}^{x_{p+1}} \left\{ p_{k-1-j}(\underbrace{v, \dots, v}_{p-n-1}, x_{p+1}, \dots, x_{k-j+n}) \right. \\ & \quad \left. - p_{k-1-j}(\underbrace{v, \dots, v}_{p-n}, x_{p+1}, \dots, x_{k-j+n-1}) \right\} dv \quad (97) \end{aligned}$$

$$- \sum_{n=p}^{k-2} \sum_{j=n}^{k-2} \binom{j}{n} (-1)^{j-n} \left\{ F^{[j+1]}(x_{n+1} + x_0 - x_p) - F^{[j+1]}(x_{n+1} + x_0 - x_{p+1}) \right\}$$

$$\left\{ p_{k-1-j}(x_{n+2}, \dots, x_{k-j+n}) - p_{k-1-j}(x_{n+1}, \dots, x_{k-j+n-1}) \right\} \quad (98)$$

$$= I_p + II_p + III_p + IV_p,$$

for  $1 \leq p \leq k-1$ , where  $I_p$ ,  $II_p$ ,  $III_p$  and  $IV_p$  are shorthand notations for the expressions given by (95), (96), (97) and (98), respectively. So we see that (90) can be represented as the sum of the terms given by (93) and (94) as well as the expressions just introduced for  $1 \leq p \leq k-1$  above. We will now continue by summing up these terms in the following way:

$$\begin{aligned} \sum_{p=1}^{k-1} I_p &= F^{[k-1]}(x_0) \sum_{n=0}^{k-2} \binom{k-1}{n} (-1)^{k-1-n} \sum_{p=n}^{k-2} \left\{ p_1(x_{p+2}) - p_1(x_{p+1}) \right\} \\ &= -F^{[k-1]}(x_0) \sum_{n=0}^{k-1} \binom{k-1}{n} (-1)^{k-1-n} p_1(x_{n+1}), \end{aligned} \quad (99)$$

where we substituted  $p \rightarrow p+1$  and then interchanged the order of summation to get the first equality. The last line was obtained after calculating the telescope-sum, splitting up the result into two summations and using the fact that

$$\sum_{n=0}^{k-2} \binom{k-1}{n} (-1)^{k-1-n} = -1.$$

If we compare (93) to (96) we notice that (93) can just as well be interpreted as a  $II_0$ . Thus, a substitution  $n \rightarrow n-1$  and similar calculations as before yield

$$\begin{aligned} \sum_{p=0}^{k-1} II_p &= \sum_{n=1}^k \binom{k-1}{n-1} (-1)^{k-n} \sum_{p=0}^{n-1} \left\{ F^{[k]}(x_n + x_0 - x_p) - F^{[k]}(x_n + x_0 - x_{p+1}) \right\} \\ &= \sum_{n=1}^k \binom{k-1}{n-1} (-1)^{k-n} F^{[k]}(x_n), \end{aligned} \quad (100)$$

as well as

$$\begin{aligned} \sum_{p=1}^{k-1} III_p &= - \sum_{n=0}^{k-2} \sum_{j=n}^{k-2} \binom{j}{n} (-1)^{j-n} F^{[j]}(x_0) \sum_{p=n}^{k-2} \int_{x_{p+1}}^{x_{p+2}} \left\{ p_{k-1-j}(\underbrace{v, \dots, v}_{p-n}, x_{p+2}, \dots, x_{k-j+n}) \right. \\ &\quad \left. - p_{k-1-j}(\underbrace{v, \dots, v}_{p-n+1}, x_{p+2}, \dots, x_{k-j+n-1}) \right\} dv \\ &= - \sum_{n=0}^{k-2} \sum_{j=n}^{k-2} \binom{j}{n} (-1)^{j-n} F^{[j]}(x_0) p_{k-j}(x_{n+1}, \dots, x_{k-j+n}). \end{aligned} \quad (101)$$

The second equality holds because the polynomials  $p_{k-1-j}$  cancel out whenever  $p - n \geq k - 1 - j$  or, equivalently, if  $p \geq k - 1 - j + n$ , and therefore

$$\begin{aligned}
 & \sum_{p=n}^{k-2} \int_{x_{p+1}}^{x_{p+2}} \left\{ p_{k-1-j}(\underbrace{v, \dots, v}_{p-n}, x_{p+2}, \dots, x_{k-j+n}) - p_{k-1-j}(\underbrace{v, \dots, v}_{p-n+1}, x_{p+2}, \dots, x_{k-j+n-1}) \right\} dv \\
 &= \sum_{p=n}^{k-2-j+n} \int_{x_{p+1}}^{x_{p+2}} \left\{ p_{k-1-j}(\underbrace{v, \dots, v}_{p-n}, x_{p+2}, \dots, x_{k-j+n}) \right. \\
 & \quad \left. - p_{k-1-j}(\underbrace{v, \dots, v}_{p-n+1}, x_{p+2}, \dots, x_{k-j+n-1}) \right\} dv \\
 &= \sum_{p=0}^{k-2-j} \int_{x_{p+n+1}}^{x_{p+n+2}} \left\{ p_{k-1-j}(\underbrace{v, \dots, v}_p, x_{p+n+2}, \dots, x_{k-j+n}) \right. \\
 & \quad \left. - p_{k-1-j}(\underbrace{v, \dots, v}_{p+1}, x_{p+n+2}, \dots, x_{k-j+n-1}) \right\} dv \\
 &= p_{k-j}(x_{n+1}, \dots, x_{k-j+n}),
 \end{aligned}$$

where the last equality follows from (19).

It can be seen that the  $IV_p$ 's show up only for  $1 \leq p \leq k - 2$ . Similarly as we did for the  $II_p$ 's above, we realize that (94) can be interpreted as a  $IV_0$ . So we include this term in the summation of the  $IV_p$ 's and obtain

$$\begin{aligned}
 \sum_{p=0}^{k-1} IV_p &= - \sum_{n=0}^{k-2} \sum_{j=n}^{k-2} \binom{j}{n} (-1)^{j-n} \sum_{p=0}^n \left\{ F^{[j+1]}(x_{n+1} + x_0 - x_p) - F^{[j+1]}(x_{n+1} + x_0 - x_{p+1}) \right\} \\
 & \quad \left\{ p_{k-1-j}(x_{n+2}, \dots, x_{k-j+n}) - p_{k-1-j}(x_{n+1}, \dots, x_{k-j+n-1}) \right\} \\
 &= - \sum_{n=1}^{k-1} \sum_{j=n}^{k-1} \binom{j-1}{n-1} (-1)^{j-n} \left\{ F^{[j]}(x_n) - F^{[j]}(x_0) \right\} \\
 & \quad \left\{ p_{k-j}(x_{n+1}, \dots, x_{k-j+n}) - p_{k-j}(x_n, \dots, x_{k-j+n-1}) \right\}. \quad (102)
 \end{aligned}$$

We can approach term (91) in an almost identical way. For  $p = k - 1$  we obtain

$$q_k(\underbrace{x_0, \dots, x_0}_k) = F^{[k-1]}(x_0) \sum_{n=0}^{k-1} \binom{k-1}{n} (-1)^{k-1-n} - \sum_{n=0}^{k-2} \sum_{j=n}^{k-2} \binom{j}{n} (-1)^{j-n} F^{[j]}(x_0)$$



$$= 0, \quad \left\{ p_{k-1-j}(x_0, \dots, x_0) - p_{k-1-j}(x_0, \dots, x_0) \right\}$$

and for  $p = 0, \dots, k-2$

$$\int_{x_p - x_0}^{x_{p+1} - x_0} q_k(\underbrace{x_0, \dots, x_0}_{p+1}, x_{p+1} - u, \dots, x_{k-1} - u) du$$

$$= \sum_{n=0}^p \binom{k-1}{n} (-1)^{k-1-n} F^{[k-1]}(x_0) \left\{ p_1(x_{p+1}) - p_1(x_p) \right\} \quad (103)$$

$$+ \sum_{n=p+1}^{k-1} \binom{k-1}{n} (-1)^{k-1-n} \left\{ F^{[k]}(x_n + x_0 - x_p) - F^{[k]}(x_n + x_0 - x_{p+1}) \right\} \quad (104)$$

$$- \sum_{n=0}^p \sum_{j=n}^{k-2} \binom{j}{n} (-1)^{j-n} F^{[j]}(x_0) \int_{x_p}^{x_{p+1}} \left\{ p_{k-1-j}(\underbrace{v, \dots, v}_{p-n}, x_{p+1}, \dots, x_{k-j+n-1}) \right. \\ \left. - p_{k-1-j}(\underbrace{v, \dots, v}_{p-n+1}, x_{p+1}, \dots, x_{k-j+n-2}) \right\} du \quad (105)$$

$$- \sum_{n=p+1}^{k-2} \sum_{j=n}^{k-2} \binom{j}{n} (-1)^{j-n} \left\{ F^{[j+1]}(x_n + x_0 - x_p) - F^{[j+1]}(x_n + x_0 - x_{p+1}) \right\} \\ \left\{ p_{k-1-j}(x_{n+1}, \dots, x_{k-j+n-1}) - p_{k-1-j}(x_n, \dots, x_{k-j+n-2}) \right\} \quad (106)$$

$$= i_p + ii_p + iii_p + iv_p,$$

with  $i_p, ii_p, iii_p$  and  $iv_p$  representing expression (103), (104), (105) and (106), respectively. Using similar calculations as before we get the following expressions for the summations of the different terms:

$$\sum_{p=0}^{k-2} i_p = -F^{[k-1]}(x_0) \sum_{n=0}^{k-1} \binom{k-1}{n} (-1)^{k-1-n} p_1(x_n), \quad (107)$$

$$\sum_{p=0}^{k-2} ii_p = - \sum_{n=0}^{k-1} \binom{k-1}{n} (-1)^{k-n} F^{[k]}(x_n), \quad (108)$$

$$\sum_{p=0}^{k-2} iii_p = - \sum_{n=0}^{k-2} \sum_{j=n}^{k-2} \binom{j}{n} (-1)^{j-n} F^{[j]}(x_0) p_{k-j}(x_n, \dots, x_{k-j+n-1}), \quad (109)$$

$$\sum_{p=0}^{k-2} iv_p = \sum_{n=1}^{k-2} \sum_{j=n+1}^{k-1} \binom{j-1}{n} (-1)^{j-n} \left\{ F^{[j]}(x_n) - F^{[j]}(x_0) \right\} \\ \left\{ p_{k-j}(x_{n+1}, \dots, x_{k-j+n}) - p_{k-j}(x_n, \dots, x_{k-j+n-1}) \right\}. \quad (110)$$

So we know that  $q_{k+1}(x_0, x_1, \dots, x_k)$  for  $k \geq 1$  is given by the sum of (99), (100), (101) and (102), subtracted by the sum of (107), (108), (109) and (110). To continue we evaluate the following differences:

$$\sum_{p=1}^{k-1} I_p - \sum_{p=0}^{k-2} i_p = - \sum_{n=0}^{k-1} \binom{k-1}{n} (-1)^{k-1-n} F^{[k-1]}(x_0) \left\{ p_1(x_{n+1}) - p_1(x_n) \right\}, \quad (111)$$

whereas

$$\sum_{p=0}^{k-1} II_p - \sum_{p=0}^{k-2} ii_p = F^{[k]}(x_k) + \sum_{n=1}^{k-1} \left\{ \binom{k-1}{n-1} + \binom{k-1}{n} \right\} (-1)^{k-n} F^{[k]}(x_n) \\ + (-1)^k F^{[k]}(x_0) \\ = \sum_{n=0}^k \binom{k}{n} (-1)^{k-n} F^{[k]}(x_n), \quad (112)$$

since  $\binom{k-1}{n-1} + \binom{k-1}{n} = \binom{k}{n}$ . Furthermore,

$$\sum_{p=1}^{k-1} III_p - \sum_{p=0}^{k-2} iii_p = - \sum_{n=0}^{k-2} \sum_{j=n}^{k-2} \binom{j}{n} (-1)^{j-n} F^{[j]}(x_0) \left\{ p_{k-j}(x_{n+1}, \dots, x_{k-j+n}) \right. \\ \left. - p_{k-j}(x_n, \dots, x_{k-j+n-1}) \right\} \quad (113)$$

and

$$\sum_{p=0}^{k-2} IV_p - \sum_{p=0}^{k-2} iv_p = - \sum_{n=0}^{k-1} \sum_{j=n}^{k-1} \binom{j}{n} (-1)^{j-n} \left\{ F^{[j]}(x_n) - F^{[j]}(x_0) \right\} \\ \left\{ p_{k-j}(x_{n+1}, \dots, x_{k-j+n}) - p_{k-j}(x_n, \dots, x_{k-j+n-1}) \right\}, \quad (114)$$

where we extended the summation to start from  $n = 0$  instead of  $n = 1$ . This does not change the overall value of (114) as all these additional terms cancel out.

Looking at the sum of (111) and (113), we realize that the result is just equal to the negative of the part of (114) involving  $F^{[j]}(x_0)$ , i.e. these parts cancel out when summed up. Thus, we are left with (112) and the remaining part of (114), and we eventually obtain that

$$q_{k+1}(x_0, x_1, \dots, x_k) = \sum_{n=0}^k \binom{k}{n} (-1)^{k-n} F^{[k]}(x_n) - \sum_{n=0}^{k-1} \sum_{j=n}^{k-1} \binom{j}{n} (-1)^{j-n} F^{[j]}(x_n)$$

$$\left\{ p_{k-j}(x_{n+1}, \dots, x_{k-j+n}) - p_{k-j}(x_n, \dots, x_{k-j+n-1}) \right\},$$

which is the statement of Theorem 4.

□

## References

- [1] Asmussen, S. (1987) *Applied Probability and Queues*. J. Wiley & Sons, Chichester.
- [2] Baccelli, F., Cohen, G., Olsder, G.J. and J.P. Quadrat (1992) *Synchronization and Linearity*. J. Wiley & Sons, Chichester.
- [3] Baccelli, F. and V. Schmidt (1996) Taylor Series Expansions for Poisson Driven (max,+)-Linear Systems. *Ann. Appl. Probab.* (to appear)
- [4] Błaszczyszyn, B. (1995) Factorial moment expansion for stochastic systems. *Stoch. Proc. Appl.* **56**, 321–335.
- [5] Błaszczyszyn, B., Frey, A. and V. Schmidt (1995) Light-traffic approximations for Markov-modulated multi-server queues. *Stochastic Models* **11**, 423–445.
- [6] Błaszczyszyn, B. and T. Rolski (1995) Expansions for Markov-modulated systems and approximations of ruin probability. *J. Appl. Probab.* **32** (to appear)
- [7] Frey, A. and V. Schmidt (1996) Taylor-series expansion for multivariate characteristics of risk processes. *Insurance: Mathematics and Economics* (to appear)
- [8] Kroese, D.P. and V. Schmidt (1996) Light-traffic analysis for queues with spatially distributed arrivals. *Math. Oper. Res.* (to appear)
- [9] Reiman, M.I. and B. Simon (1989) Open queueing systems in light traffic. *Math. Oper. Res.* **14**, 26–59.



---

Unité de recherche INRIA Lorraine, Technopôle de Nancy-Brabois, Campus scientifique,  
615 rue du Jardin Botanique, BP 101, 54600 VILLERS LÈS NANCY  
Unité de recherche INRIA Rennes, Irisa, Campus universitaire de Beaulieu, 35042 RENNES Cedex  
Unité de recherche INRIA Rhône-Alpes, 46 avenue Félix Viallet, 38031 GRENOBLE Cedex 1  
Unité de recherche INRIA Rocquencourt, Domaine de Voluceau, Rocquencourt, BP 105, 78153 LE CHESNAY Cedex  
Unité de recherche INRIA Sophia-Antipolis, 2004 route des Lucioles, BP 93, 06902 SOPHIA-ANTIPOLIS Cedex

---

Éditeur  
INRIA, Domaine de Voluceau, Rocquencourt, BP 105, 78153 LE CHESNAY Cedex (France)  
ISSN 0249-6399