



Large deviation probability and local density of sets

Philippe Barbe, Michel Broniatowski

► **To cite this version:**

Philippe Barbe, Michel Broniatowski. Large deviation probability and local density of sets. [Research Report] RR-2630, INRIA. 1995. <inria-00074057>

HAL Id: inria-00074057

<https://hal.inria.fr/inria-00074057>

Submitted on 24 May 2006

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Large deviation probability and local density of sets

Philippe BARBE, Michel BRONIATOWSKI

N° 2630

Août 1995

PROGRAMME 5

 ***rapport
de recherche***



Large deviation probability and local density of sets

Philippe BARBE, Michel BRONIATOWSKI

Programme 5 — Traitement du signal, automatique et productique

Projet Fractales

Rapport de recherche n° 2630 — Août 1995 — 19 pages

Abstract: Let X_1, X_2, \dots, X_n be n independent identically distributed real random variables and $S_n := \sum_{i=1}^n X_i$. We obtain precise asymptotics for $P(S_n \in nA)$ for rather arbitrary Borel sets A , in terms of the density of the dominating points in A .

Our result extends classical theorems in the field of large deviations for independent samples. We also obtain asymptotics for $P(S_n \in \gamma_n A)$, with $\gamma_n/n \rightarrow \infty$.

Key-words: large deviation, fractal, local density.

AMS (1991) Classification : 60F10 ; 28A80.

(Résumé : tsvp)

Philippe Barbe: CNRS Toulouse: e-mail: barbe@cict.fr

Michel Broniatowski: Université de Reims - CNRS :e-mail: mbr@ccr.jussieu.fr

Unité de recherche INRIA Rocquencourt

Domaine de Voluceau, Rocquencourt, BP 105, 78153 LE CHESNAY Cedex (France)

Téléphone : (33 1) 39 63 55 11 – Télécopie : (33 1) 39 63 53 30



Probabilité de grande déviation et dimension locale des ensembles

Résumé : Soient X_1, X_2, \dots, X_n , n variables aléatoires réelles indépendantes identiquement distribuées et $S_n := \sum_{i=1}^n X_i$. Nous obtenons des évaluations asymptotiques de $P(S_n \in n A)$ pour des ensembles Boréliens A assez généraux, en termes de la densité locale des points dominants dans A . Ces résultats généralisent des résultats classiques dans la théorie des grandes déviations pour des suites de variables indépendantes. Nous obtenons également des équivalents asymptotiques de $P(S_n \in \gamma_n A)$, lorsque $\gamma_n/n \rightarrow \infty$.

Mots-clé : grande déviation, fractale, densité locale.

AMS (1991) Classification : 60F10 ; 28A80.

1. Introduction

Let X_1, X_2, \dots, X_n be n independent and identically distributed (i.i.d) real valued random variables (r.v.'s) and let $S_n := \sum_{1 \leq i \leq n} X_i$. The main goal of this paper is to obtain precise asymptotics for $P(S_n \in n A)$ for rather arbitrary Borel sets A . This problem is strongly connected to various topics studied in the literature about large deviations and about fractal geometry.

Assume that

$$(H1) \quad X_1 \text{ is non degenerate, i.e. } P(X_1 = c) < 1 \text{ for all } c \in \mathbb{R},$$

$$(H2) \quad \phi(t) = E(e^{tX_1}) < \infty \text{ on some non-void set } \mathcal{D} \text{ with } 0 \in \text{Int}\mathcal{D}.$$

Set

$$t_0 := \sup\{t : \phi(t) < \infty\}.$$

Define the Chernoff transform

$$\Lambda(x) := \text{Sup}\{tx - \log \phi(t) ; t \in \mathbb{R}\}$$

and set

$$\Lambda(A) := \inf\{\Lambda(x) ; x \in A\}.$$

Chernoff's (1952) theorem asserts that, for any Borel set A ,

$$\begin{aligned} -\Lambda(\text{Int } A) &\leq \liminf_{n \rightarrow \infty} n^{-1} \log P(S_n \in n A) \\ &\leq \limsup_{n \rightarrow \infty} n^{-1} \log P(S_n \in n A) \leq -\Lambda(\text{cl } A). \end{aligned} \quad (1.1)$$

Thus, if $\Lambda(\text{Int } A) = \Lambda(\text{cl } A)$,

$$\lim_{n \rightarrow \infty} n^{-1} \log P(S_n \in n A) = -\Lambda(A). \quad (1.2)$$

How good is (1.1)? For instance, if $A = \{1\} \cup [2, \infty)$ and $E(X_1) = 0$, then $\Lambda(A) = \Lambda(1)$, while (1.2) holds with $\Lambda(2) = \Lambda(\text{Int } A)$. Of course this example just stresses the rather well known fact that one should take an essential infimum instead of an infimum in the definition of $\Lambda(A)$.

If we rewrite (1.2) as

$$P(S_n \in n A) = \exp(-n\Lambda(A))(1 + o(1)) \text{ as } n \rightarrow \infty, \quad (1.3)$$

it is natural to look for an equivalent of the probability itself instead of its logarithm. Further, we wish such a result to reflect some information about the fine structure of A .

Bahadur and Ranga-Rao (1960), and Petrov (1965) solve partly this problem. Consider the following hypothesis, stronger than H1,

(H3) X_1 has a continuous distribution.

Set $m(t) := (\log \phi)'(t)$ and $s^2(t) := (\log \phi)''(t)$.

Petrov's (1965) theorem asserts that, if $A = [a, \infty)$, $a > E(X_1)$, then

$$P(S_n \in n A) = P(S_n \geq n a) = \frac{e^{-n\Lambda(a)}}{\sqrt{n\Psi(a)}}(1 + o(1)), \text{ as } n \rightarrow \infty \quad (1.4)$$

where $\Psi(a) := t^* s(t^*) \sqrt{2\pi}$, and t^* is the unique root of the equation $a = m(t)$ in \mathcal{D} . For sets A more general than $[a, \infty)$, (1.4) suggests that all what really matters is the essential infimum of A . This is clearly true if the essential infimum a of A is the end point of an interval contained in A . For the multidimensional case, see Ney (1983).

What about a very discontinuous set A ? Recall that the density of a in A is defined, whenever it exists, by

$$d(a) := \lim_{\varepsilon \rightarrow 0} \frac{|A \cap [a - \varepsilon, a + \varepsilon]|}{2\varepsilon}.$$

Is (1.4) still true if $d(a) = 0$? We shall investigate the link between (1.4) and the structure of A near its essential infimum; see Falconer (1990) for the properties of d and related concepts.

We also mention other works related to (1.4). Blackwell and Hodges (1959) proved a version of (1.4) for lattice distributions. Petrov and Širikova (1973) showed that if $\phi(t) = \infty$ for any $t > 0$, then $\limsup_{n \rightarrow \infty} \rho^{-n} P(S_n \geq na) = 0$ for any $\rho \in \mathbb{R}$ (see also Steinebach (1980) and references therein). Finally, Jensen (1988) obtained uniform expansions for

$P(S_n \geq na)$ under stronger assumptions than those given above; for a survey about statistical applications of these results, see Fields and Ronchetti (1990).

Our investigation of $P(S_n \in nA)$ for rather arbitrary Borel sets A (not only closed or open) is also related to the recent interest in large deviations without topology (see Ben Arous and Ledoux (1993) for Schilder's theorem and Deheuvels and Lifshits (1993) for related investigations on Strassen's (1964) law of the iterated logarithm). The study of $P(S_n \in nA)$ is also related to the Erdős-Rényi (1970) law of large numbers (see e.g. Deheuvels, Devroye and Lynch (1986)).

We shall also study asymptotics for $P(S_n \in \gamma_n A)$ for general sequences satisfying $\gamma_n/n \rightarrow \infty$. The case $A = [a, +\infty)$ has been partly explored by Broniatowski and Mason (1994).

Section 2 presents our results, along with some remarks about a new definition of the local density of a in A . Section 3 contains the proofs.

2. Results

2.0.0

2.1 Large deviation probabilities

In the sequel we assume that (H1), (H2) and (H3) hold.

Denote α the essential infimum of A with respect to the Lebesgue measure,

$$\alpha := \text{essinf} A := \inf\{x : \text{for all } \epsilon > 0, |[x, x + \epsilon] \cap A| > 0\},$$

with $\inf \emptyset = -\infty$.

Assume

$$(H4) \quad \alpha > -\infty.$$

(H4) means that we consider rather thick sets A ; for example we do not consider Cantor type sets.

The density of α in A will not be measured in the ordinary way, but will be related to the more appropriate quantity

$$M(t) := t \int I_{A-\alpha}(y) e^{-ty} dy, \quad t > 0. \quad (2.1.1)$$

For any set A , $0 \leq M(t) \leq 1$. Notice that if $A = [\alpha, \infty)$, then $M(t) = 1$ for any $t > 0$. If there exists an interval $[\alpha, \alpha + \varepsilon] \subset A$, then $\lim_{t \rightarrow \infty} M(t) = 1$.

Here is an example for a self-similar set. Let $p > 2$ and $I_p := \left[\frac{p-1}{p}, 1 \right]$. Set $A := A_p := \bigcup_{n \in \mathbb{Z}} p^n I_p$. Then $0 = \text{essinf} A_p$ and $pA_p = A_p$. Consequently, for any $t \geq 0$, $M(tp) = M(t)$. Thus

$$\inf_{1 \leq u \leq p} M(u) = \liminf_{t \rightarrow \infty} M(t) \leq \limsup_{t \rightarrow \infty} M(t) = \sup_{1 \leq u \leq p} M(u).$$

It is convenient to define

$$M_n(t) := M(nt)/t = \int I_{A-\alpha}(y) e^{-nty} dy \quad (2.1.2)$$

and

$$\Psi_n(t) := n \log \phi(t) + \log M_n(t) - n \alpha t, \quad (2.1.3)$$

for all $t > 0$ such that $\phi(t) < \infty$.

It is well known that when (H1) and (H2) hold, $\log \phi$ is a strictly convex function. Furthermore, $m(0) = E(X_1)$ and $m(t)$ is a strictly increasing function, which we assume to satisfy:

$$(H5) \quad \lim_{t \rightarrow t_0} m(t) = +\infty.$$

See Deheuvels, Devroye and Lynch (1986) for a discussion on (H5) in connection with Petrov's (1965) theorem.

We quote some properties on the function M defined in (2.1.1). Set $\mu_n(t) := (1/n) \log M_n(t)$. For any $n \geq 1$, μ_n is a decreasing function on $[0, \infty)$, negative for n large enough. Furthermore, $\mu'_n(t) = \mu'_1(nt)$, and μ'_1 is non decreasing on $[0, \infty)$.

Let $\bar{\mu} := \lim_{t \rightarrow \infty} \mu'_1(t)$ and $\underline{\mu} := \lim_{t \rightarrow 0} \mu'_1(t)$.

Lemma 2.1.1. *Assume that (H1), (H2) and (H5) hold. Then the equation $\Psi'_n(t) = 0$ has a unique solution t_n in $(0, t_0)$ for $\alpha \in (E(X_1) + \bar{\mu}, \infty)$.*

Furthermore

(1) *If $\alpha \leq E(X_1) + \underline{\mu}$, then $\lim_{n \rightarrow \infty} t_n = 0$.*

(2) *If $\alpha > E(X_1) + \underline{\mu}$, then there exists a compact subset $K \subset (0, t_0)$ such that $t_n \in K$ for all $n \geq 1$.*

We postpone the proof to Section 3.

From now on, we suppose that $\alpha > E(X_1) + \underline{\mu}$.

Set $\psi_n(t) := \Psi''_n(t)$ and suppose that for any $\lambda > 0$,

$$(H6) \quad \lim_{n \rightarrow \infty} \sup_{|u| < \lambda} \frac{\psi_n(t_n + u/\sqrt{\psi_n(t_n)})}{\psi_n(t_n)} = 1$$

where t_n is the solution of

$$\Psi'_n(t) = 0 \tag{2.1.4}$$

in the range $(0, t_0)$.

The following result shows that (H6) is not very restrictive.

Recall that a function l is slowly varying at $+\infty$ if for any $\lambda > 0$, $\lim_{x \rightarrow +\infty} l(\lambda x)/l(x) = 1$. In this case we note $l \in \mathcal{R}_0(+\infty)$. We say that l is slowly varying at 0 if $l(1/\cdot) \in \mathcal{R}_0(+\infty)$; we denote $l \in \mathcal{R}_0(0)$. A function g is regularly varying as 0 or $+\infty$ if $g(x) = x^\rho l(x)$ for some $l \in \mathcal{R}_0(0)$ or $l \in \mathcal{R}_0(+\infty)$. We denote $g \in \mathcal{R}_\rho(0)$ or $g \in \mathcal{R}_\rho(+\infty)$; see Bingham, Goldie and Teugels (1987) for the theory of regular variation.

Lemma 2.1.2. *Assume that (H1), (H2) and (H5) hold.*

A sufficient condition for (H6) is

$$\log(M(t)/t) \in \mathcal{R}_\rho(\infty) \text{ for some } \rho \in [0, 1].$$

We also need the following condition:

$$(H7) \quad \limsup_{t \rightarrow \infty} t (\log M(t))'' < \infty.$$

A sufficient condition for (H7) is $\log(M(t)/t) \in \mathcal{R}_\rho$, for $0 \leq \rho < 1$.

We now state our main result.

Theorem. 2.1.1 *Assume that (H2) - (H7) hold. Then, for $\alpha = \text{essinf} > E(X_1) + \underline{\mu}$,*

$$P(S_n \in nA) = \frac{\phi^n(t_n) \cdot M_n(t_n) \cdot e^{-nt_n\alpha}}{\psi_n(t_n)\sqrt{2\pi}}(1 + o(1)) \text{ as } n \rightarrow \infty, \quad (2.1.5)$$

with t_n satisfying (2.1.4), provided that the function $x \mapsto P(S_n \in nA + x)$ is nonincreasing for n large enough. In particular, this last condition holds if

- (i) (Petrov): $A = (\alpha, \infty)$ or $A = [\alpha, \infty)$ (in this case (H7) is satisfied and $M_n(t) = 1/t$), or
- (ii) X_1 has a symmetric unimodal distribution or
- (iii) X_1 has a strongly unimodal distribution .

The shape of A near α is reflected in the behavior of the function $M(t)$ for large values of t . By (2.1.2) and Lemma 2.1.1, the more n is large, the more the shape of A near α is relevant in (2.1.5).

One should notice that the formula given in Theorem is the exact analogue of (1.4). Indeed, using Hölder Inequality, one easily checks that $\Psi_n(t)$ is convex.

By Lemma 2.1.1, t_n is unique in K , and t_n achieves $\sup\{n\alpha t - n \log \phi(t) + \log M_n(t)\}$; compare with the definition of $\Lambda(n\alpha)$. However, recall that the function M_n depends upon the regularity of the set A near α , while the function Λ only depends upon the distribution of X_1 .

Notice that $M_n(t)e^{-nt\alpha} = \int I_A(y)e^{-nty}dy$, from which we see that α plays no role in (2.1.5). Therefore we can substitute α by any real number γ for which $\int I_{A-\gamma}(y)e^{-ty}dy$ converges. Further, t_n defined in (2.1.4) is independent upon α . The so-called dominating point α in A can therefore be defined by $\alpha := \lim_{t \rightarrow \infty} t^{-1} \log \int I_A(y)e^{-ty}dy$.

2.2 Very large deviation probabilities

We consider asymptotics for

$$P(S_n \in \gamma_n A), \text{ with } \gamma_n/n \rightarrow +\infty.$$

For all $\alpha \in (E(X_1) + \underline{\mu}, \infty)$, let $t_n = t_n(\alpha)$ be the unique solution of the equation

$$\Psi_n(t) = 0$$

in the range $(0, t_0)$, where now

$$\Psi_n(t) = n \log \phi(t) + \log M_n(t) - \gamma_n t \alpha$$

and

$$M_n(t) = M(\gamma_n t)/t.$$

It is easy to check that $\lim_{n \rightarrow \infty} t_n = t_0$.

The following result holds:

Theorem. 2.2.1 *Assume that X_1 has a strongly unimodal distribution and (H2), (H4) - (H7) hold. With the above notation, (2.1.5) holds.*

Whenever $t_0 = +\infty$, $s^2(t)$ and $(\log M(t))''$ are regularly varying functions at $+\infty$, it is easy to characterize sequences γ_n such that (H6) holds.

2.3 Density of α in A and the function $M(t)$

In order to investigate further the role played in (2.1.5) by the regularity of the set A near its essential infimum α , we define

$$G(\epsilon) := |A \cap [\alpha, \alpha + \epsilon]|, \quad \epsilon \geq 0.$$

We interpret G as defining a measure on \mathbb{R}^+ . Shifting slightly the terminology used in the study of multifractal measures (see e.g. Falconer (1990)), we define the pointwise Hölder dimension of A at α as

$$\delta(\alpha) := \lim_{\epsilon \rightarrow 0} \frac{\log G(\epsilon)}{-\log \epsilon},$$

if the limit exists.

For instance, for the set A_p defined above, $G(\epsilon) = 1/p^k(p-1)$ if $p^{-k-1} \leq \epsilon < p^{-k}$, so that $\epsilon/(p-1) \leq G(\epsilon) \leq p\epsilon/(p-1)$ for any $\epsilon > 0$, and thus $\delta(0) = 1$. So, A_p is reasonably big around 0.

There is a strong connection between the density, the pointwise Hölder dimension and the behavior of the function $M(t)$.

Proposition 2.3.1. *i) If $M(t)/t \in \mathcal{R}_{-\rho}(\infty)$, then $0 \leq \rho \leq 1$. Moreover, $M(t) \sim c t^{-\rho+1} l(t)$ as $t \rightarrow \infty$, for some $l \in \mathcal{R}_0(\infty)$ iff $G(\varepsilon) \sim c \varepsilon^{\rho l(1/\varepsilon)}/\Gamma(1+c)$, as $\varepsilon \rightarrow 0$.*

ii) If $\log(M(t)/t) \in \mathcal{R}_{1/(1+\rho)}(\infty)$, then $\rho \geq 0$. Moreover $-\log G(\varepsilon) \sim c/\phi^-(1/\varepsilon)$ as $\varepsilon \rightarrow 0$ and for $\phi \in \mathcal{R}_{-\rho}(0^+)$ and $c > 0$ iff $-\log(M(t)/t) \sim (1+\rho) \left(\frac{\varepsilon}{\rho}\right)^{\rho/(\rho+1)} \frac{1}{\Psi^-(t)}$ as $t \rightarrow \infty$ where here $\Psi(\varepsilon) = \phi(\varepsilon)/\varepsilon \in \mathcal{R}_{-\rho-1}(0^+)$, and ϕ^- (resp. Ψ^-) denotes the generalized inverse of ϕ (resp. Ψ).

This proposition follows readily from classical Abel–Tauber theorems (see Bingham, Goldie and Teugels (1987), Ch. 2 and 4). For instance, we deduce from Proposition 2.3.1 that $G(\varepsilon) \sim \varepsilon^{\delta(\alpha)}$ (as $\varepsilon \rightarrow 0$) iff $M(t) \sim c t^{-\delta(\alpha)+1} \Gamma(1+\delta(\alpha))$ (as $t \rightarrow \infty$). Consequently, if $M_n(t) \rightarrow 1$ as $t \rightarrow \infty$ then $M(t) \sim t$ as $t \rightarrow \infty$ and $G(\varepsilon) \sim \varepsilon$ as $\varepsilon \rightarrow 0$.

3. Proofs

3.1 Proof of Lemma 2.1.1.

By the properties of the function μ_1 , $-\infty < \underline{\mu} \leq 0$. The mapping $\alpha \rightarrow t_n$ is one to one from $(E(X_1) + \bar{\mu}, \infty)$ onto $(0, t_0)$. Note that the solution t_n of (2.1.4) may not be unique on \mathcal{D} (f.i. if X_1 is normally distributed and $A = [\alpha, \infty)$).

i) Assume that $\liminf_{n \rightarrow \infty} t_n > t > 0$. For some sequence $\{k\} \subset \{n\}$, $\lim_{k \rightarrow \infty} m(t_k) + \mu'_1(k t_k) = \alpha > m(t) + \underline{\mu} > E(X_1) + \underline{\mu}$, a contradiction.

ii) If $\limsup_{n \rightarrow \infty} t_n = t_0$, then, for some $\{k\} \subset \{n\}$, $\lim_{k \rightarrow \infty} m(t_k) + \mu'_1(k t_k) = \alpha$, which implies $\lim_{k \rightarrow \infty} \mu'_1(k t_k) = -\infty$, a contradiction.

If $\liminf_{n \rightarrow \infty} t_n = 0$, then $\lim_{k \rightarrow \infty} m(t_k) + \mu'_1(k t_k) = \alpha < E(X_1)$, a contradiction. ■

3.2 Proof of Lemma 2.1.2.

The expression in (H6) is equal to

$$\frac{\mu''_1 \left(n \left(t_n + u / \sqrt{\psi_n(t_n)} \right) \right)}{\mu''_1(n t_n)} (1 + o(1)) \text{ as } n \rightarrow \infty.$$

Since t_n belongs to some compact set K (see Lemma 2.1.1) and $\lim_{n \rightarrow \infty} \psi_n(t_n) = +\infty$, the result follows from known facts on regular variation. ■

3.3 Proof of Theorem 2.1.1

The proof will be covered by several lemmas.

Lemma 3.3.1. *For any $t \in (0, t_0)$, if (H_4) holds,*

$$\int e^{tx} P(S_n \in nA + x) dx = \phi(t)^n M_{n,\gamma}(t) e^{-nt\gamma} \quad (3.3.1)$$

for all $n \geq 1$ and $\gamma \in \mathbb{R}$, where $M_{n,\gamma}(t) = (1/t) \int I_{A-\gamma}(y) e^{-nty} dy$.

Proof of Lemma 3.3.1.

Since all the functions that we integrate are nonnegative, we can use Tonelli's theorem and obtain

$$\begin{aligned} & \int e^{tx} P(S_n \in nA + x) dx = \int \int e^{tx} I_{x+nA}(u) dx P(S_n = du) \\ &= \int \int e^{tx} I_{u-nA}(x) dx P(S_n = du) = \int \int e^{t(u-ny)} I_A(y) n dy P(S_n = du) \\ &= n \int e^{tu} P(S_n = du) \int I_A(y) e^{-nty} dy = n \phi(t)^n e^{-tn\gamma} \int I_{A-\gamma}(y) e^{-tny} dy. \quad \blacksquare \end{aligned}$$

Choosing $\gamma = \alpha$, we obtain

$$\int e^{tx} P(S_n \in nA + x) dx = \phi(t)^n M_n(t) e^{-tn\alpha}. \quad (3.3.2)$$

Notice that Lemma 3.3.2 also holds in \mathbb{R}^d :

$$\int e^{\langle t, x \rangle} P(S_n \in nA + x) dx = \phi(t)^n e^{-n\langle t, \gamma \rangle} M_{n,\gamma}(t),$$

with

$$M_n(t) := n^d \int I_{A-\gamma}(y) e^{-n\langle t, y \rangle} dy$$

and assuming that $\int e^{\langle t, x \rangle} P(S_n \in nA + x) dx < +\infty$.

We now define a family of densities indexed by $t \in (0, t_0)$ namely, by (3.3.2),

$$g_n(x, t) := \frac{e^{tx} P(S_n \in nA + x)}{\phi(t)^n M_n(t) e^{-tn\alpha}}.$$

The main line of the proof consists in showing that $g_n(\cdot, t)$, suitably rescaled, converges to the Gaussian density, and then take $g_n(0, t_n)$, with t_n defined as in Lemma 2.1.1.

Notice that the Laplace transform of $g_n(\cdot, t)$ is

$$L_n(\lambda, t) := \int e^{\lambda x} g_n(x, t) dx = \frac{\phi^n(t + \lambda) M_n(t + \lambda)}{\phi^n(t) M_n(t)} e^{-n\lambda\alpha}. \quad (3.3.3)$$

We now calculate the expectation and the variance of $g_n(\cdot, t)$.

Lemma 3.3.2. *We have $c_n(t) = \int x g_n(x, t) dx = n m(t) + (\log M_n)'(t)$ and*

$$v_n^2(t) := \int (x - \mu_n(t))^2 g_n(x, t) dx = n s^2(t) + (\log M_n)''(t).$$

Proof of Lemma 3.3.2. Take the derivative at $\lambda = 0$ in (3.3.3) in order to calculate the cumulants of the density $g_n(\cdot, t_n)$ ■.

It is convenient to introduce a family of r.v.'s, say Y_n , with density $g_n(\cdot, t_n)$. Recall that we have assumed $\alpha > E(X_1) + \underline{\mu}$. Also, by Lemma 3.3.2, $E(Y_n) = c_n(t_n) = 0$.

Our next Lemma is a central limit theorem for Y_n .

Lemma 3.3.3. *Whenever (H6) holds, $Y_n/v_n(t) \xrightarrow{d} \mathcal{N}(0, 1)$ as $n \rightarrow \infty$.*

Proof of Lemma 3.3.3. Taylor's expansion at $\lambda = 0$, Lemma 3.3.2, the choice of t_n and the definition of Ψ_n yields, for some $\theta = \theta_n(\lambda) \in (0, 1)$,

$$\begin{aligned} \log E(e^{\lambda Y_n/v_n(t_n)}) &= \log L_n(\lambda/v_n(t_n), t_n) \\ &= \frac{\lambda^2}{2v_n(t_n)^2} (\log L_n(\cdot, t_n))''(\theta\lambda/v_n(t_n)) \\ &= \frac{\lambda^2 \psi_n(t_n + \theta\lambda/\sqrt{\psi_n(t_n)})}{2 \psi_n(t_n)}. \end{aligned}$$

Assuming (H6) the proof follows from classical convergence criteria (see e.g. Billingsley (1978), p. 345). ■

Having a central limit theorem for $Y_n/v_n(t_n)$, we turn it into a local central limit theorem on the density of $Y_n/v_n(t_n)$.

Let

$$h_n(x) := v_n(t_n)g_n(xv_n(t_n), t_n)$$

and its characteristic function (c.f.)

$$\hat{h}_n(u) := \int e^{iux} h_n(x) dx.$$

Lemma 3.3.4. *Assume that (H6) holds. Then there exists two positive constant C and δ such that*

$$|\hat{h}_n(\omega)| \leq e^{-C \omega^2} \quad \text{for} \quad |\omega| < \delta v_n(t_n).$$

Proof of Lemma 3.3.4. Since

$$\left| e^{ix} - \left(1 + ix - \frac{x^2}{2} \right) \right| \leq \frac{|x|^3}{6},$$

we have

$$\frac{|\phi(t_n + iu)|}{\phi(t_n)} \leq |1 - (a - ib)| + c$$

with

$$a = \frac{u^2}{2} \left(\frac{\phi''}{\phi} \right) (t_n), \quad b = u \left(\frac{\phi'}{\phi} \right) (t_n), \quad c = \frac{|u|^3}{6} \left(\frac{\phi^{(3)}}{\phi} \right) (t_n)$$

and

$$\phi^{(3)}(t) := \int |x|^3 e^{tx} dF(x)$$

where F is the d.f. of X_1 .

Let

$$A = 2a - b^2 - a^2 - c^2 - 2c\sqrt{1 + a^2 + b^2 - 2a} = u^2 s^2(t_n) - a^2 - c^2 - 2c\sqrt{1 + a^2 + b^2 - 2a}.$$

Since t_n is in a compact subset of $(0, 1)$, (Lemma 2.1.1), we have

$|a^2 + c^2 + 2c\sqrt{1 + a^2 + b^2 - 2a}| \leq c_1 u^3$ for some $c_1 > 0$ and provided $|u|$ is small enough. Therefore, in taking some $\delta > 0$ small enough, for $|u| \leq \delta$ we have

$$u^2 s^2(t_n)/2 \leq A \leq 2u^2 s^2(t_n) \leq 1/2.$$

Hence, for $|u| \leq \delta$,

$$\frac{|\phi(t_n + iu)|}{\phi(t_n)} \leq |1 - A|^{1/2} \leq 1 - A/2 \leq 1 - u^2 s^2(t_n).$$

Consequently, if $|\omega| \leq \delta v_n(t_n)$,

$$\frac{|\phi(t_n + i\omega/v_n(t_n))|^n}{\phi(t_n)^n} \leq \left(1 - \omega^2 \frac{s^2(t_n)}{v_n^2(t_n)}\right)^n.$$

If δ is small enough, the $\limsup_{n \rightarrow \infty} \delta s_n^2(t_n)/v_n^2(t_n) < 1$ and therefore, $\omega^2 s^2(t_n)/v_n(t_n) < 1$ if $|\omega| < \delta v_n(t_n)$ and so, for n large enough and $|\omega| < \delta v_n(t_n)$,

$$\frac{|\phi(t_n + i\omega/v_n(t_n))|^n}{\phi(t_n)^n} \leq \exp\left(-\omega^2 \frac{s^2(t_n)}{v_n^2(t_n)}\right). \quad \blacksquare$$

We state now the local central limit theorem.

Theorem. 3.3.1 *If (H1) – (H7) hold, then, for any $a < b$ and $x \in \mathbb{R}$,*

$$\lim_{n \rightarrow \infty} v_n(t_n) P(Y_n/v_n(t_n) \in (x + a/v_n(t_n), x + b/v_n(t_n))) = (b - a)\varphi(x),$$

where $\varphi(x) = (1/\sqrt{2\pi})e^{-x^2/2}$.

Proof of Theorem 3.3.1. Making use of the proof of Theorem 5.4 in Durrett (1991), it is enough to show that, for all θ and $x \in \mathbb{R}$,

$$v_n(t_n) E(l_\theta(Y_n - x v_n(t_n))) \rightarrow \varphi(x) \int l_\theta(y) dy \text{ as } n \rightarrow \infty, \quad (3.3.4)$$

where

$$l_\theta(y) = e^{i\theta y} l_0(y)$$

and

$$l_0(y) = \frac{1}{\pi} \frac{1 - \cos y}{y^2}$$

is the Pólya density function.

Denote \hat{l}_θ and \hat{l}_0 the Fourier transforms of l_θ and l_0 . Denote F_n the d.f. of the r.v. $Y_n - x v_n(t_n)$.

We have

$$\begin{aligned}
v_n(t_n)E(l_\theta(Y_n - xv_n(t_n))) &= v_n(t_n) \int l_\theta(y) dF_n(y) \\
&= \frac{v_n(t_n)}{2\pi} \int \int e^{-iuy} \hat{l}_\theta(u) du dF_n(y) \\
&= \frac{v_n(t_n)}{2\pi} \int e^{-iuy} dF_n(y) \int \hat{h}_\theta(u) du,
\end{aligned}$$

where we have used the inversion formula for the Fourier transform and Fubini's theorem.

Now,

$$\begin{aligned}
\int e^{-iuy} dF_n(y) &= E\left(e^{iu(Y_n - xv_n(t_n))}\right) \\
&= E\left(e^{-iuY_n}\right) e^{-iuxv_n(t_n)}.
\end{aligned}$$

Thus

$$\int e^{-iuy} dF_n(y) = e^{-iuxv_n(t_n)} \hat{h}_n(-uv_n(t_n)),$$

which yields

$$v_n(t_n)E(l_\theta(Y_n - xv_n(t_n))) = \frac{v_n(t_n)}{2\pi} \int \hat{h}_n(-uv_n(t_n)) e^{iuxv_n(t_n)} \hat{h}_\theta(u) du. \quad (3.3.5)$$

Let $M > 0$ such that $l_\theta(u) = 0$ for $u \in [-M, M]^{-c}$. Let $\delta > 0$ be as in Lemma 3.3.4, $I = [-\delta, \delta]$ and $J = [-M, M] \setminus I$. We split the integral in (3.3.5) on I and J .

$$\begin{aligned}
&\left| \frac{v_n(t_n)}{2\pi} \int_J \hat{h}_n(-uv_n(t_n)) e^{iuxv_n(t_n)} \hat{l}_\theta(u) du \right| \\
&\leq \frac{v_n(t_n)}{2\pi} \int_J |\hat{h}_n(-uv_n(t_n))| du \\
&\leq \frac{M}{\pi} v_n(t_n) \sup_{u \in J} |\hat{h}_n(-uv_n(t_n))|.
\end{aligned}$$

Moreover

$$|\hat{h}_n(-uv_n(t_n))| \leq \frac{|\phi(t_n - iu)|^n}{\phi(t)^n}.$$

Since F is continuous, there exists $\eta, 0 < \eta < 1$, such that

$$\left| \frac{v_n(t_n)}{2\pi} \int_J \hat{h}_n(-uv_n(t_n)) du \right| \leq v_n(t_n) \eta^n M / \pi, \quad (3.3.6)$$

which tends to 0 as $n \rightarrow \infty$.

We consider now

$$\frac{v_n(t_n)}{2\pi} \int_I \hat{h}_n(-uv_n(t_n)) e^{iuv_n(t_n)x} \hat{h}_\theta(u) du.$$

By Lemma 3.3.3 and 3.3.4 and the dominated convergence Theorem, we see that this expression tends to

$$\begin{aligned} \int e^{-\omega^2/2} e^{i\omega x} \hat{h}_\theta(0) d\omega &= \varphi(x) \hat{l}_\theta(0) \\ &= \varphi(x) \int l_\theta(y) dy. \end{aligned}$$

This completes the proof of Theorem 3.3.1. ■

We now complete the proof of Theorem 2.1.1.

By Theorem 3.3.1, we have, for all $\delta > 0$:

$$\frac{v_n(t_n)}{\delta} \int_0^{\delta/v_n(t_n)} h_n(x) dx = \rho(0) + \varepsilon(n, \delta), \quad (3.3.7)$$

with $\lim_{n \rightarrow \infty} \varepsilon(n, \delta) = 0$. Theorem 3.3.1 implies:

$$\frac{v_n(t_n) P(S_n \in nA)}{M_n(t_n) \phi(t_n)^n e^{-nt_n \alpha}} \frac{e^{t_n \delta}}{t_n \delta} \geq \varphi(0) + \varepsilon(n, \delta) / \delta$$

which yields

$$\frac{v_n(t_n) P(S_n \in nA)}{M_n(t_n) \phi(t_n)^n e^{-nt_n \alpha}} \geq \left(\varphi(0) + \frac{\varepsilon(n, \delta)}{\delta} \right) (1 - t_n \delta). \quad (3.3.8)$$

By Lemma 2.1.1, t_n belongs to some compact $K \subset (0, t_0)$. Taking the \liminf as $n \rightarrow \infty$ on both sides of (3.3.8) yields:

$$\liminf_{n \rightarrow \infty} \frac{v_n(t_n) P(S_n \in nA)}{M_n(t_n) \phi(t_n)^n e^{-nt_n \alpha}} \geq \varphi(0).$$

The lower bound for the \limsup is handled in the same way, using Theorem 3.3.1 with $x = 0$, $a = -\delta/v_n(t_n)$ and $b = 0$. ■

3.4 Proof of Theorem 2.2.1

Arguing as previously, we define

$$g_n(x, t) := \frac{e^{tx} P(S_n \in \gamma_n A + x)}{\phi^n(t) M_n(t) e^{-\gamma_n t \alpha}}.$$

The central limit theorem (Lemma 3.3.3) still holds whenever (H6) holds (see Lemma 2.1.2).

We cannot any longer prove the local central limit theorem 3.3.1. Instead, we prove that

Lemma 3.4.1. *For any $n \geq 1$, $g_n(\cdot, t)$ is a log-concave function.*

Proof of Lemma 3.4.1. It is enough to prove that $x \mapsto \log P(S_n \in \gamma_n A + x)$ is concave on \mathbb{R} .

The argument will go through discrete approximations. Consider first an integer-valued r.v. X with log-concave distribution, meaning that

$$P(X = m)^2 \geq P(X = m - 1)P(X = m + 1), m \in \mathbb{Z}.$$

It is known that $S_n = \sum_{i=1}^n X_i$ also possesses a log-concave distribution, where the X_i 's are n independent copies of X . Set $p_m = P(S_n = m)$, $n \in \mathbb{N}$. Consider first the case where $A = \{0, 1, 2, \dots, K\}$.

Define $\tilde{p} := (\dots, p_0, p_1, p_2, \dots)$ and $\tilde{1}_A := (\dots, 0, 1, 1, 1, \dots)$, where the first 1 in $\tilde{1}_A$ is $\tilde{1}_A(-K) = 1$. Clearly $\tilde{p} * \tilde{1}_A = \left(\sum_{i=m}^{m+K} p_i \right)_{m \in \mathbb{Z}}$. Since $\tilde{1}_A$ is log-concave, the same is $\tilde{p} * \tilde{1}_A$. Now, $(\tilde{p} * \tilde{1}_A)(m) = P(S_n \in A + m)$, which therefore is a log-concave function of m .

Consider now $C = A \cup B$ where A and B are two disjoint intervals of \mathbb{N} , say $A = \{0, 1, \dots, K\}$ and $B = \{\delta, \delta + 1, \dots, \delta + L\}$, $\delta > K$. Let $q_m := P(S_n \in A \cup B + m) = \sum_{i=m}^{m+K} p_i + \sum_{j=m+\delta}^{m+\delta+L} p_j$.

Direct calculation shows that

$$q_m^2 \geq q_{m-1} q_{m+1}$$

is equivalent to

$$\begin{aligned} & p_{m-1}p_{m+\delta+L} \left(\frac{p_m}{p_{m-1}} - \frac{p_{m+1+\delta+L}}{p_{m+\delta+L}} \right) \\ & \geq p_{m-1+\delta}p_{m+K} \left(\frac{p_{m+1+K}}{p_{m+K}} - \frac{p_{m+\delta}}{p_{m-1+\delta}} \right). \end{aligned} \quad (3.4.1)$$

Since (p_m) is log-concave,

$$\frac{p_{j+1}}{p_j} \leq \frac{p_l}{p_{l-1}} \text{ for } j > l,$$

by which (3.4.1) holds. Thus (q_m) is a log-concave sequence.

For the absolutely continuous case, it is enough to approximate the function $x \mapsto P(S_n \in \gamma_n A + x)$ by a sequence of lattice-valued log-concave functions with a decreasing grid-width, since log-concavity is preserved under simple limits. We omit the details. ■

We now complete the proof of Theorem 2.1.1.

Using Lemma 2 in Feigin and Yashchin (1983), Lemmas 3.3.3 and 3.4.1 we obtain

$$\lim_{n \rightarrow \infty} g_n(0, t_n) = \varphi(0)$$

as sought. ■

4. *

References

- [BAL93] G. Ben Arous and M. Ledoux. Schilder's large deviation principle without topology. Asymptotic Problems in Probability Theory: the Wiener functionals and asymptotics. *K.D. Elworthy, N. Ikeda, Editors. Pitman Research Notes in Mathematics Series*, 284:107–121, 1993. Longman.
- [BGT87] N. Bingham, C. Goldie, and J. Teugels. *Regular variation*. Cambridge University Press, Cambridge, 1987.
- [BH59] D. Blackwell and J.L. Hodges. The probability in the extreme tail of a convolution. *Ann. Math. Stat.*, 30:1113–1120, 1959.
- [Bill79] P. Billingsley. *Probability and Measure*. Wiley, New York, 1979.
- [BM94] M. Broniatowski and D. Mason. Extended large deviations. *Jour. Theoret. Prob.*, 7(3):647–666, 1994.
- [BRR60] R. Bahadur and R. Ranga Rao. On deviations of the sample mean. *Ann. Math. Stat.*, 31:1015–1027, 1960.
- [Che52] H. Chernoff. A measure of asymptotic efficiency for tests of a hypothesis based on the sum of observations. *Ann. Math. Stat.*, 23:493–507, 1952.
- [DDL86] P. Deheuvels, L. Devroye, and J. Lynch. Exact convergence rate in the limit theorems of Erdős-Rényi and Shepp. *Ann. Prob.*, 14:209–223, 1986.
- [DL95] P. Deheuvels and M. Lifshits. Necessary and sufficient conditions for the Strassen law of the iterated logarithm in non uniform topology. *Ann. Prob.*, 22, 1995. To appear.
- [Dur91] R. Durrett. *Probability: Theory and examples*. Wadsworth and Brooks, Pacific Grove, California, 1991.
- [ER70] P. Erdős and A. Rényi. On a new law of large numbers. *Jour. Anal. Math.*, 23:103–111, 1970.
- [Fal90] K. Falconer. *Fractal geometry*. Wiley, New York, 1990.
- [FR90] C. Fields and E. Ronchetti. Small sample asymptotics. *IMS Lecture Notes*, 13, 1990. Hayward, California.
- [FY83] P. Feigin and E. Yashchin. On a strong Tauberian result. *ZfW*, 65:35–48, 1983.
- [Jen88] J.L. Jensen. Uniform saddle point approximations. *Adv. Appl. Prob.*, 20:622–634, 1988.
- [Ney83] P. Ney. Dominating points and the asymptotics of large deviations for random walks on IR^d . *Ann. Prob.*, 11:158–167, 1983.
- [Pet65] V. Petrov. On the probabilities of large deviations for sums of independent random variables. *Th. Prob. Appl.*, 10(2):287–298, 1965.
- [PS73] V. Petrov and I.V. Sirikova. The exponential rate of convergence in the law of large numbers. *7 Math. Meh. Astronom.*, 2:155–157, 1973. Vestnik Leningrad University.

- [Ste80] J. Steinebach. Large deviation probabilities and related topics. *Carleton Math. Lecture Notes*, 28, 1980.
- [Str64] V. Strassen. An invariance principle for the law of the iterated logarithm. *ZfW*, 3:211–226, 1964.



Unité de recherche INRIA Lorraine, Technopôle de Nancy-Brabois, Campus scientifique,
615 rue du Jardin Botanique, BP 101, 54600 VILLERS LÈS NANCY
Unité de recherche INRIA Rennes, Irisa, Campus universitaire de Beaulieu, 35042 RENNES Cedex
Unité de recherche INRIA Rhône-Alpes, 46 avenue Félix Viallet, 38031 GRENOBLE Cedex 1
Unité de recherche INRIA Rocquencourt, Domaine de Voluceau, Rocquencourt, BP 105, 78153 LE CHESNAY Cedex
Unité de recherche INRIA Sophia-Antipolis, 2004 route des Lucioles, BP 93, 06902 SOPHIA-ANTIPOLIS Cedex

Éditeur

INRIA, Domaine de Voluceau, Rocquencourt, BP 105, 78153 LE CHESNAY Cedex (France)

ISSN 0249-6399