



About the rational map

Bernard Mourrain

► **To cite this version:**

| Bernard Mourrain. About the rational map. RR-2141, INRIA. 1993. <inria-00074531>

HAL Id: inria-00074531

<https://hal.inria.fr/inria-00074531>

Submitted on 24 May 2006

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

INSTITUT NATIONAL DE RECHERCHE EN INFORMATIQUE ET AUTOMATIQUE

*About the rational map
associated to a parallel robot*

Bernard Mourrain

N° 2141

Novembre 93

PROGRAMME 2

Calcul symbolique,
programmation
et génie logiciel



*Rapport
de recherche*

1994

About the rational map associated to a parallel robot

Bernard Mourrain

Programme 2 — Calcul symbolique, programmation et génie logiciel
Projet SAFIR*

Rapport de recherche n° 2141 — Novembre 93 — 41 pages

Abstract: In this paper, we consider the direct kinematic problem of a parallel robot (called the Stewart platform or left hand) from a mathematical point of view. We do not try to give real time and numerical solutions to this problem but describe tools of effective algebra, which can help us to know a little more about the geometric aspects of the question. A simple proof that the number of solutions we are expecting is actually 40 is then given. We use explicit eliminations techniques, in order to get rid of the solution at infinity and we use Bezout's theorem on surfaces with circularity to conclude. One main obstacle in the natural approach, via symbolic computations, is the great size of the polynomials appearing during the computations. We also try to show how the use of invariant theory and intrinsic approach may "factorize" the results in a more understandable way. In this text, the reader will also find some examples of successful and helpful use of a symbolic system (ie MAPLE).

Key-words: Parallel robot, rational map, Gröbner basis, resultant, component at infinity, circularity, symbolic computations

(Résumé : tsvp)

Work partially supported by PoSSo, EEC ESPRIT BRA contract 6846.

*SAFIR is a common project to INRIA (Sophia-Antipolis), Univ. De Nice-Sophia-Antipolis, CNRS

A propos de l'application rationnelle associée à un robot parallèle

Résumé : Nous considérons le problème direct d'un manipulateur parallèle (appelé plate-forme de Stewart ou main gauche) d'un point de vue algébrique. Nous n'essayons pas ici de fournir une solution en "temps réel" à ce problème, mais plutôt de décrire des outils d'algèbre effective permettant de mieux comprendre la géométrie du problème.

Nous montrons en utilisant des propriétés géométriques simples que le degré de l'application rationnelle associée à ce robot est 40 par un calcul explicite d'un résultant de 4 formes quadratiques en 4 variables et par une analyse de la "ciclicité" des surfaces ainsi obtenues.

Le principal obstacle de cette approche par le calcul formel de problèmes de mécanique est la très grande taille des polynômes manipulés. Nous essayons de convaincre le lecteur qu'une approche utilisant la théorie des invariants permet de "factoriser" les expressions tout en facilitant le suivi des calculs, notamment pour la détermination du lieu critique de l'application. Dans ce rapport sont également inclus quelques exemples d'utilisations pratiques du calcul formel, montrant comment il peut aider à mieux appréhender ces problèmes.

Mots-clé : Robot parallèle, application rationnelle, base de Gröbner, résultant, composante à l'infini, ciclicité, calcul formel

Contents

1	A rational map, its fiber, its critical locus	2
2	Some experimentations	4
2.1	Points in general position	6
2.2	The case where the base and the platform are isometric	9
2.3	Other examples	9
2.3.1	The platform is plane	9
2.3.2	The case where $Y_1 = Y_2, Y_3 = Y_4, Y_5 = Y_6$	10
2.3.3	The case where $Y_4 = Y_5 = Y_6$	10
3	The degree of the map Ψ	11
3.1	A moving tetrahedron	11
3.2	The projection of \mathcal{W}	16
3.3	Intersection with spheres	19
3.4	Yet another proof	20
3.5	The other cases	25
4	The critical locus	25
4.1	The case where $Z_1 = Z_2, Z_3 = Z_4, Z_5 = Z_6$	28
4.2	The case of arbitrary points	32
5	Conclusion	32
	Appendix A	35
	Appendix B	38
	Appendix C	40