

Variational approach of a magnetic shaping problem

Michel Crouzeix

► **To cite this version:**

Michel Crouzeix. Variational approach of a magnetic shaping problem. [Research Report] RR-1352, INRIA. 1990. <inria-00075207>

HAL Id: inria-00075207

<https://hal.inria.fr/inria-00075207>

Submitted on 24 May 2006

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



UNITÉ DE RECHERCHE
INRIA-RENNES

Institut National
de Recherche
en Informatique
et en Automatique

Domaine de Voluceau
Rocquencourt
B.P.105
78153 Le Chesnay Cedex
France
Tél.:(1) 39 63 55 11

Rapports de Recherche

N° 1352

Programme 2
Structures Nouvelles d'Ordinateurs

**VARIATIONAL APPROACH OF A
MAGNETIC SHAPING PROBLEM**

Michel CROUZEIX

Décembre 1990



Campus Universitaire de Beaulieu
35042 - RENNES CEDEX
FRANCE
Téléphone : 99.36.20.00
Télex : UNIRISA 950 473F
Télécopie : 99.38.38.32

**VARIATIONAL APPROACH
OF A MAGNETIC SHAPING PROBLEM**

**APPROCHE VARIATIONNELLE
D'UN PROBLEME DE FORMAGE ELECTROMAGNETIQUE**

Michel Crouzeix (*)

Publication interne n°557
Novembre 1990

14 pages

Programme 2

Résumé : Nous étudions un problème de formage électromagnétique de métal à partir d'une fonctionnelle d'énergie où l'on néglige la tension superficielle. Nous obtenons l'existence d'un potentiel magnétique minimisant cette fonctionnelle, et montrons qu'il est uniformément lipschitzien.

Abstract : For a problem of electromagnetic casting in absence of surface tension, we make a study from the minimization of an energy functional. We prove existence and global Lipschitz-continuity of a magnetic potential which minimizes this functional.

Key words : Free boundary, magnetic casting.

AMS (MOS) subject classification : 35R35,35A15,49A22.

(*) Mathématiques, IRISA, Université de Rennes, Campus de Beaulieu, 35042 RENNES Cedex (France).

1. Modelling.

In order to shape a tube, we have a device that we suppose to be cylindrical, vertical and infinite. Figure 1 represents a horizontal cross section of it. From the outside to the inside we have a fixed mould, then a layer of molten metal, next the vacuum and finally in the central part an inductor where a high frequency alternative current is running.




Figure 1

We assume that the conductivity of the molten metal is infinite. The surface tension on its boundary and more generally all forces in the device not produced by the electromagnetic field are neglected. We suppose we are in a perfectly cylindrical steady state. Moreover we assume that the Laplace forces acting on the inside boundary of the molten metal are strong enough to keep it away from the inductor and that the metal covers completely the wall of the exterior mould.

Let Ω be the (open) part of the horizontal plane which corresponds to the vacuum and the inductor. Its boundary $\partial\Omega$ is the interface between the vacuum and the molten metal. Let Ω^c denote the complementary set of Ω . Since we have a perfectly cylindrical steady state, the magnetic induction field \vec{B} has to be horizontal and the running current vertical. The conductivity is infinite in the molten metal, so the induced current is of pellicular type, (i.e. supported by $\partial\Omega$), and the molten metal realises a perfect shield so that $\vec{B} = 0$ outside $\bar{\Omega}$. From the Maxwell equations we have

$$(1) \quad \operatorname{div} \vec{B} = 0 \quad \text{in } \mathbb{R}^2,$$

$$(2) \quad \operatorname{curl} \vec{B} = \mu_0 j_0 \quad \text{in } \mathbb{R}^2,$$

where μ_0 is the magnetic permeability of the vacuum and j_0 denotes the (vertical component of the) current density.

Writing the equilibrium between the Laplace forces and the magnetic pressure at the boundary $\partial\Omega$ of the molten metal, we get the relation

$$(3) \quad \frac{1}{4\mu_0} |\vec{B}|^2 = \text{constant} \quad \text{on } \partial\Omega,$$

see for example [SNEYD & MOFFAT, 1982].

From equation (1), we can find a magnetic potential φ such that $\vec{B} = \vec{\text{curl}} \varphi$ in \mathbb{R}^2 . As φ is defined up to an arbitrary constant and $\vec{B} = 0$ in Ω^c , we can choose φ such that $\varphi = 0$ in Ω^c . We introduce now the functional

$$J(\varphi) = \frac{1}{2} \int_{\mathbb{R}^2} |\vec{B}|^2 dx - \int_{\mathbb{R}^2} j \varphi dx ,$$

where $j = \mu_0 j_0$ in the inductor, $j = 0$ outside, which represents the total magnetic energy of the device. Since $|\vec{B}| = |\vec{\nabla}\varphi|$, we can write

$$(4) \quad J(\varphi) = \frac{1}{2} \int_{\mathbb{R}^2} |\vec{\nabla}\varphi|^2 dx - \int_{\mathbb{R}^2} j \varphi dx .$$

We assume that j_0 is given in the conductor (i.e. j is given) and that the quantity of molten metal is fixed, which means that the area of the horizontal section corresponding to the molten metal is fixed, or equivalently the area $|\Omega|$ of Ω is equal to a fixed value γ .

In a more general context, [HENROT & PIERRE, 1989] have studied the inverse problem : for a given domain Ω , find a current density j such that (1), (2) and (3) hold ; see also [COULAUD and al.]. The direct problem : for a given current density j and a given positive number γ , find \vec{B} and Ω such that (1), (2), (3) and $|\Omega| = \gamma$ has been studied by many authors, [S & M, 1982], [GAGNOUD & SERO GUILLAUME, 1986], [HENROT and al, 1989], [DESCLOUX, 1990, 1991]. In a more general setting, they start from equations (1), (2), (3) and remark that the solutions satisfy some stationary properties of the energy functional J . In [D, 1990] some stability concepts are introduced and it is shown that the interior shaping problem can have stable solutions. In fact these results give more insight to our approach. We look for a magnetic potential which minimizes J under the constraint $|\Omega| = \gamma$ where we set $\Omega = \{x \in \mathbb{R}^2 ; \varphi(x) \neq 0\}$. Then we get easily an existence theorem for φ and if φ and Ω are smooth enough, we prove that $\vec{B} = \vec{\text{curl}} \varphi$ satisfies (1), (2), (3). The major difficulty we have to deal is the one of the regularity. We have only partially succeeded in our attempt to get such results. We show that φ is uniformly Lipschitz continuous in \mathbb{R}^2 and that Ω is an open set. In fact, at least for γ large enough, our guess is that the boundary $\partial\Omega$ and the function φ in the vacuum up to $\partial\Omega$ are analytic.

2. Existence theorem.

Let K be a compact subset of \mathbb{R}^2 such that its complementary set K^c is a connected set, $j \in L^\infty(\mathbb{R}^2)$ be a function such that

$$(5) \quad \text{support}(j) \subset K,$$

and let γ be a real number such that

$$(6) \quad \gamma > \gamma_0 = \text{area}(K).$$

The Sobolev space $H^m(\mathbb{R}^2)$ denotes the set of functions whose partial derivatives up to the order m are in $L^2(\mathbb{R}^2)$.

For a function v in $H^1(\mathbb{R}^2)$ we set $\Omega_v = \{x \in \mathbb{R}^2; v(x) \neq 0\}$, $\chi_v(x)$ the function defined by $\chi_v(x) = 1$ if $v(x) \neq 0$, $\chi_v(x) = 0$ if $v(x) = 0$, and $|\Omega_v| = \|\chi_v\| = \int_{\mathbb{R}^2} \chi_v(x) dx$. We define the two sets

$$B(\gamma) = \{w \in H^1(\mathbb{R}^2); \|\chi_w\| = \gamma\},$$

$$A(\gamma) = \{w \in H^1(\mathbb{R}^2); \|\chi_w\| \leq \gamma\}.$$

With the functional

$$J(v) = \frac{1}{2} \int_{\mathbb{R}^2} |\vec{\nabla} v|^2 dx - \int_{\mathbb{R}^2} j v dx,$$

we associate the two problems

$$(P_0) \quad \begin{cases} \text{Find } u \in B(\gamma) \text{ such that} \\ \forall v \in B(\gamma), \quad J(u) \leq J(v) \end{cases}$$

and

$$(P) \quad \begin{cases} \text{Find } u \in A(\gamma) \text{ such that} \\ \forall v \in A(\gamma), \quad J(u) \leq J(v). \end{cases}$$

From our modelling, we are concerned in the problem (P_0) but, from a mathematical point of view, the problem (P) is easier to study. Fortunately we have the relation

Lemma 1. *We assume that the situation described in (7) does not hold*

(7) *there exists $u \in H^1(\mathbb{R}^2)$ with $-\Delta u = j$ in \mathbb{R}^2 and $\|\chi_u\| < +\infty$;*
then the problems (P_0) and (P) are equivalent.

Proof. From the density of $B(\gamma)$ in $A(\gamma)$, u solution of (P_0) implies u solution of (P) . Conversely let u be a solution of (P) and $\gamma' = \|\chi_u\|$. If u were not a solution of (P_0) , we should have $\gamma' < \gamma$ and then

$$\forall \varphi \in H^1(\mathbb{R}^2) \text{ with } \|\chi_\varphi\| \leq \gamma - \gamma' , \quad J(u) \leq J(u+\varphi)$$

whence
$$\int_{\mathbb{R}^2} \vec{\nabla} u \cdot \vec{\nabla} \varphi \, dx = \int_{\mathbb{R}^2} j \varphi \, dx ,$$

so that we can conclude that (7) holds. ■

Remarks. 1) When the situation (7) occurs we have, $\forall v \in H^1(\mathbb{R}^2)$, $J(u) \leq J(v)$, moreover u is harmonic in K^c . With $\|\chi_u\| < +\infty$ that induces $u = 0$ in K^c and therefore $|\Omega_u| \leq \gamma_0$. For the physical model such a j is not suitable since we must have no contact between molten metal and the inductor.

2) When $\int_{\mathbb{R}^2} j \, dx \neq 0$, the situation (7) cannot occur.

Theorem 2. *Problem (P) admits at least one solution.*

Proof. We first remark that, from the Schwarz symmetrization principle, we have [SPERNER, 1974]

$$\forall v \in A(\gamma) , \quad \|v\|_{L^2(\mathbb{R}^2)} \leq C \gamma^{1/2} \|\vec{\nabla} v\|_{L^2(\mathbb{R}^2)} ,$$

therefore $J(v) \geq -\frac{1}{2} C^2 \gamma \|j\|_{L^2(\mathbb{R}^2)}$, and $\inf_{v \in A(\gamma)} J(v) > -\infty$. On the other hand, $A(\gamma)$ is weakly closed in $H^1(\mathbb{R}^2)$. Indeed if $v_n \in A(\gamma) \rightarrow v$ weakly in $H^1(\mathbb{R}^2)$, eventually using a subsequence, we can assume $v_n \rightarrow v$ a.e. in \mathbb{R}^2 . Then let F be a compact subset of \mathbb{R}^2 , since $\|\chi_{v_n}\| \leq \gamma$, we have

$$\int_F \frac{|v|}{\sqrt{v^2 + \varepsilon^2}} \, dx = \lim_{n \rightarrow \infty} \int_F \frac{|v_n|}{\sqrt{v_n^2 + \varepsilon^2}} \, dx \leq \gamma .$$

Therefore, taking the limit as $\varepsilon \rightarrow 0$, $\int_F \chi_v \, dx \leq \gamma$. Since this inequality holds for all compact F in \mathbb{R}^2 , we deduce $v \in A(\gamma)$. Then the theorem follows from the weak lower semi-continuity of J . ■

Remark. A similar existence result may be found in [PIRONNEAU, 1984].

Hereafter u denotes a solution of (P) . We shall see in theorem 3 that u is continuous, so that $\Omega = \Omega_u = \{x ; u(x) \neq 0\}$ is an open set. Since for all φ supported by Ω , we have $J(u) \leq J(u+\varphi)$, so

$$(8) \quad -\Delta u = j \quad \text{in } \Omega ;$$

and by the definition $u = 0$ in Ω^c .

3. A regularity result.

Let us introduce now the notations

$$(9) \quad \partial v = \frac{1}{2} \left(\frac{\partial v}{\partial x_1} - i \frac{\partial v}{\partial x_2} \right), \quad \bar{\partial} v = \frac{1}{2} \left(\frac{\partial v}{\partial x_1} + i \frac{\partial v}{\partial x_2} \right), \quad y = y_1 + i y_2,$$

(but $dy = dy_1 dy_2$ is the Lebesgue measure in \mathbb{R}^2). Let $\mathcal{D}(\mathbb{R}^2)$ denote the set of compact supported functions which are infinitely differentiable in \mathbb{R}^2 . When there is no ambiguity, we shall write

$$\int - \text{ and } \|\cdot\|_{L^p} \text{ in place of } \int_{\mathbb{R}^2} - \text{ and } \|\cdot\|_{L^p(\mathbb{R}^2)}.$$

The main result of this section is

Theorem 3. *Let u be a solution of (P). Then there exists a positive number λ such that*

$$(10) \quad \forall \varphi \in \mathcal{D}(\mathbb{R}^2), \quad 4 \int (\partial u)^2 (\bar{\partial} \varphi) dx = 2 \int j \partial u \varphi dx + \lambda \int \chi_u \partial \varphi dx.$$

Moreover we have

$$(11) \quad (\partial u)^2(x) = -\frac{1}{2\pi} \int (j \partial u)_{(x-y)} \frac{1}{y} dy - \frac{\lambda}{4\pi} \int \chi_{u(x-y)} \frac{1}{y^2} dy,$$

and the function u is uniformly Lipschitz continuous in \mathbb{R}^2 .

Proof. a) Let $\vec{\theta} \in (C^\infty(\mathbb{R}^2))^2$ be such that its derivative $D\vec{\theta} \in (L^\infty(\mathbb{R}^2))^4$ and $\int \chi_u \operatorname{div} \vec{\theta} dx > 0$. For $\delta > 0$, we consider $F_\delta(x) = x + \delta \theta(x)$ and $u_\delta = u \circ F_\delta$. Clearly if δ is small enough, F_δ is a C^∞ diffeomorphism, furthermore $u_\delta \in H^1(\mathbb{R}^2)$ and

$$\int \chi_{u_\delta}(x) dx = \int \chi_{u(y)} dF_\delta^{-1}(y) = \int \chi_{u(y)} dy - \delta \int \chi_{u(y)} \operatorname{div} \vec{\theta}(y) dy + O(\delta^2),$$

so that for $\delta > 0$ small enough, $u_\delta \in A(\gamma)$. Writing $\frac{1}{\delta} (J(u_\delta) - J(u)) \geq 0$ and taking the limit as $\delta \rightarrow 0_+$ we obtain

$$(12) \quad \int \{D\vec{\theta}(\vec{\nabla} u, \vec{\nabla} u) - \frac{1}{2} \operatorname{div} \vec{\theta} \vec{\nabla} u^2\} dx - \int j \vec{\nabla} u \cdot \vec{\theta} dx \geq 0.$$

We consider now a function $\vec{\theta}$ such that $\int \chi_u \operatorname{div} \vec{\theta} dx = 0$. We still obtain that (12) holds by using $\vec{\theta}_\varepsilon(x) = \theta(x) + \varepsilon x$ and then taking the limit as $\varepsilon \rightarrow 0_+$, but now changing $\vec{\theta}$ in $-\vec{\theta}$, we have the equality in (12).

For an arbitrary function $\vec{\theta}$, we set $\vec{\theta}_1 = \vec{\theta} - \frac{1}{2\gamma} \int \chi_u \operatorname{div} \vec{\theta} dx \vec{x}$, so that $\int \chi_u \operatorname{div} \vec{\theta}_1 dx = 0$. The equality in (12) for $\vec{\theta}_1$ leads

$$(13) \quad \int \{D \vec{\theta}(\vec{\nabla}u, \vec{\nabla}u) - \frac{1}{2} \operatorname{div} \vec{\theta} \vec{\nabla}u^2\} dx - \int j \vec{\nabla}u \cdot \vec{\theta} dx = \lambda \int \chi_u \operatorname{div} \vec{\theta} dx,$$

with

$$(14) \quad \lambda = -\frac{1}{\gamma} \int j \vec{\nabla}u \cdot \vec{x} dx.$$

We remark that λ is non negative when using (12) with $\vec{\theta} = \vec{x}$.

b) Equation (10) is exactly (13) rewritten with the notations (9) and $\varphi = \theta_1 - i \theta_2$. Using that $\int \bar{\partial}\varphi(y) (\bar{y})^{-1} dy = -\pi \varphi(0)$ and $\int \bar{\partial}\varphi(y) (\bar{y})^{-2} dy = -\pi \partial\varphi(0)$, and setting

$$(15) \quad a(x) = -\frac{1}{2\pi} \int (j \partial u)_{(x-y)} \frac{1}{\bar{y}} dy - \frac{\lambda}{4\pi} \int \chi_u(x-y) \frac{1}{\bar{y}^2} dy,$$

we deduce from (10) that

$$\forall \varphi \in \mathcal{D}(\mathbb{R}^2), \quad \int ((\partial u)^2 - a) \bar{\partial}\varphi dx = 0.$$

Therefore $(\partial u)^2 - a$ is harmonic in \mathbb{R}^2 .

We know that $u \in H^1(\mathbb{R}^2)$, thus $(\partial u)^2 \in L^1(\mathbb{R}^2)$. Since, $j \partial u \in L^1(\mathbb{R}^2) \cap L^2(\mathbb{R}^2)$ and $y \rightarrow (\bar{y})^{-1} \in L^p(\mathbb{R}^2) + L^q(\mathbb{R}^2)$ for all $1 \leq q < 2$ and $2 < p \leq +\infty$, the function

$$(16) \quad s(x) = -\frac{1}{2\pi} \int (j \partial u)_{(x-y)} (\bar{y})^{-1} dy$$

satisfies $s \in L^p$ for all $2 < p < +\infty$. We introduce now the Beurling transform T given by

$$(17) \quad Tg(x) = \frac{1}{4\pi} \int g(x-y) (\bar{y})^{-2} dy,$$

it acts from $L^p(\mathbb{R}^2)$ into $L^p(\mathbb{R}^2)$ for all $1 < p < +\infty$, (see for instance [ZYGmund, 1971]). As $\chi_u \in L^p(\mathbb{R}^2)$ for all p and $a = s - \lambda T \chi_u$, we have $a \in L^p(\mathbb{R}^2)$ for all $2 < p < +\infty$. Since the harmonic function $(\partial u)^2 - a$ belongs to $L^1(\mathbb{R}^2) + L^p(\mathbb{R}^2)$, we have $(\partial u)^2 - a = 0$ in \mathbb{R}^2 , therefore

$$(\partial u)^2 = a \in L^p(\mathbb{R}^2) \text{ for all } 2 < p < +\infty$$

and (11) is proven.

c) Now we have $j \partial u \in L^1(\mathbb{R}^2) \cap L^p(\mathbb{R}^2)$ for all $1 \leq p < +\infty$, therefore $s \in C^0(\mathbb{R}^2) \cap L^\infty(\mathbb{R}^2)$.

Let us introduce

$$(18) \quad T_\varepsilon(x) = \frac{1}{4\pi} \int_{|y| \geq \varepsilon} \chi_u(x-y) (\bar{y})^{-2} dy .$$

We admit for a while the result

Lemma 4. *There exists a constant C such that*

$$(19) \quad \forall x \in \mathbb{R}^2, \forall \varepsilon > 0, |T_\varepsilon(x)| \leq C + \|T\chi_u\|_{L^\infty(\Omega^c)} .$$

Since $\lim_{\varepsilon \rightarrow 0} T_\varepsilon = T\chi_u$ a.e. in \mathbb{R}^2 and $\partial u = 0$ in Ω^c , we have $\lambda T\chi_u = s$ in Ω^c and $\lambda T\chi_u \in L^\infty(\mathbb{R}^2)$. We deduce from (11) that $\partial u \in L^\infty(\mathbb{R}^2)$ and

$$\|\partial u\|_{L^\infty(\mathbb{R}^2)}^2 \leq 2 \|s\|_{L^\infty(\mathbb{R}^2)} + C \lambda .$$

From (14) we get $\lambda \leq \gamma^{-1} \|j \vec{\nabla} u \cdot \vec{x}\|_{L^1} \leq \|jx\|_{L^1} \|\partial u\|_{L^\infty}$

and from (16) $\|s\|_{L^\infty} \leq \frac{1}{\sqrt{2\pi}} (\|j\|_{L^\infty} + \|j\|_{L^1}) \|\partial u\|_{L^\infty}$.

Therefore we have

$$(20) \quad \|\partial u\|_{L^\infty(\mathbb{R}^2)} \leq \frac{1}{\sqrt{2\pi}} (\|j\|_{L^1} + \|j\|_{L^\infty}) + C \|jx\|_{L^\infty} .$$

Note that this estimate does not depend on γ .

d) To be complete, we have to prove Lemma 4. Without loss of generality we can suppose $x = 0$. Let $\varphi(\varepsilon) = T_\varepsilon(0)$, $D_\varepsilon = \{x \in \mathbb{R}^2; \frac{\varepsilon}{2} < |x| \leq \varepsilon\}$, $B_\varepsilon = \{x \in \mathbb{R}^2; |x| < \varepsilon\}$, $\alpha(\varepsilon) = \frac{\text{meas}(D_\varepsilon \cap \Omega^c)}{\text{meas}(D_\varepsilon)}$ and $C_0 = \|T\chi_u\|_{L^\infty(\Omega^c)}$. We have

$$\varphi(\varepsilon) - \varphi\left(\frac{\varepsilon}{2}\right) = \frac{1}{4\pi} \int_{D_\varepsilon} (1 - \chi_u(y)) (\bar{y})^{-2} dy = \frac{1}{4\pi} \int_{D_\varepsilon \cap \Omega^c} (\bar{y})^{-2} dy,$$

thus

$$(21) \quad |\varphi(\varepsilon) - \varphi\left(\frac{\varepsilon}{2}\right)| \leq \frac{3}{4} \alpha(\varepsilon).$$

On the other hand we set

$$f_1(x) = \chi_u(x) - 1, \quad f_0(x) = 1, \quad f_2(x) = 0, \quad \text{if } |x| \leq \varepsilon,$$

$$\text{and } f_1(x) = 0, \quad f_0(x) = 0, \quad f_2(x) = \chi_u(x), \quad \text{if } |x| > \varepsilon,$$

$$\text{so that } \varphi(\varepsilon) = (Tf_2)(0), \quad \chi_u = f_1 + f_2 + f_0.$$

$$\text{We can write } \varphi(\varepsilon) = T\chi_u(x) - Tf_1(x) - Tf_0(x) + (Tf_2(0) - Tf_2(x)).$$

We have $Tf_0(x) = 0$ for $|x| < \varepsilon$ (and $= -\varepsilon^2/(4\bar{x}^2)$ for $|x| > \varepsilon$), some easy computations give the (non optimal) estimate,

$$|Tf_2(0) - Tf_2(x)| \leq 2, \quad \text{for } |x| \leq \frac{\varepsilon}{2},$$

therefore we have

$$|\varphi(\varepsilon)| \leq C_0 + |Tf_1(x)| + 2, \quad \text{a.e. in } B_{\varepsilon/2} \cap \Omega^c.$$

From the continuity of T from $L^1(\mathbb{R}^2)$ into weak $L^1(\mathbb{R}^2)$, see for instance [COIFMAN & MEYER, 1978], there exists C_1 such that

$$\forall \mu > 0, \quad \mu \text{ meas}\{x \in \mathbb{R}^2, |Tf_1(x)| \geq \mu\} \leq C_1 \|f_1\|_{L^1} = C_1 \text{ meas}(B_\varepsilon \cap \Omega^c).$$

With the choice

$$\mu = |\varphi(\varepsilon)| - C_0 - 2, \quad (\text{when such a } \mu \text{ is } > 0),$$

that gives $(|\varphi(\varepsilon)| - C_0 - 2) \text{ meas}(B_{\varepsilon/2} \cap \Omega^c) \leq C_1 \text{ meas}(B_\varepsilon \cap \Omega^c)$,

$$\text{whence } |\varphi(\varepsilon)| \leq C_0 + 2 + C_1 \frac{\text{meas}(B_\varepsilon \cap \Omega^c)}{\text{meas}(B_{\varepsilon/2} \cap \Omega^c)} \leq C_0 + 2 + C_1 + C_1 \frac{\text{meas}(D_\varepsilon \cap \Omega^c)}{\text{meas}(D_{\varepsilon/2} \cap \Omega^c)},$$

therefore we have, (if $\alpha(\varepsilon/2) \neq 0$),

$$(22) \quad |\varphi(\varepsilon)| \leq C_0 + 2 + C_1 + 4C_1 \frac{\alpha(\varepsilon)}{\alpha(\varepsilon/2)}.$$

If $\alpha(\varepsilon) < 2\alpha(\varepsilon/2)$ then we have $|\varphi(\varepsilon)| \leq C_0 + 9C_1 + 2$. Otherwise there exists an integer k such that

$$\alpha(2^j \varepsilon) \leq 2^{-1} \alpha(2^{j+1} \varepsilon), \quad \text{for } j = 1, \dots, k-1 \quad \text{and} \quad \alpha(2^k \varepsilon) > 2^{-1} \alpha(2^{k+1} \varepsilon).$$

Then we have $|\varphi(2^{k+1} \varepsilon)| \leq C_0 + 9C_1 + 2$ and

$$\alpha(2^j \varepsilon) \leq 2^{j-k} \alpha(2^k \varepsilon) \leq 2^{j-k} \quad \text{for } j = 1, \dots, k.$$

From (21) we get

$$|\varphi(\varepsilon) - \varphi(\frac{\varepsilon}{2})| \leq \frac{3}{4} (\alpha(2^{k+1} \varepsilon) + \alpha(2^k \varepsilon) + \dots + \alpha(2\varepsilon)) \leq 9/4$$

so that $|\varphi(\varepsilon)| \leq C_0 + 9C_1 + 17/4$,

and (19) holds with $C = 9C_1 + 17/4$. ■

Remark. We cannot have $\lambda = 0$ in (11). Indeed $\lambda = 0$ would imply $(\partial u)^2$ is harmonic in K^c , so that $\partial u = 0$ in K^c whence $u = 0$ in K^c , which would be inconsistent with (6).

4. Conclusions.

We have got that the solution u of our minimization problem satisfies $-\Delta u = j$ in Ω . If the smoothness of Ω and u allows us to use the Green formula, we deduce from (10) that $4(\partial u)^2 = \lambda(v_1 + iv_2)^2$ on $\partial\Omega$, which means $\vec{\nabla} u = \pm \sqrt{\lambda} \vec{\nu}$; (here $\vec{\nu}$ is the unit vector normal to $\partial\Omega$). So that we get the equilibrium condition (3). With some stronger smoothness assumptions, it follows from the work of [H & P, 1989], see also [D, 1990], that u and $\partial\Omega$ have some analytic properties.

We assume now that j is fixed and satisfies $\int_{\Omega} j \, dx \neq 0$ (say $\int_{\Omega} j \, dx > 0$), that is the total current running in the inductor does not vanish. Let u_{γ} be the solution of (P), here γ appears to emphasize the dependence on γ , it is easy to show that $\lim_{\gamma \rightarrow \infty} J(u_{\gamma}) = -\infty$; we have $J(u_{\gamma}) \geq -\int_{\mathbb{R}^2} j u_{\gamma} \, dx = -\int_K j u_{\gamma} \, dx$, and from (20), $\forall x, y \in \mathbb{R}^2, |u_{\gamma}(x) - u_{\gamma}(y)| \leq c|x-y|$. This implies $\lim_{\gamma \rightarrow \infty} \min_{x \in K} \{u_{\gamma}(x)\} = +\infty$, therefore, for γ large enough, the set K is included in Ω which means for γ large enough the molten metal has no contact with the inductor. We can also show that u_{γ} is non negative in \mathbb{R}^2 .

We have proven that u is Lipschitz continuous, that gives very poor information on the smoothness on $\partial\Omega$. Note that the boundedness of ∂u implies by the lemma 4 that $|T_{\varepsilon}(x)|$ is uniformly bounded by a constant. Using the definition of T_{ε} and polar coordinates, we can easily see that $\partial\Omega$ cannot have a singularity of corner type, (except if the angle is $0, \pi$ or 2π , a cusp point is allowed).

5. References.

- COIFMAN R., MEYER Y. 1978. Au-delà des opérateurs pseudodifférentiels. *Astéristique*, 57, 76-115.
- COULAUD O., HENROT A., PIERRE M., ROCHE J.R., 1990. Numerical simulation of electromagnetic shaping. Preprint.
- DESCLOUX J., 1990. On the two dimensional magnetic shaping problem without surface tension. Report 07.90, EPFL, Dept. Math., Lausanne.
- DESCLOUX J., 1991. Stability of the solutions of the bidimensional magnetic shaping problem in absence of surface tension. Preprint.
- GAGNOUD A., SERO GUILLAUME O., 1986. Le creuset froid de lévitation : modélisation électromagnétique et application. *E.D.F Bull. Etudes et Rech.*, B, 1, 41-51.
- HENROT A., BRANCHER J.P., PIERRE M., 1989. Existence of equilibria in electromagnetic casting. *Proc. Int. Symp. Num. Mech. Eng.* 5, vol. 1, Gruber, Périaux, Shaw editions. *Computational Mechanics Publications*. Springer-Verlag, 221-230.
- HENROT A., PIERRE M., 1989. Un problème inverse en formage des métaux liquides. *M²AN*, 23, 155-177.
- PIRONNEAU O., 1984. *Optimal design for elliptic systems*. Springer-Verlag, New-York.
- SNEYD A., MOFFAT H., 1982. Fluid dynamical aspects of the levitation melting process. *J. Fluid Mech*, 117, 45-70.
- SPERNER E., 1974. Symmetrisierung für Funktionen mehrerer Variablen, *Man. Math.*, 11, 159-170.
- ZYGMUND A., 1971. *Intégrales singulières*. *Lectures notes in Math*, 204, Springer-Verlag.

LISTE DES DERNIERES PUBLICATIONS INTERNES IRISA

- PI 548 **LES PREDICATS COLLECTIFS : UN MOYEN D'EXPRESSION DU
CONTROLE DU PARALLELISME "OU" EN PROLOG**
René QUINIOU, Laurent TRILLING
Septembre 1990, 34 Pages.
- PI 549 **NORMALISATION SOUS HYPOTHESE D'ABSENCE DE LIEN
APPLICATION AU CAS NOMINAL**
François DAUDE
Septembre 1990, 42 Pages.
- PI 550 **MULTISCALE SIGNAL PROCESSING : FROM QMF TO WAVELETS**
Albert BENVENISTE
Septembre 1990, 28 Pages.
- PI 551 **ON THE TRANSITION GRAPHS OF AUTOMATA AND GRAMMARS**
Didier CAUCAL, Roland MONFORT
Septembre 1990, 46 Pages.
- PI 552 **ERREURS DE CALCUL DES ORDINATEURS**
Jocelyne ERHEL
Septembre 1990, 58 Pages.
- PI 553 **SEQUENTIAL FUNCTIONS**
Boubakar GAMATIE, Octobre 1990, 16 Pages.
- PI 554 **ANALYSE DE LA FORME D'UN COEFFICIENT D'ASSOCIATION
ENTRE VARIABLES QUALITATIVES**
Mohamed OUALI ALLAH
Octobre 1990, 26 Pages.
- PI 555 **APPROXIMATION BY NONLINEAR WAVELET NETWORKS**
Qinghua ZHANG, Albert BENVENISTE
Octobre 1990, 16 Pages.
- PI 556 **CONCEPTION ET INTEGRATION D'UN CORRELATEUR SYSTOLIQUE**
Catherine DEZAN, Eric GAUTRIN, Patrice QUINTON
Novembre 1990, 16 Pages.
- PI 557 **VARIATIONAL APPROACH OF A MAGNETIC SHAPING PROBLEM**
Michel CROUZEIX
Novembre 1990, 14 Pages.

ISSN 0249 - 6399