

Random walks in a quarter plane with zero drifts. I: ergodicity and null recurrence

Guy Fayolle, Vadim A. Malyshev, M.V. Menshikov

► **To cite this version:**

Guy Fayolle, Vadim A. Malyshev, M.V. Menshikov. Random walks in a quarter plane with zero drifts. I: ergodicity and null recurrence. RR-1314, INRIA. 1990. <inria-00075245>

HAL Id: inria-00075245

<https://hal.inria.fr/inria-00075245>

Submitted on 24 May 2006

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

INRIA

UNITÉ DE RECHERCHE
INRIA-ROCQUENCOURT

Institut National
de Recherche
en Informatique
et en Automatique

Domaine de Voluceau
Rocquencourt
B.P.105
78153 Le Chesnay Cedex
France
Tél.:(1) 39 63 55 11

Rapports de Recherche

N° 1314

Programme 3
Réseaux et Systèmes Répartis


RANDOM WALKS IN A QUARTER PLANE WITH ZERO DRIFTS. I : ERGODICITY AND NULL RECURRENCE

Guy FAYOLLE
Vadim A. MALYSHEV
Michael V. MENSHIKOV

Octobre 1990



★ R R - 1 3 1 4 ★



Random walks in a quarter plane with zero drifts. I : Ergodicity and null recurrence


G. Fayolle*, V.A. Malyshev†, M.V. Menshikov†

Abstract

In this paper, we solve the problem of non ergodicity and null recurrence for random walks in the quarter plane with zero drifts in the interior of the domain. A general criterion for null recurrence is given and then used to construct sub and supermartingales by means of Lyapounov functions, which are here functionals of quadratic forms.

*INRIA - Domaine de Voluceau, Rocquencourt - BP.105 - 78153 Le Chesnay Cedex - FRANCE.

†Moscow State University, Mechanico-Mathematical Faculty, Chair of probability, Leninskie Gori, 119899 MOSCOW - USSR



Marches aléatoires dans un quart de plan avec dérivées nulles I : Ergodicité et recurrence nulle

Guy Fayolle*, Vadim A. Malyshev†, Michael V. Menshikov†

Résumé

Dans cet article nous résolvons le problème de non ergodicité et de recurrence nulle pour des marches aléatoires dans le quart de plan, lorsque les dérivées moyennes sont nulles à l'intérieur de cette région. Un critère général de récurrence nulle est donné et utilisé pour construire des (sous) surmartingales par l'intermédiaire de fonctions de Lyapounov, qui sont ici en général des fonctionnelles de formes quadratiques.

*INRIA - Domaine de Voluceau, Rocquencourt - BP.105 - 78153 Le Chesnay Cedex - FRANCE.

†Moscow State University, Mechanico-Mathematical Faculty, Chair of probability, Leninskie Gori, 119899 MOSCOW - USSR

1 Introduction and preliminaries

We consider a discrete time homogeneous irreducible and aperiodic Markov chain $\mathcal{L} = \{\xi_n, n \geq 0\}$, with state space the lattice in the positive quarter plane $\mathbf{Z}_+^2 = \{(i, j) : i, j \geq 0 \text{ are integers}\}$, and satisfying the recursive equation

$$\xi_{n+1} = [\xi_n + \theta_{n+1}]^+,$$

where the distribution of θ_{n+1} depends only on the position of ξ_n in the following way (“maximal “ space homogeneity)

$$p\{\theta_{n+1} = (i, j) / \xi_n = (k, l)\} = \begin{cases} p_{ij}, & \text{for } k, l \geq 1, \\ p'_{ij}, & \text{for } k \geq 1, l = 0, \\ p''_{ij}, & \text{for } k = 0, l \geq 1, \\ p^0_{ij}, & \text{for } k = l = 0. \end{cases}$$

Moreover we assume, for the one step transition probabilities, the following conditions :

A (*Lower boundedness*)

$$\begin{aligned} p_{ij} &= 0, & \text{if } i < -1 & \text{ or } j < -1 ; \\ p'_{ij} &= 0, & \text{if } i < -1 & \text{ or } j < 0 ; \\ p''_{ij} &= 0, & \text{if } i < 0 & \text{ or } j < -1 . \end{aligned}$$

B (*Moment condition*)

$$E[\|\theta_{n+1}\|^3 / \xi_n = (k, l)] \leq B < \infty, \quad \forall (k, l) \in \mathbf{Z}_+^2,$$

where $\|z\|$, $z \in \mathbf{Z}_+^2$, denotes the euclidian norm and ϵ is an arbitrary but strictly positive number. In fact, as remarked later in section 3, only the existence of second moments is necessary. But the technical derivations are then more involved and will not be given here for the sake of clarity.

Notation : We shall use lower case greek letters α, β, \dots to denote arbitrary, points of \mathbf{Z}_+^2 , and then $p_{\alpha\beta}$ will mean the one step transition probabilities of the Markov \mathcal{L} . Also $\alpha > 0$ is equivalent to

$$\alpha_x > 0, \alpha_y > 0 \text{ for } \alpha = (\alpha_x, \alpha_y).$$

Define the vector

$$M(\alpha) = (M_x(\alpha), M_y(\alpha))$$

of the one step mean jumps (drifts) from the point α .

Setting

$$\alpha = (\alpha_x, \alpha_y), \quad \beta = (\beta_x, \beta_y),$$

we have

$$M_x(\alpha) = \sum_{\beta} p_{\alpha\beta}(\beta_x - \alpha_x),$$

$$M_y(\alpha) = \sum_{\beta} p_{\alpha\beta}(\beta_y - \alpha_y).$$

Condition B ensures the existence of $M(\alpha)$, for all $\alpha \in \mathbf{Z}_+^2$. By the homogeneity condition A , only 4 different drift vectors take place :

$$M(\alpha) = \begin{cases} M, & \text{for } \alpha = (\alpha_x, \alpha_y) > 0, \\ M', & \text{for } \alpha = (\alpha_x, 0), \alpha_x > 0; \\ M'', & \text{for } \alpha = (0, \alpha_y), \alpha_y > 0; \\ M_0, & \text{for } \alpha = (0, 0). \end{cases}$$

Remark :

i) all our results will be valid if one changes arbitrarily a finite number of transition probabilities.

ii) given $\xi_n = \alpha$, the components of θ_{n+1} might be taken bounded from below not by -1 , but by some arbitrary number $-K > -\infty$ provided

- first, that we keep the maximal homogeneity for the drift vectors $M(\alpha)$ introduced above (i.e. 4 of them only are different) ;
- secondly, that the second moments and the covariance of the one step jumps inside \mathbf{Z}_+^2 (i.e. from any point $\alpha > 0$) are maintained constant. These last facts will emerge more clearly in the course of the study.

The classification problem for these random walks was studied first in [1] and, in the case of bounded jumps, completely solved except for the case $M = 0$. This was generalized later, in many papers, for unbounded jumps [3, 4, 5, 11].

Nothing was precisely known for the case $M = 0$, up to now, and, in fact, this problem in many aspects is of a very different nature. In particular, intuition does not give us the evidence that the random walk can exhibit an ergodic behaviour when $M = 0$.

To the author's knowledge there exist at least 3 methods which would allow to solve some particular cases of the problem :

1. Analytic approach [8]. Now there are only some preliminary results into this direction.
2. Semi-analytic approach, using well-known explicit results about one-dimensional random walks. E.g. when the random walks inside the quarter-plane is the "composition" of independent random walks along both axis, then it is easy to prove that the mean time to reach the boundary is infinite and thus, for any parameters p'_{ij}, p''_{ij} , the chain is not ergodic when $M = 0$.
3. Method of Lyapounov functions. It seems to be the most general approach and we use it here.

One of the main differences between the cases $M \neq 0$ and $M = 0$ is the following : the case $M \neq 0$ is in a sense locally linear and $M = 0$ is locally quadratic. The local second order effects are well caught by quadratic Lyapounov functions first introduced in [5]. But the global Lyapounov function is not quadratic and this causes additional difficulties. There are some general questions about constructive criteria of ergodicity, nonergodicity etc. We tried unsuccessfully to use the existing criteria for non ergodicity. They all seem to work only either for 1-dimensional problems or for linear functions. One of our difficulties was to find a completely new one. We give it in section 2. We do not know whether it could be essentially generalized, but it allows to get the complete solution in our case.

For $M = 0$, we get the ergodicity conditions in terms of the second moments and the covariance of the one-step jumps inside \mathbf{Z}_+^2 ,

$$\lambda_x = \sum_{i,j} i^2 p_{ij} , \quad \lambda_y = \sum_{i,j} j^2 p_{ij} ,$$

$$R = \sum_{i,j} ij p_{ij} ,$$

and of the angles (counter clockwise oriented) ϕ_x, ϕ_y (Sec Figure 1).




Figure 1

Here ϕ_x is the angle between M' and the negative x-axis, ϕ_y is the angle between M'' and negative y-axis. So, if $\phi_x \neq \frac{\pi}{2}$ and $\phi_y \neq \frac{\pi}{2}$, then

$$\operatorname{tg} \phi_x = -\frac{M'_y}{M'_x} , \quad \operatorname{tg} \phi_y = -\frac{M''_x}{M''_y} .$$

Theorem 1.1 *If at least one of the following 4 conditions holds*

- (i) $\phi_x > \frac{\pi}{2}$,
- (ii) $\phi_y > \frac{\pi}{2}$,
- (iii) $\phi_x = \frac{\pi}{2}$, $\phi_y \neq 0$,
- (iv) $\phi_y = \frac{\pi}{2}$, $\phi_x \neq 0$,

then the random walk \mathcal{L} is not ergodic.

If $\phi_x, \phi_y < \frac{\pi}{2}$, then the random walk \mathcal{L} is ergodic if

$$\lambda_x \operatorname{tg} \phi_x + \lambda_y \operatorname{tg} \phi_y + 2R \equiv -\lambda_x \frac{M'_y}{M'_x} - \lambda_y \frac{M''_x}{M''_y} + 2R < 0 . \quad (1.1)$$

and non ergodic if

$$\lambda_x \operatorname{tg} \phi_x + \lambda_y \operatorname{tg} \phi_y + 2R > 0 . \quad (1.2)$$

Moreover if (1.2) holds together with $\phi_x + \phi_y < \frac{\pi}{2}$, then the random walk is null recurrent.

Remark 1 : It follows easily from the statement of the theorem that the mean first entrance time of \mathcal{L} into the boundary, when starting from some arbitrary point $\alpha > 0$ at finite distance, is finite [resp. infinite] if $R < 0$ [resp. $R > 0$], since in this case the vectors M' and M'' can be properly chosen to satisfy (1.1) [resp. (1.2)].

Remark 2 : It is clear from the formulation of the theorem that we do not consider the cases when either $\phi_x = \frac{\pi}{2}, \phi_y = 0$ or $\phi_y = \frac{\pi}{2}, \phi_x = 0$. Also, we miss the case

$$\lambda_x \operatorname{tg} \phi_x + \lambda_y \operatorname{tg} \phi_y + 2R = 0 .$$

The proof will be given in section 3.

2 General criteria for the null recurrence of countable Markov chains

Let us consider a discrete time homogeneous Markov chain \mathcal{L} with a countable state space $X = \{x_n\}, n = 1, 2, \dots$. The one step transition probabilities are denoted by q_{x_i, x_j} . \mathcal{L} is assumed to be irreducible and aperiodic. Let ξ_n be the state of \mathcal{L} at time n .

In this section, we recall some known criteria for recurrence, ergodicity and non ergodicity which will be used in the sequel. Meanwhile, theorem 2.5 is completely new and enables us to get explicit results in section 2.

Theorem 2.1 *For the irreducible Markov chain \mathcal{L} to be recurrent, it is sufficient there exists a positive function $f(x), x \in X$, such that $E[f(\xi_{m+1}) - f(\xi_m) | \xi_m = x_i] \leq 0, i \notin A, f(x_i) \rightarrow \infty$ when $i \rightarrow \infty$, and A is a finite set.*

Proof : see [7] ■

Theorem 2.2 (Foster). *An irreducible aperiodic Markov chain \mathcal{L} is ergodic if and only if there exists a positive function $f(x), x \in X, \epsilon > 0$ and a finite set A such that*

$$E(f(\xi_{m+1}) - f(\xi_m) / \xi_m = x_i) \leq -\epsilon, i \notin A,$$

$$E(f(\xi_{m+1}) / \xi_m = x_i) < \infty, i \in A.$$

Proof : See [10]. For non trivial generalisations of this criterion, see [2].

Theorem 2.3 *For an irreducible Markov chain \mathcal{L} to be non ergodic, it is sufficient that there exist a function $f(x), x \in X$, and constants c, m such that :*

1. $E(f(\xi_{n+1}) - f(\xi_n) / \xi_n = x) \geq 0$, for every n , all $x \in \{x : f(x) > c\}$, where the sets $\{x : f(x) > c\}$ and $\{x : f(x) \leq c\}$ are non empty.
2. $E(|f(\xi_{n+1}) - f(\xi_n)| / \xi_n = x) \leq m$ for every $n, x \in X$.

Under the stronger condition

$$|f(\xi_{n+1}) - f(\xi_n)| \leq \mathcal{D},$$

with probability 1, for some fixed constant $\mathcal{D}, 0 < \mathcal{D} < \infty$, this theorem was first proved in [9].

Proof : In its present form, it was apparently given in [12], by using explicitly the Markov context. It seems interesting here to point out that it is in fact a direct consequence of the following general

Theorem 2.4 *Let us consider an arbitrary sequence of random variables $(S_n), n \geq 0$. Let \mathcal{F}_n the σ -algebra generated by S_0, S_1, \dots, S_n , where S_0 will be taken constant with probability one (this does not restrict the generality). Let τ be the following \mathcal{F}_n -stopping time, C being a real constant,*

$$\tau = \inf \{n > 0, S_n < C/S_0 \geq C\}.$$

Introduce the stopped sequence

$$\tilde{S}_n = S_{n \wedge \tau} ,$$

where

$$n \wedge \tau = \begin{cases} n, & \text{if } n \leq \tau \\ \tau, & \text{if } n > \tau, \end{cases}$$

Suppose that, for $n \geq 1$,

$$E(\tilde{S}_n / \mathcal{F}_{n-1}) \geq \tilde{S}_{n-1} , \text{ a.s.} , \quad (2.1)$$

$$E(|\tilde{S}_n - \tilde{S}_{n-1}| / \mathcal{F}_{n-1}) \leq M , \text{ a.s.} , \quad (2.2)$$

Then $E(\tau) = \infty$.

Proof : For all $k \geq 1$, we get from (2.1)

$$E(|\tilde{S}_k - \tilde{S}_{k-1}|) = E(E(|\tilde{S}_k - \tilde{S}_{k-1}| / \mathcal{F}_{k-1})) \leq MP(\tau > k-1) .$$

Thus, for any $n, l, 1 \leq l \leq n$,

$$\begin{aligned} E(|\tilde{S}_n - \tilde{S}_l|) &= E\left(|\sum_{k=l+1}^n \tilde{S}_k - \tilde{S}_{k-1}|\right) \leq \sum_{k=l+1}^n E(|\tilde{S}_k - \tilde{S}_{k-1}|) . \\ &\leq M \sum_{k=l+1}^n P(\tau \geq k) , \end{aligned} \quad (2.3)$$

whence, immediately,

$$E(|\tilde{S}_n|) \leq M \sum_{k=1}^n P(\tau > k) + S_0 . \quad (2.4)$$

Assume $E(\tau) < \infty$. Then, from (2.3), (2.4) and Cauchy criterion, it follows that \tilde{S}_n is a submartingale converging almost surely (a.s) and in L_1 . [The convergence a.s. is here obvious since, by the hypothesis, $P(\tau < \infty) = 1$ and thus $\tilde{S}_n = S_{n \wedge \tau} \xrightarrow{\text{a.s.}} S_\tau$]. Thus we have

$$E(S_\tau) = \lim_{n \rightarrow \infty} E(\tilde{S}_n) \geq E(S_0) \geq C .$$

But, by the definition of τ , $E(S_\tau) < C$, which yields a contradiction. Hence $E(\tau) = \infty$ and the proof of theorem 2.4 is concluded ■

The proof of theorem 2.3 becomes now straightforward, by choosing

$$\tau = \inf \{n > 0, f(\xi_n) \leq c/f(\xi_0) > c\},$$

ξ_0 being taken constant, and

$$S_n = f(\xi_{n \wedge \tau}), S_0 = f(\xi_0).$$

The proof of theorem 2.3 is concluded. ■

One of our main results is the following

Theorem 2.5 *For an irreducible Markov chain \mathcal{L} to be null recurrent, it is sufficient that there exist two functions $f(x)$ and $\varphi(x)$, $x \in X$, and a finite subset $A \in X$, such that the following conditions hold :*

1. $f(x) \geq 0, \varphi(x) \geq 0, \forall x \in X$;
2. For some positive α, γ, ϵ , with $0 < \epsilon \leq \alpha$ and $1 \leq \alpha \leq 2$,
 $f(x) \leq \gamma(\varphi(x))^{\alpha-\epsilon}, \forall x \in X$;
3. $\varphi(x_i) \rightarrow \infty$, for $i \rightarrow \infty$,
 $\sup_{x \notin A} f(x) > \sup_{x \in A} f(x)$;
4. (a) $E[f(\xi_{n+1}) - f(\xi_n)/\xi_n = x] \geq 0, \forall x \notin A$;
(b) $E[\varphi(\xi_{n+1}) - \varphi(\xi_n)/\xi_n = x] \leq 0, \forall x \notin A$;
(c) $\sup_{x \in X} E[|\varphi(\xi_{n+1}) - \varphi(\xi_n)|^\alpha / \xi_n = x] = C < \infty$

Proof : Let us suppose that such functions exist. Conditions 1), 3) and 4b) on $\varphi(x)$ show immediately, by using theorem 4.1, that \mathcal{L} is recurrent. We shall now assume that \mathcal{L} is ergodic and then come to a contradiction, thus proving the null recurrence.

Let us denote

$$a_n = \varphi(\xi_n), b_n = f(\xi_n),$$

$$\tau = \inf \{n > 0, \xi_n \in A / \xi_0 \notin A\},$$

$$\tilde{a}_n = a_{n \wedge \tau}, \quad \tilde{b}_n = b_{n \wedge \tau}.$$

Since \mathcal{L} is assumed to be ergodic, $E(\tau) < \infty$.

It will be inconvenient throughout this study to choose ξ_0 to be a constant such that $\xi_0 \notin A$. The two following auxiliary lemmas will be useful for us.

Lemma 2.6 *Let β_n , $n \geq 1$, be a sequence of random variables such that $\beta_n \rightarrow \beta$ a.s. and*

$$E|\beta_n|^r \leq c, \quad \forall n \geq 0, \text{ for some } c, r > 0.$$

Then, for any s , $0 \leq s < r$,

$$\lim_{n \rightarrow \infty} E|\beta_n - \beta|^s \rightarrow 0,$$

and, in particular,

$$\lim_{n \rightarrow \infty} E|\beta_n|^s = E|\beta|^s$$

Proof : This is a classical result. See for instance [13]. ■

Lemma 2.7 *Let the S'_n 's of theorem 4 be now positive random variables and τ an arbitrary \mathcal{F}_n -stopping time. If, for all $n \geq 1$ and α, ϵ real numbers such that $1 \leq \alpha \leq 2$, $0 < \epsilon \leq \alpha$,*

$$E(\tilde{S}_{n+1} - \tilde{S}_n / \mathcal{F}_n) \leq 0, \quad \text{a.s.} \quad (2.5)$$

$$E(|\tilde{S}_{n+1} - \tilde{S}_n|^\alpha / \mathcal{F}_n) \leq M, \quad \text{a.s.} \quad (2.6)$$

$$E(\tau) < \infty,$$

then

$$\sup_n E(\tilde{S}_n^\alpha) < \infty, \quad (2.7)$$

$$\tilde{S}_n^{\alpha-\epsilon} \xrightarrow{L_1} S_\tau^{\alpha-\epsilon}. \quad (2.8)$$

Proof : Define

$$\Delta\tilde{S}_n = \tilde{S}_{n+1} - \tilde{S}_n .$$

The following estimate takes place, from Taylor's formula,

$$\tilde{S}_{n+1}^\alpha - \tilde{S}_n^\alpha = \alpha\Delta\tilde{S}_n(\tilde{S}_n + \theta_n\Delta\tilde{S}_n)^{\alpha-1} , \quad (2.9)$$

where $0 < \theta_n < 1$, $\forall n \geq 0$.

The right member of (2.9) can be rewritten as

$$\begin{aligned} & \alpha \tilde{S}_n^{\alpha-1} \Delta\tilde{S}_n + \alpha \tilde{S}_n^{\alpha-1} \Delta\tilde{S}_n \left[\left(1 + \frac{\theta_n \Delta\tilde{S}_n}{\tilde{S}_n}\right)^{\alpha-1} - 1 \right] \leq \\ & \alpha \tilde{S}_n^{\alpha-1} \Delta\tilde{S}_n + \alpha |\Delta\tilde{S}_n|^\alpha , \end{aligned}$$

where we have used the elementary inequalities

$$|1 + v|^q \leq 1 + v^q , \quad |1 - v|^q \geq 1 - v^q , \quad \forall q, \quad 0 \leq q \leq 1 , \quad \forall v \geq 0 .$$

Thus taking conditional expectation in (2.9) and using (2.5) and (2.6), we get

$$E[\tilde{S}_{n+1}^\alpha - \tilde{S}_n^\alpha / \mathcal{F}_n] \leq \alpha M \mathbf{1}_{\{\tau > n\}} , \text{ a.s.} \quad (2.10)$$

$$\text{where } \mathbf{1}_A = \begin{cases} 1 & \text{if } A \text{ is true,} \\ 0 & \text{otherwise.} \end{cases}$$

It follows from (2.10) that

$$E(\tilde{S}_{n+1}^\alpha) \leq \alpha M \sum_{k=0}^n P(\tau > k) + S_0^\alpha = \alpha M E(\tau) + S_0^\alpha ,$$

The finiteness of $E(\tau)$ yields (2.7). The convergence in L_1 of $\tilde{S}_n^{\alpha-\epsilon}$ to $S_\tau^{\alpha-\epsilon}$ is now a direct consequence of lemma 2.6, since \tilde{S}_n is a positive supermartingale and $\tilde{S}_n \xrightarrow{a.s.} S_\tau$.

Lemma 2.7 is proved \blacksquare

Let us return, to the proof of the theorem. Since A is a finite set,

$$\sup_{x \in A} \varphi(x) < \infty, \quad \sup_{x \in A} f(x) < \infty.$$

Since \mathcal{L} is assumed to be ergodic, $P(\tau < \infty) = 1$ and there exist two random variables \tilde{a} and \tilde{b} such that

$$\begin{aligned} \tilde{a}_n^{\alpha-\epsilon} &= \varphi^{\alpha-\epsilon}(\xi_{n \wedge \tau}) \xrightarrow{a.s.} \tilde{a}, \quad 0 \leq \tilde{a} \leq \sup_{x \in A} \varphi_x^{\alpha-\epsilon}, \\ \tilde{b}_n &= f(\xi_{n \wedge \tau}) \xrightarrow{a.s.} \tilde{b}, \quad 0 \leq \tilde{b} \leq \sup_{x \in A} f(x). \end{aligned}$$

Moreover, lemmas 2.6 and 2.7 entail that the r.v.'s $\tilde{a}_n^{\alpha-\epsilon}$ are uniformly integrable and converge to \tilde{a} in the L_1 -sense. Using condition 2) of theorem 2.5, we have

$$\tilde{b}_n = f(\xi_{n \wedge \tau}) \leq \gamma \tilde{a}_n^{\alpha-\epsilon}.$$

Thus the family $(\tilde{b}_n), n \geq 0$, dominated by a uniformly integrable family, is also uniformly integrable. This shows that \tilde{b} is the L_1 -limit of \tilde{b}_n and

$$\lim_{n \rightarrow \infty} E\tilde{b}_n = E\tilde{b} \leq \sup_{x \in A} f(x) \quad (2.11)$$

On the other hand, condition 4a) shows that \tilde{b}_n is a submartingale and

$$E[\tilde{b}_n / \xi_0 = i] \geq f(\xi_0) = f(i), \quad \forall i \notin A, \quad \forall n \geq 0. \quad (2.12)$$

From condition 3), we can choose i so that

$$f(i) > \sup_{x \in A} f(x).$$

Doing so, we get from the estimate (2.11), which does not depend on the initial position ξ_0 ,

$$\lim_{n \rightarrow \infty} E(\tilde{b}_n / \xi_0 = i) \leq \sup_{x \in A} f(x),$$

and this last inequality contradicts (2.12). Thus necessarily $E(\tau) = \infty$ and the proof of theorem 2.5 is completed. ■

3 Proof of the main result (theorem 1.1)

Let us introduce the linear function $\varphi : R_+^2 \rightarrow R_+$,

$$\varphi(x, y) = px + qy, \quad p > 0, q > 0.$$

We also shall write, for any vector $\gamma = (x, y)$

$$\varphi(\gamma) \equiv \varphi(x, y).$$

Lemma 3.1 *Let $p, q > 0$, be such that the vectors M' and M'' have the following properties (see fig. 2)*

$$\begin{cases} \varphi(M') = pM'_x + qM'_y \geq 0, \\ \varphi(M'') = pM''_x + qM''_y \geq 0. \end{cases} \quad (3.1)$$

Then, for all $(x, y) \in Z_+^2, (x, y) \neq (0, 0)$,

$$E[\varphi(\xi_{n+1})/\xi_n = (x, y)] \geq \varphi(x, y), \quad (3.2)$$

i.e. $\varphi(\xi_n)$ is a positive submartingale.

Proof: Immediate by using the linearity of φ and the fact that $M = (M_x, M_y) = (0, 0)$ for $x, y > 0$. ■

We claim that the conditions of lemma 3.1 hold in each of the cases (i)-(iv) of theorem 1.1. They are also valid in the case

$$\text{a) } \phi_x < \frac{\pi}{2}, \phi_y < \frac{\pi}{2}, \phi_x + \phi_y \geq \frac{\pi}{2},$$

which yields

$$M'_x M''_y - M'_y M''_x \leq 0. \quad (3.3)$$

Lemma 3.2 *If, for the random walk \mathcal{L} , the conditions of lemma 3.1 are fulfilled, then \mathcal{L} is not ergodic.*

Proof : It is a direct consequence of theorem 2.3. It is worth mentioning that this result holds under the mere assumption.

$$E[\|\theta_{n+1}\| / \xi_n = (x, y)] \leq K < \infty ,$$

which is weaker than the condition B stated in section 1. ■

Let us consider now the case

b) $\phi_x + \phi_y < \frac{\pi}{2}$.

Then we have the property opposite to that of lemma 3.1, since now the vectors M' and M'' look inside the simplex bounded by the two positive axis and the line $px + qy = D$. (see fig 3).




Figure 2




Figure 3

It means that there exist $p > 0, q > 0$, such that the linear function $\phi(x_t, y_t)$ be a positive supermartingale. Thus the random walk \mathcal{L} is recurrent. Our goal is to distinguish between positive and null recurrence. To that end, we introduce the following functional of quadratic form

$$f(x, y) = (ux^2 + vy^2 + xy)^\delta , \quad x, y \in \mathbb{Z}_+^2 ,$$

where $u \geq 0, v \geq 0, 0 < \delta < 1$.

We want to adjust u, v and δ to satisfy the conditions of theorem 2.5.

Define, for the sake of brevity,

$$\begin{aligned} Q(x, y) &= ux^2 + vy^2 + xy , \\ \Delta Q(x, y) &= Q(x + \theta_x, y + \theta_y) - Q(x, y) \\ &= u\theta_x^2 + v\theta_y^2 + \theta_x\theta_y + (y + 2ux)\theta_x + (x + 2vy)\theta_y , \end{aligned} \quad (3.4)$$

where (see section 1),

$$\theta_{n+1} = (\theta_x, \theta_y), \text{ given } \xi_n = (x, y) \geq 0,$$

in which case, *ad libitum*, $f(x, y)$ will be rewritten as $f(\xi_n)$.
We shall estimate the quantity

$$\begin{aligned} H(x, y) &\stackrel{\text{def}}{=} E[f(\xi_{n+1}) - f(\xi_n) / \xi_n = (x, y)] \\ &= E[(Q(x, y) + \Delta Q(x, y))^\delta - Q^\delta(x, y)] \end{aligned} \quad (3.5)$$

Lemma 3.3 *There exists $\delta > 0$ and a constant D such that*

$$H(x, y) = \delta Q^{\delta-1}(x, y)[E(\Delta Q(x, y)) + (\delta - 1)\mathcal{O}(1) + \mathfrak{o}(1)] \quad (3.6)$$

for all (x, y) , such that $(x^2 + y^2) > D^2$,

where, as usual, $|\mathcal{O}(z)| < K|z|$ and $\frac{\mathfrak{o}(z)}{z} \rightarrow 0$ as z tends to a given limit.
In particular, $\mathfrak{o}(1)$ means a function which tends to zero when $D \rightarrow \infty$.

Proof : From Taylor's formula we have

$$\begin{aligned} H(x, y) &= E[\Delta Q(x, y)(Q(x, y) + \gamma(x, y)\Delta Q(x, y))^{\delta-1}] \\ &= \delta Q^{\delta-1}(x, y)[E[\Delta Q(x, y)] + \psi(x, y)], \end{aligned} \quad (3.7)$$

where $\gamma(x, y)$ is a random variable such that $0 < \gamma(x, y) < 1$ (note that $Q + \gamma\Delta Q \geq 0$) and

$$\psi(x, y) = E[\Delta Q(x, y)\left[\left(1 + \frac{\gamma(x, y)\Delta Q(x, y)}{Q(x, y)}\right)^{\delta-1} - 1\right]].$$

It suffices to prove $\psi(x, y) = (\delta - 1)\mathcal{O}(1) + \mathfrak{o}(1)$.

i) **Case of bounded jumps.** The result is immediate from the definition (3.4), after using $|(1+z)^{\delta-1} - 1| = (\delta-1)\mathcal{O}(z)$, for z sufficiently small, since, from the boundedness of the jumps

$$\frac{\Delta^2 Q(x, y)}{Q(x, y)} < A < \infty, \quad \forall (x, y) \neq (0, 0).$$

ii) **Case of unbounded jumps.** Let us write $\psi(x, y)$ under the form

$$\psi(x, y) = \psi_1(x, y) + \psi_2(x, y),$$

with

$$\begin{aligned}\psi_1(x, y) &= E[\Delta Q(x, y)T(x, y)\mathbf{1}_{\{|\theta_x + \theta_y| \leq z\}}], \\ \psi_2(x, y) &= E[\Delta Q(x, y)T(x, y)\mathbf{1}_{\{|\theta_x + \theta_y| > z\}}],\end{aligned}\quad (3.8)$$

where z is some positive real number and

$$T(x, y) \stackrel{def}{=} \left[1 + \frac{\gamma(x, y)\Delta Q(x, y)}{Q(x, y)}\right]^{\delta-1} - 1.$$

We have, according to (3.4),

$$\Delta Q(x, y) = x(2u\theta_x + \theta_y) + y(2v\theta_y + \theta_x) + u\theta_x^2 + v\theta_y^2 + \theta_x\theta_y \quad (3.9)$$

Hence, using condition A of section 1 on the lower boundedness of jumps, we have for fixed $z > 0$

$$|\Delta Q(x, y)| \leq (ax + by)z + c^2z^2, \text{ for } |\theta_x + \theta_y| < z$$

where a, b, c are positive constants depending only u, v and it follows that for $(x^2 + y^2) > D^2, z < \sqrt{D}$,

$$\begin{aligned}|\psi_1(x, y)| &\leq E[|\Delta Q(x, y)T(x, y)| \mathbf{1}_{\{|\theta_x + \theta_y| \leq z\}}] \\ &= (1 - \delta)\mathcal{O}\left[\frac{((ax + by)z + c^2z^2)^2}{Q(x, y)}\right] = (1 - \delta)\mathcal{O}(1)\end{aligned}\quad (3.10)$$

which gives an estimate for $\psi_1(x, y)$.

On the other hand, on $\{|\theta_x + \theta_y| > z\}$, $\Delta Q(x, y)$ is positive, as emerges from (3.9), whence

$$-1 \leq T(x, y) \leq 0 \text{ and } |T(x, y)| \leq 1.$$

It follows that

$$|\psi_2(x, y)| \leq E[\Delta Q(x, y)\mathbf{1}_{\{|\theta_x + \theta_y| > z\}}].$$

Hence

$$\psi_2(x, y) \leq E[Q(\theta_x, \theta_y)\mathbf{1}_{\{|\theta_x + \theta_y| > z\}}] + (ax + by)E[|\theta_x + \theta_y| \mathbf{1}_{\{|\theta_x + \theta_y| > z\}}]. \quad (3.11)$$

To estimate the right member of (3.11), we use the following simple result, valid for any random variable X , such that $E[|X|^r] < \infty$,

$$E[|X|^r \mathbf{1}_{\{X > z\}}] = o(|z^{s-r}|), \quad \forall 0 \leq s \leq r.$$

Therefore, taking into account the moment condition B ensuring $E[|\theta_x + \theta_y|^3] < \infty$, we get

$$|\psi_2(x, y)| \leq o\left(\frac{1}{z}\right) + (ax + by) o\left(\frac{1}{z^2}\right). \quad (3.12)$$

Choosing again $z < \sqrt{D}$ in (3.12) yields

$$\psi_2(x, y) = o(1). \quad (3.13)$$

Finally, (3.10) and (3.13) together imply $|\psi(x, y)| = (1 - \delta)\mathcal{O}(1) + o(1)$ and the proof of lemma 3 is concluded ■

Remark : It is possible to refine this proof, assuming only the existence of moments of order two. We omit it to avoid unnecessary excursions, which would obscure the readability and the general ideas.

We continue with the proof of theorem 1.1.

It follows from (3.6), that one can find D and $\delta, 0 < \delta < 1$, such that, if $E[\Delta Q(x, y)] > 0$, then $H(x, y) \geq 0$, for any x, y satisfying $x^2 + y^2 > D$, or equivalently, from (3.5),

$$E[f(\xi_{n+1}) - f(\xi_n) | \xi_n = (x, y)] \geq 0.$$

But $E[\Delta Q(x, y)] > 0, x^2 + y^2 > D^2$, is equivalent, by using (3.9), to the following system of inequalities

$$\begin{cases} u_x + v\lambda_y + R > 0, \\ 2vM_y'' + M_x'' > 0, \\ 2uM_x' + M_y' > 0, \end{cases} \quad (3.14)$$

for some $u, v > 0$.

Thus if (3.14) is satisfied, there exist two functions $f(x, y) = (ux^2 + vy^2 + xy)^\delta$ and $\varphi(x, y) = px + qy$, such that, when $\phi_x + \phi_y < \frac{\pi}{2}$, the conditions 1,

3, and 4 of theorem 2.5 hold, simply taking $\alpha = 2$ in the statement of the theorem. Moreover in this case, condition 2 of theorem 2.5 becomes immediately fulfilled, since for x and y sufficiently large, $0 < \delta < 1$,

$$0 < (ux^2 + vy^2 + xy)^\delta < K(x^2 + y^2) .$$

But, since from the assumptions $M'_y \geq 0$, $M'_x < 0$, $M''_x \geq 0$, $M''_y < 0$, we conclude that (??) holds for some $u, v > 0$, if

$$-\lambda_x \frac{M'_y}{M'_x} - \lambda_y \frac{M''_x}{M''_y} + 2R > 0 ,$$

which is simply on other way of rewriting (1.2).

We have proved the “null recurrence” part of theorem 1.1.

To prove the ergodicity part of the theorem, we proceed as in [], by introducing again the quadratic form

$$Q(x, y) = ux^2 + vy^2 + xy$$

and showing that Foster’s criterion (2.2) can be satisfied, for some $u, v, \epsilon > 0$. With the notation above, we have, from (3.9),

$$\begin{aligned} K(x, y) &\stackrel{\text{def}}{=} E[Q(\xi_{n+1}) - Q(\xi_n) | \xi_n = (x, y)] \\ &= xE[2u\theta_x + \theta_y] + yE[2v\theta_y + \theta_x] + E[Q(\theta_x, \theta_y)] . \end{aligned} \quad (3.15)$$

Since $E[Q(\theta_x, \theta_y)] = \mathcal{O}(1)$, $\forall (x, y) \in \mathbf{Z}_+^2$, we get from (??), after taking into account the boundary conditions on the axes,

$$K(x, y) = \begin{cases} \lambda_x u + \lambda_y v + R, & (x, y) > 0 , \\ x(2uM'_x + M'_y) + \mathcal{O}(1), & x > 0, y = 0 , \\ y(2vM''_y + M''_x) + \mathcal{O}(1), & x = 0, y > 0 . \end{cases}$$

Thus, for some $\epsilon > 0$ and some finite subset $\mathbf{E} \in \mathbf{Z}_+^2$, we have $K(x, y) < -\epsilon$, $\forall x \notin \mathbf{E}$, provided that the following system can be satisfied, for some $u, v > 0$,

$$\begin{aligned} \lambda_x u + \lambda_y v + R &< 0 , \\ 2uM'_x + M'_y &< 0 , \\ 2vM''_y + M''_x &< 0 . \end{aligned} \quad (3.16)$$

The inequalities $M_x'' \geq 0$, $M_y'' < 0$, $M_x' \geq 0$, $M_y' < 0$, show at once that (3.16) can be satisfied for $u > 0$, $v > 0$, if (1.1) holds, that is

$$-\lambda_x \frac{M_y'}{M_x'} - \lambda_y \frac{M_x''}{M_y''} + 2R < 0 .$$


In this case the remaining conditions of Foster's criterion are clearly fulfilled and the random walk is ergodic.

The proof of theorem (1.1) is terminated. ■

References

- [1] V.A. Malyshev, Classification of two-dimensional positive random walks and almost linear semimartingales, *Dokl. Akad. Nauk. SSSR* 202 (1972), 526-528; English Transl. in *Soviet Math. Dokl.* 13 (1972)
- [2] V.A. Malyshev and M.V. Menshikov, Ergodicity, continuity and analyticity of countable Markov chains, *Trans. Moscow Math. Soc.* V.39 (1979), 2-48 ; (Transl. 1981, Issue I).
- [3] K.L. Vaninsky and B.V. Lazareva, About ergodicity and transience of homogeneous Markov chains in a positive quadrant, *Problems of Inf. Trans.* V.24 N.1 (1988), 105-110.
- [4] G. Fayolle, R. Iasnogorodski, Criteria for the non-ergodicity of stochastic processes : application to the exponential Back-off protocol. *J. Appl. Prob.* 24, (1987), 347-354.
- [5] G. Fayolle, On random walks arising in queueing systems : Ergodicity and transience via quadratic forms as Lyapounov functions - Part I, *Queueing Systems Theory and Appl.* 5, (1989), 167-184.
- [6] M.V. Menshikov, Ergodicity and transience conditions for random walks in the positive octant of space, *Dokl. Akad. Nauk SSSR* 217 (1974), 755-758. English transl. in *Soviet Math. Dokl.* 15 (1974).
- [7] S. Karlin, *A First Course in Stochastic Processes*, Academic Press, (1966).

- [8] V.A. Malyshev, *Wiener-Hopf equations, Galois Automorphisms, Random Walks in a Quarter Plane*, Moscow State Univ. (1970).
- [9] V.A. Malyshev, Doctor Thesis, Moscow (1973).
- [10] F.G. Foster, On Stochastic matrices associated with certain queueing processes, *Ann. Math. Statist.* 24 (1953), 355-360.
- [11] W.A. Rosenkrantz, Ergodicity conditions for two-dimensional Markov chains on the positive quadrant, *Prob. Theory and Related Fields*, Vol.83, N.3, (1989).
- [12] R.L. Tweedie, Criteria for classifying general Markov chains, *Adv. Appl. Prob.* 8, (1976), 737-771.
- [13] M. Loeve, *Probability Theory*, 2nd. Ed. Princeton, van Nostrand, (1960).



ISSN 0249 - 6399