



## Fairness, distances and degrees

Philippe Darondeau, Doris Nolte, Lutz Prieze, Serge Yoccoz

### ► To cite this version:

Philippe Darondeau, Doris Nolte, Lutz Prieze, Serge Yoccoz. Fairness, distances and degrees. [Research Report] RR-1199, INRIA. 1990. inria-00075359

**HAL Id: inria-00075359**

**<https://inria.hal.science/inria-00075359>**

Submitted on 24 May 2006

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



UNITÉ DE RECHERCHE  
INRIA-RENNES

Institut National  
de Recherche  
en Informatique  
et en Automatique

Domaine de Voluceau  
Rocquencourt  
BP105  
78153 Le Chesnay Cedex  
France  
Tél (1) 39 63 55 11

# Rapports de Recherche

N° 1199

*Programme 1*  
*Programmation, Calcul Symbolique*  
*et Intelligence Artificielle*

## FAIRNESS, DISTANCES AND DEGREES

**Philippe DARONDEAU**  
**Doris NOLTE**  
**Lutz PRIESE**  
**Serge YOCCOZ**

**Mars 1990**



★ RR - 1199 ★

Campus Universitaire de Beaulieu

35042-RENNES CÉDEX **Fairness, Distances and Degrees**

FRANCE

Téléphone: 99 36 20 00

Télex: UNIRISA 950 473 F

Télécopie: 99 38 38 32 Philippe Darondeau, Doris Nolte, Lutz Priese, Serge Yoccoz,

**Abstract**

*We show the identity between sets of fair computations in recursive transition graphs, sets of cluster points of finite computations for  $\Pi_1^0$  ultra-metrics refining the Baire metrics, and  $\Pi_3^0$  subsets of  $\omega^\omega$ . The results are applied to recursive marked trees, fairness definitions,  $\omega$ -regular languages, and  $\Pi_3^0$  sets.*

**Equité, distances et degrés****Résumé**

*Nous identifions les ensembles d'exécutions équitables dans des graphes de transitions récursifs, les ensembles de points d'accumulation d'exécutions finies pour des ultra-distances  $\Pi_1^0$  raffinant la distance de Baire, et les ensembles  $\Pi_3^0$  de  $\omega^\omega$ . Ces résultats sont appliqués aux arbres marqués récursifs, aux définitions de l'équité, aux langages  $\omega$ -réguliers, et aux ensembles  $\Pi_3^0$ .*

**1. Introduction**

The dynamics of a machine or program is best represented as a transition graph, defined as a set of states equipped with a set of binary transitions between states, possibly labelled on an alphabet of transition symbols [Keller]. Runs of machines or programs are identified with countable paths in the graph, called computations. Most transition graphs encountered in practice are finite if they are models of machines, e.g. finite state automata [Hopcroft and Ullman], or recursive if they are models of programming calculi, e.g. the  $\lambda$ -calculus with  $\beta$ -reduction [Barendregt] or the Calculus of Communicating Systems [Milner]. Recursively enumerable transition graphs are a borderline case. A nice example is MEJE, a synchronous process calculus with unguarded recursion [Austry-Boudol].

---

1 IRISA, Université de Rennes I, Campus de Beaulieu, 35042 Rennes cedex, France

2 Universität-Gesamthochschule, Postfach 1621, Paderborn, FRG

3 LABRI, Université de Bordeaux I, 351 ers de la Libération, 33405 Talence, France

In the case of deterministic programs, one is primarily interested in terminating computations, but infinite computations play a major role for deterministic machines, and thereby prompt the consideration of topologies. For instance, Landweber's theorem tells us that languages accepted by deterministic Büchi automata are  $G_\delta$  ( $\Pi_2^0$ ) in the topology induced by the standard ultra-metric distance on  $\omega$ -words, because  $G_\delta$ -sets coincide with Eilenberg limits of sets of finite words [Landweber][Eilenberg]. With such automata, the set of successful computations generally differs from the set of infinite computations of which it is a subset. The set of infinite computations is the derived set of the set of finite computations in the natural metric topology on computations, derived from Baire metric, and thus it is  $\Pi_1^0$ . The subset of the successful computations is specified by a  $\Pi_2^0$  predicate on infinite computations, and it is yet the derived set of the set of finite computations in another metric topology which refines the natural topology. The above situation is essentially reproduced for non deterministic machines, which accept  $\omega$ -languages located higher in Borel's hierarchy ( $\omega$ -regular languages are in the boolean closure of  $G_\delta$ ).

As regards programming calculi, a metric semantics for non deterministic recursive program schemes, based on the definition of a Baire like metric on computation trees, has been constructed in [Arnold-Nivat]. Similarly, a metric semantics for concurrent programs, based on the definition of a Baire like metric on streams, has been constructed in [de Bakker-Meyer]. That model is representative of unfairness in the following sense: the race between concurrent agents which compete for engaging themselves in an interleaved computation is totally free. The role of fairness is to place restrictions on that race, expressed by predicates on computations. The most popular forms of fairness are weak fairness and strong fairness [Francez]. In a weakly (resp. strongly) fair computation, an agent which is almost always (resp. infinitely often) enabled to act, acts infinitely often. If the set of agents is unbounded, the corresponding predicate on infinite computations is  $\Pi_2^0$  for weak fairness and  $\Pi_3^0$  for strong fairness. The set of fair computations is more complex than the set of infinite computations, which is  $\Pi_1^0$  for recursive transition graphs as the derived set of the set of finite computations in the natural topology.

More essentially, for both weak or strong fairness, the set of fair computations is the derived set of the set of finite computations in a metric topology which refines the natural topology. This central property, shown first for weak fairness in the context of a special programming language [Degano-Montanari], was established later on for strong fairness in CCS [Costa]. In the light of the papers referred to, any definition of fairness aims at restricting the convergence of sequences of finite computations which approximate infinite computations: unfair computations, which disappear from the derived set of the set of finite computations, are then discarded. The parallel with sequential machines is almost perfect, and the question arises whether one can find a counterpart to Landweber's theorem.

In this paper, we will show that any  $\Pi_3^0$  subset of  $\omega^{(n)}$  may be seen as the set of fair computations in a (fixed) recursive transition graph, for a (generic) concept of fairness based on a (varying) recursive relation which decides whether a given agent is enabled at a given step in a given computation. Furthermore, we will identify the  $\Pi_3^0$  subsets of  $\omega^{(n)}$  with the family of the derived sets of  $\omega^*$  for  $\Pi_1^0$  metrics on  $\omega^\infty$  refining the natural metric. As a result, the sets of fair computations in recursive transition graphs and the derived sets of  $\omega^*$  for  $\Pi_1^0$  metrics are the same. In addition, the correspondence between logical and metrical definitions of fairness is effective in both directions: definitions of  $\Pi_1^0$  metrics (on  $\omega^\infty$ ) are translated uniformly to definitions of fairness, and vice versa. That effective correspondence fulfills the research program which Degano and Montanari launched into. By relativizing the correspondence between  $\Pi_3^0$  subsets of  $\omega^{(n)}$  and  $\Pi_1^0$  metrics on  $\omega^\infty$ , we moreover obtain a metric characterization of the class  $F_{\sigma\delta}(\Pi_3^0)$  in the classical Borel hierarchy. This is the expected counterpart to Landweber's theorem.

The remaining sections of the paper are organized as follows. Notations and definitions are introduced in section II. The relationships between sets of fair computations and  $\Pi_3^0$  sets of functions are studied in section III. The applications are considered in section IV.

## II. Notations and definitions

In the sequel,  $\omega$  is the set of natural numbers and  $\omega^*$  is the set of finite sequences of natural numbers. We recall that  $\omega^* = \bigcup_{k=0}^{\infty} \omega^k$  is mapped recursively onto  $\omega$  by a one-one coding  $\tau^*((x_1, \dots, x_k)) = \langle x_1, \dots, x_k, k-1 \rangle + 1$ , monotonous in each one of the  $x_i$  for sequences of fixed length  $k > 0$ , where  $\tau^*(\emptyset) = 0$  [Rogers, p.71]. This coding gives rise to an isomorphism between the disjoint sums  $\omega^\infty = \omega^* \cup \omega^\omega$  and  $\Omega = \omega \cup \omega^\omega$  in the category of sets. For each  $k \geq 0$  we can therefore define a function  $[k]: \Omega \rightarrow \omega$  truncating objects at depth  $k$  according to the definitions:

$$f[0] = 0 \text{ and } f[k] = \tau^*((f(0), \dots, f(k-1))) \text{ for } f \in \omega^\omega \text{ and } k > 0,$$

$$x[0] = 0 \text{ and } x[k] = 0 \text{ for } x \in \omega \text{ and } k > 0,$$

$$x[k] = \tau^*((x_1, \dots, x_{\min(k, l)})) \text{ for } x = \tau^*((x_1, \dots, x_l)) \text{ and } k, l > 0.$$

When we deal with arithmetical functions on  $\Omega$ , and more generally with arithmetical relations on  $\Omega^k \times \omega^l$ , we define the arithmetical class of a relation  $R$  as the least upper bound (with respect to Kleene's hierarchy over  $\Sigma_n^0$ ,  $\Pi_n^0$  and  $\Delta_n^0$  classes) of the class of  $R \cap ((X_1 \times \dots \times X_l \times \omega^l) \times \omega^l)$  for variables  $X_i$  ranging over  $\{\omega, \Omega\}$ . We refer the reader to [Rogers] for a thorough presentation of the arithmetical hierarchy. Thus  $\alpha[k]$  may be considered as a recursive function from  $\Omega \times \omega$  to  $\omega$ . By way of definition,  $\alpha \wedge \beta$  is the least upper bound of the set  $\{k \mid \alpha[k] = \beta[k]\}$ .

The *natural metric* on  $\Omega$  is the ultra-metric  $\delta$  defined as  $\delta(\alpha, \beta) = 0$  if  $\alpha = \beta$ ,  $1/(1 + (\alpha \wedge \beta))$  otherwise. Because the relation  $\delta(\alpha, \beta) < 1/n$  is recursive in  $(\alpha, \beta, n)$ , we call  $\delta$  a *recursive metric*. Similarly, a metric  $d$  on  $\Omega$  is said to be  $\Pi_1^0$  (resp.  $\Sigma_2^0$ ) if the relation  $d(\alpha, \beta) < 1/n$  is  $\Pi_1^0$  (resp.  $\Sigma_2^0$ ) in  $(\alpha, \beta, n)$ . We are mainly interested in the derived sets of  $\omega$  and its recursive (or recursively enumerable) subsets for  $\Pi_1^0$  metrics  $d$  on  $\Omega$  refining  $\delta$  ( $d \geq \delta$ ). We recall from [Dugundji] that in a metric space  $(X, d)$ , the *derived set*  $A'$  (or  $A'_d$ ) of  $A$  is the set of the  $x \in X$  such that any open ball with center  $x$  and radius  $1/n$  contains at least one element  $a \in A$  distinct from  $x$ .

11.10 We will show that any  $\Pi_3^0$  subset of  $\omega^\omega$  coincides with  $\omega_d^1$  for some  $\Pi_1^0$  symmetric d on  $\Omega$ , representing a concept of effective fairness. Definitions of fairness make sense with respect to computations in transition graphs. A *transition graph*  $T$  is a quadruple  $T=(S, T, \sigma_0, \sigma_1)$  where  $S, T$  are sets of numbers (representing states and transitions) and  $\sigma_0, \sigma_1$  are mappings from  $T$  to  $S$  (indicating the source and target of transitions). A transition graph is recursive (resp. recursively enumerable) if all its components are recursive (resp.  $\Sigma_1^0$ ). For instance, the transition graph  $T_\omega=(\{\omega\}, \omega, \sigma_0, \sigma_1)$  is recursive. A *pointed transition graph*  $T_s$  is a transition graph  $T$  with an initial state  $s$ .

11.11 A *computation* in a pointed transition graph  $T_{s_0}$  is a finite or denumerable sequence of transitions  $(t_i)$  satisfying  $\sigma_0(t_i) = s_0$  if  $i=0$  and  $\sigma_0(t_i) = \sigma_1(t_{i-1})$  if  $i>0$ , and for all  $i$  in the domain of the sequence. A finite computation  $t_0 \dots t_{k-1}$  with length  $k$  is represented in  $\Omega$  by the number  $\tau^*((t_0, \dots, t_{k-1}))$ , while an infinite computation  $(t_i)_{i \in \omega}$  is represented by the function  $f(i)=t_i$ . Let  $\text{Fin}(T_s)$  resp.  $\text{Inf}(T_s)$  denote the set of the finite resp. infinite computations from  $s$  in  $T$ , then, of course  $\text{Inf}(T_s) = (\text{Fin}(T_s))'_\delta$  in the topology induced by the natural metric  $\delta$  on  $\Omega_{\text{fin}}$ .

11.12 A concept of *effective fairness* in a transition graph  $T_s$  is totally determined by an *enabling predicate* defined as a recursive relation  $E \in \omega^2$ . Intuitively,  $E(f[k], i)$  means that agent  $i$  is enabled at the  $k^{\text{th}}$  step in computation  $f$ . An *E-fair* computation is then an infinite computation in which no agent is enabled infinitely often. In formulas,  $[f \text{ is } E\text{-fair}] \Leftrightarrow [\forall i. \neg \exists^{\omega} k. E(f[k], i)]$ , or, equivalently,  $[f \text{ is } E\text{-fair}] \Leftrightarrow [\forall i. \forall^{\omega} k. \neg E(f[k], i)]$ , where  $\exists^{\omega}$  and  $\forall^{\omega}$  are the dual infinite quantifiers.

Let  $E\text{-fair}(T_s)$  denote the set of  $E$ -fair computations in transition graph  $T_s$ .

11.13 For recursive (resp. recursively enumerable) transition graphs  $T_s$ ,  $\text{Inf}(T_s)$  is  $\Pi_1^0$  (resp.  $\Pi_2^0$ ), and  $E\text{-fair}(T_s)$  is  $\Pi_3^0$ . Similar observations pertain to the usual definitions of weak fairness ( $\Pi_2^0$ ), strong and extreme fairness ( $\Pi_3^0$ ), see for instance [Harel]. We will show that all those variant definitions of fairness may in fact be reduced to  $E$ -fairness without altering the transition graphs.

### III. A connection between $\Pi_3^0$ sets of functions, $\Pi_1^0$ metrics on $\Omega$ , and E-fairness.

On support of the definitions and notations introduced in the last section, we can evolve a precise statement of the connection announced in the introduction.

**Theorem 1** For  $F \subseteq \omega^{(\omega)}$  the following assertions 1, 2, 3 and 4 are equivalent:

- (1)  $F$  is  $\Pi_3^0$  (as a set of functions),
- (2)  $F$  is the derived set of  $\omega$  for some  $\Pi_1^0$  ultra-metric on  $\Omega$ , induced from a recursive distance on  $\omega$  and refining the natural distance  $\delta$ ,
- (3)  $F$  is the derived set of a  $\Sigma_2^0$  subset of  $\omega$  for some  $\Sigma_2^0$  metric on  $\Omega$ ,
- (4)  $F$  equals E-fair( $\omega$ ) for some (recursive) enabling predicate E.

The above characterization for  $\Pi_3^0$  sets of functions is the main result of the paper. We state hereafter a variant characterization for  $\Pi_3^0$  sets of infinite computations.

**Theorem 2** For any recursively enumerable (pointed) transition graph  $T_S$  the following assertions 5, 6 and 7 are equivalent:

- (5)  $F$  is a  $\Pi_3^0$  subset of  $\text{Inf}(T_S)$ ,
- (6)  $F$  is the derived set of  $\text{Fin}(T_S)$  for some  $\Pi_1^0$  ultra-metric refining  $\delta$ ,
- (7)  $F$  equals E-fair( $T_S$ ) for some (recursive) enabling predicate E.

In view of that adapted theorem, all the classical definitions of fairness may be reduced to the universal form of E-fairness. Fundamental for the proof of both theorems is the next lemma, which points out an analogous reduction for  $\Pi_3^0$  definitions of sets of functions. This lemma extends a similar characterization for  $\Pi_3^0$  sets of numbers, established by Kreisel, Shoenfield and Wang, see [Rogers].

**Lemma 1** (normal form for  $\Pi_3^0$  sets of functions)

Any  $\Pi_3^0$  subset  $F$  of  $\omega^{(\omega)}$  may be represented in the normal form

$\{f \mid \forall i. \forall k. \neg E(f[k], i)\}$  where E is a recursive binary relation on numbers, defined uniformly from a  $\Pi_3^0$ -index of F.



proof Assume  $F = \{f \mid \forall i. \exists j. \forall k. R(f, i, j, k)\}$  for some recursive relation  $R$ .  
 (1) For all  $i$  and  $j$ , the set  $\bar{F}_{i,j}$  defined as  $\{f \mid \exists k. \neg R(f, i, j, k)\}$  is clearly equivalent to  $\{f \mid \exists k. \neg S(f[k], i, j)\}$  for some recursive relation  $S$  defined uniformly from  $R$ .  
 (2) For all  $i$ , the set  $F_i$  defined as  $\{f \mid \exists j. \forall k. S(f[k], i, j)\}$  is in turn equal to  $\{f \mid \exists j. \forall k \geq j. \neg E(f[k], i)\}$  for some recursive relation  $E$  defined uniformly from  $S$ .  
 A decision procedure defining  $E(x, i)$  is the following:

```

for k in 0:ω loop
  if  $\forall h \leq k. S(x[h], i, j)$ 
    then ( if  $x[k] = x$  then return  $\neg E(x, i)$  else  $j \leftarrow j$  )
  else ( if  $x[k] = x$  then return  $E(x, i)$  else  $j \leftarrow j+1$  ).

```

In fact,  $f \in F_i$  if and only if the following process outputs eventually always 0, which was brought to our attention by A. Louveau:

```

for k in 0:ω loop
  if  $\forall h \leq k. S(f[h], i, j)$ 
    then (output(0);  $j \leftarrow j$ )
  else (output(1);  $j \leftarrow j+1$  ).

```

Altogether,  $F = \{f \mid \forall i. \forall k. \neg E(f[k], i)\}$ , and the result is obtained.  $\square$

Let us return to the main theorem. The above lemma may be read as (1) $\Rightarrow$ (4). If we can prove (4) $\Rightarrow$ (2), the theorem will follow from (3) $\Rightarrow$ (1) and (2) $\Rightarrow$ (3), which are immediate. The remaining implication (4) $\Rightarrow$ (2) is established below.

Assume  $F = \{f \mid \forall i. \exists j. \forall k \geq j. \neg E(f[k], i)\}$ . For  $\alpha, \beta \in \Omega$  and  $j \in \omega$ , define:

$$(8) \quad \text{done}(\alpha, j) = \max\{i \leq j \mid \forall i' \leq i. \forall k \geq j. \neg E(\alpha[k], i')\},$$

(where  $\max \Phi$  equals 0 by the usual convention)

$$(9) \quad d(\alpha, \beta) = 0 \text{ if } \alpha = \beta, \text{ and otherwise}$$

$$\max(1/1 + \text{done}(\alpha, \alpha \wedge \beta), 1/1 + \text{done}(\beta, \alpha \wedge \beta)).$$

Intuitively,  $\text{done}(\alpha, j)$  is the number of the last agent whose all predecessors are *done* at stage  $j$  in computation  $\alpha$ , for they are never enabled beyond that stage.

Now  $\text{done}(\alpha, j)$  is monotonously increasing in  $j$ , thus the following equivalence holds:

$$(10) \quad (f \in F) \Leftrightarrow \lim_{j \rightarrow \omega} \text{done}(f, j) = \omega.$$

On that basis, and assuming that  $d$  is a metric refining  $\delta$ , we now prove:

$$(11) \quad F = \omega_d'.$$

Assume  $\alpha \in \omega_d'$  then by definition:  $\forall j. \exists x_j \in \omega. 0 < d(\alpha, x_j) < 1/j$ .

As  $d \geq \delta$  and  $(x_j)_j$  is a non stationary sequence of numbers, we know that  $\alpha \in \omega^\omega$ ; then  $\lim_{j \rightarrow \omega} d(\alpha, x_j) = 0 \Rightarrow \lim_{j \rightarrow \omega} \text{done}(\alpha, \alpha \wedge x_j) = \omega \Rightarrow \lim_{j \rightarrow \omega} (\alpha \wedge x_j) = \omega$ , and  $\alpha \in F$  by (10).

Assume  $f \in F$ , then by (10):  $\lim_{j \rightarrow \omega} \text{done}(f, j) = \omega$ .

$$f \wedge f[j] = j \Rightarrow d(f, f[j]) = \max(1/(1 + \text{done}(f, j)), 1/(1 + \text{done}(f[j], j)));$$

$$(j' \geq j \Rightarrow f[j] = (f[j])[j']) \Rightarrow \text{done}(f[j], j) = \max\{i \leq j \mid \forall i' \leq i. \forall j' \geq j. \neg E((f[j])[j'], i')\} \\ = \max\{i \leq j \mid \forall i' \leq i. \neg E(f[j], i')\} \geq \max\{i \leq j \mid \forall i' \leq i. \forall j' \geq j. \neg E(f[j'], i')\} = \text{done}(f, j);$$

thus  $\lim_{j \rightarrow \omega} d(f, f[j]) = 0$  entails  $f \in \omega_d'$ .  $\square$

In order to complete the proof of (4)  $\Rightarrow$  (2) and thereby establish the main theorem, we still have to show that  $d$  is a  $\Pi_1^0$  ultra-metric on  $\Omega$ , induced from a recursive distance on  $\omega$  and refining the natural metric  $\delta$ . Relation  $d(\alpha, \beta) \geq \delta(\alpha, \beta)$  is clear from the inequality  $\text{done}(\alpha, \alpha \wedge \beta) \leq \alpha \wedge \beta$ . The following lemmas 2, 3, 4 provide the rest.

**Lemma 2** For  $\alpha, \beta, \gamma \in \Omega$ :  $d(\alpha, \beta) \leq \max\{d(\alpha, \gamma), d(\beta, \gamma)\}$ .

proof The triangular inequality to be shown may be equivalently restated as

$$(12) \quad \min\{\text{done}(\alpha, \alpha \wedge \beta), \text{done}(\beta, \alpha \wedge \beta)\} \geq \\ \min\{\text{done}(\alpha, \alpha \wedge \gamma), \text{done}(\gamma, \alpha \wedge \gamma), \text{done}(\beta, \beta \wedge \gamma), \text{done}(\gamma, \gamma \wedge \beta)\}.$$

We proceed by case analysis.

case 1.  $\alpha \wedge \gamma = \beta \wedge \gamma \leq \alpha \wedge \beta$ :

$$\alpha \wedge \beta \geq \alpha \wedge \gamma \Rightarrow \text{done}(\alpha, \alpha \wedge \beta) \geq \text{done}(\alpha, \alpha \wedge \gamma),$$

$$\alpha \wedge \beta \geq \beta \wedge \gamma \Rightarrow \text{done}(\beta, \alpha \wedge \beta) \geq \text{done}(\beta, \beta \wedge \gamma).$$

case 2.  $\alpha \wedge \beta = \alpha \wedge \gamma \leq \beta \wedge \gamma$ :

$$\alpha \wedge \beta = \alpha \wedge \gamma \Rightarrow \text{done}(\alpha, \alpha \wedge \beta) \geq \text{done}(\alpha, \alpha \wedge \gamma),$$

and we claim that:

$$\text{done}(\beta, \alpha \wedge \beta) \geq \min\{\text{done}(\beta, \beta \wedge \gamma), \text{done}(\gamma, \alpha \wedge \gamma)\}.$$

Suppose the contrary, and let us search for a contradiction. Define  $\beta[k]$

$$I = I + \text{done}(\beta, \alpha \wedge \beta),$$

then one may assert:

$$(13) \quad \exists k \geq \alpha \wedge \beta. E(\beta[k], I)$$

$$\text{because } I - I = \text{done}(\beta, \alpha \wedge \beta) < I,$$

$$(14) \quad \forall k \geq \beta \wedge \gamma. \neg E(\beta[k], I)$$

$$\text{because } I \leq \text{done}(\beta, \beta \wedge \gamma),$$

$$(15) \quad \forall k \geq \alpha \wedge \gamma. \neg E(\gamma[k], I)$$

$$\text{because } I \leq \text{done}(\gamma, \alpha \wedge \gamma).$$

In view of relations  $\alpha \wedge \gamma = \alpha \wedge \beta \leq \beta \wedge \gamma$  and  $\gamma[\beta \wedge \gamma] = \beta[\beta \wedge \gamma]$ , and by (15):

$$\neg E(\beta[k], I) \text{ for any } k \text{ satisfying } \alpha \wedge \beta \leq k < \beta \wedge \gamma.$$

Hence a contradiction is reached between (13) and (14).

case 3.  $\alpha \wedge \beta = \beta \wedge \gamma \leq \alpha \wedge \gamma$ :

exchange  $\alpha$  and  $\beta$ , and proceed as in case 2.  $\square$

**Lemma 3**  $d$  is a  $\Pi_1^0$  metric on  $\Omega$ , recursive on  $\omega$ .

proof . We will give equivalent  $\Pi_1^0$  formulae for the three types of relations  $d(f, g) < 1/n$ ,  $d(f, x) < 1/n$ , and  $d(x, y) < 1/n$  (where  $f, g \in \omega^\omega$  and  $x, y, n \in \omega$ ).

By definition of  $d$ :  $d(\alpha, \beta) < 1/n \Leftrightarrow$

$$(16) \quad \alpha = \beta \vee (\text{done}(\alpha, \alpha \wedge \beta) \geq n \wedge \text{done}(\beta, \alpha \wedge \beta) \geq n).$$

By definition of  $\text{done}$ :  $\alpha = \beta \vee \text{done}(\alpha, \alpha \wedge \beta) \geq n \Leftrightarrow$

$$(17) \quad \forall j. ((\alpha[j] = \beta[j] \wedge \alpha[j+1] \neq \beta[j+1]) \rightarrow \forall i \leq n. \forall k \geq j. \neg E(\alpha[k], i)).$$

Thus  $d(\alpha, \beta) < 1/n$  is a  $\Pi_1^0$  relation.

As a matter of fact, the relations  $d(f, g) < 1/n$ ,  $d(f, x) < 1/n$ , and  $d(x, y) < 1/n$  are respectively  $\Pi_1^0$ ,  $\Pi_1^0$ , and recursive (since  $x = x[k] \Rightarrow x[k] = x[k+1]$ ).  $\square$

**Lemma 4** Let  $\alpha, \beta \in \Omega$  be such that  $\alpha \neq \beta$  then:

$$d(\alpha, \beta) < 1/n \Leftrightarrow \exists p. \forall m \geq p. d(\alpha[m], \beta[m]) < 1/n.$$

proof The above equivalence is trivially satisfied for  $\alpha \wedge \beta < n$ .

Suppose now  $\alpha \wedge \beta = I \geq n$ . By definition of  $d$  and  $\text{done}$ :

$$\begin{aligned}
(1) \quad d(\alpha, \beta) \geq 1/n &\Leftrightarrow (\text{done}(\alpha, \alpha \wedge \beta) < n \vee \text{done}(\beta, \alpha \wedge \beta) < n) \\
&\Leftrightarrow \exists i \leq n. \exists k \geq 1. (E(\alpha[k], i) \vee E(\beta[k], i)) \\
&\Leftrightarrow \exists p. \forall m \geq p. \exists i \leq n. \exists k \geq 1. \\
&\quad (E(\alpha[m], i) \vee E(\beta[m], i)) \vee (k \leq m \wedge (E(\alpha[k], i) \vee E(\beta[k], i))) \\
&\Leftrightarrow \exists p. \forall m \geq p. \exists i \leq n. \exists k \geq 1. (E(\alpha[m][k], i) \vee E(\beta[m][k], i)) \\
&\Leftrightarrow \exists p. \forall m \geq p. \exists i \leq n. \exists k \geq \min(1, m). (E(\alpha[m][k], i) \vee E(\beta[m][k], i)) \\
&\Leftrightarrow \exists p. \forall m \geq p. d(\alpha[m], \beta[m]) \geq 1/n \quad \square
\end{aligned}$$

The proof for theorem 1 is complete. We now sketch a proof for theorem 2. Let  $T_S$  be a recursively enumerable (pointed) transition graph. Since  $\text{Inf}(T_S)$  is the derived set of  $\text{Fin}(T_S)$  for the natural metric  $\delta$ , (5)  $\Rightarrow$  (6)  $\Rightarrow$  (7) by straightforward adaptation of (1)  $\Rightarrow$  (2)  $\Rightarrow$  (3). Finally, (7)  $\Rightarrow$  (5) is immediate.

As a concluding remark, let us underline the uniform construction of  $d$  from  $F$ , and conversely. Applications are discussed in the final section.

#### IV. Applications

In the above section, we have identified the  $\Pi_3^0$  subsets of  $\omega^\omega$  with the derived sets of  $\omega$  ( $\equiv \omega^*$ ) in metric topologies induced by  $\Pi_1^0$  distances on  $\Omega$  ( $\equiv \omega^* \cup \omega^\omega$ ). We will now examine applications and extensions. The main fields of application gone through are recursive marked trees, fairness expressions, and  $\omega$ -regular languages. The climax is a metric characterization of the family  $F_{\sigma\delta}(\Pi_3^0)$  in the classical Borel hierarchy, established in two different ways.

##### *Recursive marked trees*

In [Harel] was defined a language for stating properties of infinite paths in recursive marked trees. An  $\omega$ -tree  $T$  is a subset of  $\omega^*$  closed under left factors, represented by a corresponding set of numbers  $t = \{\tau^*((x_1, \dots, x_k)) \mid x_1 \dots x_k \in T\}$ .

The set of infinite paths in  $T$ , represented by the set of functions  $\{f \in \omega^\omega \mid \forall k, f[k] \in t\}$ , is  $\Pi_1^0$  if  $t$  is a recursive tree. A *recursive marked tree*  $(t, m)$  is a recursive tree  $t$  whose set of leaves is recursive and whose nodes are labelled by (possibly infinite) sets of numbers, fixed by a recursive predicate  $m \subseteq t \times \omega$ .

Infinite paths in a recursive tree may be seen as infinite computations in a recursive transition graph with a recursive set of sink states. Marks may be thought of as identifiers for computing agents and in the sequel, we shall interpret  $m(f[k], a)$  as the affirmation that agent  $a$  is *disabled* at the  $k$ th step in computation  $f$ . The complement  $\bar{m}$  of a marking predicate  $m$  is thus an enabling predicate.

Harel's language  $L$  is the union of an alternated hierarchy of languages  $L_i, L'_i$  constructed as follows. The basic language  $L_0$  provides four species of atomic formulas  $\exists a, \forall a, \exists^\omega a, \forall^\omega a$  where  $a$  ranges over the set of marks ( $a \in \omega$ ). For  $i \geq 0$ ,  $L'_i$  is the closure of  $L_i$  under finite conjunction and disjunction and under recursive  $\omega$ -conjunction, and  $L_{i+1}$  is the closure of  $L'_i$  under recursive  $\omega$ -disjunction. Formulas in  $L$  are interpreted over infinite pathes in recursive marked trees. An infinite path  $f$  in  $(t, m)$  satisfies the formula  $\exists a$ , respectively  $\exists^\omega a$ , if and only if  $m(f[k], a)$  for some  $k$ , respectively  $m(f[k], a)$  for infinitely many  $k$ . Formulas  $\forall a, \forall^\omega a$  are the duals of  $\exists a, \exists^\omega a$  and the logical connectives have the standard interpretation.

In a recursive marked tree  $(t, m)$ , the atomic formulas  $\forall a, \exists a, \forall^\omega a, \exists^\omega a$  are interpreted respectively by  $\Pi_1^0, \Sigma_1^0, \Sigma_2^0, \Pi_2^0$  sets of infinite paths, and each formula  $\varphi \in L'_0$  is therefore interpreted by a  $\Pi_3^0$  set of infinite paths. Conversely, any  $\Pi_3^0$  subset of  $\omega^\omega$  may be represented as  $\{f \mid \forall i, \forall k, m(f[k], i)\}$  for some recursive relation  $m$ , and thus coincides with the interpretation of the formula  $\bigwedge_i \forall^\omega i$  in the recursive marked tree  $(\omega, m)$ . Altogether, for  $n > 0$ , a subset of  $\omega^\omega$  is  $\Sigma_{2n+2}^0$  resp.  $\Pi_{2n+3}^0$  if and only if it coincides with the interpretation of some formula  $\varphi \in L_n$  resp.  $\varphi \in L'_n$  in some recursive marked tree  $(t, m)$ . Furthermore, the tree and the marking predicate are defined uniformly from the  $\Sigma_1^0$  or  $\Pi_1^0$  index of the set.

### *Logical expressions of fairness*

In view of the above, there are at least three equivalent ways of defining fairness in recursively enumerable transition graphs:

- i) state an arbitrary  $\Pi_3^0$  predicate acting as a filter on infinite computations,
- ii) state a recursive distance  $d$  on finite computations, refining the natural distance  $\delta$ , and define fair computations as natural limits of  $d$ -Cauchy sequences of finite computations,
- iii) state a recursive marking/enabling predicate  $m$  for finite computations, and rest on the general normal form  $F = \{f \mid \forall i. \forall k. m(f[k], i)\}$  for  $\Pi_3^0$  sets.

Furthermore, there is a uniform translation between any two types of definitions of fairness, and each type of definitions may be equipped with effective operators realizing the conjunction, disjunction, or  $\omega$ -conjunction of fairness conditions. For instance, the union of the derived sets of  $\omega$  for two different  $\Pi_1^0$  distances  $d_1$  and  $d_2$  on  $\Omega$  is again the derived set of  $\omega$  for some  $\Pi_1^0$  distance  $d_1 \vee d_2$  on  $\Omega$ . That specific property of  $\Pi_1^0$  distances is not trivial, and we have not heard about similar cases in topology.

### *$\omega$ -regular languages*

A well known theorem due to Mac Naughton states that the family of  $\omega$ -regular languages over  $A$  is the boolean closure of the family of deterministic  $\omega$ -regular languages over  $A$  [Mac Naughton]. Since any deterministic  $\omega$ -regular language is  $\Pi_2^0$ , any  $\omega$ -regular language is  $\Pi_3^0$ . Thus, any  $\omega$ -regular language over  $A$  is the derived set of  $A^*$  for some  $\Pi_1^0$  metric distance on  $A^\omega = A^* \cup A^\omega$ . This fact is not surprising, since recognition criteria used in Büchi/Müller automata are essentially fairness conditions.

### ***A metric characterization of $F_{\sigma\delta}(\Pi_3^0)$***

There exists a tight relationship between the classical Borel hierarchy of sets of functions and the effective Kleene hierarchy, see for instance [Rogers, p.356] and [Moschovakis, p.160]. Namely, if  $\Gamma$  is a (lightface) pointclass in the effective hierarchy and  $\Gamma$  is the corresponding (boldface) pointclass in the classical hierarchy, then:

$$\Gamma = \bigcup_{\varepsilon \in \omega^{(\omega)}} \Gamma[\varepsilon],$$

where a set of functions is  $\Gamma[\varepsilon]$  iff it is  $\Gamma$  *relative* to the oracle  $\varepsilon$ . More precisely,  $\Sigma_n^0[\varepsilon]$ -forms and  $\Pi_n^0[\varepsilon]$ -forms are defined in the same way as  $\Sigma_n^0$ -forms and  $\Pi_n^0$ -forms, except that recursive relations are replaced by relations *recursive in  $\varepsilon$* .

Thus, by relativizing theorem 1 to an arbitrary oracle  $\varepsilon$ , we get that a set of functions is  $\Pi_3^0$  if and only if it is the derived set of  $\omega$  for some  $\Pi_1^0$  ultra-metric on  $\Omega$ , refining the natural metric  $\delta$ . A more accurate characterization of  $\Pi_3^0$  sets may be given in terms of *inductive* distances on  $\Omega$ , defined as follows: a distance  $d$  on  $\Omega$  is inductive if it takes values in  $\{0\} \cup \{1/n \mid n \in \omega\}$ , is weakly continuous in the sense that  $d(\alpha, \beta) = \lim_k d(\alpha[k], \beta[k])$ , and has only  $\delta$ -closed  $d$ -balls.

**Theorem 3** For  $F \subseteq \omega^{(\omega)}$  the following assertions 1, 2 and 3 are equivalent:

- (1)  $F$  is a  $\Pi_3^0$  set (or  $F$  is  $F_{\sigma\delta}$ ),
- (2)  $F$  is the derived set of  $\omega$  for some inductive distance on  $\Omega$ ,
- (3)  $F$  is the derived set of  $\omega$  for some inductive ultra-metric refining  $\delta$ .

proof First of all, let us recall that the class of  $\delta$ -closed sets is just  $\Pi_1^0$ , (see [Rogers, p.342] or [Moschovakis, p.20]). The proof is then straightforward.

Suppose (1), then  $F$  is  $\Pi_3^0[\varepsilon]$  for some  $\varepsilon \in \omega^{(\omega)}$ . A remake of the proof for theorem 1 relative to  $\varepsilon$  shows that  $F$  is the derived set of  $\omega$  for some  $\Pi_1^0[\varepsilon]$  distance  $d$  on  $\Omega$ , taking values  $(1/n)$  and furthermore weakly continuous. Now, for each  $\alpha$ , each  $d$ -ball  $\{\beta \mid d(\alpha, \beta) < 1/n\}$  is  $\Pi_1^0[\gamma]$  for some  $\gamma$  determined from  $\varepsilon$  and  $\alpha$ , hence it is  $\delta$ -closed. Thus (1)  $\Rightarrow$  (2).

Suppose (2). There suffices to show that  $d(\alpha, \beta) < 1/n$  is  $\Sigma_2^0$  in  $(\alpha, \beta, n)$ , which entails (1). By definition of an inductive distance,  $d(\alpha, \beta) < 1/n \Leftrightarrow \exists i. \forall j \geq i. d(\alpha[j], \beta[j]) < 1/n$ . For  $x \in \omega$ , each  $d$ -ball  $\{\beta \mid d(x, \beta) < 1/n\}$  is closed and thus  $\Pi_1^0[\varepsilon_{x,n}]$  for some  $\varepsilon_{x,n}$  in  $\omega^\omega$ . Hence  $d(x, y) < 1/n$  is  $\Pi_1^0[\varepsilon]$  for some global oracle  $\varepsilon$  gathering the  $\varepsilon_{x,n}$ , and  $d(\alpha, \beta) < 1/n$  is  $\Sigma_2^0[\varepsilon]$  and thus  $\Sigma_2^0$  in  $(\alpha, \beta, n)$ . Thus (2)  $\Rightarrow$  (1).

Finally, (2)  $\Leftrightarrow$  (3) by the relativized version of theorem 1.  $\square$

An elegant proof for (1)  $\Rightarrow$  (3) by purely topological treatment was devised by A. Arnold, yielding an independent proof of the theorem, as (3)  $\Rightarrow$  (1) is immediate. We sketch here the construction of the distance. Let  $F$  be a  $\Pi_3^0$ -subset of  $\omega^\omega$ , thus  $F = \{f \mid \forall i. \exists j. \forall k. R(f, i, j, k)\}$  for some relation  $R$  recursive in  $\varepsilon$  for some oracle  $\varepsilon$ . Define  $F_{i,j} = \{f \mid \exists j' \leq j. \forall k. R(f, i, j', k)\}$  and  $F'_{i,j} = F_{i,j} \cup \{x \in \omega \mid \exists f. x \sqsubset f \in F_{i,j}\}$  where  $x \sqsubset f$  if  $x$  is a prefix of  $f$ , i.e.  $x = \emptyset$  or  $x = \tau^*((f(0), \dots, f(k)))$  for some  $k \geq 0$ . For each  $i \in \omega$ , the  $F'_{i,j}$  form an increasing sequence of  $\delta$ -closed subsets of  $\Omega$ . Now for any  $\alpha \in \Omega$ , define  $g_\alpha(i) = \inf\{j \mid \alpha \in F'_{i,j}\}$ , where  $\inf(\emptyset) = \infty$  and  $\infty \notin \omega$ , then  $g_\alpha(i) = \lim_k (g_{\alpha[k]}(i))$ , and  $(\alpha \in F)$  if and only if  $(\alpha \in \omega^\omega)$  and  $g_\alpha \in \omega^\omega$ . The distance  $d(\alpha, \beta)$  is finally defined as  $1/(1 + \Delta(\alpha, \beta))$  letting  $\Delta(\alpha, \alpha) = \infty$  and for  $\alpha \neq \beta$ :  $\Delta(\alpha, \beta) = \max\{n \leq \alpha \wedge \beta \mid g_\alpha[n] = g_\beta[n] = g_{\alpha \cap \beta}[n] \in \omega^n\}$  where  $\alpha \cap \beta = \alpha \upharpoonright \alpha \wedge \beta = \beta \upharpoonright \alpha \wedge \beta$ . Then  $d$  is indeed an inductive ultra-metric and  $F$  is the derived set of  $\omega$  for that metric.

**Acknowledgements** We thank A. Arnold and A. Louveau for their contributions.



## References

- [Arnold-Nivat] A. Arnold and M. Nivat. Metric interpretations of infinite trees and semantics of non deterministic recursive programs. In *Theoretical Computer Science* **11**, pp. 181-205, 1980.
- [Austry-Boudol] D. Austry and G. Boudol. Algèbre de processus et synchronisation. In *Theoretical Computer Science* **30**, pp. 91-131, 1984.
- [de Bakker-Meyer] J.W. de Bakker and J.J. Meyer. Order and metric in the stream semantics of elemental concurrency. In *Acta Informatica* **24**, pp. 491-511, 1987.
- [Barendregt] H.P. Barendregt. *The Lambda Calculus, its Syntax and Semantics*. North-Holland, 1984.
- [Costa] G. Costa. A metric characterization of fair computations in CCS. In *Lecture Notes in Computer Science* **186**, pp.239-251, 1985.
- [Degano-Montanari] P. Degano and U. Montanari. Liveness properties as convergence in metric spaces. In *STOC*, pp.31-38, 1984.
- [Dugundji] J. Dugundji. *Topology*. Allyn and Bacon, 1966.
- [Eilenberg] S. Eilenberg. *Automata, Languages, and Machines*. Academic Press, 1974.
- [Francez] N. Francez. *Fairness*. Springer, 1986.
- [Harel] D. Harel. Effective transformations on infinite trees with application to high undecidability, dominoes, and fairness. In *JACM* **33**, pp.224-248, 1986.
- [Hopcroft-Ullman] J.E. Hopcroft and J.D. Ullman. *Introduction to Automata Theory, Languages, and Computation*. Addison-Wesley, 1979.
- [Keller] R. Keller. Formal verification of parallel programs. In *CACM* **19**, pp. 371-384, 1976.
- [Landweber] L. Landweber. Decision problems for  $\omega$ -automata. In *Mathematical System Theory* **4**, pp. 376-384, 1969.
- [Milner] R. Milner. *A Calculus of Communicating Systems*. Springer, 1980.
- [MacNaughton] R. Mac Naughton. Testing and generating infinite sequences by a finite automaton. In *Information and Control* **9**, pp. 521-530, 1966.
- [Moschovakis] Y. Moschovakis. *Descriptive Set Theory*. North-Holland, 1980.
- [Rogers] H. Rogers. *Theory of recursive functions and effective computability*. Mc Graw Hill, 1967.

