



## Enumeration of bordered words

Mireille Regnier

► **To cite this version:**

| Mireille Regnier. Enumeration of bordered words. RR-0956, INRIA. 1988. <inria-00075603>

**HAL Id: inria-00075603**

**<https://hal.inria.fr/inria-00075603>**

Submitted on 24 May 2006

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# INRIA

UNITE DE RECHERCHE  
INRIA-ROCOUENCOURT

Institut National  
de Recherche  
en Informatique  
et en Automatique

Domaine de Voluceau  
Rocquencourt  
BP 105  
78153 Le Chesnay Cedex  
France  
Tel (1) 39 63 55 11

## Rapports de Recherche

N° 956

*Programme 1*

### ENUMERATION OF BORDERED WORDS

Mireille REGNIER

Décembre 1988



2959

# ENUMERATION OF BORDERED WORDS ENUMÉRATION DES MOTS À BORDS

Mireille REGNIER

**Abstract:** We consider here the family of bordered words on a  $q$ -ary alphabet, i.e. the words  $bwb$ . We also consider the  $k$ -bordered words. We enumerate such words, using generating functions, and derive asymptotic estimates by the Darboux method. In particular, we prove that the density of  $k$ -bordered words is  $\alpha_k$ ,  $\alpha_k \neq 0$ .

**Résumé:** Nous étudions dans ce papier la famille des mots à 1 bord sur un  $q$ -alphabet, i.e. les mots de la forme  $bwb$ . Nous considérons aussi les mots à  $k$  bords. Nous dénombrons ces mots, en calculant les séries génératrices associées. Puis nous obtenons des estimations asymptotiques par la méthode de Darboux. En particulier, nous prouvons que la densité des mots à  $k$  bords est non nulle.

# ENUMERATION OF BORDERED WORDS

Le langage de la vache-qui-rit †

Mireille REGNIER  
INRIA-Rocquencourt  
78 153-Le Chesnay (FRANCE)

**Abstract:** *We consider here the family of bordered words on a  $q$ -ary alphabet, i.e. the words  $bwb$ . We also consider the  $k$ -bordered words. We enumerate such words, using generating functions, and derive asymptotic estimates by the Darboux method. In particular, we prove that the density of  $k$ -bordered words is  $\alpha_k$ ,  $\alpha_k \neq 0$ .*

## I. INTRODUCTION:

This note is devoted to 1-bordered words on a  $q$ -ary alphabet  $A$ , which are to be counted. A 1-bordered word  $w$  is defined as a word:  $bw'b$  where  $b$  and  $w'$  are in  $A^*$ , and  $b$ , the border, is non empty. For example,  $w = 101.1.101$ . One can define recursively the set  $B_{k+1}$  of the  $k+1$ -bordered words:  $w$  is in  $B_{k+1}$  if  $w$  is in  $B_1$ , and if its largest side is in  $B_k$ . For example,  $w$  is in  $B_2$  as  $b = 101$  is in  $B_1$ . To count the words in sets  $B_k$ , we make use of the associated generating functions  $B_k(z)$ . The scheme is the following: we first establish functional equations satisfied by the series  $B_k(z)$ . To do so, we need a unique representation of the words in  $B_k$ . This part makes use of the general theorems in combinatorics on words. To get asymptotics for the coefficients, we do not need to solve the functional equations. We study the singularities of  $B_k(z)$ , that appear to be polar singularities. Then one can apply the Darboux theorem, and prove that the number of  $k$ -bordered words of length  $n$ ,  $b_n^k$ , satisfies  $b_n^k \sim \alpha_k \cdot q^n$ , where  $\alpha_k$  is computable, with any given precision, from the functional equation.

In Section II, we list some general theorems in combinatorics to be used in the following sections. In Section III, we introduce our techniques on the set  $B_1$ . In Section IV, we consider the more general and intricate case of the set  $B_2$ . In Section V, we extend the methods and results of Section IV to the general case of sets  $B_k$ . In Section VI, we deal with the asymptotics of  $b_n^k$ .

---

† This subtitle refers to the famous Rabier's commercial drawing: a cow with two identical © "Vache-Qui-Rit" boxes as ear-rings. Inside each cheese box, a cow with two earrings...



**Definition 4:** A word  $x \in A^*$  is said to be **primitive** if it is not a power of another word. Let  $P$  be the set of primitive words.

**Theorem B:** If

$$x^n = y^m, \quad x, y \in A^*, n, m > 0,$$

there exists a word  $z$  such that  $x, y \in z^*$ .

In particular, for each word  $w \in A^+$ , there exists a unique primitive word  $x$  such that  $w \in x^*$ .

**Theorem C:** Two words  $x, y \in A^+$  commute iff they are powers of the same word. More precisely the set of words commuting with a word  $x \in A^+$  is a monoid generated by a single primitive word.

### III 1-BORDERED WORDS:

We first state our definition of 1-bordered words.

**Definition 5:** Let  $A$  be a  $q$ -ary alphabet. Let  $B_1$  be the set of words  $w$  of the form:  $bvb$  where  $v \in A^*, b \in A^+$ . One says that  $w$  is a **bordered word** and that  $b$  is a **border** of  $w$ . One notes  $S = A^* - B_1$  its complement, the set of **unbordered words**.

**Remarks:**

- (i) The empty word and the words of length 1 are in  $S$ .
- (ii) This definition does not allow overlaps. Accordingly, 0101 is a side of 01010101 but 010101 is not.

**Example:**  $w = 1011101$  is a 1-bordered word on a binary alphabet. The words  $b_1 = 1$  and  $b_2 = 101$  are both borders of  $w$ .

We see on this example that a word in  $B_1$  may have several borders. In order to get a *unique representation* of words in  $B_1$  (and more generally in  $B_k$ ), we need to determine the relationship between different sides of a given word. This problem of deciding a unique representation is fairly general in combinatorics on words. Some interesting examples can be found in [CP88], [DA1878] or [OD85].

**Theorem 1:** Let  $b$  and  $b'$  be two borders of a same word  $w$ , with  $|b'| > |b|$ . Then there exists a primitive word  $z$  in  $A^*$ , and a factorization:  $z = ae$  such that:

$$\begin{cases} b = e.z^p, p \geq 0 \\ b' = e.z^q, q > p \end{cases}$$

**Example:** Consider the 1-bordered word  $w = 1011101$ . The words  $b' = 101$  and  $b = 1$  are two different borders. They satisfy the relationship above with:  $z = 01, e = 1, a = 0$ .

**Proof:** This is a consequence of Theorem A. As:  $w = bxb = b'x'b'$ , one has:  $b' = bu = vb$ , and  $u$  and  $v$  are conjugate. It is also worth giving a direct (and therefore intuitive) proof. We note:

$$\begin{cases} b = b_0 \\ (n-1)|u| < |b| \leq n|u|. \end{cases}$$

$$\text{If } n=1 \text{ then } \begin{cases} |b| \leq |u| = |v| \\ v = bv' \\ b' = b.(v'b) = b.z^q = e.z^p.z^q \end{cases}$$

$$\text{else } \begin{cases} b = e_u.u^{n-1} = e_z.z^p \\ b' = e.z^p.u = e.z^q \end{cases}$$

**Proposition 2:** Any word  $w$  in  $B_1$  can be written, in a single way, as:  $sw's$ , where  $s$  is in  $S^*$  and  $w'$  in  $A^*$

**Proof:** The smallest border is in  $S$ , and is the only border in  $S$ .

As a corollary of Proposition 2, we get the functional equation:

$$B_1(z) = [S(z^2) - 1].W(z) \quad (1).$$

We use the methods developed in [GJ83] and [FL84]. The concatenation  $sw'$  of two different words translates in the product of the generating functions counting these words. The repetition of the word  $s$  is taken into account by squaring  $z$  in the corresponding generating function  $S$ . Moreover, we have exactly  $q^n$  words of length  $n$ . Thus:

$$W(z) = \sum_{n \geq 0} q^n z^n = \frac{1}{1 - qz}.$$

As:  $B_1(z) + S(z) = W(z)$ , Proposition 3 follows.

**Proposition 3:** The series  $B_1$  and  $S$  satisfy the functional equations:

$$\begin{cases} S(z) = \frac{2 - S(z^2)}{1 - qz} \\ B_1(z) = \frac{1}{1 - qz} \left[ \frac{qz^2}{1 - qz^2} - B_1(z^2) \right] \end{cases}$$

**Remark:** Such an approach where combinatorial constructions translate into functional properties of generating functions is quite powerful. A general framework can be found in [FL84]. An example, related to the analysis of the Knuth-Morris-Pratt algorithm, can be found in [RE88].

In Section VI, we will derive asymptotics for the coefficients of these functions.

#### IV. 2-BORDERED WORDS:

In this Section we generalize the scheme of the previous Section. We have seen that any word in  $B_1$  can be written, in a unique manner, under the form:  $sws$ . We define here a minimal set  $G_2$  such that any word in  $B_2$  can be written, in a unique manner:  $g_2wg_2$ . Then, denoting  $G_2(z)$  (resp.  $B_2(z)$ ) the generating function counting the words in  $G_2$  (resp. in  $B_2$ ), one has:

$$B_2(z) = G_2(z^2).W(z).$$

Thus, we first state the definition of the minimal set  $G_2$ , extended for all the sets  $B_k$ . Then, we will characterize the elements of  $G_2$ . Finally, we will derive a functional equation on an associated generating function.

**Definition 5:** A bordered word  $x$  is said to be  $k$ -minimal if:

(i)  $x \in B_k$

(ii)  $w \preceq x, w \subseteq x, w \in B_k \Leftrightarrow w = x$

This subset of  $B_k$  is noted  $G_{k+1}$ .

**Example:**  $x = 1001001 \in B_1$ . As  $1001 \subseteq x$  and  $1001 \preceq x$ ,  $x \notin G_2$ , but  $1001 \in G_2$ .

**Remark:** A word in  $G_k$  has no side in  $B_{k-1}$ . And any word  $x$  in  $B_{k-1}$  contains exactly one side in  $G_k$ . Hence, we define:

**Definition 6:** We note  $g(b_{k-1})$  the unique  $k$ -minimal word such that:

$$g(b_{k-1}) \preceq b_{k-1} \text{ and } g(b_{k-1}) \subseteq b_{k-1}.$$

We can get now an equivalent to Proposition 2:

**Theorem 4:** Any  $k$ -bordered word  $b_k$  in  $B_k$  can be written, in a single way, as:

$$b_k = g_k w g_k$$

with  $g_k \in G_k$ .

**Hint:** Let  $b_{k-1}$  be the largest border of  $b_k$  and choose  $g_k = g(b_{k-1})$ .

Our aim is now to *enumerate* the set  $G_k$ . To do so, we first derive some characterizations of the words  $g_k$ . For a sake of clarity, we first consider the case of  $G_2$ .

Our characterization of  $G_2$  will rely on the Lemmas of 1-factorization and 2-factorization.

**Lemma 5 (Lemma of 1-factorization:)** Let  $x = z^*$ ,  $z \in P$ , be a word in  $L^*$ . Then, if a word  $s \in S$  is a right (resp. left) factor of  $x$ , then  $s$  is a right (resp. left) factor of  $x$ .



**Proof:** If  $s \not\subseteq z$ , then there exists  $\alpha \subseteq z$  (or  $z = \beta\alpha$ ),  $\alpha \neq \epsilon$  such that:  $s = \alpha.z^m$ ,  $m \geq 1$ . Thus:  $s = \alpha.(\beta\alpha)^{m-1}\beta.\alpha$  is not in  $S$ .

**Remark 6:** A word in  $S$  cannot overlap with himself.

As  $s = zu = vz$  implies (Theorem A):  $s = ea.e.(ae)^*$  which is not in  $S$ .

We know that any word in  $B_1$  can be written  $sas$ . Proposition 8 and Lemma L, a generalization of Lemma 5, characterize the words  $sas$  that are in  $G_2$ .

**Lemma 7 of 2-factorization** Let  $g_2 = sas$  be a word in  $G_2$ , and  $x = wsas \in z^*$ , with  $z \in P$ . Then:

$$z = \begin{cases} as \\ usas \end{cases}$$

**Example:** Let us show what happens for a word in  $B_1$ , but not in  $G_2$ , such as: 1.0100101. We have:  $s = 1$  and  $as = 0100101 \in P$ . Let  $w = 00$ . Then  $x = wsas = 00101.00101$  and  $z = 00101$ ; here  $z \neq as$  and  $sas \not\subseteq z$ .

**Proposition 8 :** Let  $H_1$  be the subset of  $B_1$ :

$$H_1 = \{sas; s \in S, as \in P\}.$$

Then:

(i)  $G_2 \subset H_1 \subset B_1$

(ii) Any  $b_1 \in B_1$  can be written:

$$\begin{cases} b_1 = s(as)^m \\ b_1 = sas.(usas)^m, usas \in P \end{cases}$$

where  $s$  and  $a$  are defined by:  $g(b_1) = sas$ .

**Proof:**

(i) From Proposition 2, one can write  $g_2 = sa's$ . If  $as \notin P$ , the Lemma of 1-factorization implies:  $g_2 = s(as)^m$ , and  $g_2$  is not minimal.

(ii) The proof of (ii) is deferred to the Appendix. It uses the important Remark 6 and Lemma 7 of 2-factorization, whose proof is also in Appendix. Note that the second form implies that  $b_1$  is in  $B_2$ , as  $sas.usas$  is.

We can turn now to the study of the generating function of 1-bordered words. We prove:

**Theorem 9:** Let  $G_2(x, z) = \sum_{g_2=sas \in G_2} x^{|s|} z^{|as|}$  be the bivariate generating function of the set  $G_2$ . It satisfies the functional equation:

$$[S(zx) - G_2(zx, z^2)].W(z) = G_2(x, z) + G_2(x, z^2) \quad (2)$$

and we have:

$$B_2(z) = G_2(z^2, z^2)W(z). \quad (3)$$

**Proof:** We define a 3-variate generating function associated to  $H_1 - B_2$ :

$$\phi(x, z, t_3) = \sum x^{|s|} z^{|as|} t_3^{|usas|} 1_{sas \in G_2} 1_{usas \in P}$$

We shall write two equations in  $\phi$  and  $G_2$  and eliminate  $\phi$  from them. We have:

$$\sum_{\substack{b_1 \in B_1 \\ b_1=sws}} x^{|s|} z^{|ws|} = S(zx)W(z).$$

From Proposition 8, this is also:

$$\begin{aligned} & \sum_{g_2 \in G_2} \left( \sum_{m \geq 1} x^{|s|} z^{m|as|} + \sum_{usas \in P} \sum_{m \geq 1} x^{|s|} z^{|as|} z^{m|usas|} \right) \\ &= \sum_{m \geq 1} G_2(x, z^m) + \sum_{m \geq 1} \phi(x, z, z^m). \end{aligned}$$

Hence:

$$S(zx)W(z) = \sum_{m \geq 1} G_2(x, z^m) + \sum_{m \geq 1} \phi(x, z, z^m) \quad (4)$$

To get the equation (5), we consider the words  $g_2 w g_2 = sas.w.sas \in B_2$ . From the Lemma 7 of 2-factorization, we get:

$$\sum x^{|s|} z^{|as|} t^{|wsas|} 1_{sas \in G_2} 1_{w \in A^*} = \sum_{m \geq 3} G_2(x, zt^m) + \sum_{m \geq 1} \phi(x, z, t^m).$$

Hence:

$$G_2(xt_3, zt_3)W(t_3) = \sum_{m \geq 3} G_2(x, zt_3^m) + \sum_{m \geq 1} \phi(x, z, t_3^m). \quad (5)$$

Eliminating  $\phi$  from (4) and (5) yields  $G_2$ .

## V. PROPERTIES OF $k$ -BORDERED WORDS

In this section, we extend the scheme of Section IV to the general case of  $k$ -bordered words. We first study the set  $G_k$ , and prove a Lemma of  $k$ -factorization. Then, we associate to  $G_k$  a  $k$ -variate

generating function  $G_k(t_k, \dots, t_1)$ . We prove that  $G_k(t_k, \dots, t_1)$  and  $G_k(z)$  satisfy equations similar to (2) and (3).

We first derive some properties of  $G_k$ .

**Proposition 10:** *Let  $g_k$  be in  $G_k$ . It can be factorized as:*

$$g_k = g_{k-1} \cdot p_{k-1}^{\theta_k} = \dots = g_i \cdot p_i^{\theta_i} = \dots = g_1 \cdot p_1^{\theta_1}$$

with  $g_i \in G_i, p_i \in P$ .

**Theorem 11 ( $k$ -overlapping):** *Let  $g_k$  be a word in  $G_k$  and  $x = z^*, z \in P$  a word in  $L^*$  such that:*

$$g_k x = y g_k, y \in L^* .$$

*Then:  $z = w g_k$  or  $z \in \{p_1, \dots, p_{k-1}\}$ .*

**Theorem 12:** *For any  $i$ , this decomposition is unique and has the following property:*

$$\text{If } g_i \subseteq p_i \text{ then } \begin{cases} \exists j \geq i : g_i \cdot p_i \in B_j - B_{j+1} \\ \theta_i = 2^{k+1-j} - 1. \end{cases}$$

$$\text{else } \begin{cases} \exists m, m \leq i : g_m \subseteq p_i = p_m . \\ \theta_m = 2^{k+1-j} - 2^{i+1-j} \end{cases}$$

**Examples:**

k=1:  $g_1 = s \in S$ .

k=2:  $g_2 = s.as, s \in S, as \in P$ .

k=3:  $g_3 = s(as)^3$  or  $sas.usas$ . We can rewrite:

$$\begin{cases} sas.(as)^2 = s.(as)^3 \\ sas.usas = s.(asusas). \end{cases}$$

k=4:  $g_4 = s(as)^7$  or  $s(as)^3.ws(as)^3$  or  $sasusas.wsasusas$  or  $sas.(usas)^3$  or  $s.(asusas)^3$ . We can rewrite, for example:

$$s.(asusas)^3 = sas.(usas.asusas.asusas) = sasusas.asasusasusas$$

We have:  $g_1 \cdot p_1 \in B_2 - B_3$  and  $g_2 \cdot p_2, g_3 \cdot p_3 \in B_3$ .

**Proof of Proposition 10:** As  $g_k \in B_{k-1}$ , its largest border  $b$  satisfies:

$$b \in B_{k-2} \subset B_{k-1} \subset \dots \subset B_1 .$$

Thus, there exists a sequence:  $g_{k-1}, g_{k-2}, \dots, g_1$  such that:

$$\begin{cases} g_1 \preceq g_2 \preceq \dots \preceq g_{k-2} \preceq g_{k-1} \preceq g_k \\ g_1 \subseteq g_2 \subseteq \dots \subseteq g_{k-2} \subseteq g_{k-1} \subseteq g_k \end{cases}$$

and we get the factorization.

**Proof of Theorem 11:** We know by Remark K that Theorem 11 holds for  $g_1$  in  $G_1$ . Assume now that it holds true for  $G_1, \dots, G_{k-1}$ . If  $z \neq wg_k$  we may assume  $x = z$ : as we have then:  $z \subset g_k$ , thus  $g_k = e_z \cdot z^p$  with  $e_z \subseteq z$  and  $p \geq 1$ . We get the equation:

$$g_k \cdot z = e_z \cdot z^{p+1} = y \cdot e_z \cdot z^p = y \cdot g_k.$$

From Remark K, we know that  $g_1 \subseteq x$ . Thus, we may define  $i, 1 \leq i \leq k-1$  by:

$$g_i \subseteq x \subset g_{i+1}.$$

We have then:  $g_{i+1}x = y'g_{i+1}$ , with  $|x| < |g_{i+1}|$ . Applying Theorem 11, we have:

$$\begin{cases} g_{i+1} = g'_m \cdot p'_m \\ x = p'_m \end{cases}$$

and

$$\left\{ \begin{array}{l} g'_m \subseteq g_{i+1} \subseteq g_k \\ g'_m \preceq g_{i+1} \preceq g_k \end{array} \right\} \Rightarrow g_m = g'_m.$$

Now, if  $i = k-1$ , we have:  $g_{k-1} \subseteq x \subseteq g_k$ . From above, we know that:  $g_k = g_m p'_m = g_m \cdot x^*$ . To prove in all other cases that  $p'_m \in \{p_1, \dots, p_{k-1}\}$ , we remark that:

$$\begin{cases} x \subseteq g_k \Rightarrow g_k = e \cdot x^p, e \subset x \\ g_i a g_i = g_i \cdot x = g_{i+1} \preceq g_k \Rightarrow g_k = g_i \cdot x^m a, a \prec x \end{cases}$$

Moreover, we have  $p \geq 1$  (as  $x \subseteq g_{k-1}$ ) and  $m \geq 1$  (as  $g_{i+1} \subseteq g_{k-1} \Rightarrow 2|g_{i+1}| \leq |g_k|$ ). Then:

$$xa = ax$$

or (Theorem D):  $a = \epsilon$  and  $g_k = g_i \cdot x^p = g_i \cdot p_i^{\theta_i}$ , hence:  $x = p_i$ .

**Proof of Theorem 12:** If  $p_i \not\subseteq g_i$  then  $|p_i| > |g_i|$ . As  $g_i \subseteq g_i \cdot p_i^{\theta_i}$ , we get  $g_i \subseteq p_i$ . Then:

$$g_i \cdot p_i = g_i \cdot w_i g_i \in B_i \subset B_{i+1} \subset \dots \subset B_j \subset B_{j+1} \subset \dots$$

The expression of  $\theta_i$  follows immediately from the remark that the largest border of  $g_i(w_i g_i)^m$ ,  $m \geq 1$  is  $g_i(w_i g_i)^{\lfloor \frac{m-1}{2} \rfloor}$ .

Now, if  $p_i \subset g_i$  the equation:  $g_k = g_i p_i^* = y g_i$  implies:

$$\begin{cases} g_i = g'_m \cdot p_m^* = g_m p_m^\theta, \theta = 2^{i+1-j} - 1 \\ p_i = p'_m. \end{cases}$$

Hence:  $g_k = g_i \cdot p_i^* = g_m \cdot p_m^\theta \cdot p_m^{\theta_i}$  with:  $\theta_i = (2^{k+1-j} - 1) - \theta$ .

We can now draw a scheme that generalizes the derivation of  $G_1$  and  $G_2$  in the previous sections. We define some multivariate generating functions.  $\psi_{i,k}(t_1, \dots, t_i, t_{i+1}, \dots, t_k)$  counts  $H_{i,k} = \{g_i \cdot p_i \mid g_i \in G_i, p_i \in P, g_i \subseteq p_i, g_i \cdot p_i \in B_k - B_{k+1}\}$ . And  $\phi_{i,k+1}(t_1, \dots, t_i, t_{i+1}, \dots, t_k)$  counts  $L_{i,k+1} = \{g_i \cdot p_i \mid g_i \in G_i, p_i \in P, g_i \subseteq p_i, g_i \cdot p_i \in B_{k+1}\}$ . One has:  $\psi_{i,j} = \phi_{i,j} - \phi_{i,j+1}$ . Then:  $G_k(t_1, \dots, t_k) = \sum_{i=1}^k \psi_{i,k}(t_1, \dots, t_k) + f(\{\phi_{i,j}\}_{1 \leq j \leq k-1})$ . For any  $k$ , counting  $\{g_i w g_i\}$  and applying Theorem 12 yields  $k$  functional equations. These equations involve:  $2k$  unknown, but dependent, functions  $(\phi_{i,k+1})_{1 \leq i \leq k+1}$  and  $(\psi_{i,k})_{1 \leq i \leq k}$ . One can derive  $(\phi_{i,k+1})_{1 \leq i \leq k+1}$  from these equations and hence:  $\psi_{i,k} = \phi_{i,k} - \phi_{i,k+1}$ . Finally, we get  $G_{k+1}$ . We can also relate these notations to the ones in Section IV:

$$\begin{aligned} \phi(t_1, t_2, t_3) &= \phi_{2,2}(t_1, t_2, t_3) \\ \phi(t_1, t_2, t_2) &= \phi_{1,2}(t_1, t_2) = \phi_{2,2}(t_1, t_2, t_3) \end{aligned}$$

We also have:  $\phi_{1,1}(t_1, t_2) = G_2(t_1, t_2)$ . Equation (4) counts  $\{g_1 w g_1\}$  and Equation (5) counts  $\{g_2 w g_2\}$ .

## VI. ASYMPTOTICS ON $k$ -BORDERED WORDS

The equations derived in the previous sections are rather involved, and cannot be solved explicitly. To derive asymptotics on the coefficients, we study the singularities of the generating functions and use the Darboux Theorem [DA1878]. Examples of this approach are developed in [ST84].

**Darboux Theorem:** *Let  $f(z)$  be some complex function, analytic for  $|z| < \rho$ , with a single singularity,  $z = \rho$ , on its circle of convergence. If it can be continued as:*

$$f(z) = g(z) \cdot \left(1 - \frac{z}{\rho}\right)^{-s} + h(z)$$

where  $g$  and  $h$  are analytic and  $s$  ranges in  $\mathcal{R} - \{0, -1, -2, \dots\}$ , then:

$$f_n = [z^n]f(z) = \rho^{-n} n^{s-1} \frac{g(s)}{\Gamma(s)} \left(1 + O\left(\frac{1}{n}\right)\right).$$

In our case,  $W(z) = \frac{1}{1-qz}$  has a unique singularity around  $z = q$ . Moreover, each  $B_k(z)$  is the product of  $W(z)$  by a, possibly intricate, generating function analytic around  $z = 1/q$  (precisely, for  $|z| \leq \frac{1}{q^2}$ ). This can be seen in Equations (1) and (2).

**Theorem 13:** Let  $B_1(z) = \sum b_n^1 z^n$  be the generating function of the bordered words on a  $q$ -ary alphabet  $A$ . Then:

$$b_n^1 \sim \alpha_1 q^{-n}$$

where:  $\alpha_1 = S(\frac{1}{q^2})$ .

**Proof:** We know that:  $W(z) = S(z) + B_1(z)$ . Thus,  $S$  and  $B_1$  are both defined at least for  $|z| < \frac{1}{q}$ . Hence  $S(z^2)$  is analytic for  $|z| < \frac{1}{\sqrt{q}}$ , notably around  $z = 1/q$ . We apply the Darboux theorem to equation (1). Note that  $S(\frac{1}{q^2})$  can be numerically computed as:

$$S(z) = 2 \sum_{k \geq 1} \frac{(-1)^k}{(1-1/q) \dots (1-q^{1-2^k})}$$

The different values of  $\alpha_1$  for  $q = 2, 3, 10$  are given in Table 2.

q	$\alpha_1$
2	0.8638659
3	0.4617496
10	0.1101101

Table 1

**Theorem 14:** Let  $B_k(z) = \sum_n b_n^k z^n$  be the generating function of the  $k$ -bordered words. Then:

$$b_n^k \sim \alpha_k q^n$$

where:  $\alpha_k = G_k(\frac{1}{q^2}, \dots, \frac{1}{q^2})$ .

**Proof:** The definition of  $G_k$  implies:

$$B_k(z) = G_k(z^2, \dots, z^2) \cdot W(z).$$

A priori,  $G_k(z, \dots, z)$  is analytic around  $z = \frac{1}{q^2}$ . It allows for the application of the Darboux theorem with  $s = 1$  and  $g(z) = G_k(z^2, \dots, z^2)$ .

## CONCLUSION

In this paper, we have considered the  $k$ -bordered words. In an algebraic part, we use general results in combinatorics on words to define a unique representation of  $k$ -bordered words. In particular, we introduce and characterize *minimal*  $k$ -bordered words. Then, we show how these constructions

translate into functional equations satisfied by the associated generating functions. Finally, we show that these equations need not to be solved (the solutions are intricate) and get directly asymptotic estimates on the number of  $k$ -bordered words. We prove that there are  $\alpha_k q^n$   $k$ -bordered words, or equivalently, that the density of the family of  $k$ -bordered words is always non-zero. The constant  $\alpha_k$  can be computed for any  $k$  from the functional equation, and is explicitly given, for various  $q$ , when  $k = 1$ . Such methods also apply to other combinatorial problems on words.

#### References:

- [CP88] M. CROCHEMORE AND D. PERRIN "Pattern Matching in Strings" in LITP (Paris) Research Report 88-5.
- [DA1878] G. DARBOUX "Mémoire sur l'approximation des fonctions de très grands nombres, et sur une classe étendue de développements en série" in *J. de Mathématiques pures et appliquées 3-ème série VI*, (1878) 1-56.
- [FL84] PH. FLAJOLET "*Mathematical Methods in the Analysis of Algorithms and Data Structures*", Lecture Notes for *A Graduate Course on Computation Theory*, Udine(Italy) 225-304(1984).
- [GJ83] I. GOULDEN AND D. JACKSON "*Combinatorial Enumerations*", Wiley, New York, (1983).
- [GO81] L.J. GUIBAS AND A.M. ODLYZKO "String overlaps, pattern matching, and nontransitive games" in *J. Comb. Theory (A)*, **30** (1981) 183-208.
- [LO83] LOTHAIRE "*Combinatorics on Words*", Addison-Wesley, Reading, Mass., (1983).
- [OD85] A.M. ODLYZKO "*Periodicities in Strings*" in *Combinatorial Algorithms on Words*, Springer NATO ASI Ser. F12, (1985) 241-254.
- [RE88] M. RÉGNIER "*Knuth-Morris-Pratt Algorithm: an Analysis*" Actes des Journées Informatique et Mathématique, Marseille, (1988).
- [ST84] J.M. STEYAERT "*Structure et Complexité des Algorithmes*" Thèse d'Etat, Paris-Orsay, (1984). ■

## APPENDIX

**Proof of Lemma 7:** Proposition E implies  $s \subseteq z$ . From Remark K:  $s = z \Rightarrow a \in z^*$ , and the minimality constraint implies  $\alpha = \epsilon$ , hence  $z = s = as$ . The case  $as \subset z \subset sas$  contradicts also Remark K. Now, if  $s \subset z \subset as$  then (minimality condition)  $as \notin z^*$ , and there exists a factorization

$$z = uv, v \preceq as, u \subseteq s \text{ or } s \subseteq u$$

From Remark K (again!):  $s \subseteq u$  and  $s \subseteq v$ . Then:  $u = bs$  and  $v$  satisfies:  $ds \preceq as$  and  $ds \subseteq as$ , hence  $d = a$ . Thus  $z = u.v = bs.as$  which contradicts our assumption ( $z \subset as$ ).

**Proof of Proposition 8:** We have already proved (i). Let  $h_1 = s'a's'$ . From Proposition E and Remark K, one has:  $s = s'$  and:  $as \preceq a's \Rightarrow as \preceq a'$ . If we note:  $a' = asb = csa$ , we have (Remark K):

$$\begin{cases} b \subseteq sa \Rightarrow b \subseteq a \\ b \not\subseteq a \Rightarrow sb \not\subseteq sa \Rightarrow sb \subseteq a. \end{cases}$$

Thus, if  $b \not\subseteq a$ , we can factorize:  $a = dsb$  and we have:  $asb = csa = cs.dsb = csd.sb$ . Hence:

$$\begin{cases} sd \subseteq a \\ ds \preceq a \end{cases} \Rightarrow \begin{cases} sds \subseteq as \subseteq sas \\ sds \preceq sa \preceq sas \end{cases} \Rightarrow d = a.$$

From this contradiction, it follows that:

$$|b| < |sa| \Rightarrow b = a.$$

If  $|b| \geq |sa|$ , then  $b = wsa$  and  $h_1 = sa's = saswsas$  and we apply the Lemma of 2-factorization. Now, let  $b_1$  be in  $B_1$ , not necessarily in  $H_1$ . Then:

$$b_1 = s.ws = s.(a's)^p = \begin{cases} s.(as)^q & \text{if } a's \in (as)^* \\ s.(as(usas)^m)^q = sas.xsas & \text{otherwise} \end{cases}$$

Applying Lemma 7 of 2-factorization in the second case yields  $sasxsas = sas(usas)^m = b_1$ .



