



# A limiting distribution for quicksort

Mireille Regnier

► **To cite this version:**

| Mireille Regnier. A limiting distribution for quicksort. RR-0739, INRIA. 1987. inria-00075813

**HAL Id: inria-00075813**

**<https://hal.inria.fr/inria-00075813>**

Submitted on 24 May 2006

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# INRIA

UNITÉ DE RECHERCHE  
INRIA-ROCCOUENCOURT

Institut National  
de Recherche  
en Informatique  
et en Automatique

Domaine de Voluceau  
Rocquencourt  
BP 105

78153 Le Chesnay Cedex  
France

Tel: (1) 39 63 55 11

## Rapports de Recherche

N° 739

### A LIMITING DISTRIBUTION FOR QUICKSORT

Mireille REGNIER

OCTOBRE 1987

# A LIMITING DISTRIBUTION FOR QUICKSORT

## DISTRIBUTION LIMITE DE QUICKSORT

**ABSTRACT:** *We establish the existence of a limiting distribution for the number of comparisons performed by quicksort, or, equivalently, for the external path length of a binary search tree. We assume a uniform distribution of the data and prove a convergence in distribution and in  $L^p$ ,  $p \geq 1$ . The proof is based on a martingale argument.*

**RESUMÉ:** Nous prouvons l'existence d'une distribution limite pour le nombre de comparaisons effectuées par quicksort, ou, de manière équivalente, pour la longueur de cheminement externe d'un arbre binaire de recherche. Sous l'hypothèse d'une distribution uniforme des données, nous prouvons la convergence en distribution et dans  $L^p$ ,  $p \geq 1$ . La preuve se fonde sur un argument de martingale.

# A LIMITING DISTRIBUTION FOR QUICKSORT

Mireille RÉGNIER

INRIA

78 153 Le Chesnay - FRANCE

**ABSTRACT:** *We establish the existence of a limiting distribution for the number of comparisons performed by quicksort, or, equivalently, for the external path length of a binary search tree. We assume a uniform distribution of the data and prove a convergence in distribution and in  $L^p$ ,  $p \geq 1$ . The proof is based on a martingale argument.*

**Keywords:** Quicksort, binary search tree, limiting distribution, martingale.

## I. INTRODUCTION

In this note, we establish the existence of limiting distributions for Quicksort and Binary Search Trees. Quicksort, discovered by Hoare [HO62], is a widely used algorithm for internal sorting [KN73],[SE77],[SE83]. Notably, it is the standard sort on Unix systems. Binary search trees are a basic data structure for sorting and searching. By a standard equivalence principle (recalled in Section II), evaluating the comparison cost of Quicksort reduces to the study of the *external path length* of a binary search tree. The first two moments, mean and variance, have been known for a long time [KN73, SE77]. More recently, a study of the moments of any order has been done. In [HE87], these moments are obtained by successive differentiation of a multivariate generating function  $C(z, q)$ . This function satisfies a simple bivariate non-linear difference-differential equation:

$$\frac{\partial}{\partial z} C(z, q) = C^2(qz, q) \quad (1)$$

Though moments of an arbitrary fixed order can be computed, it appears impossible so far to derive a general formula (a conjecture can be made, see [HE87] and our conclusion). Thus, one cannot deduce the existence of a limiting distribution from the asymptotics of the moments. In our approach, we make use of *martingale theory* [FE57,NE72] to prove the *existence of a limiting distribution* in law and in  $L^p$ . As pointed out in [HE87], this distribution is *not gaussian* since its third moment is not zero.

The plan of this note is the following. In section II, we recall briefly the Quicksort algorithm and state the relationship with Binary Search Trees. In section III, we first present some elementary

notions on martingales. Then, we associate a martingale to Quicksort and prove our convergence results. Finally, in our conclusion, we discuss the problems that remain open.

## II. QUICKSORT ALGORITHM

We first present the algorithm, and then the parameters to be analysed, and our probabilistic hypotheses.

Quicksort is a fast sorting algorithm, widely used for internal sort. The basic idea is the choice of a *partitioning element*  $K$  [SE83]. For example, let us consider the integer sequence:


45      677      98      43      42      41      60      130      32      67

and choose  $K = 67$  as the partitioning element. Scanning the left sublist from left to right, one exchanges any key greater than  $K$  with a key of the right sublist, scanned from right to left. This builds a list where  $K$  has its final position, all the keys at its left (resp. at its right) being smaller than  $K$ . The intermediate stages are:

45	32							677	67	
45	32	60					98	130	677	67
45	32	60	43	42	41	98	130	677	67	
45	32	60	43	42	41	67	130	677	98	

Then, the process can be applied recursively to both sublists (45...41) and (130...98).

These successive stages can be represented by a binary search tree. Each key is used as a partitioning element in a recursive call and the level of this call is also the depth of the node containing that key.




Now, to evaluate the cost of running Quicksort on a data set, we count the number of comparisons to be performed. Each key in a node is compared once to all the keys occurring in the

path from the root to that node. Thus, our cost is equal to the *External Path Length* of the binary search tree built. As the operations of the algorithm only depend on the relative order of the keys, this data set can also be considered as a permutation in  $\sigma_n$ . We make the standard assumption of a uniform distribution of data [KN73], [SE77], [HE87].

**Probabilistic Model:** *For data sets of size  $n$ , the  $n!$  relative orders are equally likely.*

Under this model, the analysis of Quicksort can be reduced to analyses performed on binary search trees built by *successive insertions*. This standard algorithm for binary search trees is fully described in [KN73, vol 3, 6.2.2.]. We only give here an example: the binary search tree built by successive insertions of the data listed above.



Each order of the keys is associated, for both algorithms, to a binary search tree. Although these trees are different, we have the Equivalence Principle:

**Equivalence Principle:** [KN73, vol 3, p.428] and [SE77]: *The distribution of the binary search trees associated to Quicksort when data are uniformly distributed is the same as the distribution of the binary search trees built by successive insertions of  $n$  random keys.*

This allows us to consider only the evaluation of the *External Path Length* of a binary search tree built by successive insertions. We will choose the latter approach in the following. In this case, the definition of a martingale, as well as the proof of the main properties, is more intuitive, hence easier. We shall prove:

**Theorem 1:** *There exists a random variable  $Z$  such that the internal path length (and the external path length) of a random binary search tree, once centered and normalized, converges, in law and in  $L^2$ , to  $Z$ .*

### III. THE LIMITING DISTRIBUTION

#### III.1. Elementary notions on martingales

Here, we recall very briefly some notions of probability measure, the  $\sigma$ -fields and the measurability, and the definition and elementary properties of martingales. One can refer to [FE57, vol 2, IV.3. and IV.4.] for more details about probability measure and to [FE57, vol 2, VII.9.] or [NE75] for martingale theory.

In probability theory, one deals with special subsets of the sample space, as defined below:

**Definition A:** A  $\sigma$ -field  $\mathcal{A}$  is a system of subsets of a set  $\Omega$  closed under completion and the formation of countable unions and intersections.

Random variables are real functions on  $\Omega$ , but one only uses functions for which a distribution function can be defined.

**Definition B:** Given a  $\sigma$ -field  $\mathcal{A}$  on a set  $\Omega$ , a real-valued function  $u$  on  $\Omega$  is called  $\mathcal{A}$ -measurable if for each  $t$  the set of all points  $x$  where  $u(x) \leq t$  belongs to  $\mathcal{A}$ .

The general definitions of martingales involve an increasing sequence of  $\sigma$ -fields  $(\mathcal{B}_n)$  and a sequence  $(Z_n)$  of  $\mathcal{B}_n$  measurable random variables. A particular case, that we consider here, occurs when  $(\mathcal{B}_n)$  are the  $\sigma$ -fields generated by a sequence of r.v.  $(X_n)$ , i.e. generated by the subsets  $X_n^{-1}(I)$ ,  $I \subset \mathcal{R}$ . The r.v.  $Z_n$  are  $X_n$ -measurable or even simply defined as:

$$Z_n = f_n(X_1, \dots, X_n)$$

where  $f_n$  are  $\mathcal{R}^n$ -measurable. Thus, we shall use the definition:

**Definition C:** Let  $(X_n)$  be a sequence of random variables and  $(Z_n)$  a sequence of  $\mathcal{B}_n$ -measurable random variables. The conditional expectation  $E(Z/X_1 \dots X_{n-1})$  is the  $X_n$ -measurable random variables satisfying, for every  $X_n$ -measurable random variable  $Y$ :

$$E(Z.Y) = E(E(Z/X_1 \dots X_{n-1}).Y).$$

Two important properties follow from this definition.

- (a)  $Z$  and  $E(Z/X_1 \dots X_{n-1})$  have the same expectation.
- (b) If  $Y$  is  $\mathcal{B}_n$ -measurable, then:  $E(YX/X_1 \dots X_{n-1}) = Y.E(Z/X_1 \dots X_{n-1})$ .

**Hint for (a):** Use the definition with  $Y \equiv 1$ .

**Definition D:** Let  $(X_n)$  be a sequence of random variables and  $(Z_n)$  a sequence of  $X_n$ -measurable random variables.  $(Z_n)$  is a martingale if it satisfies the property:

$$E(Z_{n+1}/X_1, \dots, X_n) = Z_n.$$

From our remark above, such variables have a constant expectation; Moreover, simple properties on the variances are sufficient to establish the existence of a limit, as stated in Theorem 2 [FE57]:

**Theorem 2:** *Let  $(Z_n)$  be an infinite martingale with  $E(Z_n^2) < C < +\infty$  for all  $n$ . There exists a random variable  $Z$  such that  $Z_n \rightarrow Z$  in law and in  $L^2$ . Furthermore  $E(Z_n) = E(Z)$  for all  $n$ .*

*Moreover, if  $(Z_n)$  satisfies:  $E(|Z_n|^p) < C_p$  for some  $p > 2$ , then  $Z_n \rightarrow Z$  in  $L^p$ .*

### III.2. A martingale associated to Quicksort

We prove here the existence of a limiting distribution for the *path length* of a binary search tree. As pointed out in Section II, this is also the number of comparisons performed when running Quicksort.

We first state our notations. The *depth of insertion* of a record is the depth of the node where this record is inserted, where we take the depth of the root to be 0. We note  $X_n$  the random variable counting the depth of insertion of a random record in a random binary search tree of size  $n - 1$ . (We have  $X_1 \equiv 0$  and  $X_2 \equiv 1$ ). The *internal path length* is then:

$$IPL_n = \sum_{i=1}^n X_i.$$

For a reason of convenience, we also define:

$$Y_n = \sum_i (X_i + 2) = IPL_n + 2n$$

the random variable representing the *external path length* of a binary search tree built on  $n$  records. Noting that the path lengths are simple  $X_n$ -mesurable functions, we are now ready to define a martingale. This is the purpose of the following Proposition 3. Then, applying Theorem 2 to this result, will yield directly our main result, Theorem 1 above.

**Proposition 3:** *The random variables*

$$Z_n = \frac{Y_n - 2(n+1)(H_{n+1} - 1)}{n+1} = \frac{IPL_n - 2(n+1)H_n + 4n}{n+1}$$

*form a martingale with a null expectation. Their variances satisfy:*

$$V_n = \left(7 - \frac{4\pi^2}{6}\right) + O\left(\frac{1}{n}\right).$$

**Proof of Proposition 3:** The proof is based on the following two lemmas.



**Lemma A:** *The conditional expectation of the random variables  $X_n$  and  $Y_n$  satisfy:*

$$\begin{cases} E(X_n/X_1 \dots X_{n-1}) = \frac{Y_{n-1}}{n} \\ E(Y_n/X_1 \dots X_{n-1}) = \frac{n+1}{n}Y_{n-1} + 2 \end{cases}$$

**Lemma B:** *The expectation of the internal and external path length satisfy:*

$$\begin{cases} E(IPL_n) = 2(n+1)H_n - 4n \\ E(Y_n) = 2(n+1)(H_{n+1} - 1) \end{cases}$$

**Proof of Lemma A:** From the definition,  $nE(X_n/X_1 \dots X_{n-1})$  is the sum of the possible values for  $n$ , in a tree built by successive insertions at depths  $X_1, \dots, X_{n-1}$ . The insertion of the  $n-1$ -st key,  $K_{n-1}$ , at depth  $X_{n-1}$ , modified the binary search tree built on the keys  $K_1, \dots, K_{n-2}$  in the following way. A leaf at depth  $X_{n-1}$  was changed to a node, where  $K_{n-1}$  was inserted, while two more leaves at depth  $X_{n-1} + 1$  were created, where the  $n$ -th key  $K_n$  may now be inserted. Thus:

$$nE(X_n/X_1 \dots X_{n-1}) = (n-1)E(X_{n-1}/X_1 \dots X_{n-2}) - X_{n-1} + 2(X_{n-1} + 1).$$

Unwinding this recurrence, with  $E(X_1) \equiv 0$ , yields:

$$nE(X_n, X_1 \dots X_{n-1}) = \sum_{i=1}^{n-1} (X_i + 2) = Y_{n-1}.$$

Now:

$$E(Y_n/X_1 \dots X_{n-1}) = E(Y_{n-1} + X_n + 2/X_1 \dots X_{n-1}) = Y_{n-1} + \frac{1}{n}Y_{n-1} + 2.$$

**Proof of Lemma B:** It follows directly from Lemma A that:

$$\frac{E(Y_n)}{n+1} = \frac{E(Y_{n-1})}{n} + \frac{2}{n+1} = 2 \sum_{i=2}^n \frac{1}{i+1} + 1 = 2(H_{n+1} - 1).$$

As an immediate consequence, one finds the martingale property for  $Z_n$ :

$$E(Z_n/X_1 \dots X_{n-1}) = E(Y_n - E(Y_n)/n + 1/X_1 \dots X_{n-1}) = \frac{Y_{n-1}}{n} + \frac{2}{n+1} - \frac{E(Y_n)}{n+1} = Z_{n-1}.$$

Now, to derive the expression of the variances, we rewrite:

$$Z_n = \frac{n}{n+1}Z_{n-1} + \frac{X_n - E(X_n)}{n+1}$$

and get:

$$E(Z_n^2) = \frac{n^2}{(n+1)^2} E(Z_{n-1}^2) + \frac{\text{Var}(X_n)}{(n+1)^2} + \frac{2n}{(n+1)^2} E(Z_{n-1}(X_n - E(X_n))) .$$

Using  $E((X_n - E(X_n))/X_1 \dots X_{n-1}) = Z_{n-1}$  and noting  $\pi_n = \frac{n+2}{n+1}$ , we get:

$$\frac{E(Z_n^2)}{\pi_n} = \frac{E(Z_{n-1}^2)}{\pi_{n-1}} + \frac{\text{Var}(X_n)}{(n+1)(n+2)} .$$

The moments of the depth of insertion  $X_n$  are well known [KN73,LO87] (the limiting distribution being gaussian). In particular:

$$\text{Var}(X_n) = 2(H_n - 1) - 4(H_n^{(2)} - 1),$$

with  $H_n^{(2)} = \sum_{i=1}^n 1/i^2$ . Summing these expressions yields:

$$E(Z_n^2) = \pi_n(7 - 4H_{n+1}^{(2)} + O(\frac{1}{n})) = 7 - 4H_{n+1}^{(2)} + O(\frac{1}{n}) .$$

This completes the proof of Proposition 3 (and hence also Theorem 1). ■

#### IV. CONCLUSION

We have been able in this note to prove the existence of a limiting distribution for the path length of a binary search tree or, equivalently, the number of comparisons performed by Quicksort. The proof uses a martingale argument and enables us to rederive, in passing, the first two moments of the distribution.

It is worth pointing out here that in [HE87], the derivation of the equation (1) gave a surprisingly simple formula for the *cumulants*  $(k_p)_{p \in \mathbb{N}}$  of the distribution, namely:

$$k_p(n) = (-1)^{p+1} 2^p (p-1)! \zeta(p) + C_p + O_p(\frac{1}{n})$$

where  $\zeta(p) = \sum_{n=1}^{\infty} \frac{1}{n^p}$ , and it is *conjectured* that such a formula, verified for  $p \leq 20$ , holds true for any  $p$ , with a uniform approximation. Solving this open problem and finding the  $C_p$  would enable us to explicitly characterize the limiting distribution for which we have given here a "non constructive" existence proof. .

#### REFERENCES:

- [BM85] F. BACCELLI AND A.M. MAKOWSKY: "Direct martingale arguments for stability: The M/G/1 case" *Systems&Control Letters*, 6 (1985) 181-186.
- [FE57] W. FELLER: *An Introduction to Probability Theory and its Applications, Vol. II*, Wiley, (1957).

- [HE87] P. HENNEQUIN: "Combinatorial Analysis of Quicksort Algorithm" to appear in *RAIRO, Th. Informatics and Applications*.
- [HO62] C.A.HOARE: "Quicksort", *Computer Journal*, 5,1 (1962).
- [KN73] D. KNUTH: *The Art of Computer Programming. Vol 3:Sorting and Searching*, Addison-Wesley, Reading, Mass.,(1973).
- [LO87] G. LOUCHARD: "Exact and Asymptotic Distributions in Digital and Binary Search Trees", to appear in *RAIRO, Th. Informatics and Applications*.
- [NE75] J. NEVEU: *Discrete-Parameter Martingale*", English Translation, North-Holland, Amsterdam, (1975).
- [SE77] R. SEDGEWICK: "The Analysis of Quicksort Programs", *Acta Informatica* 7, (1977), 327-355, and in *Quicksort*, Garland Pub.Co., New-York,(1980).

1

2

3

4

5

6

7

8

9