

# On the approximation of general shell problems by the Clough-Johnson flat plate elements. Part 3: The implementation

Michel Bernadou, Y. Ducatel

► **To cite this version:**

Michel Bernadou, Y. Ducatel. On the approximation of general shell problems by the Clough-Johnson flat plate elements. Part 3: The implementation. RR-0674, INRIA. 1987. inria-00075879

**HAL Id: inria-00075879**

**<https://hal.inria.fr/inria-00075879>**

Submitted on 24 May 2006

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# IRIA

UNITÉ DE RECHERCHE  
IRIA-ROCUENCOURT

Institut National  
de Recherche  
en Informatique  
et en Automatique

Domaine de Voluceau  
Rocquencourt  
B.P.105  
78153 Le Chesnay Cedex  
France  
Tél.(1) 39 63 55 11

## Rapports de Recherche

N° 674

**ON THE APPROXIMATION OF  
GENERAL SHELL PROBLEMS  
BY THE CLOUGH-JOHNSON  
FLAT PLATE ELEMENTS  
PART 3 : THE IMPLEMENTATION**

**Michel BERNADOU  
Yves DUCATEL**

**JUIN 1987**

ON THE APPROXIMATION OF GENERAL SHELL PROBLEMS  
BY THE CLOUGH-JOHNSON FLAT PLATE ELEMENTS

PART 3 : THE IMPLEMENTATION

Michel BERNADOU<sup>(1)</sup> - Yves DUCATEL<sup>(2)</sup>

Summary : In this series of reports, we give an account of some results obtained in the approximation of *general shell problems* by the CLOUGH-JOHNSON *flat plate elements*. The first part is concerned by the study of the *compatibility equations*. In the second part, we deliver several interesting results valid for general shells and we prove the "pseudo-convergence" of the method for a class of shallow shells. Then, this careful study allows us to introduce a perturbation of this approximation and thus, to propose a new method which is convergent for general shells. Finally, in the third part, we describe in details how to implement the CLOUGH-JOHNSON method.

SUR L'APPROXIMATION DE PROBLEMES GENERAUX DE COQUES  
PAR DES METHODES D'ELEMENTS FINIS PLATS DE CLOUGH ET JOHNSON

PARTIE 3 : IMPLEMENTATION

Résumé : Dans cette série de rapports, nous rassemblons les divers résultats obtenus dans l'approximation de *problèmes généraux de coques* par les éléments finis plats (de plaques) de CLOUGH et JOHNSON. La première partie est relative à l'étude des équations de *compatibilité*. Dans la seconde partie, nous donnons plusieurs résultats intéressants valables pour des coques générales, puis nous démontrons la pseudo-convergence de la méthode pour une classe de coques peu profondes. Cette étude détaillée nous permet alors d'introduire une perturbation de cette approximation et ainsi, de proposer une nouvelle méthode qui converge pour des coques générales. Finalement, dans la troisième partie, nous détaillons la marche à suivre pour implémenter la méthode de CLOUGH et JOHNSON.

---

(1) INRIA, Domaine de Voluceau, Rocquencourt, B.P. 105, 78153 Le Chesnay

(2) Université de Poitiers, Département de Mathématiques,  
40, avenue du Recteur Pineau, 86022 Poitiers Cedex

CONTENTS

PART 1 : COMPATIBILITY RELATIONS

- 1 - Introduction
- 2 - The continuous problem
  - 2.1 - Some notations
  - 2.2 - Variational formulation of KOITER's linear model
  - 2.3 - Existence and uniqueness theorem
- 3 - A conforming finite element method
  - 3.1 - The finite element space  $\vec{V}_h$
  - 3.2 - The discrete problem
- 4 - The discrete problem using flat plate elements and compatibility relations
  - 4.1 - The approximate middle surface  $\bar{\mathcal{I}}_h$
  - 4.2 - The discrete space  $\vec{V}_h$  using compatibility conditions
  - 4.3 - Definition of the discrete problem using flat plate elements
  - 4.4 - Another alternative : CARR's method

PART 2 : STUDY OF THE CONVERGENCE AND ERROR ESTIMATES

- 5 - The consistency errors
  - 5.1 - Abstract error estimate
  - 5.2 - Estimate of the consistency error  $|a(\vec{v}_h, \vec{w}_h) - b_h(\vec{v}_h, \vec{w}_h)|$
  - 5.3 - Estimate of the consistency error  $|f(\vec{w}_h) - g_h(\vec{w}_h)|$
  - 5.4 - The bilinear form  $b_h(.,.)$  is uniformly  $\vec{V}_h$ -elliptic
  - 5.5 - About CARR's approach
- 6 - Pseudo-convergence and error estimates ; a new convergent method
  - 6.1 - Pseudo-convergence and error estimates for shallow shells
  - 6.2 - Examples of suitable numerical integration schemes
  - 6.3 - A new convergent method for general shells

CONCLUSION


PART 3 : THE IMPLEMENTATION

- 7 - Implementation of the reduced H.C.T. triangle
  - 7.1 - Definition of the reduced H.C.T. triangle
  - 7.2 - Barycentric coordinates - Cartesian coordinates
  - 7.3 - Eccentricity parameters
  - 7.4 - Local and global degrees of freedom
  - 7.5 - Basis functions
  - 7.6 - Interpolation of a function  $v$  and partial derivatives of the interpolating function
  - 7.7 - Interpolation of the triangle of type (1)
- 8 - Implementation of the facet stiffness and second member matrices
  - 8.1 - Implementation of the facet stiffness matrix
  - 8.2 - Implementation of the facet second member matrix
  - 8.3 - Implementation of the compatibility relations
- 9 - Derivation of the global stiffness and second member matrices
  - 9.1 - Derivation of the global stiffness matrix
  - 9.2 - Derivation of the global second member.

••••

HOW TO READ THE PAPER

All the paper is divided in three parts. In the first one the reader can find all the basis. Then, *part 2 and part 3 can be read independtly* depending of the interest of everyone. In other words we can summarize these possibilities in the following "flow shart" :



ACKNOWLEDGEMENT : The Authors are indebted to F. PALMA who read the manuscript in its entirety and who suggest various improvements ; also, they gratefully appreciate the excellent typing of Mrs. DESNOUS.

7 - IMPLEMENTATION OF THE REDUCED HSIEH-CLOUGH-TOCHER TRIANGLE

Orientation : In this paragraph we record the main features of the implementation of the triangle of type 1 and of the reduced HCT triangle. For more details we refer to BERNADOU-BOISSERIE [2] and BERNADOU-HASSAN [20].

7.1 - Definition of the reduced HCT triangle

Let  $K$  be a triangle with vertices  $\Sigma_1, \Sigma_2, \Sigma_3$  and barycenter  $\Sigma$ . Let  $K_i$  be the triangles with vertices  $\Sigma, \Sigma_{i+1}, \Sigma_{i-1}$ ,  $i=1,2,3$ . Here and subsequently the indices belong to  $\{1,2,3\}$  modulo 3.

Then, the definition of the reduced HCT triangle  $(K, P_K, \Sigma_K)$  is condensed in Fig. 7.1.1. Let us record that  $P_K$  and  $\Sigma_K$  denote the discrete space and the set of degrees of freedom, respectively.




Figure 7.1.1 : The reduced HCT-triangle

7.2 - Barycentric coordinates - Cartesian coordinates

Let there be given an orthonormal fixed reference system of the plane  $\mathbb{R}^2$ ,  $(0, \vec{e}_1, \vec{e}_2)$ . The coordinates of the point P are denoted  $(\xi^1, \xi^2)$ , i.e.,

$$\vec{OP} = \xi^1 \vec{e}_1 + \xi^2 \vec{e}_2 .$$

Let K be a triangle with vertices  $\Sigma_1(\xi_1^1, \xi_1^2)$ ,  $\Sigma_2(\xi_2^1, \xi_2^2)$ ,  $\Sigma_3(\xi_3^1, \xi_3^2)$ . We assume  $\vec{\Sigma}_1 \vec{\Sigma}_2 \times \vec{\Sigma}_1 \vec{\Sigma}_3 \neq 0$  .

Then, the cartesian coordinates  $(\xi^1, \xi^2)$  of the point P are linked to the barycentric coordinates  $(\lambda_1, \lambda_2, \lambda_3)$  by the relations

$$\left. \begin{aligned} \xi^1 &= \lambda_1 \xi_1^1 + \lambda_2 \xi_2^1 + \lambda_3 \xi_3^1 , \\ \xi^2 &= \lambda_1 \xi_1^2 + \lambda_2 \xi_2^2 + \lambda_3 \xi_3^2 . \end{aligned} \right\} \quad (7.2.1)$$

Let us record that

$$1 = \lambda_1 + \lambda_2 + \lambda_3 . \quad (7.2.2)$$

Conversely,

$$\left. \begin{aligned} \lambda_1 &= \frac{(\xi^1 - \xi_2^1)(\xi_2^2 - \xi_3^2) + (\xi_3^1 - \xi_2^1)(\xi^2 - \xi_2^2)}{\Delta} \\ \lambda_2 &= \frac{(\xi^1 - \xi_3^1)(\xi_3^2 - \xi_1^2) + (\xi_1^1 - \xi_3^1)(\xi^2 - \xi_3^2)}{\Delta} \\ \lambda_3 &= \frac{(\xi^1 - \xi_1^1)(\xi_1^2 - \xi_2^2) + (\xi_2^1 - \xi_1^1)(\xi^2 - \xi_1^2)}{\Delta} \end{aligned} \right\} \quad (7.2.3)$$

where

(1)

(2)

$$\begin{aligned}
 \Delta &= \xi_1^1(\xi_2^2 - \xi_3^2) + \xi_2^1(\xi_3^2 - \xi_1^2) + \xi_3^1(\xi_1^2 - \xi_2^2) \\
 &= (\xi_1^1 - \xi_3^1)(\xi_2^2 - \xi_3^2) + (\xi_2^1 - \xi_3^1)(\xi_3^2 - \xi_1^2) \\
 &= (\overrightarrow{\Sigma_1 \Sigma_3} \times \overrightarrow{\Sigma_2 \Sigma_3}) \cdot (\vec{e}_1 \times \vec{e}_2) \neq 0 .
 \end{aligned}
 \tag{7.2.4}$$

### 7.3 - Eccentricity parameters

In order to define conveniently the orthogonal direction to the side  $\Sigma_{i-1} \Sigma_{i+1}$ , we introduce the eccentricity parameters  $\eta_i \in \mathbb{R}$ ,  $i=1,2,3$ , i.e.

$$\eta_i = 2 \frac{\overline{c_i b_i}}{\Sigma_{i-1} \Sigma_{i+1}} = \frac{(\ell_{i-1})^2 - (\ell_{i+1})^2}{\ell_i^2}, \quad i=1,2,3 .
 \tag{7.3.1}$$

By  $b_i$ ,  $c_i$ ,  $\ell_i$ , we respectively denote the midside of  $\Sigma_{i-1} \Sigma_{i+1}$ , the orthogonal projection of the point  $\Sigma_i$  on the side  $\Sigma_{i-1} \Sigma_{i+1}$  and the length of the side  $\Sigma_{i-1} \Sigma_{i+1}$ .

### 7.4 - Local and global degrees of freedom

In the definition of the degrees of freedom of the reduced HCT-triangle we have only used the geometry of the triangle. These degrees of freedom are called *local*. To assemble the contributions of each element, it is convenient to replace the direction of the sides of the triangle by the fixed directions given by the vectors  $\vec{e}_1$ ,  $\vec{e}_2$ , and next, to normalize.

Thus, to the three local degrees of freedom attached to the vertex  $\Sigma_i$ , i.e.,

$$v \rightarrow \{v(\Sigma_i), Dv(\Sigma_i)(\Sigma_{i-1} - \Sigma_i), Dv(\Sigma_i)(\Sigma_{i+1} - \Sigma_i)\}
 \tag{7.4.1}$$

we associate the three global degrees of freedom





7.5 - Basis functions

In BERNADOU-HASSAN [20], we have shown that the basis functions (shape functions) for the *general* reduced HCT-triangle are given by the figure 7.5.1 and are associated to the following interpolating function  $\pi_{K_i} v$  of a (sufficiently smooth) function  $v$  on the subtriangle  $K_i$ ,  $i=1,2,3$ .

$$\left. \begin{aligned} \pi_{K_i} v &= v(\Sigma_i) r_{i,i}^0 + v(\Sigma_{i+1}) r_{i,i+1}^0 + v(\Sigma_{i+2}) r_{i,i+2}^0 \\ &+ Dv(\Sigma_i) (\Sigma_{i+2} - \Sigma_i) r_{i,i,i+2}^1 + Dv(\Sigma_i) (\Sigma_{i+1} - \Sigma_i) r_{i,i,i+1}^1 \\ &+ Dv(\Sigma_{i+1}) (\Sigma_i - \Sigma_{i+1}) r_{i,i+1,i}^1 + Dv(\Sigma_{i+1}) (\Sigma_{i+2} - \Sigma_{i+1}) r_{i,i+1,i+2}^1 \\ &+ Dv(\Sigma_{i+2}) (\Sigma_{i+1} - \Sigma_{i+2}) r_{i,i+2,i+1}^1 + Dv(\Sigma_{i+2}) (\Sigma_i - \Sigma_{i+2}) r_{i,i+2,i}^1 \end{aligned} \right\} (7.5.1)$$

Figure 7.5.1 can be condensed as follows :

$$[R_i]_{9 \times 1} = [A_i]_{9 \times 10} [\Lambda_i]_{10 \times 1} \quad (7.5.2)$$

7.6 - Interpolation of a function  $v$  and partial derivatives of the interpolating function

Let  $\pi_{K_i} v$  be the interpolating function of a function  $v$  on the subtriangle  $K_i$ . The relations (7.4.5) (7.5.1) reveal that

$$\pi_{K_i} v = [DLLC_i(v)]_{1 \times 9} [R_i]_{9 \times 1} \quad (7.6.1)$$

or equivalently, using relations (7.4.7) and (7.5.2)

$$\pi_{K_i} v = [DLGL_i(v)]_{1 \times 9} [D_i]_{9 \times 9} [A_i]_{9 \times 10} [\Lambda_i]_{10 \times 1} \quad (7.6.2)$$

$$\begin{bmatrix} r_{i,i}^0 \\ r_{i,i+1}^0 \\ r_{i,i+2}^0 \\ r_{i,i,i+2}^1 \\ r_{i,i,i+1}^1 \\ r_{i,i+1,i}^1 \\ r_{i,i+1,i+2}^1 \\ r_{i,i+2,i+1}^1 \\ r_{i,i+2,i}^1 \end{bmatrix} = \begin{bmatrix} -\frac{1}{2}(\eta_{i+1}-\eta_{i+2}) & 0 & 0 & \frac{3}{2}(3+\eta_{i+1}) & \frac{3}{2}(3-\eta_{i+2}) & 0 & 0 & 0 & 0 \\ \frac{1}{2}(1-2\eta_{i+1}-\eta_{i+2}) & 1 & 0 & -\frac{3}{2}(1-\eta_{i+1}) & \frac{3}{2}(\eta_{i+1}+\eta_{i+2}) & 3 & 3 & 0 & 0 & 3(1-\eta_{i+1}) \\ \frac{1}{2}(1+2\eta_{i+1}+\eta_{i+2}) & 0 & 1 & -\frac{3}{2}(\eta_{i+1}+\eta_{i+2}) & -\frac{3}{2}(1+\eta_{i+1}) & 0 & 0 & 3 & 3 & 3(1+\eta_{i+1}) \\ -\frac{1}{4}(1+\eta_{i+1}) & 0 & 0 & \frac{1}{4}(5+3\eta_{i+1}) & \frac{1}{2} & 0 & 0 & 0 & 0 & 0 \\ -\frac{1}{4}(1-\eta_{i+2}) & 0 & 0 & \frac{1}{2} & \frac{1}{4}(5-3\eta_{i+2}) & 0 & 0 & 0 & 0 & 0 \\ \frac{1}{4}(1-\eta_{i+2}) & 0 & 0 & -\frac{1}{2} & -\frac{1}{4}(1-3\eta_{i+2}) & 1 & 0 & 0 & 0 & 1 \\ -\frac{1}{2}\eta_i & 0 & 0 & -\frac{1}{4}(1-3\eta_i) & \frac{1}{4}(1+3\eta_i) & 0 & 1 & 0 & 0 & \frac{1}{2}(1-3\eta_i) \\ \frac{1}{2}\eta_i & 0 & 0 & \frac{1}{4}(1-3\eta_i) & -\frac{1}{4}(1+3\eta_i) & 0 & 0 & 1 & 0 & \frac{1}{2}(1+3\eta_i) \\ \frac{1}{4}(1+\eta_{i+1}) & 0 & 0 & -\frac{1}{4}(1+3\eta_{i+1}) & -\frac{1}{2} & 0 & 0 & 0 & 1 & 1 \end{bmatrix} \begin{bmatrix} \lambda_i^3 \\ \lambda_{i+1}^3 \\ \lambda_{i+2}^3 \\ \lambda_i^2 \lambda_{i+2} \\ \lambda_i^2 \lambda_{i+1} \\ \lambda_{i+1} \lambda_i^2 \\ \lambda_{i+1} \lambda_{i+2} \\ \lambda_{i+2} \lambda_{i+1} \\ \lambda_i \lambda_{i+1} \lambda_{i+2} \end{bmatrix}$$

Figure 7.5.1 : Basis functions associated to subtriangle  $K_1$  for the reduced-HCT finite element

Then, the derivatives of the interpolating function  $\pi_{K_i} v$  are given by the expressions

$$\left. \begin{aligned} \partial_\alpha \pi_{K_i} v &= [DLGL_i(v)] [D_i] [A_i] [\partial_\alpha \Lambda_i], \quad [\partial_\alpha \Lambda_i] = \left[ \frac{\partial \Lambda_i}{\partial \xi^\alpha} \right], \\ \partial_{\alpha\beta} \pi_{K_i} v &= [DLGL_i(v)] [D_i] [A_i] [\partial_{\alpha\beta} \Lambda_i], \quad [\partial_{\alpha\beta} \Lambda_i] = \left[ \frac{\partial^2 \Lambda_i}{\partial \xi^\alpha \partial \xi^\beta} \right]. \end{aligned} \right\} (7.6.3)$$

The column matrices  $[\partial_1 \Lambda_i]$  and  $[\partial_2 \Lambda_i]$  are defined from the matrices  $\left[ \frac{\partial \Lambda_i}{\partial \lambda_j} \right]$ ,  $j=1,2,3$  (cf. tableau 7.6.1) as follows

$$\left. \begin{aligned} [\partial_1 \Lambda_i] &= \frac{1}{\Delta} \{ (\xi_2^2 - \xi_3^2) \left[ \frac{\partial \Lambda_i}{\partial \lambda_1} \right] + (\xi_3^2 - \xi_1^2) \left[ \frac{\partial \Lambda_i}{\partial \lambda_2} \right] + (\xi_1^2 - \xi_2^2) \left[ \frac{\partial \Lambda_i}{\partial \lambda_3} \right] \}, \\ [\partial_2 \Lambda_i] &= \frac{1}{\Delta} \{ (\xi_3^1 - \xi_2^1) \left[ \frac{\partial \Lambda_i}{\partial \lambda_1} \right] + (\xi_1^1 - \xi_3^1) \left[ \frac{\partial \Lambda_i}{\partial \lambda_2} \right] + (\xi_2^1 - \xi_1^1) \left[ \frac{\partial \Lambda_i}{\partial \lambda_3} \right] \}. \end{aligned} \right\} (7.6.4)$$

By analogy, we obtain  $[\partial_{\alpha\beta} \Lambda_i] = \left[ \frac{\partial^2 \Lambda_i}{\partial \xi^\alpha \partial \xi^\beta} \right]$  as :

$$\left. \begin{aligned} [\partial_{11} \Lambda_i] &= \frac{1}{\Delta^2} \{ (\xi_2^2 - \xi_3^2)^2 \left[ \frac{\partial^2 \Lambda_i}{(\partial \lambda_1)^2} \right] + (\xi_3^2 - \xi_1^2)^2 \left[ \frac{\partial^2 \Lambda_i}{(\partial \lambda_2)^2} \right] \\ &+ (\xi_1^2 - \xi_2^2)^2 \left[ \frac{\partial^2 \Lambda_i}{(\partial \lambda_3)^2} \right] + 2(\xi_2^2 - \xi_3^2)(\xi_3^2 - \xi_1^2) \left[ \frac{\partial^2 \Lambda_i}{\partial \lambda_1 \partial \lambda_2} \right] \\ &+ 2(\xi_3^2 - \xi_1^2)(\xi_1^2 - \xi_2^2) \left[ \frac{\partial^2 \Lambda_i}{\partial \lambda_2 \partial \lambda_3} \right] + 2(\xi_1^2 - \xi_2^2)(\xi_2^2 - \xi_3^2) \left[ \frac{\partial^2 \Lambda_i}{\partial \lambda_3 \partial \lambda_1} \right] \} \end{aligned} \right\} (7.6.5)$$

$$\begin{aligned}
 [\partial_{12}\Lambda_i] &= \frac{1}{\Delta} \{ (\xi_2^2 - \xi_3^2)(\xi_3^1 - \xi_2^1) \left[ \frac{\partial^2 \Lambda_i}{(\partial \lambda_1)^2} \right] + (\xi_3^2 - \xi_1^2)(\xi_1^1 - \xi_3^1) \left[ \frac{\partial^2 \Lambda_i}{(\partial \lambda_2)^2} \right] \\
 &+ (\xi_1^2 - \xi_2^2)(\xi_2^1 - \xi_1^1) \left[ \frac{\partial^2 \Lambda_i}{(\partial \lambda_3)^2} \right] \\
 &+ ((\xi_2^2 - \xi_3^2)(\xi_1^1 - \xi_3^1) + (\xi_3^2 - \xi_1^2)(\xi_3^1 - \xi_2^1)) \left[ \frac{\partial^2 \Lambda_i}{\partial \lambda_1 \partial \lambda_2} \right] \\
 &+ ((\xi_3^2 - \xi_1^2)(\xi_2^1 - \xi_1^1) + (\xi_1^2 - \xi_2^2)(\xi_1^1 - \xi_3^1)) \left[ \frac{\partial^2 \Lambda_i}{\partial \lambda_2 \partial \lambda_3} \right] \\
 &+ ((\xi_1^2 - \xi_2^2)(\xi_3^1 - \xi_2^1) + (\xi_2^2 - \xi_3^2)(\xi_2^1 - \xi_1^1)) \left[ \frac{\partial^2 \Lambda_i}{\partial \lambda_3 \partial \lambda_1} \right] \} \\
 \\ 
 [\partial_{22}\Lambda_i] &= \frac{1}{\Delta} \{ (\xi_3^1 - \xi_2^1)^2 \left[ \frac{\partial^2 \Lambda_i}{(\partial \lambda_1)^2} \right] + (\xi_1^1 - \xi_3^1)^2 \left[ \frac{\partial^2 \Lambda_i}{(\partial \lambda_2)^2} \right] \\
 &+ (\xi_2^1 - \xi_1^1)^2 \left[ \frac{\partial^2 \Lambda_i}{(\partial \lambda_3)^2} \right] + 2(\xi_3^1 - \xi_2^1)(\xi_1^1 - \xi_3^1) \left[ \frac{\partial^2 \Lambda_i}{\partial \lambda_1 \partial \lambda_2} \right] \\
 &+ 2(\xi_1^1 - \xi_3^1)(\xi_2^1 - \xi_1^1) \left[ \frac{\partial^2 \Lambda_i}{\partial \lambda_2 \partial \lambda_3} \right] + 2(\xi_2^1 - \xi_1^1)(\xi_3^1 - \xi_2^1) \left[ \frac{\partial^2 \Lambda_i}{\partial \lambda_3 \partial \lambda_1} \right] \}
 \end{aligned} \tag{7.6.5}$$

7.7 - Implementation of the triangle of type (1) (i.e.,  $P_1$ -Lagrange)

Let  $\bar{\omega}_K v$  be the interpolating function of a function  $v$  on the triangle  $K$ , using the triangle of type (1) associated to  $P_1$ -Lagrange interpolation. Similarly to (7.6.2) (7.6.3), we obtain

$$\bar{\omega}_K v = [d1g1(v)]_{1 \times 3} [\lambda]_{3 \times 1} \tag{7.7.1}$$

where

$$[\text{d}l\text{g}l(\mathbf{v})]_{1 \times 3} = [\mathbf{v}(\Sigma_1) \quad \mathbf{v}(\Sigma_2) \quad \mathbf{v}(\Sigma_3)] \quad (7.7.2)$$

$${}^t[\lambda]_{1 \times 3} = [\lambda_1 \quad \lambda_2 \quad \lambda_3] \quad (7.7.3)$$

Similarly to (7.6.4), we obtain

$${}^t[\partial_1 \lambda] = \frac{1}{\Delta} [\xi_2^2 - \xi_3^2 \quad \xi_3^2 - \xi_1^2 \quad \xi_1^2 - \xi_2^2] \quad (7.7.4)$$

$${}^t[\partial_2 \lambda] = \frac{1}{\Delta} [\xi_3^1 - \xi_2^1 \quad \xi_1^1 - \xi_3^1 \quad \xi_2^1 - \xi_1^1] \quad (7.7.5)$$

8 - IMPLEMENTATION OF THE FACET STIFFNESS AND SECOND MEMBER MATRICES

8.1 - Implementation of the facet stiffness matrix

For clarity, we begin to describe the implementation of the facet stiffness matrix directly derived from the expression (4.3.3) :

$$\begin{aligned} \tilde{a}_{hK}(\vec{u}_h, \vec{v}_h) &= \int_K \frac{Ee}{1-\nu^2} \{ (1-\nu) \tilde{\gamma}_{h\beta}^\alpha(\vec{u}_h) \tilde{\gamma}_{h\alpha}^\beta(\vec{v}_h) + \nu \tilde{\gamma}_{h\alpha}^\alpha(\vec{u}_h) \tilde{\gamma}_{h\beta}^\beta(\vec{v}_h) \\ &+ \frac{e^2}{12} [ (1-\nu) \tilde{\rho}_{h\beta}^\alpha(\vec{u}_h) \tilde{\rho}_{h\alpha}^\beta(\vec{v}_h) + \nu \tilde{\rho}_{h\alpha}^\alpha(\vec{u}_h) \tilde{\rho}_{h\beta}^\beta(\vec{v}_h) ] \} \sqrt{a_h} d\xi^1 d\xi^2 \end{aligned} \quad \left. \begin{array}{l} 4) \\ 5) \end{array} \right\} (8.1.1)$$

For the time being we leave aside the problem of compatibility relations and the only work on the facet K. Since the functions  $\vec{u}_h$  and  $\vec{v}_h$  belong to the space  $\tilde{X}_h$  - see (4.2.1) (4.2.3) (4.2.4) - the relations (7.6.2) (7.7.1) reveal that on the triangle  $K = \bigcup_{i=1}^3 K_i$  we have

$$\tilde{u}_{h\alpha} = [d_l g_l(\tilde{u}_{h\alpha})]_{1 \times 3} [\lambda]_{3 \times 1} \quad (8.1.2)$$

$$\tilde{u}_{h3} |_{K_i} = [DLGL_i(\tilde{u}_{h3})]_{1 \times 9} [D_i]_{9 \times 9} [A_i]_{9 \times 10} [\Lambda_i]_{10 \times 1} \quad (8.1.3)$$

Moreover from (8.1.2) (7.7.4) (7.7.5) on the one hand and from (8.1.3) (7.6.3) on the other hand, we derive

$$\tilde{u}_{h\alpha, \beta} = [d_l g_l(\tilde{u}_{h\alpha})]_{1 \times 3} [\partial_\beta \lambda] \quad (8.1.4)$$

$$\tilde{u}_{h3, \alpha} |_{K_i} = [DLGL_i(\tilde{u}_{h3})]_{1 \times 9} [D_i]_{9 \times 9} [A_i]_{9 \times 10} [\partial_\alpha \Lambda_i]_{10 \times 1} \quad (8.1.5)$$

$$\tilde{u}_{h3, \alpha\beta} |_{K_i} = [DLGL_i(\tilde{u}_{h3})]_{1 \times 9} [D_i]_{9 \times 9} [A_i]_{9 \times 10} [\partial_{\alpha\beta} \Lambda_i]_{10 \times 1} \quad (8.1.6)$$

Expression of  $\tilde{\gamma}_{h\beta}^\alpha(\vec{u}_h)$

From (4.1.4) (4.2.11) we obtain

2)

3)

$$\tilde{\gamma}_{h\beta}^{\alpha}(\vec{u}_h) = a_h^{\alpha\lambda} \tilde{\gamma}_{h\beta\lambda} = \frac{1}{2} a_h^{\alpha\lambda} (\tilde{u}_{h\beta,\lambda} + \tilde{u}_{h\lambda,\beta}) \quad (8.1.7)$$

or

$$\tilde{\gamma}_{h\beta}^{\alpha}(\vec{u}_h) = M_{h\beta}^{\alpha} \tilde{u}_{h12} \quad (8.1.8)$$

where

$$M_{h\beta}^{\alpha} = [0; a_h^{\alpha 1} \delta_{\beta 1} ; \frac{1}{2} a_h^{\alpha\lambda} \Delta_{\lambda\beta} ; 0 ; \frac{1}{2} a_h^{\alpha\lambda} \Delta_{\lambda\beta} ; a_h^{2\alpha} \delta_{\beta 2}] \quad (8.1.9)$$

$$\Delta_{\lambda\beta} = 0 \quad \text{if } \lambda = \beta \quad , \quad \Delta_{\lambda\beta} = 1 \quad \text{if } \lambda \neq \beta, \quad (8.1.10)$$

$${}^t\tilde{u}_{h12} = [\tilde{u}_{h1} \quad \tilde{u}_{h1,1} \quad \tilde{u}_{h1,2} \quad \tilde{u}_{h2} \quad \tilde{u}_{h2,1} \quad \tilde{u}_{h2,2}] \quad (8.1.11)$$

Let us set

$$[DG12(\vec{u}_h)]_{1 \times 6} = [d1g1(\tilde{u}_{h1}); d1g1(\tilde{u}_{h2})]_{1 \times 6} \quad (8.1.12)$$

Then, using relations (8.1.2) and (8.1.4) we can write

$${}^t\tilde{u}_{h12} = [DG12(\vec{u}_h)]_{1 \times 6} [LAMB12]_{6 \times 6} \quad (8.1.13)$$

with

$$[LAMB12]_{6 \times 6} = \left[ \begin{array}{ccc|ccc} \lambda & \partial_1 \lambda & \partial_2 \lambda & & & \\ \hline & & & \lambda & \partial_1 \lambda & \partial_2 \lambda \end{array} \right] \quad (8.1.14)$$

Hence the relation (8.1.8) can be rewritten as

$$\tilde{\gamma}_{h\beta}^{\alpha}(\vec{u}_h) = M_{h\beta}^{\alpha} {}^t[LAMB12] {}^t[DG12(\vec{u}_h)] \quad (8.1.15)$$



Expression of  $\tilde{\rho}_{h\beta}^{\alpha}(\vec{u}_h)$  :

From (4.1.4) (4.2.12) we obtain

$$\tilde{\rho}_{h\beta}^{\alpha}(\vec{u}_h) = a_h^{\alpha\lambda} \tilde{\rho}_{h\beta\lambda} = a_h^{\alpha\lambda} \tilde{v}_{h3,\lambda\beta} \quad (8.1.16)$$

or, equivalently

$$\tilde{\rho}_{h\beta}^{\alpha}(\vec{u}_h) = B_{h\beta}^{\alpha} \tilde{u}_{h3} \quad (8.1.17)$$

where

$$B_{h\beta}^{\alpha} = [0 \quad 0 \quad 0 \quad a_h^{1\alpha} \delta_{\beta 1} \quad a_h^{\alpha\lambda} \Delta_{\lambda\beta} \quad a_h^{2\alpha} \delta_{\beta 2}] \quad (8.1.18)$$

$${}^t\tilde{u}_{h3} = [\tilde{u}_{h3} \quad \tilde{u}_{h3,1} \quad \tilde{u}_{h3,2} \quad \tilde{u}_{h3,11} \quad \tilde{u}_{h3,12} \quad \tilde{u}_{h3,22}] \quad (8.1.19)$$

Let us set

$$[DA_i]_{9 \times 10} = [D_i]_{9 \times 9} [A_i]_{9 \times 10} \quad (8.1.20)$$

$$[LAMB3_i]_{10 \times 6} = [\Lambda_i \quad \partial_1 \Lambda_i \quad \partial_2 \Lambda_i \quad \partial_{11} \Lambda_i \quad \partial_{12} \Lambda_i \quad \partial_{22} \Lambda_i] \quad (8.1.21)$$

Then, the relation (8.1.19) becomes

$${}^t\tilde{u}_{h3}|_{K_i} = [DLGL_i(\tilde{u}_{h3})]_{1 \times 9} [DA_i]_{9 \times 10} [LAMB3_i]_{10 \times 6} \quad (8.1.22)$$

so that

$$\tilde{\rho}_{h\beta}^{\alpha}(\vec{u}_h)|_{K_i} = B_{h\beta}^{\alpha} {}^t[LAMB3_i] {}^t[DA_i] {}^t[DLGL_i(\tilde{u}_{h3})] \quad (8.1.23)$$

Expression of the facet stiffness matrix

Combining the relation (8.1.1) with the expressions (8.1.15) and (8.1.23), we obtain

$$\begin{aligned} \tilde{a}_{hK}(\vec{u}_h, \vec{v}_h) = & [DG12(\vec{u}_h)] \left( \int_K [LAMB12][M_h] {}^t[LAMB12] d\xi^1 d\xi^2 \right) {}^t[DG12(\vec{v}_h)] \\ & + \sum_{i=1}^3 ([DLGL_i(\vec{u}_{h3})][DA_i] \left\{ \int_{K_i} [LAMB3_i][B_h] {}^t[LAMB3_i] d\xi^1 d\xi^2 \right\} \\ & {}^t[DA_i] {}^t[DLGL_i(\vec{v}_{h3})] \end{aligned} \quad (8.1.24)$$

where

$$[M_h]_{6 \times 6} = \frac{Ee \sqrt{a_h}}{1-\nu^2} \left\{ (1-\nu) {}^tM_{h\beta}^\alpha M_{h\alpha}^\beta + \nu {}^tM_{h\alpha}^\alpha M_{h\beta}^\beta \right\} \quad (8.1.25)$$


$$[B_h]_{6 \times 6} = \frac{Ee^3 \sqrt{a_h}}{12(1-\nu^2)} \left\{ (1-\nu) {}^tB_{h\beta}^\alpha B_{h\alpha}^\beta + \nu {}^tB_{h\alpha}^\alpha B_{h\beta}^\beta \right\} . \quad (8.1.26)$$

At the beginning of this section we have mentioned that we have considered the element stiffness matrix associated to the expression (4.3.3). In fact, we consider practically the expression (4.3.14) taking into account the effect of numerical integration. With obvious notations, the relations (4.3.14) and (8.1.24) reveal that

$$\begin{aligned} a_{hK}^*(\vec{u}_h, \vec{v}_h) = & \left. \begin{aligned} & [DG12(\vec{u}_h)] \left( \sum_{\ell_1=1}^{L_1} \omega_{\ell_1, K} ([LAMB12][M_h] {}^t[LAMB12]) (b_{\ell_1, K}) {}^t[DG12(\vec{v}_h)] \right) \\ & + \sum_{i=1}^3 ([DLGL_i(\vec{u}_{h3})][DA_i] \left\{ \sum_{\ell_2=1}^{L_2} \omega_{\ell_2, K_i} ([LAMB3_i][B_h] {}^t[LAMB3_i]) \right. \\ & \left. (b_{\ell_2, K_i}) {}^t[DA_i] {}^t[DLGL_i(\vec{v}_{h3})] \right\} \end{aligned} \right\} \quad (8.1.27) \end{aligned}$$

According to (6.1.1) and (6.1.2), we can use following numerical integration schemes in order to compute respectively first and second terms of the above relation.


SCHEME 8.1.1 : integration of extension terms; exact for  $P_0$ .



$$\int_K f(x) dx \sim f(G_K) \text{meas}(K) \quad (8.1.28)$$

SCHEME 8.1.2 : integration of flexion terms; exact for  $P_2$ .

This scheme has to be used on every subtriangle  $K_i$  of triangle K.



$$\int_{K_i} f(x) dx \sim \frac{1}{3} \{f(A) + f(B) + f(C)\} \text{meas}(K_i) \quad (8.1.29)$$

## 8.2 - Implementation of the facet second member matrix

Similarly to section 8.1, we introduce the following expressions  $\tilde{f}_{hK}$  and  $f_{hK}^*$  respectively associated to the relations (4.3.4) and (4.3.15), i.e.,

$$\tilde{f}_{hK}(\vec{v}_h) = \int_K \vec{p} \cdot \vec{v}_h \sqrt{a_h} d\xi^1 d\xi^2, \quad (8.2.1)$$

$$f_{hK}^*(\vec{v}_h) = \left. \begin{aligned} & \sum_{\ell_1=1}^{L_1} \omega_{\ell_1, K} (\{\vec{p} \cdot \vec{a}_h^\alpha\} \tilde{v}_{h\alpha} \sqrt{a_h}) (b_{\ell_1, K}) \\ & + \sum_{i=1}^3 \sum_{\ell_2=1}^{L_2} \omega_{\ell_2, K_i} (\{\vec{p} \cdot \vec{a}_h^3\} \tilde{v}_{h3} \sqrt{a_h}) (b_{\ell_2, K_i}) \end{aligned} \right\} \quad (8.2.2)$$

Setting

$$\tilde{t}_{P_{h12}} = \sqrt{a_h} [\tilde{p}_h^1 \quad 0 \quad 0 \quad \tilde{p}_h^2 \quad 0 \quad 0] \quad (8.2.3)$$

$$\tilde{t}_{P_{h3}} = \sqrt{a_h} [\tilde{p}_h^3 \quad 0 \quad 0 \quad 0 \quad 0 \quad 0] \quad (8.2.4)$$

with

$$\tilde{p}_h^i = \vec{p} \cdot \vec{a}_h^i, \quad (8.2.5)$$

we derive from (8.1.11) (8.1.19)

$$\tilde{f}_{hK}(\vec{v}_h) = \int_K (\tilde{t}_{P_{h12}} \tilde{v}_{h12} + \tilde{t}_{P_{h3}} \tilde{v}_{h3}) d\xi^1 d\xi^2. \quad (8.2.6)$$

Then, the relations (8.1.13) (8.1.22) give

$$\tilde{f}_{hK}(\vec{v}_h) = \left\{ \int_K \tilde{t}_{P_{h12}} {}^t[LAMBDA12] d\xi^1 d\xi^2 \right\} {}^t[DG12(\vec{v}_h)] + \sum_{i=1}^3 \left\{ \left( \int_{K_i} \tilde{t}_{P_{h3}} {}^t[LAMBDA3_i] d\xi^1 d\xi^2 \right) {}^t[DA_i] {}^t[DLGL_i(\tilde{v}_{h3})] \right\}. \quad (8.2.7)$$

Finally, we immediately obtain

$$\begin{aligned} f_{hK}^*(\vec{v}_h) = & \left\{ \sum_{\ell_1=1}^{L_1} \omega_{\ell_1, K} ({}^t\tilde{P}_{h12} {}^t[LAMBDA12])(b_{\ell_1, K}) \right\} {}^t[DG12(\vec{v}_h)] \\ & + \sum_{i=1}^3 \left\{ \left( \sum_{\ell_2=1}^{L_2} \omega_{\ell_2, K_i} ({}^t\tilde{P}_{h3} {}^t[LAMBDA3_i])(b_{\ell_2, K_i}) \right) {}^t[DA_i] {}^t[DLGL_i(\tilde{v}_{h3})] \right\} \end{aligned} \quad (8.2.8)$$

As numerical integration schemes, we can use (8.1.28) and (8.1.29) in order to compute respectively the first and the second terms of second member of the above relation.

### 8.3 - Implementation of the compatibility relations

The relations (8.1.27) and (8.2.8) permit to compute the stiffness and second member matrices *facet by facet*. To derive the corresponding *global* stiffness and second member matrices, we have to assemble. A convenient way consists in

- (i) using the compatibility relations to obtain the expressions of  $\vec{\tilde{u}}_h$  (resp.  $\vec{\tilde{v}}_h$ ) as functions of  $\vec{u}_h$  (resp.  $\vec{v}_h$ ),
- (ii) assembling the facet contributions so that the global matrices are referred to the unknowns  $\vec{u}_h$ .

In this section we consider the point (i). The second point (ii) will be the object of next paragraph.

Let us set for any triangle  $K \in \mathcal{T}_h$  and for any  $\vec{u}_h \in \vec{X}_h$

$$\begin{aligned}
 [DG(\vec{u}_h)] = & \left[ \begin{array}{cccccc}
 u_{h1}(\Sigma_1) & u_{h1}(\Sigma_2) & u_{h1}(\Sigma_3) & u_{h2}(\Sigma_1) & u_{h2}(\Sigma_2) & u_{h2}(\Sigma_3) \\
 u_{h3}(\Sigma_1) & u_{h3}(\Sigma_2) & u_{h3}(\Sigma_3) & u_{h3,1}(\Sigma_1) & u_{h3,2}(\Sigma_1) & \\
 u_{h3,1}(\Sigma_2) & u_{h3,2}(\Sigma_2) & u_{h3,1}(\Sigma_3) & u_{h3,2}(\Sigma_3) & & 
 \end{array} \right] \cdot
 \end{aligned} \tag{8.3.1}$$

Similarly, for any  $\vec{\tilde{u}}_h \in \vec{X}_h$ , we set

$$[DG(\vec{\tilde{u}}_h)] = [DG12(\vec{\tilde{u}}_h) ; DLGL(\vec{\tilde{u}}_{h3})] \tag{8.3.2}$$

where the matrices  $[DG12(\vec{\tilde{u}}_h)]$  and  $[DLGL(\vec{\tilde{u}}_{h3})] \equiv [DLGL_1(\vec{\tilde{u}}_{h3})]$  are defined by (8.1.12) and (7.4.4), respectively. Then, using compatibility relations we want to determine the matrix  $[TILD]_{15 \times 15}$  so that, for any functions  $\vec{\tilde{u}}_h \in \vec{X}_h$  and  $\vec{u}_h \in \vec{X}_h$  in correspondence through the bijection  $F_h$  defined by theorem 4.2.1, we have :

$$[DG(\vec{\tilde{u}}_h)]_{1 \times 15} = [DG(\vec{u}_h)]_{1 \times 15} [TILD]_{15 \times 15} \tag{8.3.3}$$

From (4.2.21) (4.2.23), we obtain

$$\vec{u}_h(\Sigma_j) = \vec{u}_h(\Sigma_j) \quad , \quad j=1,2,3 .$$

Observing that on the triangle  $K$ , we have

$$\vec{u}_h(\Sigma_j) = \tilde{u}_{hi}(\Sigma_j) \vec{a}_h^i \quad (8.3.4)$$

$$\vec{u}_h(\Sigma_j) = u_{hk}(\Sigma_j) \vec{a}^k(\Sigma_j) \quad , \quad j=1,2,3, \quad (8.3.5)$$

we get

$$\tilde{u}_{hi}(\Sigma_j) = d_{hi}^k(\Sigma_j) u_{hk}(\Sigma_j) \quad , \quad j=1,2,3, \quad (8.3.6)$$

where, for any point  $M \in K$

$$d_{hi}^k(M) = \vec{a}^k(M) \cdot \vec{a}_{hi} . \quad (8.3.7)$$

Let us record that, due to the definition of the mapping  $\vec{\phi}_h|_K \in (P_1(K))^3$ , the basis vectors  $\vec{a}_h^i$  and  $\vec{a}_{hi}$  are constants on all the triangle  $K$ .

Now, let us determine  $\tilde{u}_{h3,\alpha}(\Sigma_i)$  as functions of the elements of  $[DG(\vec{u}_h)]$ . Firstly we observe that  $\tilde{u}_{h\alpha} \in \tilde{X}_{h1}$  and the relations (7.7.1) (7.7.4) (7.7.5) imply that the functions  $\tilde{u}_{h\alpha,\beta}$  are constant on all the triangle  $K$  and equal to

$$\tilde{u}_{h\alpha,\beta} = [d \lg l(\tilde{u}_{h\alpha})] [\partial_\beta \lambda] . \quad (8.3.8)$$

Hence the relation (4.2.24) give the expression of the function  $\tilde{\omega}_h^3$  which is also a constant triangle by triangle, i.e., on the triangle  $K$  :

$$\tilde{\omega}_h^3 = \frac{1}{2 \sqrt{a_h}} ([d \lg l(\tilde{u}_{h2})] [\partial_1 \lambda] - [d \lg l(\tilde{u}_{h1})] [\partial_2 \lambda]) . \quad (8.3.9)$$

Secondly, the compatibility relations (4.2.25) involve

$$\tilde{\omega}_h^\beta(\Sigma_j) \vec{a}_{h\beta} \cdot \vec{a}^\lambda(\Sigma_j) = B^\lambda(\Sigma_j) - \tilde{\omega}_h^3 \vec{a}_{h3} \cdot \vec{a}^\lambda(\Sigma_j) \quad , \quad j=1,2,3 \quad ,$$

or equivalently, by using (8.3.7)

$$d_{h\beta}^\lambda(\Sigma_j) \tilde{\omega}_h^\beta(\Sigma_j) = B^\lambda(\Sigma_j) - d_{h3}^\lambda(\Sigma_j) \tilde{\omega}_h^3 \quad , \quad j=1,2,3 \quad . \quad (8.3.10)$$

Let us denote

$$c^\lambda(\Sigma_j) = B^\lambda(\Sigma_j) - d_{h3}^\lambda(\Sigma_j) \tilde{\omega}_h^3 \quad (8.3.11)$$

$$d_h(\Sigma_j) = d_{h1}^1(\Sigma_j) d_{h2}^2(\Sigma_j) - d_{h1}^2(\Sigma_j) d_{h2}^1(\Sigma_j) \quad . \quad (8.3.12)$$

From section 4.2 we have  $d_h(\Sigma_j) \neq 0$ . Then, the relations (8.3.10) to (8.3.12) imply

$$\tilde{\omega}_h^\lambda(\Sigma_j) = \frac{1}{d_h(\Sigma_j)} e^{\lambda\nu} e_{\alpha\beta} c^\alpha(\Sigma_j) d_{h\nu}^\beta(\Sigma_j) \quad , \quad j=1,2,3 \quad .$$

Finally, from (4.2.26), we derive (compare with relation just before (5.2.45))

$$\tilde{u}_{h3,\alpha}(\Sigma_j) = \frac{\sqrt{a_h}}{d_h(\Sigma_j)} e_{\lambda\beta} c^\lambda(\Sigma_j) d_{h\alpha}^\beta(\Sigma_j) \quad , \quad j=1,2,3 \quad . \quad (8.3.13)$$

From (8.3.6) (8.3.9), we have

$$\tilde{\omega}_h^3 = \frac{1}{2\sqrt{a_h}} [DG(\vec{u}_h)]_{1 \times 15} [T^3]_{15 \times 1} \quad (8.3.14)$$

with

$$[T^3]_{15 \times 1} = \begin{bmatrix} D_{h1} & D_{h2} \\ 0 & 0 \end{bmatrix}_{15 \times 6} \begin{bmatrix} -\partial_2 \lambda \\ \partial_1 \lambda \end{bmatrix}_{6 \times 1} \quad (8.3.15)$$

and

$$[D_{hi}]_{9 \times 3} = \begin{bmatrix} d_{hi}^1(\Sigma_1) & 0 & 0 \\ 0 & d_{hi}^1(\Sigma_2) & 0 \\ 0 & 0 & d_{hi}^1(\Sigma_3) \\ d_{hi}^2(\Sigma_1) & 0 & 0 \\ 0 & d_{hi}^2(\Sigma_2) & 0 \\ 0 & 0 & d_{hi}^2(\Sigma_3) \\ d_{hi}^3(\Sigma_1) & 0 & 0 \\ 0 & d_{hi}^3(\Sigma_2) & 0 \\ 0 & 0 & d_{hi}^3(\Sigma_3) \end{bmatrix} \quad (8.3.16)$$

From (4.2.22) we have

$$B^\lambda(\Sigma_j) = [DG(\vec{u}_h)]_{1 \times 15} [T_j^\lambda]_{15 \times 1} \quad (8.3.17)$$

with

$${}^t_{[T_1^\lambda]}_{1 \times 15} = e^{\lambda\beta} \left[ \begin{array}{cccccccc} \frac{b_\beta^1}{\sqrt{a}}(\Sigma_1) & 0 & 0 & \frac{b_\beta^2}{\sqrt{a}}(\Sigma_1) & 0 & 0 & 0 & 0 & 0 \\ \frac{\delta_{1\beta}}{\sqrt{a}(\Sigma_1)} & \frac{\delta_{2\beta}}{\sqrt{a}(\Sigma_1)} & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{array} \right] \quad (8.3.18)$$

$${}^t_{[T_2^\lambda]}_{1 \times 15} = e^{\lambda\beta} \left[ \begin{array}{cccccccc} 0 & \frac{b_\beta^1}{\sqrt{a}}(\Sigma_2) & 0 & 0 & \frac{b_\beta^2}{\sqrt{a}}(\Sigma_2) & 0 & 0 & 0 & 0 \\ 0 & 0 & \frac{\delta_{1\beta}}{\sqrt{a}(\Sigma_2)} & \frac{\delta_{2\beta}}{\sqrt{a}(\Sigma_2)} & 0 & 0 & 0 & 0 & 0 \end{array} \right] \quad (8.3.19)$$



$${}^t[T_3^\lambda]_{1 \times 15} = e^{\lambda\beta} \left[ \begin{array}{cccccccc} 0 & 0 & \frac{b_\beta^1}{\sqrt{a}}(\Sigma_3) & 0 & 0 & \frac{b_\beta^2}{\sqrt{a}}(\Sigma_3) & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & \frac{\delta_{1\beta}}{\sqrt{a}(\Sigma_3)} & \frac{\delta_{2\beta}}{\sqrt{a}(\Sigma_3)} & & & \end{array} \right] . \quad (8.3.20)$$

Then, combining (8.3.6) (8.3.11) (8.3.13) (8.3.14) (8.3.16) we obtain

$$[DG(\vec{u}_h)]_{1 \times 15} = [DG(\vec{u}_h)]_{1 \times 15} [TILD]_{15 \times 15} \quad (8.3.21)$$

with

$$[TILD]_{15 \times 15} = \left[ \begin{array}{c|c|c|c|c|c|c|c|c} Dh_1 & Dh_2 & Dh_3 & R_{11} & R_{21} & R_{12} & R_{22} & R_{13} & R_{23} \\ \hline 0 & 0 & 0 & & & & & & \end{array} \right] \quad (8.3.22)$$

and

$$[R_{\alpha j}]_{15 \times 1} = \frac{d_{h\alpha}^\beta(\Sigma_j)}{d_h(\Sigma_j)} e^{\lambda\beta} \left\{ \sqrt{a}_h [T_j^\lambda]_{15 \times 1} - \frac{1}{2} d_{h3}^\lambda(\Sigma_j) [T^3]_{15 \times 1} \right\} . \quad (8.3.23)$$

9 - DERIVATION OF THE GLOBAL STIFFNESS AND SECOND MEMBER MATRICES

9.1 - Derivation of the global stiffness matrix

Using the compatibility relations obtained triangle by triangle through the relations (8.3.21), we substitute the *global* unknowns  $[DG(\vec{u}_h)]$  to the *facet* unknowns  $[DG(\vec{u}_h)]$ .

For convenience, we set

$$[J_{12}]_{15 \times 6} = \begin{bmatrix} I_6 \\ \text{-----} \\ 0_{9 \times 6} \end{bmatrix}, \quad (9.1.1)$$

$$[J_3]_{15 \times 9} = \begin{bmatrix} 0_{6 \times 9} \\ \text{-----} \\ I_9 \end{bmatrix}. \quad (9.1.2)$$

Then the relations (7.4.4) (8.1.12) (8.3.1) (8.3.2) (8.3.21) reveal that

$$[DG_{12}(\vec{u}_h)]_{1 \times 6} = [DG(\vec{u}_h)]_{1 \times 15} [TILD]_{15 \times 15} [J_{12}]_{15 \times 6} \quad (9.1.3)$$

$$[DLGL(\vec{u}_{h3})]_{1 \times 9} = [DG(\vec{u}_h)]_{1 \times 15} [TILD]_{15 \times 15} [J_3]_{15 \times 9}.$$

Observe from (7.4.4) and (8.3.2) that

$$[DLGL_i(\vec{v}_h)] = [DLGL(\vec{v}_h)][K_i]_{9 \times 9} \quad (9.1.4)$$

with

$$[K_i] = \begin{bmatrix} K_{i11} & 0 \\ 0 & K_{i22} \end{bmatrix}, \quad [K_{i11}]_{3 \times 3} = \begin{bmatrix} \delta_{1i} & \delta_{2i} & \delta_{3i} \\ \delta_{2i} & \delta_{3i} & \delta_{1i} \\ \delta_{3i} & \delta_{1i} & \delta_{2i} \end{bmatrix}$$

and

$$[K_{i22}]_{6 \times 6} = \begin{bmatrix} \delta_{1i} & 0 & \delta_{2i} & 0 & \delta_{3i} & 0 \\ 0 & \delta_{1i} & 0 & \delta_{2i} & 0 & \delta_{3i} \\ \delta_{2i} & 0 & \delta_{3i} & 0 & \delta_{1i} & 0 \\ 0 & \delta_{2i} & 0 & \delta_{3i} & 0 & \delta_{1i} \\ \delta_{3i} & 0 & \delta_{1i} & 0 & \delta_{2i} & 0 \\ 0 & \delta_{3i} & 0 & \delta_{1i} & 0 & \delta_{2i} \end{bmatrix}$$

so that

$$[DLGL_i(\vec{u}_{h3})]_{1 \times 9} = [DG(\vec{u}_h)]_{1 \times 15} [TILD]_{15 \times 15} [K_{3i}]_{15 \times 9} \quad (9.1.5)$$

with

$$[K_{3i}]_{15 \times 9} = \begin{bmatrix} 0_{6 \times 9} \\ K_i \end{bmatrix}. \quad (9.1.6)$$

From (4.3.16) and (8.1.27), we finally obtain

$$\begin{aligned} b_h(\vec{u}_h, \vec{v}_h) &= \sum_{k \in \mathcal{T}_h} \left[ [DG(\vec{u}_h)]_{1 \times 15} [TILD]_{15 \times 15} \right. \\ &\quad \left. \left\{ [J_{12}]_{15 \times 6} \left( \sum_{\ell_1=1}^{L_1} \omega_{\ell_1, K} ([LAMBDA12]_{6 \times 6} [M_h]_{6 \times 6} {}^t[LAMBDA12]) \right) \right. \right. \\ &\quad \left. \left. (b_{\ell_1, K}) {}^t[J_{12}] \right\} \right. \\ &\quad \left. + \sum_{i=1}^3 ([K_{3i}]_{15 \times 9} [DA_i]_{9 \times 10} \left\{ \sum_{\ell_2=1}^{L_2} \omega_{\ell_2, K_i} ([LAMBDA3_i]_{10 \times 6} \right. \right. \right. \\ &\quad \left. \left. [B_h]_{6 \times 6} {}^t[LAMBDA3_i]) (b_{\ell_2, K_i}) {}^t[DA_i] {}^t[K_{3i}] \right\} \right. \\ &\quad \left. {}^t[TILD] {}^t[DG(\vec{v}_h)] \right] \quad (9.1.7) \end{aligned}$$

Thus, assembling the contributions of every triangle  $K \in \mathcal{T}_h$  we get the *global stiffness matrix*.

9.2 - Derivation of the global second member

Similarly to section 9.1, the relations (4.3.17) and (8.2.8) reveal that

$$\begin{aligned}
 g_h(\vec{v}_h) = & \sum_{K \in \mathcal{T}_h} \left[ \left( \sum_{\ell_1=1}^{L_1} \omega_{\ell_1, K} \left( {}^t\tilde{P}_{h12} \quad {}^t[LAMBDA12](b_{\ell_1, K}) \right) \right) {}^t[J_{12}] \right] \\
 & + \sum_{i=1}^3 \left\{ \sum_{\ell_2=1}^{L_2} \omega_{\ell_2, K_i} \left( {}^t\tilde{P}_{h3} \quad {}^t[LAMBDA3](b_{\ell_2, K_i}) \right) \right\} {}^t[DA_i] \quad {}^t[K_{3i}] \Bigg\} \quad (9.2.1) \\
 & {}^t[TILD] \quad {}^t[DG(\vec{v}_h)] .
 \end{aligned}$$

By assembling the contributions of every triangle  $K \in \mathcal{T}_h$  we get the *global second member matrix*:

REFERENCES

- [1] BERNADOU, M. [1980] Convergence of conforming finite element methods for general shell problems. Int. J. Engng. Sc. 18, 249-276.
- [2] BERNADOU, M. ; BOISSERIE, J.M. [1982] The Finite Element Method in Thin Shell Theory : Application to Arch Dam Simulations, Birkhäuser, Boston.
- [3] CLOUGH, R.W. ; JOHNSON, C.P. [1968] A finite element approximation for the analysis of thin shells. Int. J. Solids Structures, Vol. 4.
- [4] CLOUGH, R.W. ; JOHNSON, C.P. [1970] Finite element analysis of arbitrary thin shells. Proceeding ACI Symposium on concrete thin shells, pp. 333-363, New York.
- [5] CIARLET, P.G. [1978] The Finite Element Method for Elliptic Problems. North-Holland.
- [6] CIARLET, P.G. [1976] Conforming finite element method for the shell problems, in The Mathematics of Finite Elements and Applications II (J.R. Whiteman, Ed., Academic Press, London, pp. 105-123.
- [7] JOHNSON, C.P. [1975] On finite element methods for curved shells using flat elements. In : Numerische Behandlung von Differentialgleichungen, pp. 147-154. International Series of Numerical Mathematics, Vol. 27, Basel, Stuttgart : Birkhäuser Verlag.
- [8] BERNADOU, M. ; DUCATEL, Y. [1982] Approximation of general arch problems by the straight beam elements. Numer. Math., 40, pp. 1-29.
- [9] BERNADOU, M. ; DUCATEL, Y. ; TROUVE, P. [to appear] Approximation of a circular cylindrical shell by Clough-Johnson flat plate finite elements .

- [10] ARGYRIS, J.H. ; DUNNE, C.P. ; MALEJANNAKIS, G.A. ; SCHELKLE, E. [1977] A simple triangular facet shell element with applications to linear and non-linear equilibrium and elastic stability problems. *Computer Methods in Applied Mechanics and Engineering*, 10, pp. 371-403, and 11, pp. 97-131.
- [11] DAWE, D.J. [1972] Shell analysis using a simple facet element, *Journal of strain analysis*, 7, pp. 266-270.
- [12] OLSON, M.D. ; BEARDEN, T.W. [1979] A simple flat triangular shell element revisited. *Int. J. Numer. Methods Engng. Vol. 14*, pp. 51-68.
- [13] IRONS, B.M. [1974] Un nouvel element de coques générales "semiloof", *Computing Methods in Applied Sciences and Engineering*, Part. 1, pp. 177-192, Springer-Verlag, New York.
- [14] KOITER, W.T. [1970] On the foundations of the linear theory of thin elastic shells. *Proc. Kon. Ned. Akad. Wetensch.*, B73, pp. 169-195.
- [15] BERNADOU, M. ; CIARLET, P.G. [1976] Sur l'ellipticité du modèle linéaire de coques de W.T. KOITER. "Computing Methods in Applied Sciences and Engineering". *Lecture Notes in Economic and Mathematical Systems*, Vol. 134, pp. 89-136, Springer-Verlag, Berlin.
- [16] CLOUGH, R.W. ; TOCHER, J.L. [1965] Finite element stiffness matrices for analysis of plates in bending. *Proc. Conf. Matrix Methods in Struct. Mech.*, Air Force Inst. of Techn., Wright Patterson A.F. Base, Ohio, October 1965.
- [17] BERNADOU, M. ; LALANNE, B. [1986] Sur l'approximation des coques minces par des méthodes "B-splines et éléments finis" - 1ère partie : formulation du problème et estimations d'erreur ; 2ème partie : application à la modélisation d'une pale de turbine ou d'un compresseur axial. *Rapports de Recherche INRIA*, N°s 474 et 479.

- [18] CARR, A.J. [1967] A refined finite element analysis of thin shell structures including dynamic loads, Report SESM 67-9, University of California, Berkeley.
- [19] KIKUCHI, F. [1984] Error analysis of flat plate element approximation of circular cylindrical shells, *Theoretical and Applied Mechanics*, Vol. 32, pp. 469-484, University of Tokyo Press.
- [20] GREEN, A.E. ; ZERNA, W. [1968] Theoretical Elasticity, Oxford University Press, 2nd Edition.
- [21] BERNADOU, M. ; HASSAN, K. [1981] Basis functions for general HSIEH-CLOUGH-TOCHER triangles, complete or reduced, *Int. J. Num. Math. Engng.*, 17, pp. 784-789.
- [22] BERNADOU, M. ; ODEN, J.T. [1981] An existence theorem for a class of nonlinear shallow shell problems, *J. Math. Pures et Appl.*, 60, pp. 285-308.
- [23] ZIENKIEWICZ, O.C. [1977] The Finite Element Method in Engineering Science, 3rd Edition, Mc Graw-Hill.
- [24] IRONS, B. ; AHMAD, S. [1980] Techniques of Finite Elements, J. Wiley and Sons.
- [25] GALLAGHER, R.H. [1976] Problems and progress in thin shell finite element analysis, in Finite Elements for Thin Shells and Curved Members, Chapter 1, pp. 1-14, Edited by D.G. Ashwell and R.H. Gallagher, J. Wiley and Sons, London.
- [26] KOITER, W.T. [1966] On the nonlinear theory of thin elastic shells, *Proc. Kon. Ned. Akad. Wetensch.*, B69, pp. 1-54.
- [27] STROUD, A.H. [1971] Approximate Calculation of Multiple Integrals, Prentice-Hall, Englewood Cliffs.

Imprimé en France

par

l'Institut National de Recherche en Informatique et en Automatique

